

## Artificial hollows for Carnaby's cockatoo



## How to monitor and maintain artificial hollows for Carnaby's cockatoo

It is important to monitor and maintain artificial hollows after they have been erected. Monitoring ensures that the effectiveness of the artificial hollow can be determined. It also means that problems with pest species or any maintenance requirements can be identified and resolved.

Without regular maintenance, artificial hollows are likely to fail to achieve their objective (that is, they will fail to provide nesting opportunities for threatened cockatoos). Therefore it is important to continue a regime of regular maintenance while the artificial hollow is required. It may be several (to many) decades until a natural replacement hollow is available.

Monitoring should be undertaken in order to detect:

- Use by Carnaby's cockatoo
- Maintenance requirements
- Use by other native species
- Use by pest species (e.g. feral bees, galahs, corellas etc.)



Carnaby's cockatoo female prospecting an artificial hollow.  
Photo by Rick Dawson

### How do I monitor artificial hollows?

Before undertaking monitoring of artificial hollows for Carnaby's cockatoo it is recommended that you seek advice from BirdLife Australia, the WA Museum or the Department of Parks and Wildlife. It is also important to contact Parks and Wildlife, Wildlife Licensing Section, to determine if a scientific licence is required ([wildlifelicencing@dpaw.wa.gov.au](mailto:wildlifelicencing@dpaw.wa.gov.au)).

Monitoring artificial hollows requires keen observation and naturalist skills. It is often not possible to observe evidence of breeding directly (i.e. nestlings or eggs) and inferences must be made based on observation. There are many techniques available to monitor artificial hollows. A combination of several is likely to achieve the best results.

### Looking for signs of use

Cobwebs covering the entrance to the hollow will indicate that the hollow has not been used recently. This would also apply to other light debris that may have fallen to cover the opening partially. Signs of recent use or interest in the hollow include evidence of chewing.

---

### Observing parent behaviour around the hollow

The behaviour of parent birds around a hollow will indicate an approximate age of young in the nest.

Parent behaviour	Approximate age/stage of young
Prospecting for hollow	Unborn
Male only seen out of hollow	Egg or very young nestling (< 3 - 4 weeks)
Both parents seen entering/exiting the hollow	Nestling(s) have hatched (> 3 - 4 weeks)

---

### Observing feeding flocks

Flocks of all male birds indicate that the females are incubating eggs. When flocks are mixed it suggests the birds have either not laid yet or that the nestlings have hatched and no longer require brooding (approximately 3 - 4 weeks old).

---

### Tapping

When females are sitting on eggs they will usually respond to tapping at the base of their tree (or pole) by appearing at the entrance or flying from the hollow opening. This is not a guarantee of breeding activity, but an indication that it is possibly occurring in the hollow.

---

### Observing insect activity around nest

The faecal matter produced by nestlings in a nest attracts insects, especially flies and ants. The type and number of these insects will help indicate how old any nestlings present may be. Factors such as temperature and humidity will also affect insect activity and so observations of insect activity should only be used as supporting evidence for other indications of age/use. Blowflies around a nest usually indicate that a death has occurred.

---

### Listening for nestlings

With experience it is possible to determine if one or two nestlings are present and a broad estimate of age based on the type and loudness of noises they make.

---

## **Looking inside the nest**

This can be achieved either with the aid of a telescopic pole and camera or mirror, or with the use of a ladder or other climbing equipment. This method can obtain the most detailed monitoring information for artificial hollows. However it is also the most time consuming and difficult to organise. Special equipment is likely to be needed depending on the height and positioning of artificial hollows. There are also safety issues associated with ladder or rope climbing options to reach nests to undertake observations.

---

## **How often should I monitor artificial hollows?**

The minimum frequency of monitoring and the techniques used will be determined by the aims of the monitoring and the resources available. It is important to limit disturbance to breeding birds and this should be considered when determining the techniques used and frequency.

---

## **How do I maintain artificial hollows?**

Artificial hollows require maintenance to ensure they continue to have the greatest chance of them being used by Carnaby's cockatoos. Periodic maintenance checks should be undertaken at least every two years, preferably annually. These checks should be undertaken prior to the breeding season which is between July and January with breeding occurring later in this period in southern areas. It is important to maintain a regime of regular maintenance as long as the artificial hollow is required. It may take several (to many) decades until a natural replacement hollow is available.

Maintenance checks should assess the following as a minimum:

- Condition of chewing posts (if present)
- Condition of attachment points
- Condition of hollow bases
- Stability of tree or pole used to mount the artificial hollow



Artificial hollow base needing repair.  
*Photo by Christine Groom*

## **Repairing hollows**

Any problems identified during maintenance checks should be addressed, and any repairs required done, as soon as possible. If breeding is currently occurring, maintenance may need to be delayed if it is likely to disturb the parents or nestling. Likely maintenance needs include replacement of chewing posts (frequently) or nest bases (occasionally) and repairing of any cracks (infrequently). Maintenance concerns regarding the security of attachment points or the stability of the tree or pole should be addressed as a priority for safety reasons.

For artificial hollows known to be used, spare chewing posts should be taken into the field when undertaking maintenance checks.

---

**Monitoring of artificial hollows:**

Monitoring aim	Frequency of visits	Monitoring techniques
<b>To determine possible use by Carnaby's cockatoo</b>	At least once during peak breeding season (i.e. between September and December)	<ul style="list-style-type: none"> <li>• Observing behaviour of adults around hollow</li> <li>• Tapping to see if female will flush from hollow (best undertaken between 10am and 3pm when females most likely to be sitting)</li> <li>• Listening for nestlings</li> <li>• Looking for evidence of chewing</li> <li>• Looking inside nest</li> </ul>
<b>To confirm use by Carnaby's cockatoo</b>	At least two visits during peak breeding season (i.e. between September and December)	<p>To observe at least two of the following:</p> <ul style="list-style-type: none"> <li>• Breeding behaviour of adults around hollow or evidence of chewing</li> <li>• Female flushed from hollow</li> <li>• Noises from nestlings in hollow</li> </ul> <p>Or to observe:</p> <ul style="list-style-type: none"> <li>• Nestlings or eggs in nest</li> </ul>
<b>To determine nesting success by Carnaby's cockatoo</b>	The more visits, the better. Preferably fortnightly visits between July and December. As a minimum, at least 3 visits spread throughout breeding season.	<ul style="list-style-type: none"> <li>• Looking inside nest to observe eggs or nestlings.</li> </ul>
<b>To determine use by any species</b>	As often as possible.	<ul style="list-style-type: none"> <li>• Inspection from ground as a minimum.</li> <li>• Looking inside nest for detailed observations.</li> </ul>
<b>To determine maintenance requirements</b>	At least every two years and preferably annually if hollow fitted with sacrificial chewing posts, can be longer if without.	<ul style="list-style-type: none"> <li>• A basic maintenance check can be undertaken from the ground. A ladder or elevated work platform will be required for a comprehensive check and to replace sacrificial chewing posts</li> </ul>

**Acknowledgements**

This information sheet is a joint initiative of Birdlife Australia, the Western Australian Museum and the Department of Parks and Wildlife. Many individuals have contributed to its preparation. The updated version was compiled by Rick Dawson (Department of Parks and Wildlife) with assistance from Denis Saunders.

**Other information sheets in the series: Artificial hollows for Carnaby's cockatoo**

- *How to design and place artificial hollows for Carnaby's cockatoo*
- *How to monitor and maintain artificial hollows for Carnaby's cockatoo*

Information sheets available on the *Saving Carnaby's cockatoo* webpage:

<http://www.dpaw.wa.gov.au/plants-and-animals/threatened-species-and-communities/threatened-animals/208-saving-carnaby-s-cockatoo>