



# PHOENIX

ENVIRONMENTAL SCIENCES

## Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project

Prepared for Australian Capital Equity

April 2023

Final



Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

#### Version history

Author/s	Reviewer/s	Version	Version number	Date submitted	Submitted to
F. Holmes, B. Quick	K. Crews	Draft for client comments	0.1	12-Jan-23	J. McMahon
F. Holmes		Final, client comments addressed	1.0	03-Apr-23	J. McMahon

© Phoenix Environmental Sciences Pty Ltd 2023

The use of this report is solely for the client for the purpose in which it was prepared. Phoenix Environmental Sciences accepts no responsibility for use beyond this purpose.

All rights are reserved and no part of this report may be reproduced or copied in any form without the written permission of Phoenix Environmental Sciences or the client.

Phoenix Environmental Sciences Pty Ltd  
2/3 King Edward Road OSBORNE PARK WA 6017  
P: 08 6323 5410  
E: [admin@phoenixenv.com.au](mailto:admin@phoenixenv.com.au)  
Project code: 1452-NAP-ACE-FAU

## EXECUTIVE SUMMARY

Australian Capital Equity Pty Ltd (ACE) is seeking to develop the Napier Downs Irrigation Project (the Project) located on Napier Downs Station, approximately 77 km east of Derby, Western Australia (WA). The Project will entail the development of centre irrigation pivots which will be used to produce fodder crops for cattle stocked on Napier Downs and nearby stations, with water sourced from the Grant Group Aquifer.

In March 2019 Phoenix Environmental Sciences Pty Ltd (Phoenix) was commissioned by ACE to undertake a desktop assessment of terrestrial vertebrate fauna and short-range endemic (SRE) invertebrate fauna in accordance with relevant EPA guidance. The desktop review identified a total of 42 significant vertebrate species, including 20 listed under either the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) or the *Biodiversity Conservation Act 2016* (BC Act) as Threatened or Specially Protected, 18 listed as migratory (Mig.) under both the EPBC Act and BC Act, and 9 listed as Priority by the Department of Biodiversity, Conservation and Attractions (DBCAs). None of the species identified had previously been recorded inside the study area. The closest records were from Hawkstone Creek (5.4 km east of the study area) where Northern Quoll (Endangered, EN), Gull-billed Tern (Mig.), Glossy Ibis (Mig.), Common Greenshank (Mig.), and Freshwater Crocodile (Other Specially Protected) were all recorded in 2013.

Based on the findings of the desktop assessment, Phoenix was then commissioned in August 2021 to undertake a baseline terrestrial fauna survey for the Project, which was completed between June-August 2022. This survey included both broad-scale fauna habitat mapping (within the study area plus a 1 km buffer) and sampling methods targeting several significant species identified in the previously completed desktop as potentially occurring inside the 586.5 ha study area.

The main field survey was completed between 27 June – 5 July 2022, with a secondary survey completed between 14-16 August 2022 to retrieve wet pitfall traps deployed for potential short-range endemic (SRE) invertebrate taxa. The survey involved 8 systematic trapping sites made up of Sheffield cages and Elliot box traps used to sample small mammals, as well as 2 ultrasonic recorders used to sample bats, active searches for reptiles and potential SRE taxa, avifauna surveys to sample the birds present inside the study area and the wet pitfall traps for SREs.

The survey identified 3 broad fauna habitat types based on substrate, topography, vegetation structure and composition, briefly referred to as:

- Open woodland over open shrubland over grassland
- Shrubland over grassland
- Open woodland over mixed herbs and grasses surrounding seasonally inundated depression.

The dominant fauna habitat type was the shrubland over grassland which made up over 70% of the total mapped area, while the open woodland over mixed herbs and grasses surrounding seasonally inundated depression made up less than 1% of the total mapped area.

During the survey, 114 terrestrial species were recorded, which equated to approximately 35% of the species identified in the desktop review. Additionally, 7 of the species recorded during the survey were not identified during the desktop. This was attributed to recent changes in the case of the cane toad that has moved into the region over the last decade, and incomplete contextual information due to the limited knowledge we have regarding the distribution of a number of species in the region, such as the Black Grasswren (P4) and the Northwest Kimberley Two-lined Dragon, both of which were shown in this survey to be distributed further south than previously thought.

Likelihood of occurrence was estimated for significant vertebrate species based on desktop and field data. Golden Bandicoot (VU) was the only significant species recorded inside the study area. Black Grasswren (P4) and Gouldian Finch (EN/P4) were recorded close to the study area and based on the similarity of the habitats inside and outside the study area, were considered likely to occur inside the study area. An additional 15 species identified in the desktop were assigned the category 'may possibly

occur' inside the study area, and the remaining 24 were assigned the category 'unlikely to occur' based on relevant habitat requirements.

The vertebrate faunal assemblage of the study area is broadly typical of the woodland and shrubland habitats in the region and consists mostly of species distributed throughout much of the Kimberley. The fauna habitats inside the study area are widespread across much of the region. Based on the location of the study area relative to the known distribution of Golden Bandicoot, the population is considered to be important given it is on the southern edge of the species' range.

As the habitats inside the study area are relatively continuous in terms of vegetation structure and landform attributes, it is unlikely that any invertebrate taxa are restricted to the study area. Vegetation composition was notably different in one small seasonally inundated depression; however, the topography and connectivity of the vegetation structure mean that this area was not considered to be suitable for supporting SRE taxa.

Three invertebrate taxa collected during the survey have been assigned potential SRE status, two of which are new to science (*Aname* 'MYG771' and *Cubaris* sp. indet. 'Napier') and one is from a species complex which is thought to contain multiple species. Given the relatively small size of the study area, the extent of SRE habitats it intersects, data obtained from the desktop review and the taxonomists that identified the specimens, this is likely a reflection of the limited information we have on the distribution of these invertebrates, and not a reflection of their scarcity.

## CONTENTS

Executive summary .....	ii
Contents .....	iv
1 Introduction .....	6
1.1 Background .....	6
1.2 Scope of work.....	7
1.3 Study area .....	8
2. Legislative context .....	11
2.1 Commonwealth.....	11
2.2 State .....	11
2.2.1 Threatened and Priority species .....	11
2.2.2 Critical habitat .....	12
2.2.3 Environmentally Sensitive Areas .....	12
2.2.4 Short-range endemic invertebrates .....	12
3. Existing environment .....	13
3.1 Interim Biogeographic Regionalisation of Australia.....	13
3.2 Land systems and surface geology.....	13
3.3 Climate and weather .....	17
3.4 Land use .....	18
3.5 National heritage places, Conservation reserves and ESAs .....	18
4. Methods.....	19
4.1 Desktop review .....	19
4.2 Field survey .....	20
4.2.1 Survey timing.....	20
4.2.2 Field methods.....	20
4.2.3 Survey personnel.....	27
5. Results.....	29
5.1 Desktop review .....	29
5.1.1 Vertebrate fauna .....	29
5.1.2 SRE invertebrate fauna.....	36
5.2 Field survey .....	39
5.2.1 Vertebrate fauna .....	39
5.2.2 SRE invertebrate fauna.....	49
5.3 Survey limitations.....	52
6. Discussion.....	53
6.1 Vertebrate fauna .....	53
6.2 SRE invertebrate fauna .....	54
6.3 Conclusion .....	55
References .....	56

## LIST OF FIGURES

Figure 1-1	Project location and study area .....	9
Figure 1-2	Study area in relation to previous desktop areas .....	10
Figure 3-1	Study area in relation to IBRA bioregions and subregions.....	15
Figure 3-2	Land systems and surface geology in the study area.....	16
Figure 3-3	Annual climate and weather data for Derby Airport .....	17
Figure 4-1	Survey sites.....	28
Figure 5-1	Desktop records of significant vertebrate fauna .....	35
Figure 5-2	Desktop records of SRE invertebrates.....	38
Figure 5-3	Fauna habitats and significant fauna records from the field survey.....	41
Figure 5-4	Species accumulation curve for vertebrate fauna .....	43
Figure 5-5	SRE habitats and recorded SRE taxa.....	51

## LIST OF TABLES

Table 3-1	Land systems and extent in study area .....	14
Table 4-1	Database searches conducted for the desktop review .....	19
Table 4-2	Survey dates .....	20
Table 4-3	Terrestrial fauna survey effort .....	22
Table 4-4	Short-range endemic categories .....	25
Table 4-5	Specialist taxonomists.....	26
Table 4-6	Survey personnel.....	27
Table 5-1	Significant fauna identified in the desktop review .....	30
Table 5-2	Terrestrial SRE invertebrates identified in the desktop review .....	36
Table 5-3	Extent and description of each fauna habitat in the study area.....	40
Table 5-4	Number of vertebrate species recorded in survey and desktop results.....	42
Table 5-5	Details of significant vertebrate fauna recorded during the field survey .....	44
Table 5-6	Likelihood of occurrence for significant vertebrates identified in the desktop.....	45
Table 5-7	Extent and description of each SRE habitat in the study area .....	49
Table 5-8	Specimens from SRE groups recorded in the field survey .....	50
Table 5-9	Consideration of potential survey limitations.....	52

## LIST OF APPENDICES

Appendix A	Survey site locations
Appendix B	Vertebrate fauna desktop and field survey results
Appendix C	Fauna species by site matrix

# 1 INTRODUCTION

Australian Capital Equity Pty Ltd (ACE) is seeking to develop the Napier Downs Irrigation Project (the Project) located on Napier Downs Station, approximately 77 km east of Derby, Western Australia (WA; Figure 1-1). The Project will entail the development of centre irrigation pivots which will be used to produce fodder crops for cattle stocked on Napier Downs and nearby stations. Water will be sourced from the Grant Group Aquifer.

In August 2021, Phoenix Environmental Sciences Pty Ltd (Phoenix) was commissioned by ACE to undertake a baseline terrestrial fauna survey for the Project. The purpose of the survey was to inform environmental approvals for the Project.

The study area is located in the Shire of Derby-West Kimberley and the Fitzroy Trough subregion (DAL01) of the Dampierland bioregion and the Northern Botanical Province as defined by EPA (2016b).

## 1.1 BACKGROUND

Phoenix (2019) was engaged by ACE in March 2019 to conduct a desktop assessment of terrestrial vertebrate fauna and short-range endemic (SRE) invertebrate fauna in accordance with relevant EPA guidance (Department of Mines and Petroleum 2016; EPA 2016a, d). The original desktop review was undertaken for 2 previous options for the irrigation area (Option 1 and Option 2; Figure 1-2); with a desktop addendum later prepared specifically for a third option located in Scrubby Paddock (Option 3; Figure 1-2) (Phoenix 2020).

The objective of the desktop reviews was to identify the following:

- potential significant fauna values that may be present in the study areas
- any potential values that may represent significant constraints for the Project
- proposed scope of field survey requirements for the Project.

The current study area falls partially outside the Option 3 area (Figure 1-2); however, the desktop review for Option 3 was still considered valid for the current study area.

The desktop review identified that several significant terrestrial fauna species were considered to have the potential to occur within in the study area, including 10 mammal and 16 bird species. Several SRE records were found proximal to study area, therefore, it was considered possible that SRE taxa were also present in the study area.

Based on the desktop findings, Phoenix recommended a detailed survey to verify fauna habitats and assess suitability of the study area for species identified in the desktop review (Phoenix 2020). Field verification of the potential value of the study area to SREs, particularly land snails, was also deemed required. The proposed detailed survey involved:

- detailed habitat assessment and mapping
- targeted survey for significant mammals that may occur (Bilby, Northern Quoll, Golden Bandicoot, Kimberley Brush-tailed Phascogale, Northern Brushtail Possum and Northern Short-tailed Mouse), including:
  - plot sampling for Bilby within and adjacent to the study area
  - camera trapping within the study area and along Hawkstone Creek
  - searches for signs of presence
- acoustic recordings for the significant bat species and Night Parrot
- avifauna surveys

- targeted survey for SRE invertebrates, including systematic sampling and characterisation and mapping of SRE habitats.

Discussions of desktop assessment results and proposed survey methodology with the Department of Water and Environmental Regulation (DWER) were undertaken and it was agreed that the proposed survey appeared adequate to inform environmental impact assessment for the study area (Ryan Mincham, letter dated 29/1/2021, to James McMahon).

In addition to the detailed survey carried out in the study area, additional reconnaissance surveys were undertaken at three locations within the potential zone of groundwater drawdown for the Project in late June and early July 2022. These areas included: Ngooderoodyne Spring, Hawkstone Creek and the Lennard River. Desktop findings and survey works discussed in this report were conducted for terrestrial fauna species and were specific to the study area, and the three potential groundwater drawdown sites.

## 1.2 SCOPE OF WORK

The scope of work for the baseline terrestrial fauna survey was as follows:

- Targeted detailed terrestrial fauna survey, including:
  - detailed habitat assessment and mapping within the study area; mapping within a wider 1 km buffer of the study area for context, to inform the environmental impact assessment
  - identification and mapping of habitat for significant fauna species within the study area and 1 km buffer
  - targeted survey for Threatened and Priority mammal species - Bilby *Macrotis lagotis* (Vulnerable (VU)), Northern Quoll *Dasyurus hallucatus* (Endangered), Golden Bandicoot *Isoodon auratus auratus* (VU), Northern Short-tailed Mouse *Leggadina lakedownensis* (Priority (P) 4), Kimberley Brush-tailed Phascogale *Phascogale tapoatafa kimberleyensis* (VU) and Northern Brushtail Possum *Trichosurus vulpecula arnhemensis* (VU)<sup>1</sup>, including trapping, plot sampling and camera trapping within the study area and 1 km buffer, with emphasis on the study area
  - daytime searches for signs of significant fauna activity, nocturnal spotlighting where possible
  - acoustic recordings for significant bat species – Northern Leaf-nosed Bat *Hipposideros stenotis* (P2), Ghost Bat *Macroderma gigas* (VU), Orange Leaf-nosed Bat *Rhinonictes aurantia* (P4), Bare-rumped Sheath-tailed Bat *Saccolaimus saccolaimus nudicluniatatus* (P3) and Yellow-lipped Cave Bat *Vespadelus douglasorum* (P2)
  - habitat assessment and, if required, acoustic recordings for Night Parrot *Pezoporus occidentalis* (EN Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act); Critically Endangered (CR) under the *Biodiversity Conservation Act 2016* (BC Act)
  - avifauna surveys for significant bird species within the study area and adjacent riparian habitats within 1 km buffer, and avifauna census generally

---


<sup>1</sup> Two other significant mammals identified in the desktop review – West Kimberley Black-footed Rock-wallaby *Petrogale lateralis subsp. (West Kimberley)* (VU EPBC Act; EN BC Act) and Rock Ringtail Possum *Petropseudes dahli* (P3) – were considered unlikely to occur in the study area. The survey will collect sufficient habitat data to confirm the study area is unsuitable for these species.












- Level 2 survey for SRE invertebrates, including characterisation and mapping of SRE habitats and systematic sampling in SRE prospective habitats within the study area
- searches for presence of any pools or other surface water features in the study area and adjacent toward Hawkstone Creek that may serve as refuges for the Freshwater Sawfish.
- prepare a technical report on the findings of the surveys.

### **1.3 STUDY AREA**

The study area is approximately 586.5 ha and located within Scrubby Paddock on Napier Downs pastoral lease (LPL N049855; Figure 1-1). The fauna habitat assessments and mapping encompass the study area plus an additional 1 km buffer around the study area. Including the buffer, the total mapped area (referred to hereafter as 'habitat mapping area') is approximately 1,846 ha (Figure 1-1). Additionally, basic fauna assessments were also conducted at pools and other surface water features in the area surrounding the study area.




<b>Australian Capital Equity Napier Downs Irrigation Project</b>	
Project No	1452
Date	11/01/2023
Drawn by	BK
Map author	BQ
	
	
1:750,000 (at A3) <span style="float: right;">GDA 1994 MGA Zone 51</span>	

-  Study area
-  Habitat mapping area
-  Napier Downs Station
-  Indigenous Protected Areas
-  Environmentally sensitive areas
-  DBCA managed land
-  Lakes
-  West Kimberley National Heritage Place

**Figure 1-1**  
**Project location and study area**



All information within this map is current as of 11/01/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.



<b>Australian Capital Equity Napier Downs Irrigation Project</b>		
Project No	1452	
Date	1/12/2023	
Drawn by	BK	
Map author	BQ	
1:570,200 (at A4)		GDA 1994 MGA Zone 51

- Study area
- Option 1
- Option 2
- Option 3
- Habitat mapping area
- 40 km desktop study area

**Figure 1-2**  
**Study area in relation to previous desktop areas**



All information within this map is current as of 1/12/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

## 2. LEGISLATIVE CONTEXT

The protection of fauna in WA is principally governed by 3 acts:

- Commonwealth EPBC Act
- State BC Act
- State *Environmental Protection Act 1986* (EP Act).

The BC Act came into full effect on 1 January 2019 and replaced the functions of the *Wildlife Conservation Act 1950* (WC Act).

### 2.1 COMMONWEALTH

The EPBC Act is administered by the Federal Department of Climate Change, Energy, the Environment and Water (DCCEE). The EPBC Act provides for the listing of Threatened fauna as matters of National Environmental Significance (NES). Under the EPBC Act, actions that have, or are likely to have, a significant impact on a matter of NES, require approval from the Australian Government Minister for the Environment through a formal referral process.

Conservation categories applicable to Threatened fauna species under the EPBC Act are as follows:

- Extinct (EX)<sup>2</sup> – there is no reasonable doubt that the last individual has died
- Extinct in the Wild (EW) – taxa known to survive only in captivity
- Critically Endangered (CR) – taxa facing an extremely high risk of extinction in the wild in the immediate future
- Endangered (EN) – taxa facing a very high risk of extinction in the wild in the near future
- Vulnerable (VU) – taxa facing a high risk of extinction in the wild in the medium-term
- Conservation Dependent (CD)<sup>2</sup> – taxa whose survival depends upon ongoing conservation measures; without these measures, a conservation dependent taxon would be classified as Vulnerable, Endangered or Critically Endangered.

The EPBC Act is also the enabling legislation for protection of Migratory species as matters of NES under several international agreements:

- Japan-Australia Migratory Bird Agreement (JAMBA)
- China-Australia Migratory Bird Agreement (CAMBA)
- Convention on the Conservation of Migratory Species of Wild Animals (Bonn)
- Republic of Korea-Australia Migratory Bird Agreement (ROKAMBA).

### 2.2 STATE

#### 2.2.1 Threatened and Priority species

In WA, the BC Act provides for the listing of Threatened fauna species (Government of Western Australia 2018a, b)<sup>3</sup> in the following categories:

---

<sup>2</sup> Species listed as Extinct and Conservation Dependent are not matters of NES and therefore do not trigger the EPBC Act.

<sup>3</sup> The *Wildlife Conservation (Specially Protected Fauna) Notice 2018* and the *Wildlife Conservation (Rare Flora) Notice 2018* have been transitioned under regulations 170, 171 and 172 of the *Biodiversity Conservation Regulations 2018* to be the lists of Threatened, Extinct and Specially Protected species under Part 2 of the BC Act.

- Critically Endangered (CR) – species facing an extremely high risk of extinction in the wild in the immediate future<sup>4</sup>
- Endangered (EN) – species facing a very high risk of extinction in the wild in the near future<sup>4</sup>
- Vulnerable (VU) – species facing a high risk of extinction in the wild in the medium term future<sup>4</sup>.

Species may also be listed as specially protected (SP) under the BC Act in one or more of the following categories:

- species of special conservation interest (conservation dependent fauna, CD) – species with a naturally low population, restricted natural range, of special interest to science, or subject to or recovering from a significant population decline or reduction in natural range
- migratory species (Mig.), including birds subject to international agreement
- species otherwise in need of special protection (OS).

The Department of Biodiversity, Conservation and Attractions (DBCA) administers the BC Act and maintains a non-statutory list of Priority fauna. Priority species are still considered to be of conservation significance – that is they may be Threatened – but cannot be considered for listing under the BC Act until there is adequate understanding of threat levels imposed on them. Species on the Priority fauna lists are assigned to one of four Priority (P) categories, P1 (highest) – P4 (lowest), based on level of knowledge/concern.

### 2.2.2 Critical habitat

Under the BC Act, habitat is eligible for listing as critical habitat if it is critical to the survival of a Threatened species, or a TEC and its listing is otherwise in accordance with the ministerial guidelines.

### 2.2.3 Environmentally Sensitive Areas

Under section 51B of the EP Act the Minister for Environment may declare by notice either a specified area of the State or a class of areas of the State to be ESAs. ESAs are declared in the *Environmental Protection (Environmentally Sensitive Areas) Notice 2005*, which was gazetted on 8 April 2005 (Government of Western Australia 2005).

ESAs are areas where the vegetation has high conservation value. Several types of areas are declared ESAs including:

- the area covered by vegetation within 50 metres (m) of Threatened flora, to the extent to which the vegetation is continuous with the vegetation in which the Threatened flora is located
- the area covered by a TEC
- a defined wetland (Ramsar wetlands, conservation category wetlands and nationally important wetlands) and the area within 50 m of the wetland
- Bush Forever sites.

### 2.2.4 Short-range endemic invertebrates

Short-range endemic (SRE) fauna are defined as animals that display restricted geographic distributions, nominally less than 10,000 km<sup>2</sup>, that may also be disjunct and highly localised (Harvey 2002). EPA (2016a) identifies species with restricted distributions as being significant fauna in the context of environmental impact assessments (EIA). SRE fauna need to be considered in EIA as

---

<sup>4</sup> As determined in accordance with criteria set out in the ministerial guidelines.

localised, small populations of species that are generally at greater risk of changes in conservation status due to environmental change than other, more widely distributed taxa.

Short-range endemism in terrestrial invertebrates is believed to have evolved through 2 primary processes (Harvey 2002):

- Relictual – where the drying climate reduced the area of suitable habitat available to a species, forcing a range contraction. Such habitats typically maintain historic mesic conditions (e.g. south-facing rock faces or slopes of mountains or gullies)
- Habitat speciality – where species settled in particular isolated habitat types (e.g. rocky outcrops) by means of dispersal and evolved in isolation into distinct species.

However, SRE invertebrates have also been reported in more widespread habitats such as spinifex plains or woodlands, mainly in groups with low dispersal capabilities, for example mygalomorph spiders and millipedes (see for example Car & Harvey 2014; Rix et al. 2018).

There can be uncertainty in categorising a specimen as an SRE due to several factors including poor regional survey density, lack of taxonomic research and problems of identification, i.e. specimens that may represent SREs cannot be identified to species level based on the life stage at hand. For example, in contrast to mature males, juvenile and female millipedes, mygalomorph spiders and scorpions cannot be identified to species level. Molecular techniques such as ‘barcoding’ (Hebert *et al.* 2003a; Hebert *et al.* 2003b) are routinely employed to overcome taxonomic or identification problems.

### 3. EXISTING ENVIRONMENT

#### 3.1 INTERIM BIOGEOGRAPHIC REGIONALISATION OF AUSTRALIA

The Interim Biogeographic Regionalisation of Australia (IBRA) classifies Australia’s landscapes into large ‘bioregions’ and ‘subregions’ based on climate, geology, landform, native vegetation and species information (DoEE 2016). The study area is located in the Fitzroy Trough subregion (DL1) of the Dampierland bioregion (Figure 3-1). The Fitzroy Trough subregion comprised of four basic components, described as (Graham 2001b):

- Quaternary sandplain overlying Jurassic and Mesozoic sandstones with Pindan, with hummock grasslands on hills
- Quaternary marine deposits on coastal plains, with mangal, samphire – *Sporobolus* spp. Grasslands, *Melaleuca alsophila* low forests, and *Spinifex* spp. – *Crotalaria* spp., strand communities
- Quaternary alluvial plains associated with the Permian and Mesozoic sediments of Fitzroy Trough support tree savannahs of ribbon grass (*Chrysopogon* spp.), bluegrass (*Dichanthium* spp.) and Mitchell grass (*Astrelba* spp.) scattered coolabah (*Eucalyptus microtheca*) – *Bauhinia cunninghamii*, with riparian forests of river red gum (*Eucalyptus camaldulensis*) and Cadjeput (*Melaleuca* spp.) fringe drainages
- Devonian reef limestones in the north and east supporting sparse tree steppe over lobed spinifex (*Triodia intermedia*) and limestone spinifex (*T. wiseana*) hummock grasses.

The subregion experiences a dry hot tropical and semi-arid climate with summer rainfall, with average rainfall between 500–800 mm, often, often influenced by cyclonic activity in the northwest of WA.

#### 3.2 LAND SYSTEMS AND SURFACE GEOLOGY

DPIRD undertakes land system mapping for WA using a nesting soil-landscape mapping hierarchy (Schoknecht & Payne 2011). While the primary purpose of the mapping is to inform pastoral and

agricultural land capability, it is also useful for informing biological assessments. Under this hierarchy, land systems are defined as areas with recurring patterns of landforms, soils, vegetation and drainage (Payne & Leighton 2004).



The study area intersects three land systems but falls predominantly within one of these (Table 3-1; Figure 3-2).






**Table 3-1 Land systems and extent in study area**

Land system	Description	Area (ha)	% of study area
Sisters System	Low sandy plateaux and lower slopes supporting pindan woodlands with acacias and eucalypts and curly spinifex-ribbon grass, and valley plains supporting mixed woodlands with ribbon grass.	537.0	91.6
Wanganut System	Sandplains and linear dunes supporting pindan woodlands with acacias and bloodwoods and curly spinifex- ribbon grass, and broad low-lying swales supporting bloodwood-grey box woodlands with curly spinifex-ribbon grass.	0.7	0.1
Yeeda System	Red sandplains supporting pindan vegetation with dense acacia shrubs, scattered bloodwood and grey box trees and curly spinifex and ribbon grass.	48.8	8.3
<b>Total</b>		<b>586.5</b>	<b>100</b>

According to the Surface Geology of Australia 1:1,000,000 scale, Western Australia database (Stewart *et al.* 2008), the study area intersects a single geological formation (Figure 3-2); Sand plain 38499 (Czs), which is described as ‘Sand or gravel plains; quartz sand sheets commonly with ferruginous pisoliths or pebbles, minor clay; local calcrete, laterite, silcrete, silt, clay, alluvium, colluvium, aeolian sand’.



<b>Australian Capital Equity Napier Downs Irrigation Project</b>	
Project No	1452
Date	1/11/2023
Drawn by	BK
Map author	GW
	
	
1:400,000 (at A4) <span style="float: right;">GDA 1994 MGA Zone 51</span>	


-  Study area
  -  Habitat mapping area
- Region, subregion**
-  Central Kimberley, Mount Eliza
  -  Dampierland, Fitzroy Trough
  -  Northern Kimberley, Mitchell

**Figure 3-1**  
**Study area in relation to  
 IBRA bioregions and  
 subregions**



All information within this map is current as of 1/11/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.





**Australian Capital Equity  
Napier Downs Irrigation Project**

Project No 1452  
Date 1/11/2023  
Drawn by BK  
Map author GW

0 2.5 5  
Kilometers

1:174,300 (at A4) GDA 1994 MGA Zone 51

Study area	Qa	Fossil2 System
Habitat mapping area	Qd	Gogo System
<b>Surface geology</b>	Qrlb	Neillabubica System
Czl	<b>Land system</b>	Sisters System
Czs	Alexander System	Wanganut System
DCIf	Djada System	Yeeda System
PsIj	Duffer System	

**Figure 3-2**  
**Land systems and surface geology in the study area**



All information within this map is current as of 1/11/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

### 3.3 CLIMATE AND WEATHER

The climate of the Fitzroy Trough subregion is described as dry hot tropical and semi-arid with summer rainfall. The average annual rainfall is between 500 – 800 mm (Graham 2001a). The nearest Bureau of Meteorology (BoM) weather station with comprehensive data collection and recent historic climate data is Derby Airport (no. 003032), Latitude: 17.39°S Longitude 123.68°E, located 77 km west of the study area.

Derby Airport records the highest mean maximum monthly temperature (38.3°C) in November (lowest in June, 30.8°C) and the lowest minimum mean monthly temperature (14.6°C) in July (highest in December, 26.3°C; Figure 3-3). Median annual rainfall is 706.2 mm with January and February recording the highest monthly median (172.2 and 169.0 mm respectively; Figure 3-3).

Daily mean maximum temperatures recorded at Derby Airport in the months preceding the survey were slightly above the long-term averages. Daily minimum temperatures were consistent with long-term averages, except for July when the survey took place in which the minimum temperature was 3°C cooler than the long-term average (Figure 3-3).

Records from Derby Airport indicate that most of the wet season’s rainfall was later in the season than usual, with February being the wettest month and March receiving 40% more rainfall than the long-term median. Rainfall also continued into May and June, both of which typically receive far less rainfall as the median for both those months is less than 1 mm (Figure 3-3).




Figure 3-3 Annual climate and weather data for Derby Airport (no. 003032) and mean monthly data for the 12 months preceding the survey (BoM 2022)

### 3.4 LAND USE

The dominant land uses in the Fitzroy Trough subregion are native pasture grazing, conservation reserves and unallocated crown land (Graham 2001a). The study area occurs entirely within the Napier Downs pastoral lease.

### 3.5 NATIONAL HERITAGE PLACES, CONSERVATION RESERVES AND ESAs

The study area is not situated within any conservation reserves or Environmentally Sensitive Areas (ESAs); however, an Indigenous Protected Area (IPA) called Wilinggin is located 3 km to the east (Figure 1-1). The closest conservation reserve, King Leopold Ranges Conservation Park, is situated 58 km northeast and the closest ESA is 33.5 km northeast of the study area (Figure 1-1).

The study area is situated within the West Kimberley National Heritage Place, which is listed on the National Heritage List and therefore a matter of National Environmental Significance (NES; Figure 1-1). This listing is vast in extent, covering 949.9 km<sup>2</sup> of the Kimberley region, and is recognised as nationally significant under several criteria (DoEE 2019), with many specific significant features identified, including (but not limited to):

- the King Leopold orogen, Kimberley ria coast, Lennard Shelf – for geological significance
- the Devonian Reefs, Gogo fossil sites, Dampier Coast – for evolutionary/fossil record
- northern Kimberley coast and islands, the Kimberley Plateau and the west Kimberley Devonian reefs – for their rich biodiversity
- vine thickets – for endemic invertebrates
- river systems (the Drysdale, Prince Regent, Roe, Moran, Carson, Isdell, Mitchell and King Edward Rivers) – as refuges for freshwater fish species
- Roebuck Bay – for Migratory shorebird habitat
- Kimberley coast from the Buccaneer Archipelago to King George River, Mitchell River National Park, King George Falls, King George River, Geiki Gorge Conservation Park, Geikie Gorge National Park, Windjana Gorge National Park, King Leopold Ranges and the Kimberley coast from the Buccaneer Archipelago to King George River – for aesthetic landscape values
- numerous indigenous heritage sites of national significance.

While the study area is situated over the King Leopold Orogen geological province; it does not intersect any of the other specific features described in the West Kimberley National Heritage Place; the Monsoon vine thickets and Camaenid land snails of limestone ranges (Napier Range) Priority Ecological Community (PEC) is the closest, located approximately 20.8 km to the northeast.

## 4. METHODS

The terrestrial fauna survey was conducted in accordance with relevant survey guidelines and guidance, including:

- *EPA Environmental Factor Guideline: Terrestrial fauna* (EPA 2016a)
- *EPA Technical Guidance: Sampling of short range endemic invertebrate fauna* (EPA 2016c)
- *EPA Technical Guidance: Terrestrial vertebrate fauna surveys for environmental impact assessment* (EPA 2020)
- *Interim guideline for preliminary surveys of Night Parrot (*Pezoporus occidentalis*) in Western Australia* (DPaW 2017)
- *Approved conservation advice for *Pezoporus occidentalis* (Night Parrot)* (DSEWPac 2008)
- *EPBC Act referral guideline for the endangered Northern Quoll *Dasyurus hallucatus** (DoE 2016)
- *Guideline for the survey and relocation of bilby in Western Australia (draft)* (DBCA 2018)
- *Survey guidelines for Australia's threatened mammals. Guidelines for detecting mammals listed as threatened under the EPBC Act* (DSEWPac 2011a)
- *Survey guidelines for Australia's threatened birds. Guidelines for detecting birds listed as threatened under the EPBC Act* (DSEWPac 2010)
- *Survey guidelines for Australia's threatened reptiles. Guidelines for detecting reptiles listed as threatened under the EPBC Act* (DSEWPac 2011b).

### 4.1 DESKTOP REVIEW

The results of the previous desktop assessments (Phoenix 2019, 2020) were used to inform this survey. The study area lies inside the 40 km buffer applied for database searches conducted for the Phoenix (2019) desktop assessment. Subsequently, the results of the database searches were applicable to the current study area. Database search buffers are therefore not of a precise length around the perimeter of the study area.

Searches of several biological databases were undertaken to identify and prepare lists of significant fauna that may occur within the study area (Table 4-1). The desktop results from the previous assessment for Option 3 (Phoenix 2020) were updated for the current study area, and these results are presented in this report.

**Table 4-1 Database searches conducted for the desktop review**

Database	Target group/s	Search coordinates and extent
Protected Matters Search Tool (DCCEE 2022a)	EPBC Act Threatened flora, fauna, and ecological communities	Centre point of study area with a 40 km buffer
DBCA Threatened and Priority Fauna Database (DBCA 2019b)	Threatened and Priority fauna	Centre point of Option 1 and Option 2 plus a 40 km buffer
DBCA Threatened and Priority Ecological Communities Database (DBCA 2019c)	TECs and PECs	Centre point of Option 1 and Option 2 plus a 40 km buffer
DBCA NatureMap Database (DBCA 2019a)	Flora and fauna records	Centre point of Option 1 and Option 2 plus a 40 km buffer
BirdLife Birddata (Birdlife Australia 2019)	Avifauna	Centre point of Option 1 and Option 2 plus a 40 km buffer

Database	Target group/s	Search coordinates and extent
WA Museum Arachnid and Myriapod Database, Mollusca Database (WAM 2019)	Arachnid, myriapod and mollusc SREs	Centre point of Option 1 and Option 2 plus a 100 km buffer

## 4.2 FIELD SURVEY

### 4.2.1 Survey timing

Field survey dates are provided in Table 4-2.

**Table 4-2 Survey dates**

Survey type	Season	Dates
Field survey (trip 1) – Detailed and reconnaissance survey	Winter	27 June – 5 July 2022
Field survey (trip 2) – SRE pit trap retrieval	Winter	14 – 16 August 2022

### 4.2.2 Field methods

The terrestrial fauna survey comprised of a combination of reconnaissance assessment, systematic sampling for fauna assemblage and targeted methods for significant fauna considered to have potential to occur, as identified in the desktop review (Phoenix 2020).

Field methods for the fauna survey of the study area included:

- initial reconnaissance (see 4.2.2.1)
- habitat assessments and mapping (see 4.2.2.2)
- active diurnal and nocturnal searches (see 4.2.2.3)
- systematic trapping (4.2.2.4)
- targeted survey for Bilby (4.2.2.5)
- bat echolocation recordings (4.2.2.6)
- habitat assessment for Night Parrot (see 4.2.2.7)
- avifauna surveys (4.2.2.8)
- opportunistic records (see 4.2.2.9)
- SRE invertebrate sampling (4.2.2.11)
- SRE potential habitat rating (4.2.2.12).

A total of 9 survey sites were sampled within the study area, and 7 sites outside the study area at pools and other surface water features within 20 km of the study area (Figure 4-1; Appendix A).

#### 4.2.2.1 Initial reconnaissance

An initial reconnaissance assessment was conducted throughout the study area at the start of the survey to identify suitable habitat resources for significant fauna species, including shelter sites such as tree hollows and tree species (for arboreal mammals), rocky habitats, caves, and water resources. Information gathered in the reconnaissance was used to refine and finalise the survey plan, methods, and site locations.

#### 4.2.2.2 Habitat assessment and mapping

Initial habitat characterisation was undertaken using various remote geographical tools, including aerial photography (Google Earth®), land system maps and topographic maps. Habitats with the potential to support significant terrestrial fauna species were identified based on known habitats of such species within the Dampierland bioregion. Tentative sites were selected for the terrestrial fauna survey to represent all habitat types. Final survey site selection was conducted after ground-truthing of site characteristics.

At the broadest scale, site selection considered aspect, topography, and land systems. At the finer scale, consideration was given to proximity to water bodies (drainage lines and creek), vegetation complexes and condition and soil type. Sites were primarily chosen to represent the best example of distinct habitats within the broader habitat associations of the study area with a focus on species of conservation significance identified in the desktop review. Habitat descriptions and characteristics were recorded at all survey sites within the study area, and at pools and other surface water features in the surrounding area (Figure 4-1; Table 4-3).

Mapping of broad fauna habitats was undertaken based on mapped vegetation types, site-based habitat assessments and additional habitat data points collected during the field survey. Mapping within a 1 km buffer was extrapolated from the detailed mapping within the study area. Mapping of habitat for significant fauna was derived for broad fauna habitat classifications.

**Table 4-3 Terrestrial fauna survey effort**

Site name	Site type	Birding (hours)	Large cage trap (trap nights)	Large Elliot trap (trap nights)	Active searching (hours)	Ultrasonic recording (nights)	Wet pitfall trap (trap nights)
SITE01	Systematic fauna site	2.5	35	35	1.5	3	198
SITE02	Systematic fauna site	1	35	35	1.4		198
SITE03	Systematic fauna site	1	35	35	2		
SITE04	Systematic fauna site	1	35	35	2		198
SITE05	Systematic fauna site	0.6	35	35	1.5		
SITE06	Systematic fauna site	1	35	35	1.5		
SITE07	Systematic fauna site	1.5	35	35	2		
SITE08	Systematic fauna site	2	35	35	2.2	3	198
SITE09	Opportunistic fauna site	1.5	7	7	1		
SITE10	Regional fauna site	0.5			1		
SITE11	Regional fauna site	0.5			1		
SITE12	Regional fauna site	0.7			1		
SITE13	Regional fauna site	0.5			1		
SITE14	Regional fauna site	0.5			2		
SITE15	Regional fauna site	1			3		
SITE16	Regional fauna site	2			3		

#### 4.2.2.3 Active diurnal and nocturnal searches

Active searches were undertaken at all 16 fauna sites (Figure 4-1). Active searches primarily targeted diurnal herpetofauna and mammals from direct sightings and secondary evidence. Searches focused primarily on significant species identified in the desktop review as potentially occurring within the study area, including Northern Quoll, Golden Bandicoot, Northern-short-tailed Mouse, Bilby, Western Kimberley Black-footed Rock-wallaby, Rock Ringtail Possum, Kimberley Brush-tailed Phascogale and Northern Brushtail Possum.

Searches were undertaken in any observable microhabitats considered likely to support mammals, reptiles, and amphibians. Techniques included: raking leaf and bark litter, overturning logs, searching beneath the bark of trees, investigating dead trees and logs, investigating burrows, investigating infrastructure ruins or disused building materials and identifying any secondary evidence including tracks, diggings, scats, fur or sloughs (shed skins), predation or feeding sites, and fauna constructed structures such as pebble mounds or nests. A minimum of 1 person hour was spent active searching at each site for a total of 27.1 hours over the duration of the field survey (Table 4-3).

#### 4.2.2.4 Systematic trapping

Eight systematic trapping sites were established to capture terrestrial mammals, reptiles, and amphibians (Figure 4-1). Each site contained 5 Sheffield cage traps (60 cm x 20 cm x 20 cm), and 5 aluminium box Elliott traps spaced approximately 20 m apart in a straight line. The 2 trap types were alternated across the line, and all traps were baited with a universal bait mixture consisting of oats, peanut butter, and sardines.

Elliott and Sheffield traps were shrouded with reflective closed cell insulation (R2.5 rated) to provide shade and protection for any captured animals. All traps were given as much shade as possible under/around vegetation.

Traps were open for 7 consecutive nights and checked within 3 hours of sunrise each day. Baits were removed and replaced every second day. The total vertebrate trapping effort for the 8 systematic trapping sites during the surveys was 280 trap-nights (Table 4-3), where a trap-night is defined as one trap remaining open for one night.

In addition to the 8 systematic trapping sites, an opportunistic ninth site (SITE09) containing just one Sheffield and one Elliott was set up near a large termite mound close to where the field team camped each night. The traps were set in the same manner as the systematic trapping sites and were deployed for the same duration.

#### 4.2.2.5 Targeted survey for Bilby

Suitability of habitat for Bilby was assessed at all sites based on substrate, vegetation structure and density. In the event suitable habitat was found, standardised plot surveys would be conducted over 2 ha plots (~142 m x 142 m) to systematically search for signs of Bilby presence including tracks, scats, foraging diggings and/or burrows. No suitable habitat was found, and no plot surveys were conducted.

#### 4.2.2.6 Bat echolocation recordings

Song Meter SM2 recording devices were used to record bat echolocation calls at 2 sites during the field survey (SITE01 and SITE08; Figure 4-1). Recording devices were deployed for 3 nights of continuous recording between 30 minutes prior to sunset to 30 minutes after sunrise (Table 4-3). Devices were aimed at a 45° angle to the ground. The Song Meters were positioned in areas of habitat



likely to have increased insect activity and provide foraging areas, movement corridors or potential roosting sites for bats.

#### 4.2.2.7 Habitat assessment for Night Parrot

Habitat assessments for Night Parrot was conducted throughout the study area. Song Meter SM4 acoustic recording devices were to be deployed in areas identified as providing suitable roosting habitat as defined by DPaW (2017), i.e. old spinifex grassland with ring-forming hummocks greater than 50 cm in height. No suitable habitat was identified across the study area and so no recording devices were deployed.

#### 4.2.2.8 Avifauna surveys

Avifauna surveys were undertaken at each systematic site (Figure 4-1; Table 4-3). Each avifauna survey encompassed an area of 2 ha and was confined to a single habitat type. All avifauna surveys were a minimum of 20 minutes. These surveys were undertaken throughout the day with a focus on periods of higher activity around sunrise and sunset. Surveys consisted of bird recordings from visual sightings and call recognition. A total of 17.8 person hours of avifauna surveys was undertaken during the field survey (Table 4-3). Additional avifauna observations were also recorded opportunistically while other field work was completed, including observations made during travel and active searches.

#### 4.2.2.9 Opportunistic records

All vertebrate fauna observed while traversing the study area and sites surrounding the study area were recorded.

#### 4.2.2.10 Analysis of survey completeness

A species accumulation curve was produced on a samples and abundance basis to obtain an estimate of survey completeness (i.e. whether the collection adequately represents the vertebrate fauna assemblage of the study area) for the systematic methods (vertebrate fauna trapping using Sheffield cage and Elliott aluminium box traps and avifauna surveys at the 8 systematic trapping sites) completed within the study area. All sample types were aggregated by day and no data transformation was undertaken.

#### 4.2.2.11 SRE invertebrate sampling

Sampling for SRE invertebrates was conducted at 4 systematic SRE sites across the study area (Figure 4-1), in areas identified as suitable habitat for SREs. Sampling comprised the following methods:

- wet pit trapping
- active foraging.

Four wet pitfall trapping sites were established, each comprising of 5 one litre plastic containers with a 70 mm diameter dug in flush with the surface in suitable microhabitats at each site. Pit traps were half-filled with a 50:50 mixture of propylene glycol and alcohol. All traps were covered with a plastic lid elevated 25 mm above the trap with wooden blocks to minimise by-catch of vertebrates where possible. Traps remained open following the setup and were retrieved 6 weeks later.

Active foraging for SRE invertebrate groups comprised inspection of logs, larger plant debris, the underside of bark of larger trees and the underside of rocks. Methodical searches were conducted amongst the leaf litter of shade-bearing tall shrubs and trees, including raking of litter, and spinifex

bases were inspected thoroughly. Rocks and rock crevices were inspected, particularly for pseudoscorpions.

A standardised approach was undertaken whereby each site was sampled for one person hour (concurrently with active searches for vertebrate fauna), a total search effort of approximately 27.1 hours (Table 4-3). Trapdoor spider burrows identified during the searches were excavated if they were considered inhabited. Excavation involved removing soil from around the burrow to carefully expose the burrow chamber and remove the spider.

#### 4.2.2.12 SRE potential habitat rating

Fauna habitat was assessed for its potential to support endemic SRE species and communities. Potential SRE habitat was rated as follows:

- High – defined/known areas of habitat that contain elements that often give rise to specialisation or dependency in invertebrate fauna, such as aspect (e.g. south-facing slopes, geological features (e.g. granite), soil types that retain water (e.g. clay, loam). These habitats may also include habitat isolates which have the capacity to restrict dispersal.
- Low – areas of largely in-tact native vegetation that occur broadly across the landscape, are less incised and typically link more restricted habitats. This may include land that was cleared but has since been rehabilitated or is in the process of being rehabilitated.
- None – land that has been previously cleared for other uses that no longer contains native vegetation.

#### 4.2.2.13 SRE status rating

Currently, there is no accepted system to determine the likelihood that a species is an SRE. The WA Museum applies 3 categories: confirmed, potential, and widespread. Confirmed SREs are taxa for which the distribution is known to be less than 10,000 km<sup>2</sup>, the taxonomy is well known, and the group is well represented in collections and/ or via comprehensive sampling (WAM 2013). Potential SREs include those taxa for which there is incomplete knowledge of the geographic distribution of the group and its taxonomy, and the group is not well represented in collections. Phoenix applies 4 categories based on the WA Museum criteria (Table 4-4).

**Table 4-4 Short-range endemic categories**

SRE category	Criteria
Confirmed	Distribution <10,000 km <sup>2</sup> . Taxonomy of the group is well known (but not necessarily published); group is well represented in collections, in particular from the region in question; high levels of endemism exist in documented species; inference is often possible from immature specimens.
Potential	Distribution <10,000 km <sup>2</sup> . Taxonomically poorly resolved group; patchy distribution, often common in certain micro-habitats, but no other regional records; congeners (= species in the same genus) both widespread and restricted in distribution.
Widespread	Distribution >10,000 km <sup>2</sup> .
Uncertain	Taxonomy cannot be resolved to species level (i.e. indeterminate species designations due to sex, life stage or damage) and therefore species distribution remains uncertain).

#### 4.2.2.14 SRE Taxonomy

Initial higher-level (class, order, family) identifications of specimens are undertaken by Phoenix staff in Phoenix' invertebrate laboratory. Final specials designations are allocated using specialist morphological and/or molecular sequencing (Table 4-5).

Where possible identifications are on compared with reference material from the WA Museum and/or taxonomist reference collections.

**Table 4-5 Specialist taxonomists**

Person	Title	Taxa
Dr Erich S. Volschenk	Taxonomic consultant, Alacran	Scorpiones, Pseudoscorpiones
Dr Simon Judd	Taxonomic consultant	Isopoda
Dr Mark Harvey	Taxonomist, WA Museum	Mygalomorph spiders

Genomic analysis was undertaken for all specimens for which morphological identification didn't provide sufficient taxonomic resolution. A total of 45 specimens were sent for molecular analyses, comprising 11 mygalomorph spiders, 2 scorpions, 8 pseudoscorpions, and 3 isopods. Of these, only one species of spider, one species of isopod, 2 species of spider and 3 species of pseudoscorpion were successfully sequenced. Tissue from each specimen was obtained in Phoenix' laboratory and sequenced by Genotyping Australia.

Sequences were edited and analysed using Genius 2022.2. Sequences for comparison were sourced from GenBank (Benson *et al.* 2012) and Phoenix's DNA database using the megablast search function in Geneious. For each sequence, the most similar ten matches were retrieved. In cases where the retrieved sequences represented a species more than twice, then the two longest sequences were retained, and the shorter conspecific sequences discarded. Where megablast results yielded families differing from the morphological assessment, then additional sequences were obtained from GenBank, representing the morphological taxonomic assessment. If all the resulting blast sequences represented organisms from a different taxonomic class, sequences were discarded as likely contamination.

SRE specimens collected during the survey have been lodged with the WA Museum.

#### 4.2.2.15 Likelihood of occurrence assessment

Following the field survey, the likelihood of occurrence for each significant fauna species identified in the desktop review was assessed and assigned to one of four ratings:


- recorded – species recorded within the study area by previous or current survey
- likely – study area within current known range of species, suitable habitat within the study area and home range of species intersects study area based on known records
- possible – study area within current known range of species, suitable habitat within the study area and home range of species does not intersect study area based on known records
- unlikely – study area outside current known range of species or no suitable habitat present in study area.

### 4.2.3 Survey personnel

The personnel involved in the surveys are listed in Table 4-6. All survey work was carried out under relevant licences issued by DBCA under the BC Act, and DPIRD under the *Animal Welfare Act 2002* (AW Act; Table 4-6).

**Table 4-6 Survey personnel**

Name	Permit	Qualifications	Role/s
Floyd Holmes	Fauna taking (biological assessment) licence no. BA27000620	PhD, Hons (Biological Sciences), BSc (Physics & Conservation Biology)	Project manager, field survey, reporting
Simon Pynt	TFA (authorisation to take or disturb threatened species) no. 2022-0030 Scientific Use Licence no. U304/2022-2024	BSc (Zoology)	Field survey
Brooke Quick		BSc (Environmental Science)	Reporting, taxonomy & lab work
Lachlan Petersen		BSc (Environmental Science)	Taxonomy & lab work



Australian Capital Equity Napier Downes Irrigation Project		
Project No	1452	
Date	1/11/2023	
Drawn by	BK	
Map author	BQ	
1:303,400 (at A4)		GDA 1984 MGA Zone 51

- Study area
- Habitat mapping area
- Major watercourses
- Sites outside study area
- Sites inside study area

**Figure 4-1**  
**Terrestrial fauna survey sites**



All information within this map is current as of 1/11/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

## **5. RESULTS**

### **5.1 DESKTOP REVIEW**

#### **5.1.1 Vertebrate fauna**

A total of 42 significant species were identified in the desktop review, comprising 20 species listed under the EPBC Act and/or BC Act as Threatened (CR, EN, VU) or Specially Protected (OS) (Table 5-1; Figure 5-1). A further 18 species are listed as Migratory under the EPBC Act and BC Act, and 9 species are listed as Priority by the DBCA (Table 5-1). This assemblage included those species where suitable habitats may be present within the search extent and is not indicated by definitive records of the species.

No desktop records were returned for any significant fauna species within the study area; however, 5 significant species were recorded in 2013 from a flood plain site on the eastern side of Hawkstone Creek, 5.4 km east of the study area; Northern Quoll (EN), Gull-billed Tern (Mig.), Glossy Ibis (Mig.), Common Greenshank (Mig.) and Freshwater Crocodile (OS).

Table 5-1 Significant fauna identified in the desktop review

Species	Status <sup>1</sup>	Proximity to study area <sup>2</sup>	Habitat
<b>Reptiles</b>			
<i>Crocodylus johnstoni</i> Freshwater Crocodile	OS (BC Act)	5.4 km E	Freshwater Crocodiles inhabit various freshwater environments such as rivers, creeks, pools, billabongs, lagoons, and swamps. During the wet seasons as habitats become inundated with water, crocodiles can move throughout flood plains. During the dry season, crocodiles congregate to larger and deeper water bodies and will usually return to the same water body between years (Australian Museum 2020a).
<i>Crocodylus porosus</i> Salt-water Crocodile	OS (BC Act)	*	The Salt-water Crocodile is found in Australian coastal waters, estuaries, lakes, inland swamps and marshes (Webb <i>et al.</i> 1987). Despite the species' common name, the Salt-water Crocodile can persist in freshwater bodies (DCCEEW 2022b).
<b>Birds</b>			
<i>Actitis hypoleucos</i> Common Sandpiper	Mig (EPBC & BC Act)	42.4 km SE	Found across all Australian states, the Common Sandpiper never occurs in large flocks, mostly singly. In WA, the species is mostly coastal with some inland records (Geering <i>et al.</i> 2007). They are found across a wide range of wetlands: small ponds, large inlets and mudflats where they forage on the shore usually close to vegetation (DoEE 2020).
<i>Apus pacificus</i> Fork-tailed Swift	Mig (EPBC & BC Act)	*	The Fork-tailed Swift is a non-breeding visitor to all states and territories of Australia (Higgins 1999). It occurs in a wide range of dry or open habitats across most of WA and is uncommon to moderately common in the north-west (DoEE 2020).
<i>Botaurus poiciloptilus</i> Australasian Bittern	EN (EPBC & BC Act)	59.4 km SE	Found in freshwater or brackish swamps with dense vegetation (McKilligan 2005).
<i>Calidris acuminata</i> Sharp-tailed Sandpiper	Mig (EPBC & BC Act)	31.8 km S	Occurs on saline wetlands such as coastal lagoons, estuaries, bays, swamps, lakes, inundated grasslands, saltmarshes, river pools, creeks, floodplains and artificial wetlands (DCCEEW 2022b).
<i>Calidris ferruginea</i> Curlew Sandpiper	CR/Mig (EPBC Act) CR (BC Act)	*	Mainly occur on intertidal mudflats in sheltered coastal areas, also around non-tidal swamps, lakes, and lagoons near the coast. Less often inland around ephemeral and permanent lakes and waterholes, usually with bare edges of mud or sand (DCCEEW 2022b).
<i>Calidris melanotos</i> Pectoral Sandpiper	Mig (EPBC & BC Act)	*	Found in wetlands, inland as well as on the coast. Occurs on shallow fresh to saline wetlands, usually coastal or near-coastal but occasionally further inland. Prefers wetlands that have open fringing mudflats and low, emergent or fringing vegetation (DCCEEW 2022b).

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	Status <sup>1</sup>	Proximity to study area <sup>2</sup>	Habitat
<i>Cecropis daurica</i> Red-rumped Swallow	Mig (EPBC & BC Act)	*	Irregular visitor to Australia during its non-breeding seasons. Builds mud nests in mountains, sea cliffs and on buildings and other structures. Forages over open areas (Johnstone <i>et al.</i> 2013).
<i>Charadrius veredus</i> Oriental Plover	Mig (EPBC & BC Act)	24.5 km S	Inhabits coastal habitats such as estuarine mudflats and sandbanks, sandy or rocky ocean beaches or nearby reefs, or in near-coastal grasslands, before dispersing further inland where they inhabit flat, open, semi-arid or arid grasslands (DCCEEW 2022b).
<i>Erythrotriorchis radiatus</i> Red Goshawk	VU (EPBC & BC Act)	*	In the Kimberley area, Red Goshawks are most often found in extensive open forest, open woodlands and riparian vegetation dominated by mature <i>Eucalyptus tetradonta</i> , <i>E. miniata</i> (woollybutt), and <i>Melaleuca leucadendron</i> (cadjeputs) (Queensland Department of Environment and Resource Management 2012).
<i>Erythrura gouldiae</i> Gouldian Finch	EN (EPBC Act) P4 (DBCA)	30.4 km E	In WA, the species only occurs in the Kimberley. It inhabits open woodlands with favoured annual and perennial grasses (especially sorghum), a nearby source of surface water and, in the breeding season, unburnt hollow-bearing <i>Eucalyptus</i> trees (DoEE 2020).
<i>Falco hypoleucos</i> Grey Falcon	VU (EPBC & BC Act)	60.4 km SE	The Grey Falcon is a widespread but rare species inhabiting much of the semi-arid interior of Australia. Its distribution is centred on inland drainage systems. It has a large foraging range extending from timbered plains, such as <i>Acacia</i> shrublands, into open grasslands (Garnett & Crowley 2000b).
<i>Falco peregrinus</i> Peregrine Falcon	OS (BC Act)	27.8 km SW	Preferred habitat includes cliffs and wooded watercourses. Nesting occurs mainly on cliff ledges, granite outcrops, quarries and in trees with old raven or Wedge-tailed Eagle nests (Johnstone & Storr 1998).
<i>Gelochelidon nilotica</i> Gull-billed Tern	Mig (EPBC & BC Act)	5.4 km E	Occur in freshwater swamps, salt lakes, beaches, mudflats and sewage farms, and are rarely found over the ocean (DCCEEW 2022b).
<i>Glareola maldivarum</i> Oriental Pratincole	Mig (EPBC & BC Act)	*	In Australia, inhabits open plains, floodplains and grasslands, often with extensive bare areas (DCCEEW 2022b).
<i>Hirundo rustica</i> Barn Swallow	Mig (EPBC & BC Act)	*	Occurs in open country in coastal lowlands, an uncommon visitor to Australia (DCCEEW 2022b).
<i>Motacilla cinerea</i> Grey Wagtail	Mig (EPBC & BC Act)	*	Vagrant visitor to Australia that inhabits fast flowing streams and rivers (IUCN 2019).
<i>Motacilla flava</i> Yellow Wagtail	Mig (EPBC & BC Act)	*	Uncommon but regular visitor to the Kimberley; primarily inhabits a range of damp or wet habitats with low vegetation including damp meadows, marshes, waterside pastures, and sewage farms (IUCN 2019; Johnstone <i>et al.</i> 2013).



Species	Status <sup>1</sup>	Proximity to study area <sup>2</sup>	Habitat
<i>Numenius madagascariensis</i> Eastern Curlew	CR/Mig (EPBC Act) CR (BC Act)	*	Occurs mainly on intertidal mudflats, on exposed seagrass beds or mudflats (Geering <i>et al.</i> 2007). Also utilises sand spits of estuaries, mangroves, lake shores and ocean beaches.
<i>Pandion cristatus</i> Osprey	Mig (EPBC & BC Act)	23.0 km SSE	Mostly found in coastal areas, where they require extensive areas of water for foraging (DCCEE 2022b).
<i>Pezoporus occidentalis</i> Night Parrot	EN (EPBC Act) CR (BC Act)	*	Rare and cryptic species most frequently associated with <i>Triodia</i> hummock grass and chenopod shrubs (samphire, saltbush etc), particularly where these occur together as a mosaic or along a boundary (ecotone). Suitability of habitat is thought to depend on particular stages of regeneration after fire (DoEE 2020).
<i>Plegadis falcinellus</i> Glossy Ibis	Mig (EPBC & BC Act)	5.4 km E	Predominantly inhabits terrestrial wetlands, foraging in shallow water over soft substrate or on grassy or muddy verges of wetlands providing a variety of water depths. Inland, freshwater wetlands are preferred, especially permanent or ephemeral waterbodies on floodplains and shallow swamps with abundant aquatic flora (Johnstone <i>et al.</i> 2013; Marchant & Higgins 1990)
<i>Polytelis alexandrae</i> Princess Parrot	VU (EPBC Act) P4 (DBCA)	*	Princess Parrots inhabit sandy deserts where they feed on seeds and flowers (Garnett & Crowley 2000a). They also occur in riverine/ littoral areas, or open savannah woodlands and shrublands that usually consist of scattered strands of <i>Eucalyptus</i> , <i>Casuarina</i> or <i>Allocasuarina</i> trees; an understory of shrubs and groundcover dominated by <i>Triodia</i> species (DCCEE 2022b).
<i>Rostratula australis</i> Australian Painted Snipe	EN (EPBC & BC Act)	*	Generally, inhabits shallow terrestrial freshwater (occasionally brackish) wetlands, including temporary and permanent lakes, swamps and claypans. Typical sites include those with rank emergent tussocks of grass, sedges, rushes or reeds, or samphire; often with scattered clumps of <i>Muehlenbeckia</i> (lignum) or canegrass or sometimes <i>Melaleuca</i> (tea-tree) (DoEE 2020).
<i>Tringa glareola</i> Wood Sandpiper	Mig (EPBC & BC Act)	23.5 km S	Prefers the shallows of wooded lakes or swamps with trees. It also inhabits freshwater swamps, lakes, flooded pastures and occasionally, mangroves (Morcombe 2004).
<i>Tringa nebularia</i> Common Greenshank	Mig (EPBC & BC Act)	5.4 km E	Mostly on the coast but sometimes inland; uses permanent and ephemeral terrestrial wetlands, including rivers and creeks (DCCEE 2022b).
<i>Tringa stagnatilis</i> Marsh Sandpiper	Mig (EPBC & BC Act)	30.1 km SE	Inhabits coastal and inland wetlands, estuarine and mangrove mudflats, beaches, swamps, lakes and several other types of wetlands (Morcombe 2004).


Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	Status <sup>1</sup>	Proximity to study area <sup>2</sup>	Habitat
<i>Tyto novaehollandiae kimberli</i> Masked Owl	VU (EPBC Act) P1 (DBCAs)	62.1 km SE	In northern Australia, the Masked Owl has been recorded from riparian forest, rainforest, open forest, <i>Melaleuca</i> swamps and the edges of mangroves, as well as along the margins of sugar cane fields (DoEE 2020).
<b>Mammals</b>			
<i>Dasyurus hallucatus</i> Northern Quoll	EN (EPBC & BC Act)	5.4 km E	Most commonly found in rocky areas, with rugged rocky habitats such as gorges, gullies, escarpments, boulder fields and small caves critical for denning and shelter (DCCEEW 2022b).
<i>Hipposideros stenotis</i> Northern Leaf-nosed Bat	P2 (DBCAs)	31.2 km N	The Northern Leaf-nosed Bat is a secretive bat, with most records of this species being from cracks and caves in the western escarpment of the Arnhem Land plateau, in sandstone cliffs, gorges and water holes bordered by paperbark trees. The species has been recorded in WA, and NW Queensland, as well as a range of offshore islands (Menkhorst & Knight 2011).
<i>Isoodon auratus auratus</i> Golden Bandicoot	VU (EPBC & BC Act)	*	The Golden Bandicoot previously occurred throughout central Australia but is now restricted to the Kimberley (offshore islands and the mainland) and Marchinbar Island (offshore Arnhem Land). During the day it rests in dense vegetation or other shelter (DCCEEW 2022b).
<i>Leggadina lakedownsensis</i> Northern Short-tailed Mouse / Lakeland Downs Mouse	P4 (DBCAs)	25.8 km E	Occurs on sandy soils and cracking clays in WA. Occupies a diverse range of environments from the monsoon tropical coast to semiarid climates, including spinifex and tussock grasslands, samphire and sedgeland, <i>Acacia</i> shrublands, tropical <i>Eucalyptus</i> and <i>Melaleuca</i> woodlands and stony ranges (DEC 2012; Moro & Kutt 2008).
<i>Macrodarma gigas</i> Ghost Bat	VU (EPBC & BC Act)	31.2 km N	Australia's only carnivorous bat (Van Dyck & Strahan 2008), and the largest microbat. Ghost bats are dependent on diurnal roosts in caves, crevices, deep overhangs, and artificial roosts such as underground mines (Hourigan 2011a). They are known to forage in productive plain areas with thin mature woodland over patchy tussock or hummock grass, or waterholes and riparian zones (Bullen 2021).
<i>Macrotis lagotis</i> Bilby	VU (EPBC & BC Act)	65.8 km SE	Bilby prefers hummock grassland in plains and alluvial areas, open tussock grassland on uplands and hills, and mulga woodland/shrubland on ridges and rises (DCCEEW 2022b). Areas where it is now regionally extinct include open and exposed habitat types.
<i>Petrogale lateralis subsp.</i> (West Kimberley) West Kimberley Black-footed Rock-wallaby	VU (EPBC Act) EN (BC Act)	*	The species is patchily distributed across the western half of Australia. Populations are scattered and restricted to sites with suitable rocky habitat with caves and crevices (DCCEEW 2022b).

Species	Status <sup>1</sup>	Proximity to study area <sup>2</sup>	Habitat
<i>Petropseudes dahlii</i> Rock Ringtail Possum	P3 (DBCA)	23.4 km E	Rock Ringtail Possums are found in northern Australia, where they shelter in rocky or sandstone outcrops during the day. At night they forage in trees close to their dens. These rocky areas are usually surrounded by flat, lowland areas (Strahan 1995).
<i>Phascogale tapoatafa kimberleyensis</i> Kimberley Brush-tailed Phascogale	VU (EPBC & BC Act)	*	Inhabits bunch grasslands with a savanna woodland structure, dominated by <i>Eucalyptus</i> and <i>Corymbia</i> species with old growth and dead trees to provide nesting hollows (Threatened Species Scientific Committee 2017).
<i>Rhinoicteris aurantia</i> Orange Leaf-nosed Bat	P4 (DBCA)	31.5 km N	This distinctive, bright orange bat occurs across northern Australia, as well as a distinct form from the Kimberley. Roosting in colonies from 20 to 20,000 this species emerges at dusk to feed on a range of invertebrates, such as termites, flies, wasps, and cockroaches, and fly with a fast zig-zag pattern, usually within 1 m of the ground, foraging over open grasslands and sparse tree and shrub savannah (Hourigan 2011b).
<i>Saccolaimus saccolaimus nudicliviatus</i> Bare-rumped Sheath-tailed Bat	VU (EPBC) P3 (DBCA)	*	The Bare-rumped Sheath-tail Bat occurs mostly in lowland areas, typically in a range of woodland, forest and open environments. In Australia, they roost in tree hollows and forage over habitat edges such as the edges of forest clearings or rainforests (DCCEEW 2022b).
<i>Trichosurus vulpecula arnhemensis</i> Northern Brushtail Possum	VU (EPBC & BC Act)	65.2 km SE	Occurs discontinuously from the Gulf of Carpentaria hinterland, NT (Woinarski J. et al. 2011) to the Kimberley region (Kerle and How 2008, in (Van Dyck & Strahan 2008). Occurs mainly in tall eucalypt open forests, particularly where the understorey includes some shrubs that bear fleshy fruits, also in some mangrove communities and rainforests (Friend & Taylor 1985; Menkhorst & Woinarski 1992; Woinarski J. et al. 2011). Needs hollow bearing trees.
<i>Vespardelus douglasorum</i> Yellow-lipped Cave Bat	P2 (DBCA)	31.3 km N	Occur in regions with high rainfall (over 800 mm), in tropical woodlands along waterways and in adjacent open woodlands. Yellow-lipped Cave Bats roost in sandstone and limestone caves (Australian Museum 2020b).

<sup>1</sup> CR – Critically Endangered; EN – Endangered; VU – Vulnerable; OS – Specially Protected; Mig – Migratory; P1–4 – Priority 1–4.

<sup>2</sup> \* – EPBC Protected Matters Search does not return species record locations and may include instances where suitable habitat may occur, but the species has not necessarily been observed.



Australian Capital Equity Napier Downes Irrigation Project		
Project No	1452	
Date	1/11/2023	
Drawn by	BK	
Map author	BQ	
1:600,000 (at A4)		GDA 1994 MGA Zone 51

Study area  
 Desktop study area

**Figure 5-1**

**Desktop records of significant vertebrate fauna**

**PHOENIX**  
ENVIRONMENTAL SCIENCES

All information within this map is current as of 1/11/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

8125280  
8103540  
8082000  
8060360  
8038720  
8017080  
7995440

### 5.1.2 SRE invertebrate fauna

Four terrestrial invertebrates listed as Threatened under the BC Act and 5 Priority species were identified in the desktop review (Table 5-2). All are molluscs (land snails) in the family Camaenidae and are Confirmed SREs; the records are associated with rocky habitats of the surrounding ranges, mainly Napier Range (Figure 5-2). Two PECs are associated with the SRE invertebrates: Invertebrate community of Napier Range Cave, located 27 km east of the study area, and Monsoon vine thickets and Camaenid land snails of limestone ranges, 20 km northeast (Figure 5-2).

Records of a further 30 confirmed (all land snails) and 13 potential terrestrial SRE species were identified through the WA Museum database searches (Figure 5-2; Figure 5-2). None of these are from within the study area but there is one record of an opilione (Assamiidae sp., potential SRE) 1.7 km west of the study area. No habitat information is provided for this record but the site is within the same broad vegetation association (Shepherd *et al.* 2002) as the study area, adjacent to a drainage line.

The SREs have been collected from a wide range of habitat types, including rocky habitats (rocky outcrops, limestone outcrops, rock piles, boulders, rubble, rock crevices, rock/scree slopes, rocky gullies, bases of escarpments and cliffs), on plains under spinifex, open woodlands, caves/cave entrances, embankments, vine thickets and on roots, creek beds, trunk and/or branches of trees (boabs).


Habitat of the study area is unlikely to support any of the Threatened or Priority camaenids or the invertebrate associated PECs. However, there is potential for other SRE taxa to occur, based on the habitat descriptions above.



**Table 5-2 Terrestrial SRE invertebrates identified in the desktop review**






Higher taxon, Family	Species	Conservation status	SRE status
<b>Arachnida - Mygalomorphae (trapdoor spiders)</b>			
Euagridae	<i>Cethegus</i> `sp. nov.`		Potential
Halonoproctidae	<i>Conothele</i> `MYG542`		Potential
Halonoproctidae	<i>Conothele</i> sp.		Potential
Idiopidae	<i>Idiosoma</i> `occidentalis sp. group`		Potential
<b>Arachnida - Araneomorphae (modern spiders)</b>			
Selenopidae	<i>Karaops jenniferae</i>		Potential
Sparassidae	<i>Heteropoda cavernicola</i>		Potential
<b>Arachnida - Opiliones (harvestmen)</b>			
Assamiidae	<i>Dampetrus</i> sp.		Potential
Assamiidae	Assamiidae sp.		Potential
<b>Arachnida - Pseudoscorpiones (pseudoscorpions)</b>			
Chthoniidae	<i>Austrochthonius</i> `minutissimus`		Potential
<b>Diplopoda (millipedes)</b>			
Paradoxosomatidae	<i>Helicopodosoma</i> `Mt Hart`		Potential
<b>Malacostraca - Isopoda (isopods)</b>			
Armadillidae	<i>Kimberleydillo waldockae</i>		Potential
<b>Gastropoda - Pulmonata (land snails)</b>			
Camaenidae	<i>Amplirhagada carinata</i>		Confirmed









**Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity**

Higher taxon, Family	Species	Conservation status	SRE status
Camaenidae	<i>Amplirhagada napierana</i>		Confirmed
Camaenidae	<i>Amplirhagada (Tenuiragada) percita</i>		Confirmed
Camaenidae	<i>Kendrickia ignivenatus</i>		Confirmed
Camaenidae	<i>Kimboraga mccorryi</i>		Confirmed
Camaenidae	<i>Kimboraga micromphala</i>	P2 (DBCA)	Confirmed
Camaenidae	<i>Kimboraga yammerana</i>	P1 (DBCA)	Confirmed
Camaenidae	<i>Mouldingia occidentalis</i>	CR (BC Act)	Confirmed
Camaenidae	<i>Rhagada basedowana</i>		Confirmed
Camaenidae	<i>Rhagada gatta</i>		Confirmed
Camaenidae	<i>Rhagada construa</i>		Confirmed
Camaenidae	<i>Rhagada gibbensis</i>	P1 (DBCA)	Confirmed
Camaenidae	<i>Rhagada mimika</i>		Confirmed
Camaenidae	<i>Rhagada sutra</i>		Confirmed
Camaenidae	<i>Kimberleytrachia (Torresitrachia) crawfordi</i>		Confirmed
Camaenidae	<i>Westraltrachia alterna</i>	VU (BC Act)	Confirmed
Camaenidae	<i>Westraltrachia (Parrhagada) commoda (P. ferrosa)</i>		Confirmed
Camaenidae	<i>Westraltrachia cunicula</i>		Confirmed
Camaenidae	<i>Westraltrachia derbyi (Trachia orthocheila)</i>		Confirmed
Camaenidae	<i>Westraltrachia (Trachia) froggatti froggatti</i>		Confirmed
Camaenidae	<i>Westraltrachia froggatti complanata</i>		Confirmed
Camaenidae	<i>Westraltrachia inopinata</i>	VU (BC Act)	Confirmed
Camaenidae	<i>Westraltrachia instita</i>		Confirmed
Camaenidae	<i>Westraltrachia lievreana</i>	P2 (DBCA)	Confirmed
Camaenidae	<i>Westraltrachia limbana</i>		Confirmed
Camaenidae	<i>Westraltrachia rotunda</i>		Confirmed
Camaenidae	<i>Westraltrachia sp.1</i>		Potential
Camaenidae	<i>Westraltrachia sp.2</i>		Potential
Camaenidae	<i>Westraltrachia subtila</i>	P2 (DBCA)	Confirmed
Camaenidae	<i>Westraltrachia tropida</i>		Confirmed
Camaenidae	<i>Westraltrachia turbinata</i>	VU (BC Act)	Confirmed
Camaenidae	<i>Westraltrachia woodwardi (Parrhagada detecta)</i>		Confirmed




<b>Australian Capital Equity</b> <b>Napier Downs Irrigation Project</b>	
Project No	1452
Date	1/12/2023
Drawn by	FK
Map author	BQ
	
	
1:1,263,800 (at A4)      GDA 1994 MGA Zone 51	

- Study area**
-  Option 1
  -  Study area
  -  Option 2
  -  Habitat mapping area
  -  100 km buffer

- SRE status**
-  Araneomorph spider, Potential
  -  Harvestman spider, Potential
  -  Land snail, Confirmed
  -  Land snail, Potential
  -  Millipede, Potential
  -  Mygalomorph spider, Potential
  -  Mygalomorph spider, uncertain
  -  Pseudoscorpion, Potential

**Figure 5-2**

**Desktop records of SRE invertebrates**



All information within this map is current as of 1/12/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

## 5.2 FIELD SURVEY

### 5.2.1 Vertebrate fauna

#### 5.2.1.1 Habitats

Three broad fauna habitat types were identified in the habitat mapping area (Table 5-3; Figure 5-3). These habitats comprised of the following:

- shrubland over grassland
- open woodland over open shrubland over grassland
- open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression.




The shrubland over grassland habitat type made up most of the habitat mapping area (1,354.5 ha; 70.8%) and surrounded one small seasonally inundated depression (4.3 ha; 0.2%) located inside the study area. The seasonally inundated depression comprised of open woodland (shrubby regrowth) over mixed herbs and grasses. The shrubland habitat type contained fewer large trees and fallen timber than the open woodland habitat, and typically formed dense mid and lower story vegetation cover. It contained the majority of *Isoodon* (bandicoot) diggings (see section 5.2.1.3 for more information).


The seasonally inundated depression likely provides a valuable source of water during periods where surface water is present. The overall species diversity at the site located in the open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression was the second highest of all 16 fauna sites, largely due to the large number of bird species observed at this site.

The open woodland over open shrubland over grassland covered the remaining 26.4% (487.7 ha) of the habitat mapping area and was found widely fringing drainage lines located outside the perimeter of the habitat mapping area, in areas such as Hawkstone Creek. This habitat type had the highest abundance of fallen logs and tree hollows suitable for fauna refugia.











**Table 5-3 Extent and description of each fauna habitat in the study area**

Habitat type	Site/s	Extent in habitat mapping area	Extent in study area	Representative photograph
<b>Open woodland over open shrubland over grassland</b> Mid <i>Corymbia</i> , <i>Eucalyptus</i> and <i>Acacia</i> open woodland over mixed open shrubland over <i>Sorghum stipoidesum</i> , <i>Chrysopogon</i> and <i>Triodia</i> grassland.	SITE03, SITE04, Opp-Isoodon-01, Opp-Isoodon-03, Opp-Isoodon-09, Opp-Isoodon-10,	487.7 ha 26.4%	75.9 ha 13%	
<b>Shrubland over grassland</b> Sparse <i>Corymbia</i> and <i>Eucalyptus</i> open woodland over mixed open shrubland over <i>Sorghum</i> grassland.	SITE02, SITE05, SITE06, SITE07, SITE08, SITE09, Opp-02, Opp-03, Opp-08, Opp-09, Opp-15, Opp-16, Opp-Isoodon-02, Opp-Isoodon-04, Opp-Isoodon-05, Opp-Isoodon-06, Opp-Isoodon-07, Opp-Isoodon-08, Opp-Isoodon-11, Opp-Isoodon-12, Opp-Isoodon-13,	1,354.5 ha 70.8%	506.2 ha 86.3%	
<b>Open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression</b> Open <i>Melaleuca</i> and <i>Corymbia</i> woodland (primarily shrubby regrowth) over mixed herbs and grasses.	SITE01	4.3 ha 0.2%	4.3 ha 0.7%	





**Fauna habitats**

-  open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression
-  open woodland over open shrubland over grassland
-  shrubland over grassland

-  Study area
-  Habitat mapping area
- Species, status**
-  *Amytomis housei*, P4 (DBC list)
-  *Erythrura gouldiae*, P4 (DBC list)
-  *Isoodon auratus auratus*, VU (EPBC & BC Acts)

**Figure 5-3**  
**Fauna habitats and significant fauna records from the field survey**



Australian Capital Equity Napier Downs Irrigation Project		
Project No	1452	
Date	1/12/2023	
Drawn by	BK	
Map author	GW	
		
1:44,549 (at A4)		GDA 1994 MGA Zone 51

All information within this map is current as of 1/12/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

### 5.2.1.2 Assemblage

A total of 114 terrestrial vertebrate species were recorded in the study area and 1 km buffer during the field survey (Table 5-4). This equated to approximately one third of the 330 species identified in the desktop review, including 5 introduced species (Table 5-4).

**Table 5-4 Number of vertebrate species recorded in survey and desktop results**

Group	No. species identified in desktop review	No. species recorded in survey	% recorded
Amphibians	16	3 (inc. 1 introduced)	18.75
Reptiles	48	6	12.5
Birds	223	89	40
Mammals	43 (inc. 4 introduced)	16 (inc. 3 introduced)	37
<b>Total</b>	<b>330</b>	<b>114</b>	<b>35</b>


Of the 3 species of amphibians recorded during the survey, only the Northern Laughing Tree Frog had been previously identified in the desktop review. The other two included one native species, the Northern Toadlet, and one introduced species, the Cane Toad. It is likely that the absence of these species from the desktop is due to the scarcity of survey work in the area, and the recent movement of Cane Toads, as the population has expanded its distribution with a western direction out of the Northern Territory into WA. None of the amphibians identified in the desktop or field survey are listed as significant species.

Reptiles recorded during the survey included one species of snake, the Olive Python, and 5 species of lizard, all skinks from the family Scincidae. The only reptiles of significance identified in the desktop were the two crocodile species the Australian Freshwater Crocodile and the Salt-water Crocodile (listed as OS under the BC Act), neither of which were detected during the field survey.

The bird assemblage was the most comprehensively catalogued, with 40% of the species identified in the desktop review being recorded during the field survey. Many of the bird species identified in the desktop but not the field survey are unlikely to occur in the study area as they are associated with aquatic habitats, such as marine and riparian environments, that are absent from the study area. This includes various ducks and other waterbirds and 21 species of migratory birds.

Mammals were the second most comprehensively catalogued taxonomic group, with 37% of the species identified in the desktop being recorded during the field survey. The only significant vertebrate identified inside the study area was the Golden Bandicoot (VU under the BC and EPBC Acts). Three of the 4 introduced mammals identified in the desktop were also identified inside the study area, including European Cattle, Dog, and Cat.

A species accumulation curve by trapping night confirms that the survey adequately captured the fauna richness of the study area at the time the survey was conducted (Figure 5-4).





**Figure 5-4 Species accumulation curve for vertebrate fauna**

### 5.2.1.3 Significant vertebrate fauna

One Threatened, one Priority, and one Threatened / Priority species of vertebrate fauna were recorded during the survey, Golden Bandicoot (VU), Black Grasswren (P4) and Gouldian Finch (EN & P4) (Table 5-5; Figure 5-3). Black Grasswren and Gouldian Finch were both recorded outside the study area; however, due to the proximity of records and the presence of suitable habitat, it is possible the species also occur inside the study area. Black Grasswren was not identified in the desktop assessment, however, they were observed opportunistically during the survey along an access track 4.5 km to the south of the study area while the team were on route to Long Pool. Gouldian Finches were observed coming in to drink at a cattle watering station located 2.5 km to the north of the study area.

Golden Bandicoot were detected at one of the trapping sites (Site 08) inside the study area. Additionally, secondary evidence of bandicoot activity (digging) was recorded at an additional 15 sites across the study area and adjacent areas of open woodland and shrubland. As there are two species of bandicoot known to occur inside the study area based on the trapping data (Golden and Northern Brown Bandicoot), it was not possible to determine whether the secondary evidence was the significant Golden Bandicoot, or the Northern Brown Bandicoot which is not listed. It is likely that both species forage inside the study area and the surrounding area.

**Table 5-5 Details of significant vertebrate fauna recorded during the field survey**

Species	Distribution and ecology	Survey records	Photograph
<i>Isoodon auratus auratus</i> Golden Bandicoot (VU under EPBC and BC Act)	Previously occurred throughout central Australia but is now restricted to the Kimberley (offshore islands and the mainland) and Marchinbar Island (offshore Arnhem Land) (DCCEE 2022b). Golden Bandicoot can occupy a wide range of habitats including: Hummock and tussock grasslands on sand-dunes and sand plains in the arid zone, acacia and eucalypt woodlands in tropical semi-arid the zone, vine thickets, heath and woodlands in rugged sandstone, and volcanic country in the subhumid tropics (Palmer <i>et al.</i> 2003).	One male in breeding condition, and one female with 2 juvenile offspring were recorded on separate days at Site 08. Additionally, secondary evidence of <i>Isoodon sp.</i> diggings were recorded across the study area in additional 15 locations. These diggings were likely made by both Golden Bandicoot and the similar Northern Brown Bandicoot which was also captured at Site 08.	
<i>Amytornis housei</i> ; Black Grasswren (P4, DBCA list)	Known to occur to the north of the study area at Prince Regent National Park and the Mitchell Plateau, in habitats containing large rocky sandstone boulders and dense hummock and spinifex grasslands.	One bird of unknown gender was observed opportunistically while travelling through the hummock grassland to the south of the study area.	No photo taken
<i>Erythrura gouldiae</i> Gouldian Finch EN (EN under EPBC Act, P4, DBCA list)	In WA, the species only occurs in the Kimberley. It inhabits open woodlands with favoured annual and perennial grasses (especially Sorghum), a nearby source of surface water and, in the breeding season, unburnt hollow-bearing <i>Eucalyptus</i> trees (DoEE 2020).	Three Gouldian Finches were observed drinking in a mixed flock at a cattle watering station 2.5 km north of the study area.	

The likelihood of occurrence assessment (section 4.2.2.14) for the remaining significant species identified in the desktop review (section 5.1.1) determined two were likely to occur in the study area, 15 may possibly occur and the remaining 24 are unlikely to occur (Table 5-6).

Table 5-6 Likelihood of occurrence for significant vertebrates identified in the desktop

Species	Status <sup>1</sup>	Likelihood of occurrence	Habitat in the study area
<b>Reptiles</b>			
<i>Crocodylus johnstoni</i> Freshwater Crocodile	OS (BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Crocodylus porosus</i> Salt-water Crocodile	OS (BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<b>Birds</b>			
<i>Actitis hypoleucos</i> Common Sandpiper	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Apus pacificus</i> Fork-tailed Swift	Mig (EPBC & BC Act)	Possible	Not limited by habitat type and therefore a possible visitor to the study area.
<i>Botaurus poiciloptilus</i> Australasian Bittern	EN (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Calidris acuminata</i> Sharp-tailed Sandpiper	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Calidris ferruginea</i> Curlew Sandpiper	CR/Mig (EPBC Act) VU/Mig (BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Calidris melanotos</i> Pectoral Sandpiper	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Cecropis daurica</i> Red-rumped Swallow	Mig (EPBC & BC Act)	Unlikely	Rare Australian vagrant occasionally recorded in Northern Australia.
<i>Charadrius veredus</i> Oriental Plover	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Erythrotriorchis radiatus</i> Red Goshawk	VU (EPBC & BC Act)	Possible	Due to large foraging ranges, Red Goshawk may occur within the study area to forage. Typically associated with river systems in Northern Australia. The red goshawk rarely breeds in areas with fragmented native vegetation (Aumann & Baker-Gabb 1991) The stick nests, in which 1–2 eggs are laid, are restricted to trees that are taller than 20 m and within 1 km of a watercourse or wetland (Aumann & Baker-Gabb 1991).

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	Status <sup>1</sup>	Likelihood of occurrence	Habitat in the study area
<i>Erythrura gouldiae</i> Gouldian Finch	P4 (BC Act)	Likely	Suitable breeding habitat in the Eucalypt woodland stands, and access to surface water in the seasonally inundated depression. Gouldian Finch were also recorded approximately 3 km to the north of the study area during the survey.
	EN (EPBC Act)		
<i>Falco hypoleucos</i> Grey Falcon	VU (EPBC & BC Act)	Possible	Due to large foraging ranges, Grey Falcon may occur within the study area to forage and nest. Typically forages in open habitats, and nests in tall trees and infrastructure towers.
<i>Falco peregrinus</i> Peregrine Falcon	OS (BC Act)	Possible	Due to large foraging ranges, Peregrine Falcon may occur within the study area to forage. Typically nests on cliffs and in trees.
<i>Gelochelidon nilotica</i> Gull-billed Tern	Mig (EPBC & BC Act)	Unlikely	Absence of suitable coastal/marine habitats. Rarely observed in inland environments.
<i>Glaucala maldivarum</i> Oriental Pratincole	Mig (EPBC & BC Act)	Possible	Not limited by habitat type and therefore a possible visitor to the study area. Typically associated with open habitats.
<i>Hirundo rustica</i> Barn Swallow	Mig (EPBC & BC Act)	Possible	Not limited by habitat type and therefore a possible visitor to the study area.
<i>Motacilla cinerea</i> Grey Wagtail	Mig (EPBC & BC Act)	Unlikely	Absence of suitable open grassland or riparian habitat within the study area.
<i>Motacilla flava</i> Yellow Wagtail	Mig (EPBC & BC Act)	Unlikely	Absence of suitable open grassland or riparian habitat within the study area.
<i>Numenius madagascariensis</i> Eastern Curlew	CR/Mig (EPC Act)	Unlikely	Absence of suitable coastal/marine habitats. Rarely observed in inland environments.
<i>Pandion cristatus</i> Osprey	VU/Mig (BC Act)	Unlikely	Absence of suitable coastal/marine habitats. Typically associated with coastal areas.
	Mig (EPBC & BC Act)		
<i>Pezoporus occidentalis</i> Night Parrot	EN (EPBC Act)	Unlikely	Absence of old growth spinifex within the study area.
	CR (BC Act)		
<i>Plegadis falcinellus</i> Glossy Ibis	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
	VU (EPBC Act)		
<i>Polytelis alexandrae</i> Princess Parrot	P4 (DBCA)	Unlikely	Typically occurs in arid environments in central Australia.
<i>Rostratula australis</i> Australian Painted Snipe	EN (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	Status <sup>1</sup>	Likelihood of occurrence	Habitat in the study area
<i>Tringa glareola</i> Wood Sandpiper	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Tringa nebularia</i> Common Greenshank	Mig (EPBC & BC Act)	Unlikely	No suitable coastal, drainage, wetland or salt lake habitat in the study area.
<i>Tringa stagnatilis</i> Marsh Sandpiper	Mig (EPBC & BC Act)	Unlikely	Absence of suitable riparian/river habitat within the study area.
<i>Tyto novaehollandiae</i> Kimberli Masked Owl	VU (EPBC Act)	Possible	Nests in a variety of habitats, provided there are large hollows present. Open woodland habitats inside the study area may contain suitable breeding habitat.
<b>Mammals</b>			
<i>Dasyurus hallucatus</i> Northern Quoll	EN (EPBC & BC Act)	Possible	Previously recorded 5.4 km east of the study area in riparian habitat. No suitable rocky denning habitat within study area. Study area may provide foraging habitat.
<i>Hipposideros aaurantia</i> Northern Leaf-nosed Bat	P2 (DBCA)	Possible	No suitable breeding or roosting habitat, but whole of study area contains potential foraging habitat.
<i>Leggadina lakedownsensis</i> Northern Short-tailed Mouse	P4 (DBCA)	Likely	Occurs across a diverse range of environments from the monsoon tropical coast to semi-arid climates including grasslands, shrublands, and woodlands.
<i>Macrodarma gigas</i> Ghost Bat	VU (EPBC & BC Act)	Possible	No suitable roost habitat, but whole of study area contains potential foraging habitat
<i>Macrotis lagotis</i> Bilby	VU (EPBC & BC Act)	Possible	Occurs across a range of vegetation types including grassland, shrubland and woodland habitats like those found inside the study area. However, the study area is located beyond the northern extent of their known distribution.
<i>Petrogale lateralis</i> subsp. (West Kimberley) West Kimberley Black-footed Rock-wallaby	VU (EPBC Act)	Unlikely	Absence of suitable rocky habitats with complex caves or crevices, and permanent water.
<i>Petropseudes dahli</i> Rock Ringtail Possum	EN (BC Act)		
	P3 (DBCA)	Unlikely	Absence of suitable rocky habitats with large boulders and deeply fissured rocks.



Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	Status <sup>1</sup>	Likelihood of occurrence	Habitat in the study area
<i>Phascogale tapoatafa</i> kimberleyensis Kimberley Brush-tailed Phascogale	VU (EPBC & BC Act)	Possible	Restricted to eucalypt forests where it is highly arboreal, although has been recorded foraging on the ground as well as in trees. Requires tree hollows for nesting and refuge during the day. May occur in the open woodland habitats.
<i>Rhinoicteris aurantia</i> Orange Leaf-nosed Bat	P4 (DBCA)	Possible	No suitable breeding or roosting habitat, but whole of study area contains potential foraging habitat.
<i>Saccolaimus saccolaimus nudicinctatus</i> Bare-rumped Sheath-tailed Bat	P3 (DBCA)	Possible	Forages over the tops of trees of a variety of forest types like those present in the open woodland habitats. Typically roosts in caves which are absent in the study area but can also utilise tree hollows for roosting habitat which is present in the woodland areas.
	VU (BC Act)		
<i>Trichosurus vulpecula arnhemensis</i> Northern Brushtail Possum	VU (EPBC & BC Act)	Possible	Typically occur in wooded areas in the tropical regions of Northern Australia. Displays a low specificity for habitat selection occurring in both rural and urban areas.
<i>Vespadelus douglasorum</i> Yellow-lipped Cave Bat	P2(DBCA)	Unlikely	Roosts in caves and forages over streams, neither of which are present inside the study area.

<sup>1</sup> CR – Critically Endangered; EN – Endangered; VU – Vulnerable; OS – Specially Protected; Mig – Migratory; P1-4 – Priority 1-4.

## 5.2.2 SRE invertebrate fauna

### 5.2.2.1 Habitats

The 3 fauna habitats mapped in the study area (section 5.2.1.1) were all classified as Low potential habitat for SRE invertebrates (Table 5-7; Figure 5-5). None of these represent High potential SRE habitats due to the lack of isolating habitat features such as rocky outbreaks or drainage systems. With the exception of the seasonally inundated depression which comprised a unique flora assemblage not seen elsewhere in the study area, the native flora species recorded extend throughout the study area and much of the surrounding landscape, and do not impose barriers for fauna dispersal.

**Table 5-7 Extent and description of each SRE habitat in the study area**

Habitat code	Habitat type	Sites	SRE habitat rating	Extent in study area and % of study area
1	Open woodland over open shrubland over grassland	SITE04	Low	553.2 ha 28.9%
2	Shrubland over grassland	SITE02, SITE08	Low	1,354.5 ha 70.8%
3	Open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression	SITE01	Low	4.3 ha 0.3%

### 5.2.2.2 SRE records

A total of 45 specimens from four SRE groups were collected within the study area (Figure 5-5; Table 5-8). These comprise of 3 species of pseudoscorpions, 2 species of mygalomorph spider, and one species of scorpion and isopod.

Of these, 3 are Potential SREs, 2 are widespread species and 2 are of Uncertain SRE status due to poor taxonomic resolution. The 3 Potential SREs are:

- *Aname* 'MYG771' – a mygalomorph spider collected from all 4 survey sites (in and out of the study area) and in all 3 habitat types. This species has not been recorded previously.
- *Cubaris* sp. indet. – an isopod (slater) collected from 3 sites (in and out of the study area) within 2 habitats (Shrubland over grassland and Open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression).
- *Lychas* 'annulatus group' – a scorpion which belongs to a species complex has a distribution throughout Australia. The specimens collected from the survey are similar in appearance to the widespread 'central deserts form' (pers. comm. Dr. Erich Volschenk) and were collected from one site in one habitat (Open woodland over open shrubland over grassland).

**Table 5-8 Specimens from SRE groups recorded in the field survey**

Group	Taxa	Sites	Habitats	No. specimens	SRE status
Mygalomorph spider	<i>Aname</i> 'MYG771'	SITE01, SITE02, SITE04, SITE08	1, 2, 3	10	Potential
	<i>Selenotholus</i> 'MYG381'	SITE08	2	1	Widespread
Scorpion	<i>Lychas</i> 'annulatus grp'	SITE04	1	2	Potential
Pseudoscorpion	<i>Austrohorus</i> sp. indet.	SITE02, SITE04	1, 2	2	Uncertain
	<i>Beierolpium</i> 'sp. 8/4'	SITE04	1	1	Widespread
	<i>Indolpium</i> sp. indet.	SITE01, SITE02, SITE04, SITE08	1, 2, 3	20	Uncertain
Isopod (slater)	<i>Cubaris</i> sp. indet. 'Napier'	SITE01, SITE02, SITE08	2, 3	9	Potential



**Fauna habitats**

- open woodland (shrubby regrowth) over mixed herbs and grasses surrounding a seasonally inundated depression
- open woodland over open shrubland over grassland
- shrubland over grassland



**Australian Capital Equity  
Napier Downes Irrigation Project**

Project No 1452  
 Date 1/12/2023  
 Drawn by BK  
 Map author AJ



0 0.5 1  
 Kilometers  
 1:30,723 (at A4) GDA 1994 MGA Zone 51

- Study area
- Habitat mapping area
- Species, SRE status**
- Indolpium* sp. indet, Uncertain
- Cubaris* sp. indet, Potential SRE
- Aname* 'MYG771', Potential SRE
- Selenotholus* 'MYG381', Widespread
- Lychas* 'annulatus grp', Potential SRE
- Beierolpium* 'sp. 8/4', Potential SRE
- Austrohorus* sp. indet, Uncertain

- Isopod**
- Indolpium* sp. indet, Uncertain
- Cubaris* sp. indet, Potential SRE
- Aname* 'MYG771', Potential SRE
- Selenotholus* 'MYG381', Widespread
- Mygalomorph spider**
- Aname* 'MYG771', Potential SRE
- Selenotholus* 'MYG381', Widespread
- Pseudoscorpion**
- Austrohorus* sp. indet, Uncertain
- Beierolpium* 'sp. 8/4', Potential SRE

**Figure 5-5**  
**SRE habitats and recorded SRE taxa**



All information within this map is current as of 1/12/2023. This product is subject to COPYRIGHT and is property of Phoenix Environmental Sciences (Phoenix). While Phoenix has taken care to ensure the accuracy of this product, Phoenix make no representations or warranties about its accuracy, completeness or suitability for any particular purpose.

### 5.3 SURVEY LIMITATIONS

The limitations of the terrestrial fauna survey have been considered in accordance with EPA (2016b, d) (Table 5-9).

**Table 5-9 Consideration of potential survey limitations**

Limitations	Comments
Availability of contextual information at a regional and local scale	Limited contextual information was available at both the regional and local scale.
Competency/experience of the team carrying out the survey	Field surveys were led by a zoologist with over 10 years' experience conducting surveys in WA who has conducted numerous surveys in the Kimberley bioregion.
Scope and completeness	The scope was completed.
Proportion of flora and fauna recorded and/or collected, any identification issues	All recorded fauna identified, except some invertebrates with indeterminate morphology. Approximately one third of the vertebrate species identified in the desktop review were detected. Assemblage results were unsurprising, and not considered a limitation, due to the limited habitat diversity in the study area compared with the wider desktop study area. The SRE assemblage recorded during the survey was expected both in terms of taxa and abundance, including both known and previously unknown taxa.
Access within the study area	The study area was fully accessible at the time of the survey.
Timing, rainfall, season	Timing was not optimal as it was outside of the wet season. This was due to restricted access into the study area caused by late wet season rainfall, delaying the start of the survey by several months.
Disturbance that may have affected the results of the survey	There was no disturbance that would have affected the results of the survey.

## 6. DISCUSSION

The surveys have documented a diverse assemblage of vertebrate fauna species across the study area, including several that were not on any of the database searches conducted for the desktop assessment. This is partially due to the limited availability of survey data, and highly diverse assemblage of fauna for this area, including seasonal migrants and occasional vagrant species as well as residents.

The fauna assemblage in the Kimberley has been surveyed far less thoroughly than areas such as the Pilbara and Goldfields of WA in recent years. As such, it is unsurprising that several of the species detected during this survey did not come up in the desktop review. Additionally, as the desktop review included numerous habitat types that were absent from the study area, many of the species identified in the desktop were absent from the field survey.

Although habitat quality has been impacted through human activities involving agriculture and groundwater modification, as well as predation and poisoning by introduced species such as cats and cane toads, the regions vertebrate communities remain relatively intact and do not appear to be undergoing a high extinction rate.

The most notable species detected during the field survey that were not identified in the desktop assessment were the invasive Cane Toad, which is a recent addition to the assemblage as the population has dispersed out of the NT and into WA, and the significant Black Grasswren (P4), which is known to occur in isolated locations to the north of the study area. Far less is known about the invertebrate fauna of the region, particularly in those groups with limited dispersal abilities resulting in the precautionary SRE status of many taxa.

### 6.1 VERTEBRATE FAUNA

The habitat types identified and mapped in the study area match those in the surrounding vicinity (Table 5-3, Figure 5-3) and are widespread in the Kimberley region. Restricted habitats in the region that would provide core habitat for significant vertebrates include major drainages, rocky cliffs and gorges, and sand dunes, none of which occur within the study area.

The areas containing open woodland over mixed herbs and grasses surrounding a seasonally inundated depression have a relatively restricted distribution within the landscape, and represent habitat for significant vertebrates such as the Gouldian Finch, as well as invertebrates (see next section).

The desktop review identified 333 vertebrate taxa potentially occurring in the vicinity of the study area, of which 42 are listed in significant categories (section 5.1.1, Table 5-1). The surveys recorded 108 terrestrial vertebrate species, 8 of which were not identified in the desktop review. Three significant vertebrate taxa were recorded during the surveys; Golden Bandicoot (VU), Black Grasswren (P4) and Gouldian Finch (P4/EN), however only the Golden Bandicoot was recorded inside the study area.

Additionally, 5 significant species were recorded in 2013 from a flood plain site on the eastern side of Hawkstone Creek, 5.4 km east of the study area; Northern Quoll (EN), Gull-billed Tern (Mig.), Glossy Ibis (Mig.), Common Greenshank (Mig.) and Freshwater Crocodile (OS). With the exception of the Northern Quoll, all of the species from Hawkstone Creek are unlikely to occur in the study area as they are all strongly associated with aquatic habitats.

No vertebrate species are restricted (endemic) to the study area or the Fitzroy Trough subregion, and each of the significant species that may occur has its main area of distribution outside the subregion. For example, Gouldian Finch require eucalypt tree hollows for breeding, *Sorghum spp.* and similar seeds for feeding, and water within about 4 km of the nest site (Tidemann 1996) which is present only in the woodland areas of the study area, and is present across much of the Kimberley region.

Similarly, Golden Bandicoot historically occupied much of northern Australia from central WA to western QLD and a number of islands along the Kimberley and Pilbara coast, spanning a diverse range of habitat types. The species has severely declined since European settlement, which has been attributed to the introduction of predators such as feral cats, and changes to fire regimes causing habitat loss and degradation (Ringma *et al.* 2018). The study area is located towards the south-western extent of the Golden Bandicoots currently known distribution. As such, this population is considered important based on its location relative to the currently known distribution of Golden Bandicoot.

Two Golden Bandicoot were captured during the survey, one male and one female. Both were in breeding condition, and the female had two juveniles still attached. Additionally, the extensive bandicoot digging observed across the study area indicates that Golden Bandicoot are likely resident in the area and use the available habitats for foraging and are reproducing in and around the study area. The shrubland habitat contained the highest number of bandicoot diggings. The shrubland habitats may provide the best protection from predators, as they typically had the densest understory of the habitat types present inside the study area, and likely contained an abundance of arthropods, small vertebrates, and plant materials for the bandicoots to eat. This habitat type is relatively common in the region, making up approximately 86% of the study area, and 70% of the mapped area.

Northern Quoll have been recorded approximately 5.4 km to the east of the study area at a nearby riparian habitat. It is possible they may also move through the study area as it would likely provide suitable foraging habitat, however it is unlikely the Northern Quoll would breed inside the study area as there is no rocky habitats containing caves suitable for denning habitat. None of the habitat inside the study area would be considered critical habitat for Northern Quoll, but all habitats present may be used for dispersal and foraging.

## 6.2 SRE INVERTEBRATE FAUNA

The habitats within the study are considered to be relatively continuous and contain overlapping vegetation structure, and landform attributes. Aside from one habitat displaying a higher incidence of moisture/inundation (at site O1), this was not considered enough to create a habitat isolate for SREs. The Yeeda land system, which occupies the majority of the study area and includes all four survey sites, extends uninterrupted 66 km to the west, and is also well represented immediately east of the Hawkstone River and south of the Lennard River. The larger rivers, and the Napier Ranges to the north, may be potential barriers to dispersal, along with any major changes to vegetation and soils.

Three potential SREs were recorded from the study area, of which two are new to science (*Aname* 'MYG771' and *Cubaris* sp. indet. 'Napier') and one is from a species complex which is thought to contain multiple species:

- *Aname* 'MYG771' – Several other undescribed species of *Aname* are known from the Kimberley region (Pers. comm Dr. Mark Harvey, WA Museum); however, this is the first record of *Aname* 'MYG771' and is therefore currently only known from the study area. Considering this mygalomorph species was collected from widespread and unrestrictive habitat types, it is likely to occur more broadly in the general area within similar habitats
- *Cubaris* sp. indet. 'Napier' – *Cubaris* isopods are from the Armadillidae family of isopods which are uncommon in the Kimberly region. Several highly endemic taxa have been recorded from the region e.g. *Filippinodillo kimberleyensis*, *Acanthodillo agasketos* (both from one location in wet pitfall traps) (Lewis 1998), and *Kimberleydillo waldockensis* (one location – cave entrance) (Dalens 1993), indicating this species is likely to also be endemic. Given this species was recorded from 3 locations within 2 habitat types, including a widespread habitat, it is likely to occur more broadly in the general area within similar habitats.
- *Lychas* 'annulatus group' – Many morphologically distinct species are known from this scorpion group, but some display high levels of cryptic speciation and thus taxonomy of this

group is not well resolved. Given this taxon was collected from a widespread habitat type, it is likely to occur more broadly in the general area within similar habitat.

The 2 uncertain SREs are both pseudoscorpions from genera which have generally poor taxonomic resolution. *Indolpium* and *Austrohorus*, both from the Olpiidae family of pseudoscorpions occur widely throughout WA; however, few species have been described. The two widespread species, *Beierolpium* 'sp. 8/4' and *Selenothelus* 'MYG381' are known to occur in arid and northern Australia.

The habitats within the study area are not considered suitable for the confirmed SREs returned in the desktop review due to absence of typical isolated habitat types these are known from (e.g. rocky outcrops, limestone outcrops, rock piles, boulders, rubble, rock crevices, rock/scree slopes, rocky gullies, bases of escarpments and cliffs), cave entrances, embarkments, vine thickets and on roots, creek beds, trunk and/or branches of trees (boabs).

### 6.3 CONCLUSION

The faunal assemblage of the study area is broadly typical of the woodland and shrubland habitats in the region and consists mostly of species with large distributions across northern WA and the NT. All the fauna habitats present inside the study area are relatively common in the surrounding region.

The Golden Bandicoot was the only significant vertebrate species recorded inside the study area and was also recorded as several sites to the south of the study area, which was unsurprising given the small size of the study area, and the habitats inside the study area being relatively common in the surrounding area.

The SRE invertebrate records from the survey comprise of a mixture of new, potentially range restricted SRE invertebrates, and known widespread species. The habitats in the study area are of only low habitat potential for SREs and do not contain any distinct isolating features that may create physical barriers to invertebrate dispersal. It is highly unlikely that any SRE taxa would be restricted to the study area.



## REFERENCES

- Aumann, T. & Baker-Gabb, D. J. 1991. *The ecology and status of the Red Goshawk in northern Australia*. Royal Australasian Ornithologists Union.
- Australian Museum. 2020a. *Freshwater Crocodile*. Australian Museum. Available at: <https://australian.museum/learn/animals/reptiles/freshwater-crocodile/>
- Australian Museum. 2020b. *Yellow-lipped Bat*. Australian Museum. Available at: <https://australian.museum/learn/animals/bats/yellow-lipped-bat/> (accessed 8/9/22).
- Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D., Ostell, J. & Sayers, E. W. 2012. GenBank. *Nucleic Acids Research* **41**: D36–D42.
- Birdlife Australia. 2019. *Birddata*. Birdlife Australia, Calton, VIC. Available at: <https://birddata.birdlife.org.au/>
- BoM. 2022. *Climate statistics for Australian locations*. Commonwealth of Australia, Bureau of Meteorology. Available at: <http://www.bom.gov.au/climate/data/>
- Bullen, R. D. 2021. *A review of ghost bat ecology, threats and survey requirements*. Prepared for the Department of Agriculture, Water and Environment.
- Car, C. A. & Harvey, M. S. 2014. The millipede genus *Antichiropus* (Diplopoda: Polydesmida: Paradoxosomatidae), part 2: species of the Great Western Woodlands region of Western Australia. *Records of the Western Australian Museum* **29**: 20–77.
- Dalens, H. 1993. Two new genera of terrestrial isopods (Crustacea: Isopoda: Oniscidea) from north-western Western Australia. *Records of the Western Australian Museum* **16**: 257–267.
- DBCA. 2018. *Guideline for the survey and relocation of bilby in Western Australia (draft)*. Department of Biodiversity, Conservation and Attractions, Perth, WA.
- DBCA. 2019a. *NatureMap*. Department of Biodiversity, Conservation and Attractions. Available at: <https://naturemap.dpaw.wa.gov.au/default.aspx>
- DBCA. 2019b. *Threatened and Priority Fauna database search*. Department of Biodiversity, Conservation and Attractions, Kensington, WA.
- DBCA. 2019c. *Threatened Flora, Fauna and Ecological Communities database searches*. Department of Biodiversity, Conservation and Attractions, Kensington, WA.
- DCCEEW. 2022a. *Protected Matters Search Tool*. Available at: [pmst.awe.gov.au](http://pmst.awe.gov.au)
- DCCEEW. 2022b. *Species Profile and Threats Database*. Department of Agriculture, Water and Environment, Canberra, ACT. Available at: <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>
- DEC. 2012. *Lakeland Downs Short-tailed Mouse *Leggadina lakedownensis* (Watts, 1976)*. Department of Environment and Conservation, Kensington, WA.
- Department of Mines and Petroleum. 2016. *Guidelines for Mining Proposals in Western Australia*. Department of Mines and Petroleum, Government of Western Australia, Perth, WA. Available at: <http://www.dmp.wa.gov.au/Documents/Environment/ENV-MEB-213.pdf>
- DoE. 2016. *EPBC Act referral guideline for the endangered Northern Quoll *Dasyurus hallucatus**. Department of the Environment, Canberra, ACT. Available at: <http://www.environment.gov.au/biodiversity/threatened/publications/referral-guideline-northern-quoll>
- DoEE. 2016. *Maps: Australia's bioregions (IBRA)*. Department of the Environment and Energy, Canberra, ACT. Available at: <http://www.environment.gov.au/topics/land/national-reserve-system/science-maps-and-data/australias-bioregions-ibra>
- DoEE. 2019. *Australia's National Heritage List*. Department of the Environment and Energy, Canberra, ACT. Available at: <https://www.environment.gov.au/heritage/places/national-heritage-list>
- DoEE. 2020. *Species Profile and Threats Database*. Department of the Environment and Energy, Australian Government, Canberra, ACT. Available at: <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>

- DPaW. 2017. *Interim guideline for preliminary surveys of Night Parrot (Pezoporus occidentalis) in Western Australia*. Department of Parks and Wildlife, Kensington, WA. Available at: [https://www.dpaw.wa.gov.au/images/documents/plants-animals/animals/interim\\_guideline\\_for\\_night\\_parrot\\_survey.pdf](https://www.dpaw.wa.gov.au/images/documents/plants-animals/animals/interim_guideline_for_night_parrot_survey.pdf)
- DSEWPaC. 2008. *Approved conservation advice for Pezoporus occidentalis (Night Parrot)*. Department of Sustainability, Environment, Water, Population and Communities, Parkes, ACT. A statement for the purposes of approved conservation advice (s266B of the Environment Protection and Biodiversity Conservation Act 1999).
- DSEWPaC. 2010. *Survey guidelines for Australia's threatened birds. Guidelines for detecting birds listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999*. Department of Sustainability, Environment, Water, Population and Communities, Parkes, ACT.
- DSEWPaC. 2011a. *Survey guidelines for Australia's threatened mammals. Guidelines for detecting mammals listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999*. Department of Sustainability, Environment, Water, Population and Communities, Parkes, ACT.
- DSEWPaC. 2011b. *Survey guidelines for Australia's threatened reptiles. Guidelines for detecting reptiles listed as threatened under the Environmental Protection and Biodiversity Conservation Act 1999*. Department of Sustainability, Environment, Water, Population and Communities, Parkes, ACT.
- EPA. 2016a. *Environmental Factor Guideline: Terrestrial fauna*. Environmental Protection Authority, Perth, WA. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/Guideline-Terrestrial-Fauna-131216\\_3.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Guideline-Terrestrial-Fauna-131216_3.pdf) (accessed 20 December 2016).
- EPA. 2016b. *Technical Guidance: Flora and vegetation surveys for Environmental Impact Assessment*. Environmental Protection Authority, Perth, WA. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/EPA%20Technical%20Guidance%20-%20Flora%20and%20Vegetation%20survey\\_Dec13.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA%20Technical%20Guidance%20-%20Flora%20and%20Vegetation%20survey_Dec13.pdf) (accessed 20 December 2016).
- EPA. 2016c. *Technical Guidance: Sampling of short range endemic invertebrate fauna*. Environmental Protection Authority, Perth, WA. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/Tech%20guidance-%20Sampling-SREs-Dec-2016.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Tech%20guidance-%20Sampling-SREs-Dec-2016.pdf) (accessed 20 December 2016).
- EPA. 2016d. *Technical Guidance: Terrestrial fauna surveys*. Environmental Protection Authority, Perth, WA. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/Tech%20guidance-%20Terrestrial%20Fauna%20Surveys-Dec-2016.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Tech%20guidance-%20Terrestrial%20Fauna%20Surveys-Dec-2016.pdf) (accessed 10 December 2016).
- EPA. 2020. *Technical Guidance: Terrestrial vertebrate fauna surveys for environmental impact assessment*. Environmental Protection Authority, Perth, WA. Available at: [https://epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/EPA-Technical-Guidance-Vertebrate-Fauna-Surveys.pdf](https://epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA-Technical-Guidance-Vertebrate-Fauna-Surveys.pdf)
- Friend, G. & Taylor, J. 1985. Habitat preferences of small mammals in tropical open-forest of the Northern Territory. *Australian Journal of Ecology* **10**: 173-185.
- Garnett, S. T. & Crowley, G. M. 2000a. *The action plan for Australian birds 2000*. Birds Australia, Environment Australia, Canberra, ACT. Available at: [www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/](http://www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/)
- Garnett, S. T. & Crowley, G. M. 2000b. *The action plan for Australian birds 2000. List of extinct, threatened and near threatened Australian birds*. Birds Australia, Environment Australia. Available at: [www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/ts-list.html](http://www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/ts-list.html) (accessed 26 Jul 2010).
- Geering, A., Agnew, L. & Harding, S. 2007. *Shorebirds of Australia*. CSIRO Publishing, Collingwood, Vic.

- Government of Western Australia. 2005. *Environmental Protection (Environmentally Sensitive Areas) Notice 2005*. Government of Western Australia, Perth, Western Australia.
- Government of Western Australia. 2018a. *Wildlife Conservation Act 1950 Wildlife Conservation (Rare Flora) Notice 2018*. Government Gazette, WA. Government of Western Australia, Perth, WA.
- Government of Western Australia. 2018b. *Wildlife Conservation Act 1950, Wildlife Conservation (Specially Protected Fauna) Notice 2018*. Government Gazette, WA, Perth, WA.
- Graham, G. 2001a. Dampierland 1 (DL1—Fitzroy Trough subregion). In: May, J. E. & McKenzie, N. L. (eds) *A biodiversity audit of Western Australia's 53 biogeographical subregions in 2002*. Department of Environment and Conservation, Perth, WA, pp. 170–178.
- Graham, G. 2001b. Great Sandy Desert 1 (GSD1—McLarty subregion). In: May, J. E. & McKenzie, N. L. (eds) *A biodiversity audit of Western Australia's 53 biogeographical subregions in 2002*. Department of Environment and Conservation, Perth, WA, pp. 326–331.
- Harvey, M. S. 2002. Short-range endemism among the Australian fauna: some examples from non-marine environments. *Invertebrate Systematics* **16**: 555–570.
- Hebert, P. D. N., A., C., Ball, S. L. & de Waard, J. R. 2003a. Biological identifications through DNA barcodes. *Proceedings of the Royal Society London B* **270**: 313–321.
- Hebert, P. D. N., Ratnasingham, S. & de Waard, J. R. 2003b. Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society London B, Supplement* **270**: 96–99.
- Higgins, P. J. (ed.) 1999. *Handbook of Australian, New Zealand and Antarctic birds. Volume 4: Parrots to Dollarbird*. Oxford University Press Melbourne, Vic.
- Hourigan, C. 2011a. *Ghost bat, Macroderma gigas. Targeted species survey guidelines*. Queensland Herbarium, Department of Environment and Science, Brisbane, Queensland. Available at: [https://www.qld.gov.au/data/assets/pdf\\_file/0018/67140/ghost-bat.pdf](https://www.qld.gov.au/data/assets/pdf_file/0018/67140/ghost-bat.pdf)
- Hourigan, C. 2011b. *Orange leaf-nosed bat, Rhinonicteris aurantia. Targeted species survey guidelines*. Queensland Herbarium, Department of Environment and Science, Brisbane, Queensland. Available at: [https://www.qld.gov.au/data/assets/pdf\\_file/0021/68007/orange-leafnosed-bat.pdf](https://www.qld.gov.au/data/assets/pdf_file/0021/68007/orange-leafnosed-bat.pdf)
- IUCN. 2019. *The IUCN Red List of Threatened Species*.
- Johnstone, R. E., Burbidge, A. H. & Darnell, J. C. 2013. Birds of the Pilbara region, including seas and offshore islands, Western Australia: distribution, status and historical changes. *Records of the Western Australian Museum, Supplement* **78**: 343–441.
- Johnstone, R. E. & Storr, G. M. 1998. *Handbook of Western Australian birds. Volume 1: Non-passerines (Emu to Dollarbird)*. Western Australian Museum, Perth, WA.
- Lewis, F. 1998. New genera and species of terrestrial isopods (Crustacea: Oniscidea) from Australia. *Journal of Natural History* **32**: 701–732.
- Marchant, S. & Higgins, P. J. (eds). 1990. *Handbook of Australian, New Zealand and Antarctic birds. Volume 1: Ratites to ducks*. Oxford University Press, Melbourne, Vic.
- McKilligan, N. (ed.) 2005. *Hérons, egrets and bitterns*. CSIRO Publishing, Collingwood, Vic.
- Menkhorst, K. & Woinarski, J. 1992. Distribution of mammals in monsoon rainforests of the Northern Territory. *Wildlife Research* **19**: 295–316.
- Menkhorst, P. W. & Knight, F. 2011. *A field guide to the mammals of Australia. 3rd edition*. Oxford University Press, Oxford (UK).
- Morcombe, M. 2004. *Field guide to Australian birds. Complete compact edition*. Steve Parish Publishing, Archerfield, QLD.
- Moro, D. & Kutt, A. S. 2008. Northern Short-tailed Mouse, *Leggadina lakedonensis*. In: Van Dyck, S. & Strahan, R. (eds) *The mammals of Australia*. Reed New Holland, Sydney, NSW, pp. 583–584.
- Palmer, C., Taylor, R. & Burbidge, A. 2003. *Recovery plan for the Golden Bandicoot Isoodon auratus and Golden-backed Tree-rat Mesembriomys macrurus 2004 - 2009*. Northern Territory Department of Infrastructure, Planning and Environment, Darwin.

- Payne, A. L. & Leighton, K. A. 2004. Land systems. In: van Vreeswyk, A. M. E., Payne, A. L., Leighton, K. A. & Hennig, P. (eds) *Technical Bulletin 9. An inventory and condition survey of the Pilbara region, Western Australia*. Department of Agriculture, Government of Western Australia, South Perth, WA, pp. 175–384.
- Phoenix. 2019. *Environmental desktop review for the Napier Downs Irrigation Project*. Phoenix Environmental Sciences Pty Ltd, Balcatta, WA. Unpublished report prepared for Australian Capital Equity Pty Ltd.
- Phoenix. 2020. *Environmental desktop review for the Napier Downs Irrigation Project – Report Addendum*. Phoenix Environmental Sciences Pty Ltd, Osborne Park, WA. Unpublished report prepared for Australian Capital Equity Pty Ltd.
- Queensland Department of Environment and Resource Management. 2012. *National recovery plan for the Red Goshawk *Erythrotriorchis radiatus**. Queensland Department of Environment and Resource Management, Brisbane, QLD.
- Ringma, J., Legge, S., J., W., Radford, J., Wintle, B. & Bode, M. 2018. Australia's mammal fauna requires a strategic and enhanced network of predator-free havens. *Nature Ecology and Evolution* **2**: 410-411.
- Rix, M. G., Huey, J. A., Cooper, S. J. B., Austin, A. D. & Harvey, M. S. 2018. Conservation systematics of the shield-backed trapdoor spiders of the *nigrum*-group (Mygalomorphae, Idiopidae, *Idiosoma*): integrative taxonomy reveals a diverse and threatened fauna from south-western Australia. *Zookeys* **756**: 1–121 <http://dx.doi.org/10.3897/zookeys.756.24397>.
- Schoknecht, N. R. & Payne, A. L. 2011. *Land systems of the Kimberley region, Western Australia*. Department of Agriculture and Food, Western Australia, Perth.
- Shepherd, D. P., Beeston, G. R. & Hopkins, A. J. M. 2002. *Native vegetation in Western Australia. Extent, type and status*. Department of Agriculture, South Perth, WA. Resource Management Technical Report 249.
- Stewart, A. J., Sweet, I. P., Needham, R. S., Raymond, O. L., Whitaker, A. J., Liu, S. F., Phillips, D., Retter, A. J., Connolly, D. P. & Stewart, G. 2008. *Surface geology of Australia 1:1,000,000 scale, Western Australia [Digital Dataset]*. The Commonwealth of Australia, Geoscience Australia, Canberra.
- Strahan, R. 1995. *Mammals of Australia*. Smithsonian Institution Press, Washington, DC.
- Threatened Species Scientific Committee. 2017. *Conservation Advice, *Phascogale tapoatafa kimberleyensis*, Kimberley brush-tailed phascogale*. Threatened Species Scientific Committee, Canberra, ACT.
- Tidemann, S. C. 1996. Causes of the decline of the Gouldian Finch *Erythrura gouldiae*. *Bird Conservation International* **6**: 49–61.
- Van Dyck, S. & Strahan, R. 2008. *The mammals of Australia*. New Holland Publishers, Sydney, NSW.
- WAM. 2013. *WAM short-range endemic categories and sub-categories*. Western Australian Museum, Welshpool.
- WAM. 2019. *WA Museum Arachnology/Myriapodology, Crustacea and Mollusca database*, Welshpool, WA.
- Webb, G., Whitehead, P. & Manolis, S. 1987. Crocodile management in the Northern Territory of Australia. In: GJW, W., Manolis, S. & Whitehead, P. J. (eds) *Wildlife management: Crocodiles and Alligators*. Surrey Beatty & Sons, Sydney, pp. 107-124.
- Woinarski J., Ward S., Mahney T., Bradley J., Brennan K., Ziembicki M. & A., F. 2011. The mammal fauna of the Sir Edward Pellew Islands, Northern Territory: refuge and death-trap. *Wildlife Research* **38**: 307-322.

**Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity**

**Appendix A Survey site locations**

Site name	Latitude	Longitude
SITE01	-17.2174	124.3829
SITE02	-17.2208	124.3884
SITE03	-17.2235	124.38
SITE04	-17.2279	124.3729
SITE05	-17.222	124.3937
SITE06	-17.2225	124.3976
SITE07	-17.2309	124.3965
SITE08	-17.2322	124.3932
SITE09	-17.22	124.3869
SITE10	-17.1944	124.4045
SITE11	-17.1932	124.3327
SITE12	-17.1371	124.471
SITE13	-17.196	124.258
SITE14	-17.3169	124.6778
SITE15	-17.1376	124.4851
SITE16	-17.2687	124.4025
Opp-Isoodon-01	-17.2253	124.3775
Opp-Isoodon-02	-17.2246	124.3841
Opp-Isoodon-03	-17.2264	124.3774
Opp-Isoodon-04	-17.219	124.3802
Opp-Isoodon-05	-17.2262	124.3811
Opp-Isoodon-06	-17.2262	124.3801
Opp-Isoodon-07	-17.2265	124.3795
Opp-Isoodon-08	-17.2249	124.3804
Opp-Isoodon-09	-17.2265	124.3774
Opp-Isoodon-10	-17.225	124.376
Opp-Isoodon-11	-17.2223	124.3953
Opp-Isoodon-12	-17.2205	124.3865
Opp-Isoodon-13	-17.2333	124.3949

Site name	Latitude	Longitude
Opp-01	-17.3037	124.6909
Opp-02	-17.2266	124.4007
Opp-03	-17.2334	124.3953
Opp-04	-17.1371	124.4721
Opp-05	-17.2911	124.6634
Opp-06	-17.1369	124.4716
Opp-07	-17.1769	124.561
Opp-08	-17.2232	124.4034
Opp-09	-17.2228	124.398
Opp-10	-17.2439	124.3873
Opp-11	-17.1849	124.4335
Opp-12	-17.1393	124.5001
Opp-13	-17.2476	124.6234
Opp-14	-17.292	124.7723
Opp-15	-17.2237	124.403
Opp-16	-17.2264	124.4009
Opp-17	-17.1427	124.5303
Opp-18	-17.2463	124.6211
Opp-19	-17.302	124.6736
Opp-20	-17.1778	124.4392
Opp-21	-17.2715	124.398
Opp-22	-17.2232	124.379

Appendix B Vertebrate fauna desktop and field survey results

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
<b>FISHES (1)</b>										
Pristidae	Pristis pristis	Freshwater Sawfish	MI & P3	VU & MI				•		
<b>FROGS (18)</b>										
Bufoidea	Rhinella marina	Cane Toad								•
Limnodynastidae	Notaden nichollsi	Desert Spadefoot					•			
Limnodynastidae	Platyplectrum ornatum	Ornate Burrowing Frog				•	•			
Myobatrachidae	Uperoleia aspera	Derby Toadlet				•	•			
Myobatrachidae	Uperoleia borealis	Northern Toadlet								•
Myobatrachidae	Uperoleia lithomoda	Stonemason Toadlet					•			
Myobatrachidae	Uperoleia mjobergii	West Kimberley Toadlet					•			
Pelodyadidae	Cyclorana australis	Giant Frog					•			
Pelodyadidae	Cyclorana cryptotis	Hidden-ear Frog				•	•			
Pelodyadidae	Cyclorana longipes	Long-footed Frog				•	•			
Pelodyadidae	Litoria caerulea	Green Tree Frog					•			
Pelodyadidae	Litoria inermis	Bumpy Rocket Frog					•			
Pelodyadidae	Litoria meiriana	Rockhole Frog					•			
Pelodyadidae	Litoria pallida	Pale Rocket Frog				•	•			
Pelodyadidae	Litoria rothii	Northern Laughing Tree Frog				•	•			
Pelodyadidae	Litoria rubella	Little Red Tree Frog				•	•			
Pelodyadidae	Litoria splendida	Splendid Tree Frog					•			
Pelodyadidae	Litoria tornieri	Black-shinned Rocket Frog				•				
<b>REPTILES (52)</b>										
Agamidae	Lophognathus gilberti	Top End Ta-Ta Dragon					•			
Agamidae	Chlamydosaurus kingii	Frill-necked Lizard				•	•			
Agamidae	Ctenophorus caudicinctus	Ring-tailed Dragon					•			

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Agamidae	Ctenophorus nuchalis	Central Notted Dragon					•			
Agamidae	Diporiphora lalliae	Northern Deserts Dragon					•			
Agamidae	Diporiphora magna	Yellow-sided Two-lined Dragon				•	•			
Agamidae	Diporiphora margaretae	Northwest Kimberley Two-lined Dragon								•
Agamidae	Diporiphora pindan	Pindan Dragon					•			
Agamidae	Pogona minor	Northwestern Bearded Dragon					•			
Boidae	Liasis olivaceus	Olive Python								•
Carphodactylidae	Nephrurus sheai	Northern Knob-tailed Gecko					•			
Colubridae	Boiga irregularis	Brown Tree Snake					•			
Crocodylidae	Crocodylidae	Australian freshwater crocodile	OS		•		•			
Crocodylidae	Crocodylus porosus	Salt-water Crocodile				•		•		
Diplodactylidae	Crenodactylus ocellatus	Clawless Gecko					•			
Diplodactylidae	Lucasium stenodactylus	Western Sandplain Gecko				•				
Diplodactylidae	Strophurus ciliaris	Northern Spiny-tailed Gecko				•	•			
Diplodactylidae	Strophurus taeniatus	Northern Phasmid Gecko					•			
Elapidae	Furina ornata	Moon Snake					•			
Gekkonidae	Gehyra montium	Central Rock Dtella				•				
Gekkonidae	Gehyra nana	Northern Spotted Rock Dtella					•			
Gekkonidae	Gehyra occidentalis	Kimberley Plateau Dtella				•	•			
Gekkonidae	Gehyra pilbara	Pilbara Dtella					•			
Gekkonidae	Gehyra punctata	Spotted Dtella					•			
Gekkonidae	Heteronotia binoei	Bynoe's Gecko					•			
Gekkonidae	Heteronotia planiceps	North-west Prickly Gecko				•	•			
Pygopodidae	Delma borea	Rusty-topped Delma				•	•			
Pygopodidae	Delma nasuta	Sharp-snouted Delma					•			
Pygopodidae	Lialis burtonis	Burton's Legless Lizard					•			

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Pythonidae	<i>Aspidites melanocephalus</i>	Black-headed Python					•			
Scincidae	<i>Carlia munda</i>	Shaded-litter Rainbow Skink					•			
Scincidae	<i>Cryptoblepharus megastictus</i>	Blotched Snake-eyed Skink					•			
Scincidae	<i>Cryptoblepharus metallicus</i>	Metallic Snake-eyed Skink					•			
Scincidae	<i>Cryptoblepharus ruber</i>	Tawny Snake-eyed Skink				•				
Scincidae	<i>Cryptoblepharus tythos</i>	Pygmy Snake-Eyed Skink					•			
Scincidae	<i>Ctenotus inornatus</i>	Plain Ctenotus				•				
Scincidae	<i>Ctenotus serventyi</i>	North-western Sandy-loam Ctenotus					•			
Scincidae	<i>Cyclodomorphus melanops</i>	Slender Blue-tongue					•			
Scincidae	<i>Eremiascincus isolepis</i>	Northern Bar-lipped Skink					•			•
Scincidae	<i>Lerista bipes</i>	North-western Sandslider								•
Scincidae	<i>Lerista borealis</i>	Northern Slider					•			
Scincidae	<i>Lerista greeri</i>	South-eastern Kimberley Sand Slider				•				
Scincidae	<i>Menetia greyii</i>	Common Dwarf Skink					•			
Scincidae	<i>Morethia ruficauda</i>	Lined Firetail Skink					•			
Scincidae	<i>Proablepharus tenuis</i>	Slender Snake-eyed Skink					•			
Scincidae	<i>Tiliqua scincoides</i>	Eastern Blue-tongue				•				
Scincidae	<i>Tiliqua scincoides</i>	Common Blue-tongue					•			
Varanidae	<i>Varanus acanthurus</i>	Spiny-tailed Monitor					•			
Varanidae	<i>Varanus gouldii</i>	Bungarra or Sand Monitor					•			
Varanidae	<i>Varanus panoptes</i>	Yellow-spotted Monitor					•			
Varanidae	<i>Varanus scalaris</i>	Spotted Tree Monitor				•				
Varanidae	<i>Varanus storri</i>	Storr's Monitor				•				



Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
<b>Birds (221)</b>										
Acanthizidae	<i>Gerygone olivacea</i>	White-throated Gerygone				•	•		•	
Acanthizidae	<i>Smicronis brevirostris</i>	Weebill				•	•		•	•
Acanthizidae	<i>Gerygone chloronota</i>	Green-backed Gerygone				•	•		•	
Accipitridae	<i>Aquila audax</i>	Wedge-tailed Eagle				•	•		•	•
Accipitridae	<i>Circus approximans</i>	Swamp Harrier				•	•		•	•
Accipitridae	<i>Circus assimilis</i>	Spotted Harrier				•	•		•	
Accipitridae	<i>Erythroriorchis radiatus</i>	Red Goshawk						•		
Accipitridae	<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle				•			•	
Accipitridae	<i>Haliastur sphenurus</i>	Whistling Kite				•	•		•	•
Accipitridae	<i>Hamirostra isura</i>	Square-tailed Kite				•			•	
Accipitridae	<i>Hamirostra melanosternon</i>	Black-breasted Buzzard				•	•		•	•
Accipitridae	<i>Accipiter cirrocephalus</i>	Collared Sparrowhawk				•	•		•	•
Accipitridae	<i>Accipiter fasciatus</i>	Brown Goshawk				•	•		•	•
Accipitridae	<i>Accipiter novaehollandiae</i>	Grey Goshawk							•	
Accipitridae	<i>Elanus caeruleus</i>	Black-shouldered Kite					•		•	
Accipitridae	<i>Milvus migrans</i>	Black Kite				•	•		•	
Accipitridae	<i>Hieraetus morphnoides</i>	Little Eagle				•	•		•	•
Acrocephalidae	<i>Acrocephalus orientalis</i>	Oriental Reed Warbler						•		
Aegothelidae	<i>Aegotheles cristatus</i>	Australian Owllet-nightjar				•	•		•	
Alaudidae	<i>Mirafra javanica</i>	Horsfield's Bushlark				•	•		•	
Alcedinidae	<i>Dacelo leachii</i>	Blue-winged Kookaburra				•	•		•	•
Alcedinidae	<i>Todiramphus sanctus</i>	Sacred Kingfisher				•	•		•	•
Alcedinidae	<i>Todiramphus pyrrophygius</i>	Red-backed Kingfisher				•	•		•	•
Alcedinidae	<i>Ceyx azureus</i>	Azure Kingfisher				•	•		•	•

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Anatidae	Anas gracilis	Grey Teal				•	•		•	
Anatidae	Anas superciliosa	Pacific Black Duck				•	•		•	•
Anatidae	Aythya australis	Hardhead				•	•		•	•
Anatidae	Cygnus atratus	Black Swan				•	•			
Anatidae	Dendrocygna eytoni	Plumed Whistling Duck				•	•		•	
Anatidae	Malacorhynchus membranaceus	Pink-eared Duck				•	•		•	
Anatidae	Nettapus pulchellus	Green Pygmy-goose				•	•		•	
Anatidae	Stictonetta naevosa	Freckled Duck				•				
Anhingiidae	Anhinga novaehollandiae	Australasian Darter				•	•		•	
Anseranatidae	Anseranas semipalmata	Maggie Goose				•			•	•
Apodidae	Apus pacificus	Fork-tailed Swift						•		
Ardeidae	Ardea novaehollandiae	White-faced Heron				•	•		•	•
Ardeidae	Ardea pacifica	White-necked Heron				•	•		•	•
Ardeidae	Ardea alba	Great Egret				•				•
Ardeidae	Ardea garzetta	Little Egret				•	•		•	
Ardeidae	Ardea ibis	Cattle Egret				•				•
Ardeidae	Ardea intermedia	Intermediate Egret				•	•		•	•
Ardeidae	Ixobrychus flavicollis	Black Bittern				•			•	
Ardeidae	Nycticorax caledonicus	Rufous Night Heron				•	•		•	•
Ardeidae	Botaurus poicilopilus	Australasian Bittern	EN	EN	•					
Ardeidae	Ardea modesta	Great Egret				•	•		•	
Artamidae	Artamus minor	Little Woodswallow				•	•		•	•
Artamidae	Artamus personatus	Masked Woodswallow				•	•		•	•
Artamidae	Cracticus nigrogularis	Pied Butcherbird				•	•		•	•
Artamidae	Artamus cinereus	Black-faced Woodswallow				•	•		•	•
Artamidae	Artamus leucorhynchus	White-breasted Woodswallow				•	•		•	•

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Artamidae	<i>Cracticus tibicen</i>	Australian Magpie				•	•		•	•
Artamidae	<i>Cracticus torquatus</i>	Grey Butcherbird					•		•	
Burhinidae	<i>Burhinus grallarius</i>	Bush Stone-curlew					•		•	•
Cacatuidae	<i>Nymphicus hollandicus</i>	Cockatiel				•	•		•	
Cacatuidae	<i>Cacatua galerita</i>	Sulphur-crested Cockatoo				•	•		•	•
Cacatuidae	<i>Cacatua roseicapilla</i>	Galah				•	•		•	•
Cacatuidae	<i>Cacatua sanguinea</i>	Little Corella				•	•		•	•
Cacatuidae	<i>Calyptorhynchus banksii</i>	Red-tailed Black-Cockatoo				•	•		•	•
Campephagidae	<i>Coracina maxima</i>	Ground Cuckoo-shrike							•	
Campephagidae	<i>Lalage tricolor</i>	White-winged Triller				•	•		•	•
Campephagidae	<i>Coracina novaehollandiae</i>	Black-faced Cuckoo-shrike				•	•		•	•
Campephagidae	<i>Coracina papuensis</i>	White-bellied Cuckoo-shrike				•	•		•	
Campephagidae	<i>Lalage leucomela</i>	Varied Triller					•		•	
Caprimulgidae	<i>Eurostopus argus</i>	Spotted Nightjar				•	•		•	•
Casuariidae	<i>Dromaius novaehollandiae</i>	Emu				•				
Charadriidae	<i>Charadrius veredus</i>	Oriental Plover	MI	MI	•	•	•	•	•	
Charadriidae	<i>Erythronyx cinctus</i>	Red-kneed Dotterel				•	•		•	
Charadriidae	<i>Charadrius leschenaultii</i>	Greater Sand Plover				•		•		
Charadriidae	<i>Vanellus miles</i>	Masked Lapwing				•	•		•	•
Charadriidae	<i>Eisayornis melanops</i>	Black-fronted Dotterel				•	•		•	•
Ciconiidae	<i>Ephippiorhynchus asiaticus</i>	Black-necked Stork				•	•		•	•
Cisticolidae	<i>Cisticola exilis</i>	Golden-headed Cisticola				•	•		•	
Climacteridae	<i>Climacteris melanurus</i>	Black-tailed Treecreeper				•			•	
Columbidae	<i>Chalcophaps indica subsp. longirostris</i>	Emerald Dove							•	

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Columbidae	<i>Geopelia cuneata</i>	Diamond Dove				•	•		•	•
Columbidae	<i>Geopelia humeralis</i>	Bar-shouldered Dove				•	•		•	•
Columbidae	<i>Geopelia striata</i> subsp. <i>placida</i>	Peaceful Dove				•	•		•	
Columbidae	<i>Geophaps plumifera</i>	Spinifex Pigeon					•		•	
Columbidae	<i>Ocyphaps lophotes</i>	Crested Pigeon				•	•		•	•
Columbidae	<i>Phaps chalcoptera</i>	Common Bronzewing				•			•	
Columbidae	<i>Petrophassa albipennis</i>	White-quilled Rock Pigeon							•	
Columbidae	<i>Geophaps smithii</i>	Partridge Pigeon						•		
Coraciidae	<i>Eurystomus orientalis</i>	Dollarbird				•	•		•	
Corvidae	<i>Corvus bennetti</i>	Little Crow					•		•	
Corvidae	<i>Corvus orru</i>	Torresian Crow				•	•		•	•
Cuculidae	<i>Chrysococcyx basalis</i>	Horsfield's Bronze Cuckoo				•	•		•	•
Cuculidae	<i>Chrysococcyx osculans</i>	Black-eared Cuckoo				•			•	
Cuculidae	<i>Cacomantis variolosus</i>	Brush Cuckoo				•	•		•	
Cuculidae	<i>Centropus phasianinus</i>	Pheasant Coucal				•	•		•	
Cuculidae	<i>Eudynamys scolopacea</i>	Common Koel					•			
Cuculidae	<i>Scythrops novaehollandiae</i>	Channel-billed Cuckoo					•		•	
Cuculidae	<i>Cacomantis pallidus</i>	Pallid Cuckoo				•	•		•	•
Cuculidae	<i>Cuculus optatus</i>	Oriental Cuckoo						•		
Cuculidae	<i>Eudynamys orientalis</i>	Pacific Koel					•		•	
Dicaeidae	<i>Dicaeum hirundinaceum</i>	Mistletoebird				•	•		•	
Estrilidae	<i>Emblema pictum</i>	Painted Finch					•		•	
Estrilidae	<i>Erythrura gouldiae</i>	Gouldian Finch	P4	EN	•	•	•	•	•	•
Estrilidae	<i>Heteromunia pectoralis</i>	Pictorella Mannikin				•	•		•	
Estrilidae	<i>Poephila acuticauda</i>	Long-tailed Finch				•	•		•	•

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Estrilidae	<i>Neochmia phaeton</i>	Crimson Finch				•			•	•
Estrilidae	<i>Neochmia ruficauda</i>	Star Finch				•	•		•	•
Estrilidae	<i>Poephila personata</i>	Masked Finch							•	•
Estrilidae	<i>Taeniopygia guttata</i>	Zebra Finch				•	•		•	•
Estrilidae	<i>Taeniopygia bichenovii</i>	Double-barred Finch				•	•		•	•
Falconidae	<i>Falco hypoleucos</i>	Grey Falcon	VU		•	•		•	•	
Falconidae	<i>Falco subniger</i>	Black Falcon				•	•		•	
Falconidae	<i>Falco berigora</i>	Brown Falcon				•	•		•	•
Falconidae	<i>Falco cenchroides</i>	Australian Kestrel				•	•		•	•
Falconidae	<i>Falco longipennis</i>	Australian Hobby				•			•	•
Falconidae	<i>Falco peregrinus</i>	Peregrine Falcon	OS		•	•	•		•	
Falconidae	<i>Falcunculus frontatus</i>	Crested Shrike-tit						•		
Glareolidae	<i>Glareola maldivarum</i>	Oriental Pratincole						•		
Gruidae	<i>Grus rubicunda</i>	Brolga				•	•		•	•
Gruidae	<i>Grus antigone</i>	Sarus Crane				•				
Hirundinidae	<i>Hirundo rustica</i>	Barn Swallow						•		
Hirundinidae	<i>Cecropis daurica</i>	Red-rumped Swallow						•		
Hirundinidae	<i>Petrochelidon ariel</i>	Fairy Martin				•	•		•	•
Hirundinidae	<i>Petrochelidon nigricans</i>	Tree Martin				•	•		•	•
Jacaniidae	<i>Irediparra gallinacea</i>	Comb-crested Jacana							•	
Laridae	<i>Sterna hybrida</i>	Whiskered Tern				•			•	
Laridae	<i>Gelochelidon nilotica</i>	Gull-billed Tern	MI	MI	•	•				
Locustellidae	<i>Cincloramphus mathewsi</i>	Rufous Songlark				•			•	
Locustellidae	<i>Megalurus timoriensis</i>	Tawny Grassbird					•		•	•
Maluridae	<i>Malurus coronatus</i>	Purple-crowned Fairy-wren				•				
Maluridae	<i>Malurus lamberti</i>	Variegated Fairy-wren				•	•		•	•

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Maluridae	Malurus melanocephalus	Red-backed Fairy-wren				•	•		•	•
Maluridae	Amytornis housei	Black Grasswren	P4							•
Meliphagidae	Cissomela pectoralis	Banded Honeyeater				•	•		•	•
Meliphagidae	Conopophila rufogularis	Rufous-throated Honeyeater				•	•		•	
Meliphagidae	Manorina flavigula	Yellow-throated Miner				•	•		•	
Meliphagidae	Melithreptus albogularis	White-throated Honeyeater				•	•		•	•
Meliphagidae	Ramsayornis fasciatus	Bar-breasted Honeyeater				•	•		•	
Meliphagidae	Lichmera indistincta	Brown Honeyeater				•	•		•	•
Meliphagidae	Melithreptus cyanotis	Blue-faced Honeyeater							•	
Meliphagidae	Melithreptus gularis	Black-chinned Honeyeater				•	•		•	•
Meliphagidae	Myzomela erythrocephala	Red-headed Honeyeater							•	
Meliphagidae	Philemon argenticeps	Silver-crowned Friarbird				•	•		•	
Meliphagidae	Philemon citreogularis	Little Friarbird				•	•		•	•
Meliphagidae	Gavicalis virescens	Singing Honeyeater				•	•		•	•
Meliphagidae	Ptilotula flavescens	Yellow-tinted Honeyeater				•	•		•	•
Meliphagidae	Stomiopera unicolor	White-gaped Honeyeater				•	•		•	•
Meliphagidae	Ptilotula plumula	Grey-fronted Honeyeater				•	•		•	
Meropidae	Merops ornatus	Rainbow Bee-eater				•	•		•	•
Monarchidae	Grallina cyanoleuca	Magpie-lark				•	•		•	•
Monarchidae	Myiagra inquieta	Restless Flycatcher				•	•		•	•
Monarchidae	Myiagra rubecula	Leaden Flycatcher				•	•		•	
Monarchidae	Myiagra ruficollis	Broad-billed Flycatcher							•	
Motacillidae	Anthus australis	Australian Pipit				•			•	
Motacillidae	Motacilla flava	Yellow Wagtail						•		
Motacillidae	Motacilla cinerea	Grey Wagtail						•		
Neosittidae	Daphoenositta chrysoptera	Varied Sittella				•	•		•	

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Oriolidae	Oriolus sagittatus	Olive-backed Oriole				•	•		•	•
Oriolidae	Sphecotheres vieilloti	Australasian Figbird							•	
Otididae	Ardeotis australis	Australian Bustard				•	•		•	•
Pachycephalidae	Colluricincla woodwardi	Sandstone Shrike-thrush				•	•		•	
Pachycephalidae	Colluricincla harmonica	Grey Shrike-thrush				•	•		•	•
Pachycephalidae	Colluricincla megarhyncha	Little Shrike-thrush				•			•	
Pachycephalidae	Pachycephala rufiventris	Rufous Whistler				•	•		•	•
Pandionidae	Pandion haliaetus	Osprey	MI	MI	•			•	•	
Pardalotidae	Pardalotus rubricatus	Red-browed Pardalote				•	•		•	
Pardalotidae	Pardalotus striatus	Striated Pardalote				•	•		•	•
Pelecanidae	Pelecanus conspicillatus	Australian Pelican				•	•		•	
Petroicidae	Microeca fascians	Jacky Winter				•	•		•	
Petroicidae	Microeca flavigaster	Lemon-breasted Flycatcher					•		•	
Petroicidae	Poecilodryas superciliosa	White-browed Robin				•				
Petroicidae	Melanodryas cucullata	Hooded Robin							•	
Petroicidae	Poecilodryas cerviniventris	Buff-sided Robin					•		•	
Phalacrocoracidae	Phalacrocorax sulcirostris	Little Black Cormorant				•	•		•	
Phalacrocoracidae	Phalacrocorax carbo	Great Cormorant				•	•		•	
Phalacrocoracidae	Phalacrocorax melanoleucos	Little Pied Cormorant				•	•		•	•
Phalacrocoracidae	Phalacrocorax varius	Pied Cormorant				•	•		•	
Phasianidae	Coturnix ypsilophora	Brown Quail				•	•		•	•
Podargidae	Podargus strigoides	Tawny Frogmouth				•	•		•	
Podicipedidae	Poliiocephalus poliocephalus	Hoary-headed Grebe				•	•		•	

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Podicipedidae	<i>Tachybaptus novaehollandiae</i>	Australasian Grebe				•	•		•	•
Pomatostomidae	<i>Pomatostomus temporalis</i>	Grey-crowned Babbler				•	•		•	•
Psittaculidae	<i>Aprosmictus erythropterus</i>	Red-winged Parrot				•	•		•	•
Psittaculidae	<i>Melopsittacus undulatus</i>	Budgerigar				•	•		•	
Psittaculidae	<i>Neophema bourkii</i>	Bourke's Parrot				•	•			
Psittaculidae	<i>Pezoporus occidentalis</i>	Night Parrot						•		
Psittaculidae	<i>Platycercus venustus</i>	Northern Rosella				•	•		•	
Psittaculidae	<i>Polytelis alexandrae</i>	Princess Parrot						•		
Psittaculidae	<i>Trichoglossus rubitorquus</i>	Red-collared Lorikeet				•	•		•	•
Psittaculidae	<i>Trichoglossus versicolor</i>	Varied Lorikeet				•	•		•	
Psittaculidae	<i>Trichoglossus haematodus</i>	Rainbow Lorikeet				•	•			
Ptilonorhynchidae	<i>Ptilonorhynchus nuchalis</i>	Great Bowerbird				•	•		•	•
Rallidae	<i>Fulica atra</i>	Eurasian Coot				•	•		•	
Rallidae	<i>Gallirallus philippensis</i>	Buff-banded Rail							•	
Rallidae	<i>Amaurornis moluccana</i> subsp. <i>ruficrissa</i>	Pale-vented Bush-hen							•	
Recurvirostridae	<i>Himantopus himantopus</i>	Black-winged Stilt				•	•		•	
Rhipiduridae	<i>Rhipidura fuliginosa</i>	Grey Fantail							•	
Rhipiduridae	<i>Rhipidura leucophrys</i>	Willie Wagtail				•	•		•	•
Rhipiduridae	<i>Rhipidura rufiventris</i>	Northern Fantail				•			•	
Rhipiduridae	<i>Rhipidura albiscapa</i>	Grey Fantail					•			
Rhipiduridae	<i>Rostratula australis</i>	Australian Painted Snipe						•		
Scolopacidae	<i>Calidris acuminata</i>	Sharp-tailed Sandpiper	MI	MI	•	•			•	•
Scolopacidae	<i>Calidris ferruginea</i>	Curlew Sandpiper						•		•



Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Scolopacidae	Calidris melanotos	Pectoral Sandpiper				•		•		
Scolopacidae	Calidris ruficollis	Red-necked Stint				•				
Scolopacidae	Gallinago hardwickii	Latham's Snipe				•				
Scolopacidae	Numenius madagascariensis	Eastern Curlew						•		
Scolopacidae	Tringa glareola	Wood Sandpiper	MI	MI	•	•	•		•	
Scolopacidae	Tringa nebularia	Common Greenshank	MI	MI	•	•	•			
Scolopacidae	Tringa stagnatilis	Marsh Sandpiper	MI	MI	•	•	•		•	
Scolopacidae	Calidris canutus	Red Knot						•		
Scolopacidae	Limosa lapponica	Bar-tailed Godwit						•		
Scolopacidae	Actitis hypoleucos	Common Sandpiper			•	•		•	•	
Strigidae	Ninox rufa subsp. rufa	Rufous Owl					•			
Strigidae	Ninox connivens	Barking Owl				•	•		•	
Strigidae	Ninox novaeseelandiae	Boobook Owl				•				
Strigidae	Ninox boobook	Australian Boobook							•	
Threskiornithidae	Platalea flavipes	Yellow-billed Spoonbill				•	•			
Threskiornithidae	Platalea regia	Royal Spoonbill				•	•		•	
Threskiornithidae	Plegadis falcinellus	Glossy Ibis	MI	MI	•	•	•		•	
Threskiornithidae	Threskiornis spinicollis	Straw-necked Ibis				•	•		•	•
Threskiornithidae	Threskiornis moluccus	Australian White Ibis				•			•	
Turnicidae	Turnix pyrrhothorax	Red-chested Button-quail					•		•	
Turnicidae	Turnix velox	Little Button-quail					•		•	
Turnicidae	Turnix maculosa	Red-backed Button-quail							•	
Tytonidae	Tyto alba	Barn Owl							•	
Tytonidae	Tyto novaehollandiae	Masked Owl						•	•	
<b>Mammals (46)</b>										
Bovidae	Bos taurus	European Cattle					•			•

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Canidae	Canis familiaris	Dog								•
Dasyuridae	Dasyurus hallucatus	Northern Quoll	EN	EN	•		•	•		
Dasyuridae	Phascogale tapoatafa	Brush-tailed Phascogale						•		
Dasyuridae	Pseudantechinus ningbing	Ningbing Pseudantechinus					•			•
Emballonuridae	Saccolaimus flaviventris	Yellow-bellied Sheath-tailed Bat				•	•			
Emballonuridae	Saccolaimus saccolaimus	Bare-rumped Sheath-tail-bat						•		
Emballonuridae	Taphozous georgianus	Common Sheath-tailed Bat				•	•			
Equidae	Equus caballus	Horse					•			
Felidae	Felis catus	Cat					•			•
Hipposideridae	Hipposideros ater	Dusky Leafnosed-bat					•			
Hipposideridae	Hipposideros stenotis	northern leaf-nosed bat	P2		•		•			
Macropodidae	Macropus agilis	Agile Wallaby					•			
Macropodidae	Macropus antilopinus	Antilopine Wallaroo					•			
Macropodidae	Macropus robustus	Euro					•			•
Macropodidae	Macropus rufus	Red Kangaroo					•			
Macropodidae	Notamacropus agilis	Agile Wallaby				•				
Macropodidae	Onychogalea unguifera	Northern Nailtail Wallaby				•	•			•
Macropodidae	Petrogale brachyotis	Short-eared Rock-wallaby				•	•			
Macropodidae	Petrogale concinna	Nabarlek						•		
Macropodidae	Petrogale lateralis	Black-footed Rock-wallaby						•		
Megadermatidae	Macroderma gigas	Ghost Bat	VU	VU	•		•	•		
Miniopteridae	Miniopterus orianae	Northern Bent-winged Bat or Large Bent-winged Bat					•			•
Molossidae	Chaerephon jobensis	Greater Northern Freetail-bat				•	•			•
Muridae	Leggadina lakedownensis	Northern Short-tailed Mouse, Lakeland Downs mouse, kerakenga	P4		•	•	•			

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Family	Species	Common name	BC Act	EPBC Act	TPFA	ALA	Nature Map	PMST	Birdlife Bird Data	This Survey
Muridae	<i>Pseudomys delicatulus</i>	Delicate Mouse					•			•
Muridae	<i>Pseudomys nanus</i>	Western Chestnut Mouse				•	•			•
Muridae	<i>Rattus tunneyi</i>	Pale Field-rat				•	•			•
Muridae	<i>Rattus villosissimus</i>	Long-haired Rat								•
Muridae	<i>Zyomys argurus</i>	Common Rock-rat				•	•			
Peramelidae	<i>Isoodon auratus</i>	Golden Bandicoot	VU	VU				•		•
Peramelidae	<i>Isoodon macrourus</i>	Northern Brown Bandicoot								•
Phalangeridae	<i>Trichosurus vulpecula arnhemensis</i>	Northern Brushtail Possum	VU	VU	•			•		
Pseudocheiridae	<i>Petropseudes dahli</i>	Rock Ringtail Possum	P3		•	•	•			
Pteropodidae	<i>Pteropus alecto</i>	Black Flying-fox				•	•			
Pteropodidae	<i>Pteropus scapulatus</i>	Little Red Flying-fox				•				
Rhinonycteridae	<i>Rhinonictis aurantia</i>	Orange leaf-nosed bat	P4		•		•			
Tachyglossidae	<i>Tachyglossus aculeatus</i>	Short-beaked Echidna					•			
Thylacomyidae	<i>Macrotis lagotis</i>	Bilby	VU	VU	•			•		
Vespertilionidae	<i>Chalinolobus gouldii</i>	Gould's Wattled Bat				•	•			•
Vespertilionidae	<i>Chalinolobus nigrogriseus</i>	Hoary Wattled Bat				•	•			•
Vespertilionidae	<i>Nyctophilus arnhemensis</i>	Arnhem Land Long-eared Bat					•			
Vespertilionidae	<i>Nyctophilus walkeri</i>	Pygmy Long-eared Bat					•			
Vespertilionidae	<i>Scotorepens greyii</i>	Little Broad-nosed Bat					•			
Vespertilionidae	<i>Vespardelus caurinus</i>	Western Cave Bat				•	•			
Vespertilionidae	<i>Vespardelus douglasorum</i>	Yellow-lipped Cave Bat	P2		•	•	•			

Appendix C Fauna species by site matrix

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total															
<b>AMPHIBIANS</b>																																																						
<i>Rhinella marina</i>															1																											1												
<i>Litoria rothii</i>									3																																				3									
<i>Uperoleia borealis</i>	1																																												1									
<b>REPTILES</b>																																																						
<i>Liasis olivaceus</i>																1																															1							
<i>Ctenotus piankai</i>		1																																														1						
<i>Eremiascincus isolepis</i>								1																																									1					
<i>Lerista bipes</i>	9			1				1																																								11						
<i>Morethia ruficauda</i>								2	1																																								3					
<i>Diporiphora margaretae</i>		1																																														1						
<i>Tiliqua scincoides</i>			1																																															1				
<b>BIRDS</b>																																																						
<i>Anas superciliosa</i>										42																																									42			
<i>Aythya australis</i>										1																																										1		
<i>Anseranas semipalmata</i>									2																																											2		
<i>Eurostodopus argus</i>																																																					3	
<i>Burhinus grallarius</i>																																																					2	
<i>Elsyaornis melanops</i>										2																																											4	

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total			
<i>Vanellus miles</i>																													5											5		
<i>Ardea ibis</i>								1		1																															1	
<i>Ardea intermedia</i>								2																																		2
<i>Ardea modesta</i>															1																										1	
<i>Ardea novaehollandiae</i>								1		1			1																												3	
<i>Ardea pacifica</i>								1		3			1		2																											7
<i>Nycticorax caledonicus</i>													1																												1	
<i>Ephippiorhynchus asiaticus</i>																												3													3	
<i>Threskiornis spinicollis</i>								1		2																															3	
<i>Geopelia cuneata</i>	6										1																														7	
<i>Geopelia humeralis</i>	2						1		10	1	1	1	1	1	1	1																									17	
<i>Ocyphaps lophotes</i>											1																														1	
<i>Ceyx azureus</i>															1																										1	
<i>Dacelo leachii</i>											1			1																											2	
<i>Todiramphus pyrrhopygius</i>				1																																					1	
<i>Todiramphus sanctus</i>													1	1	1																										2	
<i>Merops ornatus</i>	2			1	2	1	2	1	5						1	1																									16	
<i>Cacomantis pallidus</i>											1																														1	
<i>Chrysococcyx basalis</i>	1			1		1	1	1	1	1																															6	

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total			
<i>Accipiter cirrocephalus</i>	1			1	2	1				1																														5		
<i>Accipiter fasciatus</i>		1																																							1	
<i>Aquila audax</i>	1																																								1	
<i>Circus approximans</i>										1																															1	
<i>Haliastur sphenurus</i>								1		1	1	1	2	2	1	2																									7	
<i>Hamirostra melanosternon</i>													1			3																									4	
<i>Hieraaetus morphnoides</i>															1																										1	
<i>Falco berigora</i>										1	1	1								1																					2	
<i>Falco cenchroides</i>																												1													1	
<i>Falco longipennis</i>										1																															1	
<i>Coturnix ypsilophora</i>	2	3									1																														6	
<i>Grus rubicunda</i>										1												3																				4
<i>Ardeotis australis</i>										1					1	2											1															5
<i>Smicromis brevirostris</i>															1																											1
<i>Artamus cinereus</i>							1							15							3																				19	
<i>Artamus leucorynchus</i>																																			3						3	
<i>Artamus minor</i>									1					6	1																				1						9	
<i>Artamus personatus</i>								3																																		3
<i>Coracina novaehollandiae</i>	1	1	1					1																																		4

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total				
<i>Lalage tricolor</i>				1																																				1			
<i>Corvus orru</i>				1	1	1		12		1																															16		
<i>Cracticus nigrogularis</i>			1				1																																		2		
<i>Cracticus tibicen</i>																																									1		
<i>Grallina cyanoleuca</i>	2	2			2				1	1	10	1	1	1	1	2																									12		
<i>Myiagra inquieta</i>				1							1	1	1	1								1																			5		
<i>Rhipidura leucophrys</i>	1				1	1	1	1	1	1	3	1	1	1	1	1																									13		
<i>Erythrura gouldiae</i>									3																																3		
<i>Neochmia phaeton</i>																																		2							2		
<i>Neochmia ruficauda</i>																																		3							3		
<i>Poephila acuticauda</i>							3	30	3	30	1																															74	
<i>Poephila personata</i>							1		1																																		4
<i>Taeniopygia bichenovii</i>										10	1	1	1																					1								13	
<i>Taeniopygia guttata</i>					30	2					1																															33	
<i>Petrochelidon ariel</i>								6		6			5																														11
<i>Petrochelidon nigricans</i>															4																					2						6	
<i>Amytornis housei</i>																																											1
<i>Malurus lamberti</i>	3																																									3	
<i>Malurus melanocephalus</i>	3													3	1																												7

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total	
<i>Cissomela pectoralis</i>						2															1																		3	
<i>Gavialis virescens</i>	1	3	1	1	1	2	1	1	1				1																											13
<i>Lichmera indistincta</i>	2	2	1	6	1	1	1	2																																16
<i>Manorina flavigula</i>																																							5	
<i>Melithreptus albogularis</i>			1																																				1	
<i>Melithreptus brevirostris</i>											6																												6	
<i>Philemon argenticeps</i>	1																																						1	
<i>Philemon citreogularis</i>	1	2	1	1			1	1	1				1																										8	
<i>Ptilotula flavescens</i>	1	2	3					2					5	1	2																									16
<i>Stomiopera unicolor</i>													1	2																									3	
<i>Oriolus sagittatus</i>																																							2	
<i>Colluricincla harmonica</i>	1						1																																2	
<i>Pachycephala rufiventris</i>	1			1			1						1	3																						1			8	
<i>Pardalotus striatus</i>	5			1												2																							8	
<i>Pomatostomus temporalis</i>	1	1	4	2	1	1	1	1					2																										12	
<i>Ptilonorhynchus nuchalis</i>													1																										5	
<i>Megalururus timoriensis</i>							1	1																															2	
<i>Phalacrocorax melanoleucos</i>											2																												2	



Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total		
<i>Tachybaptus novaehollandiae</i>										47																														47	
<i>Aprosmictus erythropterus</i>					4	2					1		5																												12
<i>Cacatua galerita</i>													1																												1
<i>Cacatua roseicapilla</i>	2		1		8	1	1	40							1																									54	
<i>Cacatua sanguinea</i>	1							2					4																											7	
<i>Calyptorhynchus banksii</i>			1					4								2																								13	
<i>Melopsittacus undulatus</i>								20																																20	
<i>Trichoglossus rubritorquis</i>		4	3	2																																				21	
<b>MAMMALS</b>																																									
<i>Bos taurus</i>	1	3	1	1	1	1	1	1					1			1									5																15
<i>Canis familiaris</i>		1				1										1																									4
<i>Felis catus</i>								1								1																									3
<i>Pseudantechinus ningbing</i>									1																																1
<i>Onychogalea unguifera</i>	1			1																																					4
<i>Osphranter robustus</i>				1										1																											2
<i>Isodon auratus / macrourus</i>			1	3																																					4
<i>Isodon auratus</i>																																									1
<i>Isodon macrourus</i>								1																																	1
<i>Pseudomys delicatulus</i>	2	1	1	1	1	1	1	2	1																																10

Baseline Terrestrial Fauna Survey for the Napier Downs Irrigation Project  
Prepared for Australian Capital Equity

Species	SITE01	SITE02	SITE03	SITE04	SITE05	SITE06	SITE07	SITE08	SITE09	SITE10	SITE11	SITE12	SITE13	SITE14	SITE15	SITE16	Opp-01	Opp-02	Opp-03	Opp-04	Opp-05	Opp-06	Opp-07	Opp-08	Opp-09	Opp-10	Opp-11	Opp-12	Opp-13	Opp-14	Opp-15	Opp-16	Opp-17	Opp-18	Opp-19	Opp-20	Opp-21	Opp-22	Grand Total				
<i>Pseudomys nanus</i>					1	1	1	1																																	4		
<i>Rattus tunneyi</i>						1	1																																				2
<i>Rattus villosissimus</i>	1				1	1	1	1																																			5