

Vertebrate Fauna Survey – Yanchep Lagoon, Yanchep



Version 2. February 2020

Prepared for:
City of Wanneroo
Locked Bag 1
Wanneroo, WA 6946

By: Terrestrial Ecosystems 10 Houston Place Mt Claremont WA 6010

RECORD OF DISTRIBUTION

| No. of copies | Report File Name | Report Status | Date | Prepared for: | Initials |
|---------------|---------------------|---------------|------------------|------------------|----------|
| Electronic | 2019-0054-006-gt-V1 | Draft | 8 January 2020 | City of Wanneroo | GT/ST |
| Electronic | 2019-0054-006-gt-V2 | Final | 12 February 2020 | City of Wanneroo | GT/ST |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

Suggested citation: Terrestrial Ecosystems (2020) *Vertebrate Fauna Survey – Yanchep Lagoon, Yanchep*. Unpublished report for the City of Wanneroo, Perth.

DISCLAIMER

This document is prepared in accordance with and subject to an agreement between Terrestrial Ecosystems and the client, City of Wanneroo. It has been prepared and is restricted to those issues that have been raised by the client in its engagement of Terrestrial Ecosystems and prepared using the standard of skill and care ordinarily exercised by environmental scientists in the preparation of such reports.

Persons or agencies that rely on or use this document for purposes or reasons other than those agreed by Terrestrial Ecosystems and its client without first obtaining prior consent, do so at their own risk and Terrestrial Ecosystems denies all liability in tort, contract or otherwise for any loss, damage or injury of any kind whatsoever (whether in negligence or otherwise) that may be suffered as a consequence.

Front Cover: Fauna habitat in the project area



Contents

| 1 | Inti | roduction | 1 |
|---|------|--|----------------------------|
| | 1.1 | Background | 1 |
| | 1.2 | Project objectives and scope of works | 1 |
| 2 | Exi | isting Environment | 2 |
| | 2.1 | Location of project area | 2 2 2 2 2 3 |
| | 2.2 | Landforms and soils | 2 |
| | 2.3 | Land use history | 2 |
| | 2.4 | Climate | 2 |
| | 2.5 | Regional biological fauna context of project area | 3 |
| | 2.5 | .1 Fauna species at risk | 4 |
| 3 | Me | thodology | 5 5 5 7 |
| | 3.1 | Database searches | 5 |
| | 3.2 | Site inspection and fauna habitat assessment | 5 |
| | 3.3 | Survey and reporting staff | 7 |
| | 3.4 | Taxonomy and nomenclature | 7 |
| | 3.5 | Limitations | 7 |
| 4 | Res | sults | 9 |
| | 4.1 | Fauna habitat | 9 |
| | 4.1 | .1 Feral pests | 10 |
| | 4.2 | Bioregional vertebrate fauna assemblage | 11 |
| | 4.3 | Conservation significant fauna | 13 |
| 5 | Dis | cussion | 20 |
| | 5.1 | Adequacy of the fauna survey data for fauna habitats represented in the project area | 20 |
| | 5.1 | .1 Amphibians | 20 |
| | 5.1 | .2 Reptiles | 20 |
| | 5.1 | .3 Birds | 20 |
| | 5.1 | .4 Mammals | 21 |
| 6 | Sur | nmary | 22 |
| 7 | Ref | ferences | 23 |



Chart

1. Climate averages for Lancelin

Plates

- 1. Coastal low heath on sand and karst limestone
- 2. Coastal low heath on sand and karst limestone
- 3. Mixed open and closed shrubland and heath on sand and limestone
- 4. Mixed open and closed shrubland and heath on sand and limestone
- 5. Highly disturbed, cleared habitat or planted vegetation
- 6. Highly disturbed, cleared habitat or planted vegetation
- 7. Replanted vegetation
- 8. Replanted vegetation
- 9. Anthropogenic disturbance
- 10. Anthropogenic disturbance

Tables

- 1. Fauna survey limitations and constraints
- 2. Habitat types
- 3. Birds potentially found near of the project area
- 4. Amphibians potentially found near of the project area
- 5. Mammals potentially found near the project area
- 6. Reptiles potentially found near the project area
- 7. Assessment of the potential presence of a conservation significant fauna species in the project area

Figures

- 1. Regional location
- 2. Habitat assessment locations

Appendices

- A. Results of the *EPBC Act* national protected matters search
- B. Vertebrate fauna recorded in biological surveys in the region
- C. Definitions of Significant Fauna under the WA *Biodiversity Conservation Act 2016* and Priority Species
- D. Fauna Habitat Assessment Results



EXECUTIVE SUMMARY

The City of Wanneroo requested a Level 1 vertebrate fauna survey for an area of 33ha, comprising of Newman Park 'A' Reserve foreshore area, and the adjacent land surrounded by Brazier Road, Capricorn Esplanade and Wilkie Avenue. This area includes the bushland on the western end of Yanchep Beach Rd and includes bituminised roads, car parks, built infrastructure and access tracks to the beach.

This Level 1 vertebrate fauna survey included a review of the available literature to determine vertebrate fauna species potentially in the project area and a site visit to determine major fauna habitats and their condition as the basis for an impact assessment which is contained in a separate report.

The assessed area included three broad fauna habitats in the project area:

- coastal low heath on sand and karst limestone;
- mixed open and closed shrubland and heath on sand and limestone; and
- highly disturbed, cleared habitat or planted vegetation.

Some of the site is highly disturbed, cleared, contains buildings and bituminised car parks and provides no habitat value for vertebrate fauna. Some parts of the project area have a relatively steep slope and there are sections with replanted vegetation.

It is probable that Quenda (Priority 4) and Black-striped Snake (Priority 3) are present in the project area. The Osprey (Migratory) will be regularly seen flying over the site but there is a very low probability that the Peregrine Falcon would be seen near project area. There was evidence of rabbits, cats and foxes in the project area.



1 INTRODUCTION

1.1 Background

The City of Wanneroo requested a Level 1 vertebrate fauna survey for an area of 33ha, comprising of Newman Park 'A' Reserve foreshore area, and the adjacent land surrounded by Brazier Road, Capricorn Esplanade and Wilkie Avenue. This area includes the bushland on the western end of Yanchep Beach Rd and includes bituminised roads, car parks, built infrastructure and access tracks to the beach (Figure 1).

The City of Wanneroo requested a Level 1 vertebrate fauna assessment as part of investigating opportunities to enhance the Yanchep Lagoon Precinct through the implementation of community events, infrastructure and development opportunities.

1.2 Project objectives and scope of works

Terrestrial Ecosystems was commissioned by the City of Wanneroo to undertake a Level 1 vertebrate fauna survey of the project area to support an environmental impact assessment which has been provided as a separate report (Terrestrial Ecosystems and One Tree Botanical 2020). The methodology broadly follows that described in the Environmental Protection Authority (EPA) *Technical Guidance Terrestrial Fauna Surveys* (EPA 2016b) and the *Technical Guidance - Sampling methods for terrestrial vertebrate fauna* (EPA 2016a).

A Level 1 fauna survey involves undertaking a desktop review and reconnaissance site visit. The objectives of this fauna survey were to:

- provide an indication of the vertebrate fauna assemblage (reptiles, amphibians, mammals and birds) in and near the project area, so that potential impacts on the fauna and fauna assemblage might be adequately assessed; and
- describe the major vertebrate fauna habitats present.

To achieve these objectives, Terrestrial Ecosystems:

- reviewed Terrestrial Ecosystems' database [includes Atlas of Living Australia and Department of Biodiversity, Conservation and Attractions (DBCA) records in NatureMap] to identify potential vertebrate fauna within the area;
- searched the DBCA's NatureMap for Threatened and Priority Species;
- searched the Commonwealth Governments database of fauna of national environmental significance to identify species potentially occurring within the area that are protected under the *Environment Protection and Biodiversity Conservation (EPBC) Act 1999* or international migratory bird agreements (JAMBA/CAMBA);
- undertook a site reconnaissance survey;
- reviewed previous fauna surveys conducted near the project area in similar habitat types; and
- discussed the likelihood of *EPBC Act 1999* and *Biodiversity Conservation (BC) Act 2016* listed species being present in the project area.

2 EXISTING ENVIRONMENT

2.1 Location of project area

The project area is within the Swan Coastal Plain 2 (SWA2) Interim Biogeographic Regionalisation of Australia (IBRA) subregion. This subregion is a low lying coastal plain, once vegetated by Banksia and Tuart on sandy soils, with *Casuarina obesa* on outwash plains and paperbark in swampy areas (Mitchell *et al.* 2002). The subregion is part of the West Botanical Province which has high species richness and diversity in flora and vegetation. The area is located within Bush Forever (Site 397) which has been identified as regionally significant bushland for protection by reservation or within a statutory planning framework (Department of Planning 2000).

2.2 Landforms and soils

The project area is in the Quindalup Dunes, which includes calcareous sands formed into parabolic dunes and beach ridge plains (Churchward and McArthur 1980, Gozzard 2007). These dunes are Holocene in age (McArthur and Bettenay 1974).

2.3 Land use history

The dominant land uses in the IBRA subregion are urban, rural residential, industrial, cultivation, forestry plantations, grazing and conservation areas. The greater Perth metropolitan area now extends almost from Mandurah to Alkimos, with towns further north at Yanchep and Two Rocks and east over the Darling Scarp. The project area has older established urban development on the southern boundary and more recent estates to the east and the north, with the coastal vegetation of the project area extending north almost to Two Rocks town site.

2.4 Climate

The project area is characterised as warm Mediterranean (Mitchell *et al.* 2002). Lancelin, which is approximately 70km to the north, and similarly situated on the coast, has an annual rainfall of approximately 853mm, although this varies considerably from year-to-year. The highest mean maximum and minimum temperatures in Lancelin are in January to March (Bureau of Meteorology 2019). The lowest mean daily maximum and minimum temperatures occur in July (Chart 1). Rainfall predominantly occurs between May and August and winter rains result from low pressure cells moving in an easterly direction.




Chart 1. Climatic averages for Lancelin



2.5 Regional biological fauna context of project area

The frogs, reptiles, mammals and birds in the vicinity of the project area have been surveyed for other environmental assessments and research purposes and are therefore known. Fauna surveys and assessments undertaken in the vicinity of the project area that have been reviewed for this assessment include:

- Alan Tingay & Associates (1991) Response to Draft north-west corridor structure plan, Department of Planning & Urban Development (February 1991), Yanchep Structure Plan, Vertebrate Fauna Survey. Perth.
- Alan Tingay & Associates (1996) Alkimos Eglington Vertebrate Fauna Survey, October 1996. Perth.
- Alan Tingay & Associates (1998) Yanchep Sun City, Environmental Assessment for the Rezoning of lots 201 and 202 Breakwater Drive, Two Rocks to Rural Community. Perth.
- Alan Tingay & Associates (1999a) Pt Lot 2 Burns Beach, Vertebrate Fauna. Perth.
- Alan Tingay & Associates (1999b) Shire of Wanneroo, Town Planning Scheme No. 1, Amendment 787, Yanchep Two Rocks, Environmental Review. Perth.
- Alan Tingay & Associates (1999c) Shire of Wanneroo, Town Planning Scheme No. 1, Amendment 837 Yanchep / Two Rocks, Environmental Review. Perth.
- Alan Tingay & Associates (2002) Review of Two Rocks Yanchep Foreshore Management Plan. Perth.
- ATA Environmental (1991) Yanchep Structure Plan Vertebrate Fauna Survey, Unpublished report for Tokyu Corporation, Perth.
- ATA Environmental (2005). *Metropolitan Region Scheme Amendment 1029/33. Alkimos-Eglinton Flora, Vegetation and Fauna Baseline Information. Interpretation Report.* Perth.
- ATA Environmental (2007) *Vertebrate Fauna Assessment St Andrews Estate* (*Southern Precinct*), *Yanchep*, Unpublished report for Yanchep Sun City Pty Ltd, Perth.
- ATA Environmental (2008) Vertebrate Fauna Assessment Lot 3 Romeo Road, Alkimos, Unpublished report for Northern Corridor Developments Limited. Perth.
- ATA Environmental (2008) Vertebrate Fauna Assessment Lot 3 Romeo Road, Alkimos. Perth.
- Bamford Consulting Ecologists (1998) Report on a Vertebrate Survey at Burns Beach. Perth.
- Bamford Consulting Ecologists (2005) *Alkimos Proposed Wastewater Treatment Plant: Fauna Assessment*, Unpublished report for Water Corporation, Perth.
- Bamford Consulting Ecologists (2005) *Alkimos Proposed Wastewater Treatment Plant: Fauna Assessment*. Perth.
- Biota Environmental Sciences (2000) Lot 52 Burns Beach Road Fauna Survey. Perth.
- Department of Conservation and Land Management. (1993) Fauna studies in water supply Reserve 34537, adjacent to Neerabup National Park. Perth.
- ENV Australia (2006) Lots 1005 and 1006 Alkimos Fauna Habitat Assessment. Perth.
- GHD (2014a) Mitchell Freeway extension: Burns Beach Rd to Romeo Rd Level 2 Flora & Level 1 Fauna Assessment. Unpublished report for Main Roads Western Australia, Perth.
- GHD (2014b) *Neerabup Road Extension Level 2 Fauna Survey*, Unpublished report for Main Roads Western Australia. Perth.
- Gole, C.A. (2003) Bird Survey in selected Perth Metropolitan Reserves. A Joint Biodiversity Conservation Project between Birds Australia WA and Perth Biodiversity Project, Unpublished report Birds Australia and Perth Biodiversity Project, Perth.
- Harvey, M. S., Dell, J., How, R. A. and Waldock, J. M. (1997) *Ground Fauna of Bushland Remnants on the Ridge Hill Shelf and Pinjarra Plain Landforms, Perth*, Report to the Australian Heritage commission NEP Grant N95/49.
- Kitchener, D.J., Chapman, A. and Barron, G. (1978). Mammals of the Northern Swan Coastal Plain. *Faunal Studies of the Northern Swan Coastal Plain*. Unpublished report for the Western Australian Museum and Department of Conservation and Environment. Perth.
- Ninox Wildlife Consulting (1990) Eglinton Beach Resort an appraisal of the vertebrate fauna. Perth.
- Storr, G.M., Harold G. and Barron, G. (1978a) The amphibians and reptiles of the northern Swan Coastal Plain. *Faunal Studies of the Northern Swan Coastal Plain*. Western Australian Museum, unpublished report, Perth.
- Storr, G.M., Johnstone, R.E. and Harold, G. (1978b) Birds of the northern Swan Coastal Plain, Western Australia. *Faunal Studies of the Northern Swan Coastal Plain*. Western Australian Museum, Unpublished report. Perth.
- Terrestrial Ecosystems (2005) *Terrestrial Vertebrate Fauna Species Likely to be found in the Alkimos-Eglinton Area, with a comment on Significant Fauna Species and the Impacts of the Proposed Disturbance*. Unpublished report for ATA Environmental, Dilhorn House, 2 Bulwer St, Perth, WA, 6000. Perth.
- Terrestrial Ecosystems (2018) City of Wanneroo Black-Cockatoo Habitat Survey. Perth.



- Terrestrial Ecosystems (2020a) *Vertebrate Fauna Survey Two Rocks Beach Access, Two Rocks*. Unpublished report for the City of Wanneroo.
- Terrestrial Ecosystems (2020b) *Vertebrate Fauna Survey Lot 211 Quinns Road, Mindarie*. Unpublished report for the City of Wanneroo.
- Valentine, I.E., Wilson, B.A., Reaveley, A., Huang, N., Johnson, B. and Brown, P.R. (2009) Patterns of Ground-dwelling Vertebrate Biodiversity in the Gnangara Sustainability Strategy Study Area, Unpublished report for the Department of Environment and Conservation. Perth.
- Western Australian Museum (1978). Faunal Studies of the Northern Swan Coastal Plain. Western Australian Museum. Perth.

Data in the Atlas of Living Australia and Western Australian Museum has also been added to the information contained in Appendix B, and the compilation of the species lists for the project area.

Few of these reports contain survey data for vertebrate fauna on the dune system close to the coast where there are few substantive trees. Reports providing useful data included ATA Environmental (1991, 2007, 2008), Biota Environmental Sciences (2000), Valentine *et al.* (2009), GHD (2014b), Gole (2003) and Bamford Consulting Ecologists (2005). Data from these reports are provided in Appendix B along with data from the Western Australian Museum, NatureMap and Atlas of Living Australia. Some of the data from consultants' fauna surveys are deposited in government databases, so it is possible there is some duplication of data in Appendix B.

2.5.1 Fauna species at risk

Mitchell *et al.* (2002) reported multiple vertebrate fauna species at risk in the subregion. However, some of these species have not been recorded near the project area for many years (e.g. *Myrmecobius fasciatus, Pseudocheirus occidentalis, Setonix brachyurus*), although, species such as *Calyptorhynchus latirostris, Calyptorhynchus banksii naso, Isoodon fusciventer* and *Neelaps calonotos* are still present, and regularly encountered. There is a very low probability that the Peregrine Falcon would be seen in the bioregion.



3 METHODOLOGY

3.1 Database searches

A review of the *EPBC* list of protected species was undertaken to identify species of conservation interest to the Commonwealth Government. The search area was a linear shaped polygon along the coastal strip using the following coordinates 31.46962°S 115.56826°E, 31.46962°S 115.583 E, 31.64574°S 115.69925°E, 31.64574°S 115.68169°E (Appendix A). In addition, a desktop search of the Terrestrial Ecosystems' fauna survey database was used to develop an appreciation of the vertebrate fauna assemblages in relevant sections of the bioregion near the project area. The DBCA threatened and priority species database was searched via the records in NatureMap.

Other more general texts were also used to provide supplementary information on vertebrate fauna in the bioregion, including Tyler *et al.* (2000) for frogs; Storr *et al.* (1983, 1990, 1999, 2002) for reptiles; Johnstone and Storr (1998, 2004) for birds; and Van Dyck and Strahan (2008) for mammals.

Collectively these sources of information were used to create lists of species expected to utilise the project area and broader subregion. It should be noted that these lists will include species that have been recorded in the general region but are possibly vagrants and they will not generally be found in the project area due to a lack of suitable habitat (e.g. water and shore birds). Vagrants can be recorded almost anywhere. Many of the records are historical and the species is no longer present in the area (e.g. Malleefowl). Many of the bird, mammal, reptile and amphibian species have specific habitat requirements that may be present in the general area but not in the project area. Also, the ecology of many of these species is often not well understood and it can sometimes be difficult to indicate those species whose specific habitat requirements are not present in the project area. Therefore, many species will be included in the lists produced from database searches but will not be present in the actual project area.

There are errors in most databases, including NatureMap, Atlas of Living Australia and the WAM collection. These errors occur because of a misidentification of individuals, taxonomic name changes and incorrect coordinates being entered into the database. Terrestrial Ecosystems was unable to verify the primary records, so it has used the information provided. Obvious errors have been removed but readers should appreciate that species lists and fauna surveys reported in the appendices may include these errors.

3.2 Site inspection and fauna habitat assessment

A site visit was undertaken on 29 August 2019 to assess fauna habitat types and condition in the project area. This information included a description of the habitat structure, habitat condition, landform, soils and vegetation and time since last fire.

The fauna habitat assessment had two foci:

- assessing fauna habitat types and their condition; and
- assessing the possible presence of and recording evidence of conservation significant fauna.

Dr Scott Thompson, who undertook the site assessment, stopped at multiple locations within the project area and recorded a suite of data about the fauna habitat and its condition. This information included a description of the habitat structure, habitat condition, landform, soils and vegetation and time since last fire. The following data were assessed at each location as part of the habitat assessment:

| Observer's name | |
|--|---------------------------------|
| Coordinates of the location as UTM (GDA94) | |
| Fire history – options | |
| o > 5 years | |
| o 1-5 years | |
| o < 1 year | |
| Landform – options | |
| o Beach | Lower slope |
| Clay plain | Mid slope |
| o Cliff | o Ridge |
| Creek line | o River |
| o Dam | Rocky outcrop / breakaway |



| Drainage line | Salt lake |
|--------------------------------------|---------------------------------|
| o Dune crest | Sand dune |
| Dune slope | Sand plain |
| o Dune swale | Stony plain |
| Escarpment | o Swamp |
| o Flat | Undulating |
| o Gorge | Upper slope |
| o Gully | Wetland |
| o Intertidal / mangrove | Water hole |
| Lake / lake edge | |
| YY 1 * | |

Habitat quality – options

- o *High quality fauna habitat* These areas closely approximate the vegetation mix and quality that would have been in the area prior to any disturbance. The habitat has connectivity with other habitats and is likely to contain the most natural vertebrate fauna assemblage.
- Very good fauna habitat These areas show minimal signs of disturbance (e.g. grazing, clearing, fragmentation, weeds) and generally retain many of the characteristics of the habitat if it had not been disturbed. The habitat has connectivity with other habitats and fauna assemblages in these areas are likely to be minimally affected by disturbance.
- Good fauna habitat These areas showed signs of disturbance (e.g. grazing, clearing, fragmentation, weeds) but generally retain many of the characteristics of the habitat if it had not been disturbed. The habitat has connectivity with other habitats and fauna assemblages in these areas are likely to be affected by disturbance.
- O Disturbed fauna habitat— These areas showed signs of significant disturbance. Many of the trees, shrubs and undergrowth are cleared. These areas may be in the early succession and regeneration stages. Areas may show signs of significant grazing, containing weeds or have been damaged by vehicle or machinery. Habitats are fragmented or have limited connectivity with other fauna habitats. Fauna assemblages in these areas are likely to differ significantly from what might be expected in the area had the disturbance not occurred.
- Highly degraded fauna habitat These areas often have a significant loss of vegetation, an abundance of weeds, and a large number of vehicle tracks or are completely cleared. Limited or no fauna habitat connectivity. Fauna assemblages in these areas are likely to be significantly different to what might have been in the area pre-disturbance.

| Habitat | Habitat structure – combined into habitat description | | | |
|---------|---|---|---|--|
| | Upper stratum | | | |
| 0 | Tall open woodland | 0 | Scattered tall trees | |
| 0 | Tall woodland | 0 | Scattered trees | |
| 0 | Open woodland | 0 | Scattered low trees | |
| 0 | Woodland | 0 | Low closed forest | |
| 0 | Open forest | 0 | Low open forest | |
| 0 | Closed forest | 0 | Low woodland | |
| 0 | Tall closed forest | 0 | Low open woodland | |
| 0 | Tall open forest | | | |
| Middle | e stratum | | | |
| 0 | Shrubland | 0 | Open heath | |
| 0 | Tall shrubland | 0 | Low closed heath | |
| 0 | Tall open shrubland | 0 | Low open heath | |
| 0 | Low shrubland | 0 | Tall closed scrub | |
| 0 | Scattered low shrubs | 0 | Tall open scrub | |
| 0 | Low open shrubland | 0 | Scattered tall shrubs | |
| 0 | Scattered tall shrubs | 0 | Open shrubland | |
| 0 | Closed heath | 0 | Scattered shrubs | |
| Lower | stratum | | | |
| 0 | Closed hummock grassland | 0 | Closed tussock grassland / sedgeland / herbland | |
| 0 | Mid-dense hummock grassland | 0 | Tussock grass land / sedgeland / herbland | |
| 0 | Hummock grassland | 0 | Open tussock grassland / sedgeland / herbland | |
| 0 | Open hummock grassland | 0 | Scattered tussock / grasses / sedges / herbs | |
| 0 | Scattered hummock grassland | 0 | Very open tussock grassland / herbland | |
| Soil Ty | Soil Type – options | | | |
| 0 | Sand | 0 | Silty loam | |



| 0 | Loamy sand | 0 | Sand clay loam |
|--------------------------|------------------|---|-------------------|
| 0 | Clayey sand | 0 | Clay |
| 0 | Clay loam | 0 | Peat / organic |
| 0 | Silty clay loam | 0 | Stony |
| 0 | Sandy loam | | |
| Soil co | lour - options | | |
| 0 | Black | 0 | Red |
| 0 | Brown | 0 | White |
| 0 | Grey | 0 | Yellow |
| 0 | Orange | | |
| Surface stones – options | | | |
| 0 | None | 0 | Boulders (>250mm) |
| 0 | Pebbles (0-50mm) | 0 | Rocks |
| 0 | Cobbles (51-250) | | |

3.3 Survey and reporting staff

Dr Scott Thompson undertook the site investigation and fauna habitat assessment. Drs Graham and Scott Thompson prepared the report and Dr Scott Thompson reviewed the report before it was sent to the client. Both senior scientists have appropriate relevant post-graduate qualifications, extensive experience in conducting fauna assessments on the Swan Coastal Plain, have published research articles on biodiversity, fauna assemblages, conservation significant species, trapping techniques and temporal variations in trapped fauna assemblages and are therefore appropriately trained and experienced for the task of preparing this assessment.

3.4 Taxonomy and nomenclature

Taxonomy and nomenclature for fauna species used in this report are generally based on the WA Museum species list except for bats, which follow Churchill (2008) and birds which follow Christidis and Boles (2008). Terrestrial Ecosystems' has presumed that the identifications referred to in the appendices or in reports used to provide local and regional comparative data are correct and we have only corrected obvious records where the nomenclature was known to be incorrect.

3.5 Limitations

This Level 1 fauna risk assessment is based on information contained in the Commonwealth Government database and other published and unpublished fauna survey data for the bioregion and a site visit. It is acknowledged that multiple surveys conducted in different seasons, repeated over several years are necessary to fully appreciate the fauna assemblage in the project area.

The EPA's (2016a) *Technical Guidance Terrestrial Fauna Surveys* suggested that fauna surveys may be limited by many variables. Limitations associated with each of these variables are assessed in Table 1.



Table 1. Fauna survey limitations and constraints

| Table 1. Fauna survey limitations and constraints | | | |
|--|--|--|--|
| Possible limitations | Constraint (yes/no); significant, moderate or negligible | Comment | |
| Competency and experience of the consultant carrying out this assessment | No | The environmental scientists that undertook the site assessment, drafted and reviewed this report are familiar with the vertebrate fauna of this bioregion. | |
| Scope | No | All aspects of the scope of works have been addressed. | |
| Proportion of fauna identified, recorded and/or collected | No | Not applicable. | |
| Accuracy of previous survey work | Yes, negligible | Terrestrial Ecosystems has reported fauna survey data recorded by various authors but is not able to vouch for the accuracy of much of this information. It is acknowledged that the taxonomy of Western Australian vertebrates is continually being revised and the nomenclature of some of the species listed in the appendices may have changed since publication by the authors. | |
| Sources of information | Yes, negligible | Vertebrate fauna information was available from on-line databases and unpublished and published reports of surveys conducted in the bioregion in a variety of habitat types. Many of these surveys employed a low level of trapping effort which significantly impacts on the capacity of these data to represent the fauna assemblages in the areas surveyed. | |
| Proportion of the task achieved | No | All tasks completed. | |
| Timing/weather/ season/ cycle | N/A | Weather was fine during the site visit. | |
| Disturbances which affected results of the survey | No | Disturbance areas throughout the project area have been factored into this assessment. | |
| Intensity of survey effort | N/A | | |
| Completeness | No | All aspects of this assessment have been completed. | |
| Resources | No | Adequate resources were available. | |
| Remoteness and/or access problems | No | All areas could be accessed. | |
| Availability of contextual information on the region | No | Fauna survey data are scant for the coastal dune habitats, and specifically fauna habitats accessed in the project area. | |



4 RESULTS

4.1 Fauna habitat

One hundred and twenty seven habitat assessments were completed in the project area (Figure 2). There are three broad fauna habitats in the project area (Table 2), excluding developed areas (e.g. roads, housing lots, etc). Some of the site is highly disturbed or cleared and provides no habitat value for vertebrate fauna and there were multiple areas that have been replanted. Some sections of the project area are on a steep slope.

Table 2. Habitat types

| Habitat category | Description | Area (ha) |
|--------------------------------------|--|-----------|
| Coastal low heath on sand and | Low coastal heath on unconsolidated sandy low primary | 3.58 |
| karst limestone | dunes. The quality of fauna habitat was variable. | |
| Mixed open and closed shrubland | Mixed open or closed shrubs and heath on taller | 2.74 |
| and heath on sand and limestone | unconsolidated sandy dunes. The quality of fauna habitat | |
| | was variable. | |
| Highly disturbed, cleared habitat or | These are generally area that have little vegetation, or the | 18.78 |
| planted vegetation | native vegetation has largely been removed and the area | |
| | supports weeds, or is a grassed area. | |
| Developed areas, roads, house lots, | | 7.90 |
| etc | | |

Plates 1-6 provide representative images of the fauna habitat types, Plates 7 and 8 show replanted vegetation and Plates 9 and 10 provide evidence of anthropogenic disturbance.



Plate 1. Coastal low heath on sand and karst limestone



Plate 2. Coastal low heath on sand and karst limestone



Plate 3. Mixed open and closed shrubland and heath on sand and limestone



Plate 4. Mixed open and closed shrubland and heath on sand and limestone







Plate 6. Highly disturbed, cleared habitat or planted vegetation



Plate 7. Replanted vegetation




Plate 8. Replanted vegetation



Plate 9. Anthropogenic disturbance



Plate 10. Anthropogenic disturbance

The condition of the fauna habitat varied from high quality, particularly in areas where the dense vegetation inhibits human access, to areas that are highly degraded. Highly degraded areas included beach access tracks, areas cleared of native vegetation, bituminised roads and sealed carparks.

4.1.1 Feral pests

There was evidence to indicate a high density of rabbits (e.g. scats and diggings), and cats (tracks) and foxes (tracks) in the project area.



4.2 Bioregional vertebrate fauna assemblage

Appendix B provides a summary of the fauna survey data that are available near the project area. There are appreciable differences in the recorded fauna assemblages within and among fauna surveys shown in Appendix B. These differences are partially due to the low survey effort deployed by some of the surveys and they also reflect variations in soils and vegetation as well as temporal variations in the fauna assemblages.

Tables 3-6 provide a list of vertebrate species potentially found near the project area that have been compiled based on the fauna survey report results shown in Appendix B.

Table 3. Birds potentially found near the project area

| Family | Species | Common Name | Family | Species | Common Name |
|-------------------|---------------------------------|---------------------------|-----------------|-------------------------------|-----------------------------|
| Accipitridae | Accipiter cirrocephalus | Collared Sparrowhawk | Campephagidae | Coracina novaehollandiae | Black-faced Cuckoo-shrike |
| | Accipiter fasciatus | Brown Goshawk | | Lalage tricolor | White-winged Triller |
| | Aquila audax | Wedge-tailed Eagle | Corvidae | Corvus coronoides | Australian Raven |
| | Circus approximans | Swamp Harrier | Hirundinidae | Cheramoeca leucosterna | White-backed Swallow |
| | Elanus axillaris | Black-shouldered Kite | | Hirundo neoxena | Welcome Swallow |
| | Haliastur sphenurus | Whistling Kite | | Petrochelidon nigricans | Tree Martin |
| | Hieraaetus morphnoides | Little Eagle | Maluridae | Malurus lamberti | Variegated Fairy-wren |
| | Lophoictinia isura | Square-tailed Kite | | Malurus leucopterus | White-winged Fairy-wren |
| Anatidae | Tadorna tadornoides | Australian Shelduck | | Malurus splendens | Splendid Fairy-wren |
| Podargidae | Podargus strigoides | Tawny Frogmouth | | Stipiturus malachurus | Southern Emu-wren |
| Casuariidae | Dromaius novaehollandiae | Emu | Megaluridae | Cincloramphus mathewsi | Rufous Songlark |
| Laridae | Chroicocephalus novaehollandiae | Silver Gull | | Acanthorhynchus superciliosus | Western Spinebill |
| Threskiornithidae | Threskiornis spinicollis | Straw-necked Ibis | | Anthochaera carunculata | Red Wattlebird |
| Columbidae | Columba livia | Domestic Pigeon | | Anthochaera chrysoptera | Little Wattlebird |
| | Ocyphaps lophotes | Crested Pigeon | | Anthochaera lunulata | Western Little Wattlebird |
| | Phaps chalcoptera | Common Bronzewing | | Epthianura albifrons | White-fronted Chat |
| | Phaps elegans | Brush Bronzewing | | Gavicalis virescens | Singing Honeyeater |
| | Spilopelia senegalensis | Laughing Turtle-dove | | Gliciphila melanops | Tawny-crowned Honeyeater |
| | Spilpopelia chinensis | Spotted Turtle-dove | | Lichenostomus ornatus | Yellow-plumed Honeyeater |
| Halcyonidae | Dacelo novaeguineae | Laughing Kookaburra | | Lichmera indistincta | Brown Honeyeater |
| | Todiramphus sanctus | Sacred Kingfisher | | Manorina flavigula | Yellow-throated Miner |
| Meropidae | Merops ornatus | Rainbow Bee-eater | | Melithreptus lunatus | White-naped Honeyeater |
| Cuculidae | Cacomantis flabelliformis | Fan-tailed Cuckoo | | Sugomel nigrum | Black Honeyeater |
| | Chalcites basalis | Horsfield's Bronze-cuckoo | | Phylidonyris niger | White-cheeked Honeyeater |
| | Chalcites lucidus | Shining Bronze-cuckoo | Meliphagidae | Phylidonyris novaehollandiae | New Holland Honeyeater |
| | Heteroscenes pallidus | Pallid Cuckoo | Monarchidae | Grallina cyanoleuca | Magpie-lark |
| Falconidae | Falco berigora | Brown Falcon | Motacillidae | Anthus novaeseelandiae | Australasian Pipit |
| Falconidae | Falco cenchroides | Nankeen Kestrel | Nectariniidae | Dicaeum hirundinaceum | Mistletoe Bird |
| | Falco longipennis | Australian Hobby | Pachycephalidae | Colluricincla harmonica | Grey Shrike-thrush |
| | Falco peregrinus | Peregrine Falcon | 7 1 | Pachycephala pectoralis | Golden Whistler |
| Phasianidae | Coturnix pectoralis | Stubble Quail | | Pachycephala rufiventris | Rufous Whistler |
| Otididae | Ardeotis australis | Australian Bustard | Pardalotidae | Pardalotus punctatus | Spotted Pardalote |
| Rallidae | Porzana tabuensis | Spotless Crake | | Pardalotus striatus | Striated Pardalote |
| Acanthizidae | Acanthiza apicalis | Inland Thornbill | Petroicidae | Eopsaltria georgiana | White-breasted Robin |
| | Acanthiza chrysorrhoa | Yellow-rumped Thornbill | | Petroica boodang | Scarlet Robin |
| | Acanthiza inornata | Western Thornbill | Rhipiduridae | Rhipidura albiscapa | Grey Fantail |
| | Gerygone fusca | Western Gerygone | 1 | Rhipidura leucophrys | Willie Wagtail |
| | Sericornis frontalis | White-browed Scrubwren | Timaliidae | Zosterops lateralis | Silvereye |
| | Smicrornis brevirostris | Weebill | Cacatuidae | Cacatua sanguinea | Little Corella |
| Acrocephalidae | Acrocephalus australis | Australian Reed-warbler | Cucumanduc | Calyptorhynchus banksii naso | Forest Red-tailed Cockatoo |
| Artamidae | Artamus cinereus | Black-faced Woodswallow | | Calyptorhynchus latirostris | Carnaby's Cockatoo |
| | Artamus cyanopterus | Dusky Woodswallow | | Eolophus roseicapilla | Galah |
| | Artamus personatus | Masked Woodswallow | Psittacidae | Barnardius zonarius | Australian Ringneck |
| | Cracticus nigrogularis | Pied Butcherbird | - Sittle Falle | Neophema elegans | Elegant Parrot |
| | Cracticus torquatus | Grey Butcherbird | | Trichoglossus haematodus | Rainbow Lorikeet |
| | Gymnorhina tibicen | Australian Magpie | Strigidae | Ninox boobook | Southern Boobook |



Table 4. Amphibians potentially found near the project area

| Family | Species | Common Name |
|-----------------|------------------------|--------------------|
| Hylidae | Litoria moorei | Motorbike Frog |
| Limnodynastidae | Heleioporus eyrei | Moaning Frog |
| | Limnodynastes dorsalis | Western Banjo Frog |

| Family | Species | Common Name |
|----------------|------------------------|--------------------|
| Myobatrachidae | Crinia insignifera | Squelching Froglet |
| | Myobatrachus gouldii | Turtle Frog |
| | Pseudophryne guentheri | Gunther's Toadlet |

Table 5. Mammals potentially found near the project area

| Family | Species | Common Name |
|--------------|-------------------------|-----------------------|
| Canidae | Vulpes vulpes | Red Fox |
| Felidae | Felis catus | Cat |
| Dasyuridae | Sminthopsis fuliginosus | Grey-bellied Dunnart |
| Macropodidae | Macropus fuliginosus | Western Grey Kangaroo |
| | Notamacropus irma | Western Brush Wallaby |
| Tarsipedidae | Tarsipes rostratus | Honey Possum |

| Family | Species | Common Name |
|------------------|-----------------------|---------------------|
| Leporidae | Oryctolagus cuniculus | Rabbit |
| Peramelidae | Isoodon fusciventer | Quenda |
| Muridae | Mus musculus | House Mouse |
| | Rattus fuscipes | Bush Rat |
| | Rattus rattus | Black Rat |
| Vespertilionidae | Vespadelus regulus | Southern Forest Bat |

Table 6. Reptiles potentially found near the project area

| Family | Species | Common Name |
|-----------------|------------------------------|-------------------------------------|
| Agamidae | Ctenophorus adelaidensis | Western Heath Dragon |
| | Pogona minor | Dwarf Bearded Dragon |
| Diplodactylidae | Crenadactylus ocellatus | Clawless Gecko |
| | Diplodactylus polyophthalmus | Speckled Stone Gecko |
| | Strophurus elderi | Jewelled Gecko |
| | Strophurus spinigerus | South-western Spiny-tailed Gecko |
| Elapidae | Brachyurophis fasciolata | Narrow-banded Burrowing Snake |
| | Brachyurophis semifasciata | Half-girdled Snake |
| | Demansia psammophis | Yellow-faced Whipsnake |
| | Echiopsis curta | Bardick |
| | Neelaps bimaculatus | Black-naped Burrowing Snake |
| | Neelaps calonotus | Black-striped Burrowing Snake |
| | Parasuta gouldii | Gould's Snake |
| | Pseudonaja affinis | Dugite |
| | Pseudonaja mengdeni | Western Brown Snake |
| | Simoselaps bertholdi | Jan's Banded Snake |
| | Simoselaps littoralis | West Coast Banded Snake |
| Gekkonidae | Christinus marmoratus | Marbled Gecko |
| Pygopodidae | Aprasia repens | Southwest Sandplain Worm Lizard |
| | Delma concinna | Javelin Lizard |
| | Delma fraseri | Fraser's Delma |
| | Delma grayii | Side-barred Delma |
| | Lialis burtonis | Burton's Legless Lizard |
| | Pletholax gracilis | Keeled Legless Lizard |
| | Pygopus lepidopodus | Common Scaly-foot |
| Pythonidae | Morelia spilota | Carpet Python |
| Scincidae | Acritoscincus trilineatus | Western Three-lined Skink |
| | Cryptoblepharus buchananii | Buchanan's Snake-eyed Skink |
| | Ctenotus australis | Western Limestone Ctenotus |
| | Ctenotus fallens | West-coast Laterite Ctenotus |
| | Cyclodomorphus celatus | Western Slender Bluetongue |
| | Egernia kingii | King's Skink |
| | Egernia napoleonis | Southwestern Crevice Skink |
| | Hemiergis quadrilineatum | Two-toed Earless Skink |

| Family | Species | Common Name |
|-------------|-------------------------|---------------------------------------|
| | Lerista distinguenda | South-western Orange-tailed Slider |
| | Lerista elegans | West Coast Four-toed Lerista |
| | Lerista lineopunctulata | Dotted-line Robust Slider |
| | Lerista praepedita | Blunt-tailed West-coast Slider |
| | Menetia greyii | Common Dwarf Skink |
| | Morethia lineoocellata | Pale-flecked Morethia |
| | Morethia obscura | Shrubland Pale-flecked Morethia |
| | Tiliqua occipitalis | Western Blue-tongued Lizard |
| | Tiliqua rugosa | Bobtail |
| Typhlopidae | Anilios australis | Austral Blind Snake |
| | Anilios pinguis | Rotund Blind Snake |
| Varanidae | Varanus gouldii | Gould's Goanna |
| | Varanus tristis | Black-headed Monitor |



These lists include species commonly found in Banksia and Tuart woodlands on the inland side of the coastal dunes, so there are multiple species shown in these lists that are unlikely to be recorded in the coastal dunes, although they may infrequently be recorded as vagrants, particularly for the avian species.

4.3 Conservation significant fauna

Conservation significant fauna are protected by the Commonwealth *EPBC Act 1999*, and this list includes species covered by international treaties such as the Japan-Australia Migratory Bird Agreement (JAMBA) and China-Australia Migratory Bird Agreement (CAMBA) and the Western Australia (WA) *BC Act 2016*. The *BC Act 2016* provides for the publishing of the *Wildlife Conservation (Specially Protected Fauna) Notice* that lists species under multiple categories. In addition, DBCA maintains a list of fauna that require monitoring under four priorities based on the current knowledge of their distribution, abundance and threatening processes. The *EPBC Act 1999* and *BC Act 2016* imply legislative requirements for the management of anthropogenic impacts to minimise the effects of disturbances on species and their habitats. Priority species have no statutory protection, other than the DBCA wishes to monitor potential impacts on these species. Environmental consultants and proponents of developments are encouraged to avoid and minimise impacts on these species. Definitions of the significant fauna under the *BC Act 2016* are provided in Appendix C.

The fauna species that have special status in either State or Commonwealth government legislation or are on the DBCA Priority species list and are potentially present in the vicinity of the project area are listed in Table 7. Although they were recorded in the search of the MNES online database, migratory species that typically would be found around the edge of salt lakes, clay pans, estuaries and marshes have been excluded from Table 7 as there is no suitable habitat nearby.

Threatened and conservation significant waders and shorebirds that utilise the beaches along the edge of the ocean or are marine migratory species, or marine turtles that were identified in the MNES online search have not been included in this assessment as the project area does not include habitat in which they will forage or nest.

Two threatened species of fauna and two migratory species of birds were identified under the *EPBC Act 1999* as potentially occurring in the project area or surrounds. There is one Schedule 7 species as listed under the *BC Act 2016*, and two species listed on the DBCA's Threatened and Priority Fauna List that potentially occur in the project area or surrounds. The following is an assessment of the likelihood of each of the species listed in Table 7 being found in the project area.

Table 7. Assessment of the potential presence of a conservation significant fauna species in the project area

| Species | DBCA Schedule / Priority | Status under Commonwealth EPBC Act | Comment on the potential presence of a species |
|---|--------------------------------|--|--|
| Woylie | Critically | Endangered | Locally extinct from this area. |
| Bettongia penicillata | Endangered | | |
| Western Ringtail Possum | Critically | Critically | Locally extinct from this area. |
| Pseudocheirus occidentalis | Endangered | Endangered | |
| Australasian Bittern | Endangarad | Endangarad | Not present in the project area due to a lack |
| Botaurus poiciloptilus | Endangered | Endangered | of suitable habitat. |
| Carnaby's Black-Cockatoo Calyptorhynchus latirostris | Endangered | Endangered | Flies over the project area, but a lack of suitable feeding, roosting and nesting resources means any visits will be infrequent. |
| Forest Red-tailed Black-Cockatoo Calyptorhynchus banksii naso | Vulnerable | Vulnerable | Flies over the project area, but a lack of suitable feeding, roosting and nesting resources means any visits will be infrequent. |



| Species | DBCA Schedule / Priority | Status under Commonwealth EPBC Act | Comment on the potential presence of a species |
|---|--------------------------------|--|--|
| Malleefowl Leipoa ocellata | Vulnerable | Vulnerable | Locally extinct from this area. |
| Chuditch Dasyurus geoffroii | Vulnerable | Vulnerable | Locally extinct from this area. |
| Balston's Pygmy Perch Nannatherina balstoni | | Vulnerable | Not present in the project area due to a lack of suitable habitat. |
| Lancelin Island Skink Ctenotus lancelini | Vulnerable | Vulnerable | Highly unlikely to be in the project area. |
| Fork-tailed Swift Apus pacificus | Migratory | Migratory | May infrequently be seen flying in the region. |
| Grey Wagtail Motacilla cinerea | Migratory | Migratory | Highly unlikely to be seen in the project area. |
| Osprey Pandion haliaetus | Migratory | Migratory | Regularly seen flying over the project area but there are no roosting trees, so it is unlikely to roost in the project area. |
| Quenda <i>Isoodon fusciventer</i> | P4 | | Potentially in the project area. |
| Black-striped Snake Neelaps calonotos | P4 | | Potentially in the project area. |
| Peregrine Falcon Falco peregrinus | OS | | May very infrequently be seen in the project area. |

Results of the Commonwealth EPBC Act 1999 protected matters database search are provided in Appendix A.

Woylie (*Bettongia penicillata*) – Critically endangered under the *BC Act* 2016 and endangered under the *EPBC Act* 1999

The Brush-tailed Bettong or Woylie is a small (1-1.6kg) mammal that has a preference for open forests and woodlands, with clumped low understorey of tussock grasses or clumped low woody scrub (Christensen 2000). Woinarski *et al.* (2014) reported a population reduction of greater than 90% in the last 10 years.

It has not been recorded near the project area for many years, so it is Terrestrial Ecosystems' view that it is not present in the project area.

Western Ringtail Possum (*Pseudocheirus occidentalis*) – Critically endangered under the *BC Act* 2016 and *EPBC Act* 1999

The Western Ringtail Possum is an arboreal mammal with a body weight between 820–1020g. This species is regularly encountered in urban development and disturbed areas throughout its distribution which has contracted from what appears to have been a patchy distribution covering the south west of Western Australia from south-east of Geraldton to the Nullarbor with the most inland recordings coming from the Tuatanning Nature Reserve (de Tores *et al.* 1995). Its distribution encompassed a variety of vegetation types including coastal Peppermint (*Agonis flexuosa*), and Peppermint/Tuart (*Eucalyptus gomphocephala*) associations, Eucalypt and Casuarina (*Allocasuarina huegeliana*) woodlands, and mallee heath from the Hampton Tableland (Baynes 1987). It is now almost exclusively restricted to the coastal Peppermint woodland and coastal Peppermint/Tuart associations from the Australind-Eaton area to Two Peoples Bay. The only known natural extant inland populations are in the lower Collie River Valley, Perup Nature Reserve, around Albany and surrounding forest block near Manjimup. DBCA has translocated individuals to Yalgorup National Park and there are isolated populations south of Mandurah.

Factors thought to have contributed to this species decline include habitat loss, modification or fragmentation, changing fire regimes, disease, competition and predation by introduced predators (Clarke *et al.* 2008, Department of Environment Water Heritage and the Arts 2008).



It has not been recorded near the project area for many years, so it is Terrestrial Ecosystems' view that it is not present in the project area.

Australasian Bittern (Botaurus poiciloptilus) Endangered under the BC Act 2016 and EPBC Act 1999

The Australasian Bittern has a distribution from Moora through much of the south-west and east to Mt Arid; however, it is rarely recorded. It is almost always found in dense *Typha*, *Baumea* and sedges in freshwater or brackish swamps (Johnstone and Storr 1998). Garnett *et al.* (2011) reported its population across Australia as less than 2,000 and in decline. Most of the Western Australian records come from Lake Muir.

It has not been recorded in the vicinity of the project area in other fauna surveys. It is highly unlikely to be present in or near the project are due to a lack of semi-permanent water on the very sandy soils.

Carnaby's Black-Cockatoo (Calyptorhynchus latirostris) - Vulnerable under the BC Act 2016 and EPBC Act 1999

Carnaby's Black-Cockatoo (*Calyptorhynchus latirostris*) is a large, pied, cockatoo. Garnett *et al.* (2011) and the DSEWPaC (2011) reported that Carnaby's Black-Cockatoo inhabits the south-west of Western Australia, from Kalbarri to as east on the south coast as Esperance. It breeds inland and moves to the coastal areas when chicks have fledged (Saunders *et al.* 1985). Carnaby's Black-Cockatoos are highly gregarious, usually seen in trios, small parties or large flocks (up to 5000 birds)(Perry 1948). These flocks usually contain males, females and immature birds.

Carnaby's Black-Cockatoos are partly migratory and partly sedentary (Higgins 1999). In the drier regions of their geographic range where most of the native vegetation has been cleared (e.g. wheatbelt), Carnaby's Black-Cockatoos are postnuptial migrants (Saunders 1980, Saunders and Ingram 1995). After breeding, individuals in these areas migrate to feed in higher rainfall areas including the Swan Coastal Plain, and to a lesser extent, forests dominated by *E. marginata* (Jarrah), *C. calophylla* (Marri) and *E. diversicolor* (Karri; Saunders 1980). On the Swan Coastal Plain, Carnaby's Black-Cockatoos have been recorded foraging in most suburbs and in pine plantations within the greater Perth metropolitan area (Perry 1948). Vagrants have been recorded on Rottnest Island (Winnett 1989) and Garden Island (Wykes *et al.* 1999). These later two sightings clearly indicate that Carnaby's Black-Cockatoo will fly considerable distances over non-vegetated areas to forage.

Garnett et al. (2011) estimated there were between 10,000 and 60,000 birds in the population.

Saunders (1980) recorded non-breeding cockatoos at Coomallo Creek foraging within a 50km radius of their breeding area, whereas, cockatoos at Manmanning moved a much greater distance to the coastal plain during their non-breeding season. These data suggest that Carnaby's Black-Cockatoo move from areas where there is little food to southern and western coastal areas where food is presumably more plentiful during summer and autumn (Davies 1966, Saunders 1980).

Carnaby's Black-Cockatoo breed between July and November mostly in eucalypt woodland (Saunders 1980, 1986). Carnaby's Black-Cockatoo nest in tree hollows that are created by fire, fungi, termites or old age, with hollows between 2.5 and 12m above the ground (Saunders 1979, Higgins 1999). Hollows are large, ranging from 10 to over 250cm in depth (Higgins 1999). These hollows are usually in live or dead smooth-barked *Eucalyptus salmonophloia* (Salmon Gum) or *Eucalyptus wandoo* (Wandoo). However, Carnaby's Black-Cockatoo will also nest in *E. longicornis* (Red Morrell), *E. loxophleba* (York Gum), *E. gomphocephala* (Tuart), *E. rudis* (Flooded Gum), *E. salubris* (Gimlet), *E. occidentalis* (Swamp Yate) and *C. calophylla* (Higgins 1999, Cale 2003). When breeding, they most often forage in the surrounding shrubland and kwongan heath (Higgins 1999). On the Swan Coastal Plain, breeding could occur in *E. gomphocephala*, *E. rudis*, *E. occidentalis* and *C. calophylla*. Adults return to the same breeding area each year (Saunders 1977) and some use the same tree hollow for many years in succession to raise their chicks, others shift their nests among a number of trees in the same area (Saunders and Ingram 1998).

At Coomallo Creek, Carnaby's Black-Cockatoo travelled on average 1.4km from their nests to forage, whereas at Manmanning they foraged more widely and travelled an average of 2.5km from their nest to forage (Saunders 1980). At Manmanning, road and railway reserves were extensively used for foraging, presumably as this was the closest



food source to their nests. The availability of food near the nest influenced the time spent incubating eggs and fledging body mass (Saunders 1980). At Manmanning, Carnaby's Black-Cockatoo traversed agricultural land to forage in remnant plots of uncleared land.

Saunders (1980) reported Carnaby's Black-Cockatoo at Coomallo Creek (breeding area) foraged mostly on native plants, with the only exception being *Erodium* sp.. Higgins (1999) reported the habitat of Carnaby's Black-Cockatoo was uncleared or remnant woodlands dominated by *Eucalyptus*, particularly *E. wandoo* and *E. salmonophloia* and often in shrubland or kwongan heathland dominated by *Hakea*, *Dryandra*, *Banksia* and *Grevillea* and seasonally in *Pinus* plantations and less often in *C. calophylla*, *E. diversicolor* or *E. marginata*.

Carnaby's Black-Cockatoo is unlikely to forage in the project area due to a lack of suitable feeding resources and the lack of tall trees means it will not roost or nest in this area.

Forest Red-tailed Black-Cockatoo (Calyptorhynchus banksii naso) - Vulnerable under the BC Act 2016 and EPBC Act 1999

The Forest Red-tailed Black-Cockatoo is one of three large black-cockatoos found in Western Australia. *Calyptorhynchus banksii naso* frequents the humid to sub-humid south-west of Western Australia from Gingin in the north, to Albany in the south and west to Cape Leeuwin and Bunbury (Department of Sustainability Environment Water Population and Communities 2011). It was mostly seen in the hills, but small numbers of birds were seen at Mundijong, Baldivis, Karnup, Stakehill, Pinjarra, Coolup and in the Lake Clifton area (Johnstone *et al.* 2011). In 2011, there was an increase in the number of Forest Red-tailed Black-Cockatoo on the coastal strip north from Rockingham to the northern metropolitan suburbs. The reason for the recent increase in abundance is unknown.

Forest Red-tailed Black-Cockatoo nest hollows have been recorded between 6.5 and 33m above the ground, with entrance sizes ranging from 10 x 12cm to 44 x 150cm and a depth of 0.3-8.2m (Johnstone *et al.* 2013a, b). Breeding occurs in all months, but peaks in April-June and August-October with an incubation period of 29-31 days. A female broods her hatchling for the first 3-10 days after hatching and then leaves the nest each day at dawn and returns to feed the chick at dusk. Hatchlings are fully feathered at about 48 days. The majority of nests are in Marri, but they have also been recorded in Jarrah, Blackbutt, Bullich and Wandoo. Nest sites are often clustered in an area.

Johnstone and Kirkby (2011) reported the Forest Red-tailed Black-Cockatoo to feed mostly on seeds from *C. calophylla*, *E. marginata*, but also on *Allocasuarina fraseriana* (Sheoak), *Persoonia longifolia* (Snottygobble), *Eucalyptus patens* (Blackbutt) and introduced species such as *M. azedarach* (Cape Lilac) and *Corymbia citriodora* (Lemon-scented Gum).

Loss of breeding habitat in the form of suitable hollows and adequate feeding resources in the vicinity of nesting hollows to enable adults to feed chicks is a primary threat. Abbott (1998) reported that trees within its known breeding distribution was not a factor in limiting breeding. He estimated there were about 15,000 birds and Garnett *et al.* (2011) thought about 10% of these birds bred each year. Competition for nesting hollows by other cockatoos, Wood Ducks, Galahs and feral Honey Bees appears to also be a significant threat (Garnett *et al.* 2011).

The Forest Red-tailed Black-Cockatoo is unlikely to forage in the project area due to a lack of suitable feeding resources and the lack of tall trees means it will not roost or nest in this area.

Malleefowl (*Leipoa ocellata*) – Vulnerable under the *BC Act* 2016 and *EPBC Act* 1999

Malleefowl are large, ground-dwelling birds that rarely fly unless alarmed or are perching for the night. Historically, Malleefowl have been found in mallee regions of southern Australia from approximately the 26th parallel of latitude southwards. Prior to vegetation clearing for agriculture, Malleefowl were abundant in the WA Wheatbelt. Vegetation clearing for agriculture also opened adjacent bushland to predators, and in the south-west of WA, Malleefowl often only persist in isolated remnant patches of native vegetation. Sheep and other herbivores (e.g. goats, kangaroos) grazing in remnant vegetation removes or thins the undergrowth, and they also compete with



Malleefowl for herbaceous foods and can cause changes to the structure and floristic diversity of foraging habitats (Benshemesh 2007).

Malleefowl and their eggs are vulnerable to predation by foxes, and newly hatched chicks are vulnerable to foxes, cats and raptors (Priddel and Wheeler 1990, 1997, Benshemesh and Burton 1999, Benshemesh 2007, Lewis and Hines 2014). Their abundance in the Goldfields is low and they are sparsely distributed, favouring those areas that are more densely vegetated. Malleefowl build distinctive nests that comprise a large mound of soil/rock covering a central core of leaf litter. These nest mounds range in diameter but can span more than five metres and may be up to one metre high. Malleefowl are generally monogamous and once breeding commences they pair for life. The presence of nest mounds provides an indication of the presence of Malleefowl in the area.

Malleefowl has not been observed in the bioregion for many decades and it is not present in or near the project area.

Chuditch (Dasyurus geoffroii) – Vulnerable under the BC Act 2016 and EPBC Act 1999.

The Chuditch is the largest extant carnivorous marsupial in WA. It is usually active from dusk to dawn. Formally known from over 70% of Australia, the Chuditch now has a patchy distribution throughout the Jarrah forest and mixed Karri/Marri/Jarrah forest of south-west WA and other isolated areas. Chuditch are solitary animals for most of their life and den in hollow logs, burrows, culverts, etc. and have also been recorded in tree hollows and rock cavities. Chuditch are opportunistic feeders, and forage primarily on the ground at night. Their diet can include other mammals, birds, lizards, bird and reptile eggs but the majority is a mixture of large invertebrates (e.g. spiders, scorpions and crickets).

Chuditch have not been recorded in or near the project area for many years, so it is highly unlikely to be present.

Balston's Pygmy Perch (Nannatherina balstoni) - Vulnerable under the BC Act 2016 and EPBC Act 1999.

Balston's Pygmy Perch is the rarest of all the endemic freshwater fishes of south-western Australia. It inhabits coastal streams, lakes, ponds and swamps, where the water is dark and acidic (pH as low as 3.0) and has a large seasonal fluctuation in temperature (11-30°C). It is often found in association with tall sedge thickets.

Balston's Pygmy Perch is not present in the project area due to a lack of permanent freshwater.

Lancelin Island Skink (Ctenotus lancelini) - Vulnerable under the BC Act 2016 and EPBC Act 1999.

The approved conservation advice (Anon 2008) for the Lancelin Island Skink indicates that this small grey-brown lizards with indistinct streaks running along its back from the neck to the base of the tail is known only from Lancelin Island off the Western Australian coast (Pearson and Jones 2000). It typically uses all vegetation types on the island, but favours slopes facing north to north-east; areas protected from the prevailing south-westerly winds.

Having not been found on the mainland, except for a single individual near Lancelin, it is highly unlikely that the Lancelin Island Skink would be recorded in the project area.

Fork-tailed Swift (Apus pacificus) - Migratory species under the EPBC Act 1999 and BC Act 2016

This species breeds in the northeast and mid-east Asia and winters in Australia and southern New Guinea. It is a visitor to most parts of Western Australia, beginning to arrive in the Kimberley in late September, in the Pilbara in November and in the southwest land division in mid-December, and leaving by late April. The Fork-tailed Swift is an almost exclusively aerial species, foraging and sleeping on the wing. It rarely comes to ground, usually only for breeding. It is common in the Kimberley, uncommon to moderately common near northwest, west and southeast coasts and rare to scarce elsewhere. It is rarely seen in the Goldfields.



Terrestrial Ecosystems' assessment is that the Fork-tailed Swift may very infrequently be seen flying over the project area, however, the Fork-tailed Swift is essentially an aerial species and would be highly unlikely to land in the project area.

Grey Wagtail (Motacilla cinerea) - Migratory species under the EPBC Act 1999 and BC Act 2016

The Grey Wagtail is a small yellow breasted bird with a grey back and head. Johnstone and Storr (2004) reported this migratory species as breeding in Palearctic from western Europe and north-west Africa to eastern Asia and wintering in Africa, south-east Asia, Indonesia, the Philippines, New Guinea and Australia. Its preferred habitat in Australia is banks and rocks in fast-running fresh water including rivers, streams and creeks where it feeds on insects. The Atlas of Living Australia records two sightings on the south-coast of Western Australia and none around the project area.

It is highly unlikely to be seen in the project area due to a lack of records and suitable habitat.

Osprey (Pandion haliaetus) – Migratory under the EPBC Act 1999

The Osprey is a large raptor that is mostly found in coastal areas, off-shore islands and the lower sections of rivers. It mainly feeds on fish, sea-snakes and large lizards. This species is a regular coastal visitor and likely to be recorded flying over the project area; however, due to a lack of trees is unlikely to roost on site.

The Osprey is seen along the coastal area searching for food in the shallow water. It nests on tall structures, rock outcrops and large trees, none of which are in the project area.

Quenda (Isoodon fusciventer) - Priority 4 species with the DBCA

Quenda prefer dense scrub (up to one metre high), with swampy vegetation but are found in a variety of other habitats. They will often feed in adjacent forest and woodland that is open grassland, pasture and crop land lying close to dense cover.

Quenda have been recorded as far north as Two Rocks in the DBCA threatened species database, and Dr Scott Thompson caught them near the old Club Capricorn Resort. It is possible that Quenda are present in very low densities in areas that provide suitable habitat. Given the abundance of foxes and cats along the coastal zone, Quenda are only going to survive in areas of dense undergrowth which provide some protection from these predators.

Black-striped Snake (Neelaps calonotus) - Priority 3 with DBCA

This species occurs on dunes and sand-plains vegetated with heaths and eucalypt/banksia woodlands. It feeds largely on skinks and its distribution is restricted and threatened by urban development. In its natural undisturbed state, the project area would provide habitat for the Black-striped Snake. The DBCA threatened species database has records of this snake around Mindarie, and the Atlas of Living Australia records one south of Lancelin, so it is feasible that they are in the project area.

Peregrine Falcon (Falco peregrinus) – Other specially protected fauna under the BC Act 2016

The Peregrine Falcon is uncommon, although widespread throughout much of Australia excluding the extremely dry areas and has a wide and patchy distribution. It favours hilly or mountainous country and open woodlands and may be an occasional visitor to the project area. Nesting sites include ledges along cliffs, granite outcrops and quarries, hollow trees near wetlands and old nests of other large bird species. There is no evidence to suggest any change in status in the last 50 years.



The Atlas of Living Australia contains records of this species around Joondalup and Lancelin, so it is possible that they are infrequently seen in the project area, however, the habitat in the project area is atypical for this species. The Peregrine Falcon will not rely on this site for continued survival in the region.



5 DISCUSSION

5.1 Adequacy of the fauna survey data for fauna habitats represented in the project area

The EPA's (2016a) *Technical Guidance on Terrestrial Fauna* indicated that a Level 2 fauna assessment is required for a disturbance area of in excess of 10ha in this bioregion. The project area is larger than 10ha, but it contains large areas of disturbed habitat, reducing the area of native vegetation and high quality habitat for native fauna.

Level 2 fauna surveys are typically not undertaken now on the Swan Coastal Plain as the EPA considers the vertebrate fauna in this IBRA subregion to be well known. Even if such a survey was undertaken, it is unlikely to provide new species, in particular a conservation significant species that have not previously been identified for this area that would alter the assessment of potential impacts, however, as with all surveys, until it is completed the outcome is unknown.

5.1.1 Amphibians

Amphibians typically found on the Swan Coastal Plain are listed in Table 4. The lack of permanent freshwater means that only those species able to survive away from permanent water on very porous sandy soil (e.g. potentially *Heleioporus eyrei*, *Limnodynastes dorsalis* and *Myobatrachus gouldii*) are likely to be present in the project area. Frogs in this area are normally only detected immediately after rainfall. All three species have a wide-spread distribution and are abundant. There are no conservation significant amphibians in the Swan Coastal IBRA subregion.

5.1.2 Reptiles

Reptile species richness in the project area will be comparable with other sandy dune habitats in the subregion. The list provided in Appendix A represents species likely to be found over a large area of diverse habitat types. Sandy dune habitats are likely to have a restricted reptile fauna assemblage and these areas would typically support 8-15 species of reptiles, but many of these would be in low abundance (see Table 6). The herpetofauna assemblage in the project area are likely to be similar to that in the adjacent areas.

The Black-striped Snake, which is listed as a Priority 4 species, is found in Banksia woodlands and sandy areas in the Perth region. There are records of this snake in the vicinity of the project area, so it is potentially present in the project area.

The Lancelin Island Skink is essentially only known from an off-shore island, and the single record on the mainland probably represents a translocation from the island. It is highly unlikely that this species is present in the project area.

5.1.3 Birds

Avian species richness on the Swan Coastal Plain is influenced by rainfall, urban disturbance and vegetation clearing, and in coastal areas the migratory shore birds and waders that may be seasonally present along the beaches. The list provided in Table 3 represents species likely to be found over a large area of diverse habitat types. Coastal dunes with limited variation in the vegetation structure and diverseness in habitats would typically support between 20-40 species of birds, but many of these would be in very low numbers (see Appendix A).

The Peregrine Falcon, which is a Schedule 7 species under the *BC Act*, will normally have a very large home range and is unlikely to be seen on the relatively flat coastal dunes.



5.1.4 Mammals

The diversity of small terrestrial mammals potentially caught in the project area would be low given the lack of diversity on the sandy coastal dunes. Although, records of Chuditch (*D. geoffroii*), Woylie (*B. penicillata*) and Western Ringtail Possums (*P. occidentalis*) are present in the MNES database search for the area (Appendix B), they are no longer present in the region, having been predated on by foxes and cats and lost due to vegetation clearing and urban development many years ago.

Quenda (*I. fusciventer*) maybe present on the sections of dense vegetation on the sandy dunes in the vicinity of the project area. This species is likely to be in low abundance, as it is predated on by foxes and feral cats which are present in the area.

It was noted during the site visit that there was an abundance of rabbits, and moderate density of cats and foxes utilising the project area.



6 SUMMARY

The City of Wanneroo has requested a Level 1 vertebrate fauna risk assessment as part of its investigation of opportunities to enhance the Yanchep Lagoon Precinct.

Terrestrial Ecosystems undertook a Level 1 vertebrate fauna survey of the 33ha project area, which included a review of the available literature to determine vertebrate fauna species potentially in the project area and a site visit to determine major fauna habitats and their condition as the basis for an impact assessment which is contained in a separate report (Terrestrial Ecosystems and One Tree Botanical 2020).

The total assessed area is 33ha and there are three broad fauna habitats in the project area:

- coastal low heath on sand and karst limestone;
- mixed open and closed shrubland and heath on sand and limestone; and
- highly disturbed, cleared habitat or planted vegetation.

Some of the site is highly disturbed, cleared, contains buildings and bituminised car parks and provides no habitat value for vertebrate fauna. Some parts of the project area have a relatively steep slope and there are sections with replanted vegetation.

It is probable that Quenda (Priority 4) and Black-striped Snake (Priority 3) are present in the project area. The Osprey (Migratory) will be regularly seen flying over the site but there is a very low probability that the Peregrine Falcon would be seen near project area. There was evidence of rabbits, cats and foxes in the project area.



7 REFERENCES

Abbott, I. 1998. Conservation of the forest red-tailed black cockatoo, a hollow-dependent species, in the eucalypt forests of Western Australia. Forest Ecology and Management **109**:175-185.

Alan Tingay & Associates. 1991. Response to Draft north-west corridor structure plan, Department of Planning & Urban Development (February 1991), Yanchep Structure Plan, Vertebrate Fauna Survey.

Alan Tingay & Associates. 1996. Alkimos - Eglinton Vertebrate Fauna Survey, October 1996. Perth.

Alan Tingay & Associates. 1998. Yanchep Sun City, Environmental Assessment for the Rezoning of lots 201 and 202 Breakwater Drive, Two Rocks to Rural Community. Perth.

Alan Tingay & Associates. 1999a. Pt Lot 2 Burns Beach, Vertebrate Fauna. Perth.

Alan Tingay & Associates. 1999b. Shire of Wanneroo, Town Planning Scheme No. 1, Amendment 787, Yanchep - Two Rocks, Environmental Review. Perth.

Alan Tingay & Associates. 1999c. Shire of Wanneroo, Town Planning Scheme No. 1, Amendment 837 - Yanchep / Two Rocks, Environmental Review. Perth.

Alan Tingay & Associates. 2002. Review of Two Rocks - Yanchep Foreshore Management Plan. Perth.

Anon. 2008, Approved Conservation Advice for Ctenotus lancelini (Lancelin Island Skink), Canberra.

ATA Environmental. 1991. Yanchep Structure Plan Vertebrate Fauna Survey. Unpublished report for Tokyu Corporation, Perth.

ATA Environmental. 2005. Metropolitan Region Scheme Amendment 1029/33. Alkimos-Eglinton Flora, Vegetation and Fauna Baseline Information. Interpretation Report.

ATA Environmental. 2007. Vertebrate Fauna Assessment St Andrews Estate (Southern Precinct), Yanchep.

ATA Environmental. 2008. Vertebrate Fauna Assessment Lot 3 Romeo Road, Alkimos.

Bamford Consulting Ecologists. 1998. Report on a Vertebrate Survey at Burns Beach. Perth.

Bamford Consulting Ecologists. 2005. Alkimos Proposed Wastewater Treatment Plant: Fauna Assessment. Perth.

Baynes, A. 1987. The original mammal fauna of the Nullarbor and southern peripheral regions: evidence from skeletal remains in superficial cave deposits. Pages 139-152 *in* A. C. Robinson and N. L. McKenzie, editors. A Biological Survey of the Nullarbor region of south and western Australia in 1984. Department of Environment and Planning, Adelaide.

Benshemesh, J. 2007. National Recovery Plan for Malleefowl. South Australia.

Benshemesh, J., and P. Burton. 1999. Fox predation on Malleefowl three years after the spread of RCD in Victoria. Unpublished report for Parks Victoria and Department of Natural Resources and Environment, Mildura.

Bureau of Meteorology. 2019. Lancelin - Climate Statistics for Australian locations. (http://www.bom.gov.au/climate/averages/tables/cw 009114.shtml

Biota Environmental Sciences. 2000. Lot 52 Burns Beach Road Fauna Survey. Perth.

Cale, B. 2003. Carnaby's Black-Cockatoo (*Calyptorhynchus latirostris*) Recovery Plan. Department of Conservation and Land Management, Perth.

Christensen, P. 2000. Brush-tailed Bettong. Pages 292-293 in R. Strahan, editor. Mammals of Australia. Reed, Sydney.

Christidis, L., and W. E. Boles. 2008. Systematics and Taxonomy of Australian Birds. CSIRO Publishing, Collingwood, Victoria.

Churchill, S. 2008. Australian Bats. Jacana Books, Crows Nest, NSW.

Churchward, H. M., and W. M. McArthur. 1980. Atlas of Natural Resources: Darling Systems, Western Australia. Clarke, J., K. Warren, I. Robertson, M. Calver, and P. de Tores. 2008. Health and survival of translocated western ringtail possums. Australian Wildlife Management Society 21st Annual Conference 24-27 November, Fremantle, Western Australia.

Davies, S. J. J. F. 1966. The movements of the White-tailed Black Cockatoo (*Calyptorhynchus baudinii*) in southwestern Australia. The Western Australian Naturalist **10**:33-42.

Department of Conservation and Land Management. 1993. Fauna studies in water supply Reserve 34537, adjacent to Neerabup National Park. Perth.

Department of Environment Water Heritage and the Arts. 2008. Background Paper to EPBC Act Policy Statement 3.10 - Nationally Threatened Species and Ecological Communities. Significant impact guidelines for the vulnerable western ringtail possum (*Pseudocheirus occidentalis*) in the southern Swan Coastal Plain, Western Australia., DEWHA, Canberra.

Department of Planning. 2000. Bush Forever Volume 1 Policies, Principles and Processes. Perth.

Department of Sustainability Environment Water Population and Communities. 2011. Environment Protection and Biodiversity Conservation Act 1999 draft referral guidelines for three threatened black cockatoo species: Carnaby's cockatoo (endangered) *Calyptorhynchus latirostris* Baudin's cockatoo (vulnerable) *Calyptorhynchus baudinii* Forest red-tailed black cockatoo (vulnerable) *Calyptorhynchus banksii naso*. Canberra.

ENV Australia. 2006. Lots 1005 and 1006 Alkimos Fauna Habitat Assessment. Perth.



- Environmental Protection Authority. 2016a. Technical Guidance Sampling methods for terrestrial vertebrate fauna. Environmental Protection Authority, Perth.
- Environmental Protection Authority. 2016b. Technical Guidance Terrestrial Fauna Surveys. Perth.
- Garnett, S. T., J. K. Szabo, and G. Dutson. 2011. The Action Plan for Australian Birds 2010. CSIRO, Collingwood, Melbourne.
- GHD. 2014a. Mitchell Freeway extension: Burns Beach Rd to Romeo Rd Level 2 Flora & Level 1 Fauna Assessment.
- GHD. 2014b. Neerabup Road Extension Level 2 Fauna Survey. Perth.
- Gole, C. A. 2003. Bird Survey in selected Perth Metropolitan Reserves. A Joint Biodiversity Conservation Project between Birds Australia WA and Perth Biodiversity Project. Perth.
- Gozzard, J. R. 2007. Geology and Landforms of the Perth Region. Western Australia Geological Survey, Perth.
- Harvey, M. S., J. Dell, R. A. How, and J. M. Waldock. 1997. Ground Fauna of Bushland Remnants on the Ridge Hill Shelf and Pinjarra Plain Landforms, Perth, Report to the Australian Heritage commission NEP Grant N95/49.
- Higgins, P. J. 1999. Handbook of Australian, New Zealand and Antarctic Birds Volume 4 Parrots to Dollardbird.
- Johnstone, R. E., C. Johnstone, and T. Kirkby. 2011. Carnaby's Cockatoo (*Calyptorhynchus latirostris*), Baudin's Cockatoo (*Calyptorhynchus baudinii*) and Forest Red-tail Black Cockatoo (*Calyptorhynchus banksii naso*) on the Swan Coastal Plain (Lancelin-Dunsborough), Western Australia. Studies on distribution, status, breeding, food, movement and historical changes. Perth.
- Johnstone, R. E., T. Kirkby, and K. Sarti. 2013a. The breeding biology of the Forest Red-tailed Black Cockatoo Calyptorhynchus banksii naso Gould in south-western Australia. I. Characteristics of nest trees and nest hollows. Pacific Conservation Biology **19**:121-142.
- Johnstone, R. E., T. Kirkby, and K. Sarti. 2013b. The breeding biology of the Forest Red-tailed Black Cockatoo Calyptorhynchus banksii naso Gould in south-western Australia. II. Breeding behaviour and diet. Pacific Conservation Biology 19:143-155.
- Johnstone, R. E., and G. M. Storr. 1998. Handbook of Western Australian Birds. Volume I Non-Passerines (Emu to Dollarbird). Western Australian Museum, Perth.
- Johnstone, R. E., and G. M. Storr. 2004. Handbook of Western Australian Birds, Volume II Passerines (Bluewinged Pitta to Goldfinch). Western Australian Museum, Perth.
- Kitchener, D. J., A. Chapman, and G. Barron. 1978. Mammals of the Northern Swan Coastal Plain. Faunal Studies of the Northern Swan Coastal Plain. Unpublished report for the Western Australian Museum and Department of Conservation and Environment., Perth.
- Lewis, M., and M. Hines. 2014. Malleefowl activity at nesting sites increase fox and other feral animal visitation rates. Pages 242-247 Proceedings of the 5th National Malleefowl Forum 2014.
- McArthur, W. M., and E. Bettenay. 1974. The Development and Distribution of the Soils on the Swan Coastal Plain, Western Australia. CSIRO, Melbourne.
- Mitchell, D., K. Williams, and A. Desmond. 2002. Swan Coastal Plain 2 (SWA2 Swan Coastal Plain subregion). Pages 606-623 A Biodiversity Audit of Western Australia's 53 Biogeographic Subregions in 2002. Department of Conservation and Land Management, Perth.
- Ninox Wildlife Consulting. 1990. Eglinton Beach Resort an appraisal of the vertebrate fauna.
- Pearson, D., and B. Jones. 2000. Lancelin Island Skink Recovery Plan. Department of Conservation and Land Management, Perth.
- Perry, D. H. 1948. Black Cockatoos and pine plantations. The Western Australian Naturalist 1:133-135.
- Priddel, D., and R. Wheeler. 1990. Survival of Malleefowl *Leipoa ocellata* chicks in the absence of ground-dwelling predators. Emu **90**:81-87.
- Priddel, D., and R. Wheeler. 1997. Efficacy of fox control in reducing the mortality of released captive-reared Malleefowl, *Leipoa ocellata*. Wildlife Research **24**:469-482.
- Saunders, D. A. 1979. The availability of tree hollows for use as nest sites by White-tailed Black Cockatoos. Australian Wildlife Research 6:205-216.
- Saunders, D. A. 1980. Food and movement of the Short-billed form of the White-tailed Black Cockatoo. Australian Wildlife Research 7:257-269.
- Saunders, D. A. 1986. Breeding season, nesting success and nestling growth in Carnaby's Cockatoo, *Calyptorhynchus funereus latirostris*, over 16 years at Coomallo Creek, and a method for assessing the viability of populations in other areas. Australian Wildlife Research **13**:261-273.
- Saunders, D. A., and J. A. Ingram. 1995. Birds of Southwestern Australia: An Atlas of Changes in the Distribution and Abundance of the Wheatbelt avifauna. Surrey Beatty, Sydney.
- Saunders, D. A., and J. A. Ingram. 1998. Twenty-eight years of monitoring a breeding population of Carnaby's Cockatoo. Pacific Conservation Biology **4**:261-270.
- Saunders, D. A., I. Rowley, and G. T. Smith. 1985. The effects of clearing for agriculture on the distribution of cockatoos in the southwest of Western Australia.in A. Keast, H. F. Recher, H. A. Ford, and D. A.



- Saunders, editors. Birds of Eucalypt Forest and Woodlands: Ecology, Conservation, Management. RAOU and Surrey Beatty and Sons, Melbourne and Chipping North.
- Storr, G. M., G. Harold, and G. Barron. 1978a. The amphibians and reptiles of the northern Swan Coastal Plain. Faunal Studies of the Northern Swan Coastal Plain. Western Australian Museum, unpublished report,
- Storr, G. M., R. E. Johnstone, and G. Harold. 1978b. Birds of the northern Swan Coastal Plain, Western Australia. Faunal Studies of the Northern Swan Coastal Plain. Western Australian Museum, Unpublished report, Perth
- Storr, G. M., L. A. Smith, and R. E. Johnstone. 1983. Lizards of Western Australia. II: Dragons and Monitors. Western Australian Museum, Perth, Western Australia.
- Storr, G. M., L. A. Smith, and R. E. Johnstone. 1990. Lizards of Western Australia. III: Geckos and Pygopods. Western Australian Museum, Perth.
- Storr, G. M., L. A. Smith, and R. E. Johnstone. 1999. Lizards of Western Australia. I: Skinks. Western Australian Museum, Perth.
- Storr, G. M., L. A. Smith, and R. E. Johnstone. 2002. Snakes of Western Australia. Western Australian Museum,
- Terrestrial Ecosystems. 2005. Terrestrial Vertebrate Fauna Species Likely to be found in the Alkimos-Eglinton Area, with a comment on Significant Fauna Species and the Impacts of the Proposed Disturbance. Unpublished report for ATA Environmental, Dilhorn House, 2 Bulwer St, Perth, WA, 6000., Perth.
- Terrestrial Ecosystems. 2018. City of Wanneroo Black-Cockatoo Habitat Survey. Perth.
- Terrestrial Ecosystems. 2020a. Vertebrate Fauna Survey Two Rocks Beach Access, Two Rocks. Perth. Terrestrial Ecosystems. 2020b. Vertebrate Fauna Survey Lot 211 Quinns Road, Mindarie. Perth.
- Terrestrial Ecosystems, and One Tree Botanical. 2020. Flora, Vegetation and Vertebrate Fauna Environmental Impact Assessment - Yanchep Lagoon, Yanchep. Perth.
- Tyler, M. J., L. A. Smith, and R. E. Johnstone. 2000. Frogs of Western Australia. Western Australian Museum, Perth.
- Valentine, I. E., B. A. Wilson, A. Reaveley, N. Huang, B. Johnson, and P. R. Brown. 2009. Patterns of Grounddwelling Vertebrate Biodiversity in the Gnangara Sustainability Strategy Study Area. Perth.
- Van Dyck, S., and R. Strahan. 2008. The Mammals of Australia. Reed New Holland, Sydney.
- Western Australian Museum. 1978. Faunal Studies of the Northern Swan Coastal Plain. Western Australian Museum, Perth.
- Winnett, S. 1989. White-tailed Black Cockatoos on Rottnest Island. The Western Australian Naturalist 18:64.
- Woinarski, J. C. Z., A. A. Burbidge, and P. L. Harrison. 2014. The Action Plan for Australian Mammals 2012. CSRIO Publishing, Melbourne.
- Wykes, B. J., D. Pearson, and J. Majer. 1999. Fauna Survey of Garden Island, WA, 1996-1997. HMAS Stirling, Environmental Working Paper No. 12, Perth.




Figures
Vertebrate Fauna Survey - Yanchep Lagoon, Yanchep





Appendix A Results of the *EPBC Act* Protected Matters Search

Vertebrate Fauna Survey - Yanchep Lagoon, Yanchep



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about <u>Environment Assessments</u> and the EPBC Act including significance guidelines, forms and application process details.

Report created: 31/12/19 14:00:39

Summary

Details

Matters of NES

Other Matters Protected by the EPBC Act

Extra Information


Caveat

<u>Acknowledgements</u>



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

Coordinates
Buffer: 1.0Km



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the <u>Administrative Guidelines on Significance</u>.

| World Heritage Properties: | None |
|---|------|
| National Heritage Places: | None |
| Wetlands of International Importance: | None |
| Great Barrier Reef Marine Park: | None |
| Commonwealth Marine Area: | None |
| Listed Threatened Ecological Communities: | 2 |
| Listed Threatened Species: | 40 |
| Listed Migratory Species: | 41 |

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at http://www.environment.gov.au/heritage

A <u>permit</u> may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

| Commonwealth Land: | 1 |
|------------------------------------|------|
| Commonwealth Heritage Places: | None |
| Listed Marine Species: | 65 |
| Whales and Other Cetaceans: | 12 |
| Critical Habitats: | None |
| Commonwealth Reserves Terrestrial: | None |
| Australian Marine Parks: | None |

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

| State and Territory Reserves: | None |
|----------------------------------|------|
| Regional Forest Agreements: | None |
| Invasive Species: | 34 |
| Nationally Important Wetlands: | None |
| Key Ecological Features (Marine) | None |

Details

Matters of National Environmental Significance

Listed Threatened Ecological Communities

| Listed Threatened Ecological Communities | | [Resource information] |
|---|--------------------------|--|
| For threatened ecological communities where the distriplans, State vegetation maps, remote sensing imagery community distributions are less well known, existing vegetation produce indicative distribution maps. | and other sources. Where | threatened ecological |
| Name | Status | Type of Presence |
| Banksia Woodlands of the Swan Coastal Plain ecological community | Endangered | Community likely to occur within area |
| Tuart (Eucalyptus gomphocephala) Woodlands and Forests of the Swan Coastal Plain ecological community | Critically Endangered | Community likely to occur within area |
| Listed Threatened Species | | [Resource Information] |
| Name | Status | Type of Presence |
| Birds | | • • |
| Anous tenuirostris melanops | | |
| Australian Lesser Noddy [26000] | Vulnerable | Species or species habitat may occur within area |
| Calidris canutus | | |
| Red Knot, Knot [855] | Endangered | Species or species habitat known to occur within area |
| <u>Calidris ferruginea</u> | | |
| Curlew Sandpiper [856] | Critically Endangered | Species or species habitat may occur within area |
| Calyptorhynchus banksii naso Forest Red-tailed Black-Cockatoo, Karrak [67034] | Vulnerable | Species or species habitat likely to occur within area |
| Calyptorhynchus latirostris | | |
| Carnaby's Cockatoo, Short-billed Black-Cockatoo [59523] | Endangered | Species or species habitat known to occur within area |
| Diomedea amsterdamensis | | |
| Amsterdam Albatross [64405] | Endangered | Species or species habitat may occur within area |
| Diomedea epomophora | | |
| Southern Royal Albatross [89221] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| <u>Diomedea exulans</u> | | |
| Wandering Albatross [89223] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| <u>Diomedea sanfordi</u> | | |
| Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Leipoa ocellata Malleefowl [934] | Vulnerable | Species or species habitat likely to occur within area |
| | | |

[Resource Information]

| Name | Status | Type of Presence |
|---|-----------------------|--|
| Limosa lapponica baueri Bar-tailed Godwit (baueri), Western Alaskan Bar-tailed Godwit [86380] | Vulnerable | Species or species habitat may occur within area |
| Limosa lapponica menzbieri Northern Siberian Bar-tailed Godwit, Bar-tailed Godwit (menzbieri) [86432] | Critically Endangered | Species or species habitat may occur within area |
| Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat may occur within area |
| Pachyptila turtur subantarctica Fairy Prion (southern) [64445] | Vulnerable | Species or species habitat known to occur within area |
| Phoebetria fusca Sooty Albatross [1075] | Vulnerable | Species or species habitat may occur within area |
| Rostratula australis Australian Painted Snipe [77037] | Endangered | Species or species habitat likely to occur within area |
| Sternula nereis nereis Australian Fairy Tern [82950] | Vulnerable | Foraging, feeding or related behaviour known to occur within area |
| Thalassarche cauta cauta Shy Albatross [82345] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche cauta steadi White-capped Albatross [82344] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within area |
| Insects | | |
| Hesperocolletes douglasi Douglas' Broad-headed Bee, Rottnest Bee [66734] | Critically Endangered | Species or species habitat may occur within area |
| Mammals | | |
| Balaenoptera musculus Blue Whale [36] | Endangered | Species or species habitat likely to occur within area |
| Dasyurus geoffroii Chuditch, Western Quoll [330] | Vulnerable | Species or species habitat likely to occur within area |
| Eubalaena australis Southern Right Whale [40] | Endangered | Breeding known to occur within area |
| Megaptera novaeangliae Humpback Whale [38] | Vulnerable | Species or species habitat known to occur within area |

| Name | Status | Type of Presence |
|---|---------------------------|--|
| Neophoca cinerea Australian Sea-lion, Australian Sea Lion [22] | Vulnerable | Species or species habitat known to occur within area |
| Plants | | |
| <u>Diuris micrantha</u> Dwarf Bee-orchid [55082] | Vulnerable | Species or species habitat likely to occur within area |
| <u>Drakaea elastica</u> Glossy-leafed Hammer Orchid, Glossy-leaved Hammer Orchid, Warty Hammer Orchid [16753] | Endangered | Species or species habitat likely to occur within area |
| Eucalyptus argutifolia Yanchep Mallee, Wabling Hill Mallee [24263] | Vulnerable | Species or species habitat likely to occur within area |
| Lepidosperma rostratum Beaked Lepidosperma [14152] | Endangered | Species or species habitat likely to occur within area |
| Reptiles | | |
| Caretta caretta Loggerhead Turtle [1763] | Endangered | Species or species habitat known to occur within area |
| Chelonia mydas Green Turtle [1765] | Vulnerable | Species or species habitat known to occur within area |
| Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat known to occur within area |
| Natator depressus Flatback Turtle [59257] | Vulnerable | Species or species habitat known to occur within area |
| Sharks | | |
| Carcharias taurus (west coast population) Grey Nurse Shark (west coast population) [68752] | Vulnerable | Species or species habitat known to occur within area |
| Carcharodon carcharias White Shark, Great White Shark [64470] | Vulnerable | Species or species habitat known to occur within area |
| Rhincodon typus Whale Shark [66680] | Vulnerable | Species or species habitat may occur within area |
| Listed Migratory Species | | [Resource Information] |
| * Species is listed under a different scientific name on | the EPBC Act - Threatened | |
| Name Migratory Marine Birds | Threatened | Type of Presence |
| Anous stolidus | | |
| Common Noddy [825] | | Species or species habitat likely to occur within area |
| Apus pacificus Fork-tailed Swift [678] | | Species or species habitat likely to occur within area |
| Ardenna carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [82404] | | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea amsterdamensis Amsterdam Albatross [64405] | Endangered | Species or species habitat may occur within area |

| Name | Threatened | Type of Presence |
|---|-------------|--|
| Diomedea epomophora Southern Royal Albatross [89221] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea exulans Wandering Albatross [89223] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea sanfordi Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Hydroprogne caspia Caspian Tern [808] | | Foraging, feeding or related behaviour known to occur within area |
| Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Onychoprion anaethetus Bridled Tern [82845] Phoebetria fusca | | Breeding known to occur within area |
| Sooty Albatross [1075] | Vulnerable | Species or species habitat may occur within area |
| Sterna dougallii Roseate Tern [817] | | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche cauta Shy Albatross [89224] | Vulnerable* | Species or species habitat may occur within area |
| Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche steadi White-capped Albatross [64462] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| Migratory Marine Species | | within area |
| Balaena glacialis australis Southern Right Whale [75529] Balaenoptera edeni | Endangered* | Breeding known to occur within area |
| Bryde's Whale [35] | | Species or species habitat may occur within area |
| Balaenoptera musculus Blue Whale [36] | Endangered | Species or species habitat likely to occur within area |
| Caperea marginata Pygmy Right Whale [39] | | Species or species habitat may occur within area |
| Carcharodon carcharias White Shark, Great White Shark [64470] | Vulnerable | Species or species habitat known to occur within area |
| Caretta caretta Loggerhead Turtle [1763] | Endangered | Species or species |

| Name | Threatened | Type of Presence |
|--|-----------------------|---|
| | | habitat known to occur |
| <u>Chelonia mydas</u> | | within area |
| Green Turtle [1765] | Vulnerable | Species or species habitat |
| | | known to occur within area |
| <u>Dermochelys coriacea</u> | | |
| Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat known to occur within area |
| | | known to occur within area |
| Lamna nasus Darbaaria, Maakaral Chark [92299] | | Charies or angeles habitat |
| Porbeagle, Mackerel Shark [83288] | | Species or species habitat may occur within area |
| | | , |
| Manta alfredi Reef Manta Ray, Coastal Manta Ray, Inshore Manta | | Species or species habitat |
| Ray, Prince Alfred's Ray, Resident Manta Ray [84994] | | may occur within area |
| Manta birostris | | |
| Giant Manta Ray, Chevron Manta Ray, Pacific Manta | | Species or species habitat |
| Ray, Pelagic Manta Ray, Oceanic Manta Ray [84995] | | may occur within area |
| Megaptera novaeangliae | | |
| Humpback Whale [38] | Vulnerable | Species or species habitat known to occur within area |
| | | known to occur within area |
| Natator depressus Flotback Turtle [50257] | Vulnerable | Chasias ar angeige habitat |
| Flatback Turtle [59257] | vuinerable | Species or species habitat known to occur within area |
| Orainua araa | | |
| Orcinus orca Killer Whale, Orca [46] | | Species or species habitat |
| | | may occur within area |
| Rhincodon typus | | |
| Whale Shark [66680] | Vulnerable | Species or species habitat |
| | | may occur within area |
| Migratory Terrestrial Species | | |
| Motacilla cinerea Grey Wagtail [642] | | Species or species habitat |
| Croy Wagtan [042] | | may occur within area |
| Migratory Wetlands Species | | |
| Actitis hypoleucos | | |
| Common Sandpiper [59309] | | Species or species habitat |
| | | may occur within area |
| Calidris acuminata | | On a sing on an arise helitat |
| Sharp-tailed Sandpiper [874] | | Species or species habitat may occur within area |
| | | |
| Calidris canutus Red Knot, Knot [855] | Endangered | Species or species habitat |
| | 9 | known to occur within area |
| <u>Calidris ferruginea</u> | | |
| Curlew Sandpiper [856] | Critically Endangered | Species or species habitat |
| | | may occur within area |
| <u>Calidris melanotos</u> | | |
| Pectoral Sandpiper [858] | | Species or species habitat may occur within area |
| | | a, oodar within area |
| <u>Limosa lapponica</u> Bar-tailed Godwit [844] | | Species or species habitat |
| Dai tanoa Coawit [OTT] | | may occur within area |
| Numenius madagascariensis | | |
| Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat |
| | | may occur within area |
| | | |

| Name | Threatened | Type of Presence |
|-------------------|------------|--|
| Pandion haliaetus | | |
| Osprey [952] | | Species or species habitat may occur within area |

Other Matters Protected by the EPBC Act

Commonwealth Land [Resource Information]

The Commonwealth area listed below may indicate the presence of Commonwealth land in this vicinity. Due to the unreliability of the data source, all proposals should be checked as to whether it impacts on a Commonwealth area, before making a definitive decision. Contact the State or Territory government land department for further information.

Name

Commonwealth Land -

| Listed Marine Species | | [Resource Information] |
|--|-----------------------------|--|
| * Species is listed under a different scientific nam | ne on the EPBC Act - Threat | ened Species list. |
| Name | Threatened | Type of Presence |
| Birds | | |
| Actitis hypoleucos | | |
| Common Sandpiper [59309] | | Species or species habitat may occur within area |

Anous stolidus

Common Noddy [825] Species or species habitat likely to occur within area

Anous tenuirostris melanops

Australian Lesser Noddy [26000] Vulnerable Species or species habitat

may occur within area

Apus pacificus

Fork-tailed Swift [678] Species or species habitat

likely to occur within area

Ardea alba

Great Egret, White Egret [59541] Breeding known to occur

within area

Ardea ibis

Cattle Egret [59542] Species or species habitat

may occur within area

Calidris acuminata

Sharp-tailed Sandpiper [874] Species or species habitat

may occur within area

Calidris canutus

Red Knot, Knot [855] Endangered Species or species habitat

known to occur within area

Calidris ferruginea

Curlew Sandpiper [856] Critically Endangered Species or species habitat

may occur within area

| Name | Threatened | Type of Presence |
|---|-----------------------|--|
| <u>Calidris melanotos</u> | | |
| Pectoral Sandpiper [858] | | Species or species habitat may occur within area |
| Diomedea amsterdamensis | | |
| Amsterdam Albatross [64405] | Endangered | Species or species habitat may occur within area |
| <u>Diomedea epomophora</u> | | |
| Southern Royal Albatross [89221] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| <u>Diomedea exulans</u> | Mada a salida | |
| Wandering Albatross [89223] Diomedea sanfordi | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Northern Royal Albatross [64456] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Haliaeetus leucogaster | | |
| White-bellied Sea-Eagle [943] | | Species or species habitat likely to occur within area |
| Larus novaehollandiae | | |
| Silver Gull [810] Larus pacificus | | Breeding known to occur within area |
| Pacific Gull [811] | | Foraging, feeding or related behaviour may occur within area |
| <u>Limosa lapponica</u> | | |
| Bar-tailed Godwit [844] | | Species or species habitat may occur within area |
| Macronectes giganteus | | |
| Southern Giant-Petrel, Southern Giant Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli | | |
| Northern Giant Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Merops ornatus | | |
| Rainbow Bee-eater [670] | | Species or species habitat may occur within area |
| Motacilla cinerea | | |
| Grey Wagtail [642] | | Species or species habitat may occur within area |
| Numenius madagascariensis | | |
| Eastern Curlew, Far Eastern Curlew [847] | Critically Endangered | Species or species habitat may occur within area |
| Pachyptila turtur | | |
| Fairy Prion [1066] | | Species or species habitat known to occur within area |
| Pandion haliaetus | | |
| Osprey [952] | | Species or species habitat may occur within area |
| Phoebetria fusca | | |
| Sooty Albatross [1075] | Vulnerable | Species or species habitat may occur within area |
| Puffinus assimilis | | |
| Little Shearwater [59363] | | Foraging, feeding or related behaviour known to occur within area |
| Puffinus carneipes Floch footed Shorweter, Flochy footed | | Egraciae fooding or |
| Flesh-footed Shearwater, Fleshy-footed | | Foraging, feeding or |

| Name | Threatened | Type of Presence |
|--|--------------|--|
| Shearwater [1043] | Tilledictied | related behaviour likely to |
| | | occur within area |
| Rostratula benghalensis (sensu lato) | | |
| Painted Snipe [889] | Endangered* | Species or species habitat |
| | | likely to occur within area |
| Sterna anaethetus | | |
| Bridled Tern [814] | | Breeding known to occur |
| | | within area |
| Sterna caspia | | |
| Caspian Tern [59467] | | Foraging, feeding or related |
| | | behaviour known to occur |
| Sterna dougallii | | within area |
| Roseate Tern [817] | | Foraging, feeding or related |
| | | behaviour likely to occur |
| | | within area |
| <u>Thalassarche cauta</u> | | |
| Shy Albatross [89224] | Vulnerable* | Species or species habitat |
| | | may occur within area |
| Thalassarche impavida | | |
| Campbell Albatross, Campbell Black-browed Albatross | Vulnerable | Species or species habitat |
| [64459] | | may occur within area |
| | | · |
| <u>Thalassarche melanophris</u> | | |
| Black-browed Albatross [66472] | Vulnerable | Species or species habitat |
| | | may occur within area |
| Thalassarche steadi | | |
| White-capped Albatross [64462] | Vulnerable* | Foraging, feeding or related |
| | | behaviour likely to occur |
| | | within area |
| Fish | | |
| Acentronura australe Southorn Pygmy Pincherse [66185] | | Species or species habitat |
| Southern Pygmy Pipehorse [66185] | | Species or species habitat may occur within area |
| | | may boodi within area |
| Campichthys galei | | |
| Gale's Pipefish [66191] | | Species or species habitat |
| | | may occur within area |
| Choeroichthys suillus | | |
| Pig-snouted Pipefish [66198] | | Species or species habitat |
| | | openies of species flabitat |
| | | · |
| | | may occur within area |
| Halicampus brocki | | may occur within area |
| Halicampus brocki Brock's Pipefish [66219] | | may occur within area Species or species habitat |
| • | | may occur within area |
| Brock's Pipefish [66219] | | may occur within area Species or species habitat |
| Brock's Pipefish [66219] Hippocampus angustus | | may occur within area Species or species habitat may occur within area |
| Brock's Pipefish [66219] | | may occur within area Species or species habitat |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] | | Species or species habitat may occur within area Species or species habitat |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps | | Species or species habitat may occur within area Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse | | Species or species habitat may occur within area Species or species habitat may occur within area Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps | | Species or species habitat may occur within area Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse | | Species or species habitat may occur within area Species or species habitat may occur within area Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] | | Species or species habitat may occur within area Species or species habitat may occur within area Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus | | Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] | | Species or species habitat may occur within area Species or species habitat |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus | | Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] | | Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus | | Species or species habitat may occur within area |
| Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus | | Species or species habitat may occur within area |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus Prophet's Pipefish [66250] | | Species or species habitat may occur within area Species or species habitat may occur within area |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus Prophet's Pipefish [66250] | | Species or species habitat may occur within area |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus Prophet's Pipefish [66250] Maroubra perserrata Sawtooth Pipefish [66252] | | Species or species habitat may occur within area Species or species habitat may occur within area |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus Prophet's Pipefish [66250] Maroubra perserrata Sawtooth Pipefish [66252] | | Species or species habitat may occur within area Species or species habitat may occur within area |
| Brock's Pipefish [66219] Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234] Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235] Hippocampus subelongatus West Australian Seahorse [66722] Lissocampus fatiloquus Prophet's Pipefish [66250] Maroubra perserrata Sawtooth Pipefish [66252] | | Species or species habitat may occur within area Species or species habitat may occur within area |

| Name | Threatened | Type of Presence |
|--|-------------|---|
| | | area |
| Nannocampus subosseus Bonyhead Pipefish, Bony-headed Pipefish [66264] | | Species or species habitat may occur within area |
| Phycodurus eques | | |
| Leafy Seadragon [66267] | | Species or species habitat may occur within area |
| Phyllopteryx taeniolatus | | |
| Common Seadragon, Weedy Seadragon [66268] | | Species or species habitat may occur within area |
| Pugnaso curtirostris | | |
| Pugnose Pipefish, Pug-nosed Pipefish [66269] | | Species or species habitat may occur within area |
| Solegnathus lettiensis | | |
| Gunther's Pipehorse, Indonesian Pipefish [66273] | | Species or species habitat may occur within area |
| Stigmatopora argus | | |
| Spotted Pipefish, Gulf Pipefish, Peacock Pipefish [66276] | | Species or species habitat may occur within area |
| Stigmatopora nigra | | |
| Widebody Pipefish, Wide-bodied Pipefish, Black Pipefish [66277] | | Species or species habitat may occur within area |
| Syngnathoides biaculeatus | | |
| Double-end Pipehorse, Double-ended Pipehorse, Alligator Pipefish [66279] | | Species or species habitat may occur within area |
| <u>Urocampus carinirostris</u> | | |
| Hairy Pipefish [66282] | | Species or species habitat may occur within area |
| Vanacampus margaritifer | | |
| Mother-of-pearl Pipefish [66283] | | Species or species habitat may occur within area |
| Mammals | | |
| Arctocephalus forsteri | | |
| Long-nosed Fur-seal, New Zealand Fur-seal [20] | | Species or species habitat may occur within area |
| Neophoca cinerea | | |
| Australian Sea-lion, Australian Sea Lion [22] | Vulnerable | Species or species habitat known to occur within area |
| Reptiles | | |
| Aipysurus pooleorum Charle Boy Cooppeles [66061] | | Chasias ar angeige habitat |
| Shark Bay Seasnake [66061] | | Species or species habitat may occur within area |
| Caretta caretta Loggerhead Turtle [1763] | Endangered | Species or species habitat known to occur within area |
| Cholonia mudas | | |
| <u>Chelonia mydas</u> Green Turtle [1765] | Vulnerable | Species or species habitat |
| | vuirierable | known to occur within area |
| <u>Dermochelys coriacea</u> Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Species or species habitat known to occur within area |
| Disteira kingii | | |
| Spectacled Seasnake [1123] | | Species or species habitat may occur within area |
| Natator depressus | | |
| Flatback Turtle [59257] | Vulnerable | Species or species |
| | | |

| Polamic platurus | | witnin area |
|---|------------|--|
| Pelamis platurus Yellow-bellied Seasnake [1091] | | Species or species habitat may occur within area |
| Whales and other Cetaceans | | [Resource Information] |
| Name | Status | Type of Presence |
| Mammals | | |
| Balaenoptera acutorostrata | | |
| Minke Whale [33] | | Species or species habitat may occur within area |
| Balaenoptera edeni | | |
| Bryde's Whale [35] | | Species or species habitat may occur within area |
| Balaenoptera musculus | | |
| Blue Whale [36] | Endangered | Species or species habitat likely to occur within area |
| Caperea marginata | | |
| Pygmy Right Whale [39] | | Species or species habitat may occur within area |
| Delphinus delphis | | |
| Common Dophin, Short-beaked Common Dolphin [60] | | Species or species habitat may occur within area |
| Eubalaena australis | | |
| Southern Right Whale [40] | Endangered | Breeding known to occur within area |
| Grampus griseus | | |
| Risso's Dolphin, Grampus [64] | | Species or species habitat may occur within area |
| Megaptera novaeangliae | | |
| Humpback Whale [38] | Vulnerable | Species or species habitat known to occur within area |
| Orcinus orca | | |
| Killer Whale, Orca [46] | | Species or species habitat may occur within area |
| Stenella attenuata | | |
| Spotted Dolphin, Pantropical Spotted Dolphin [51] | | Species or species habitat may occur within area |
| Tursiops aduncus | | |
| Indian Ocean Bottlenose Dolphin, Spotted Bottlenose Dolphin [68418] | | Species or species habitat likely to occur within area |
| Tursiops truncatus s. str. | | |
| Bottlenose Dolphin [68417] | | Species or species habitat may occur within area |
| | | |

Threatened

Type of Presence

within area

habitat known to occur

Extra Information

Name

Invasive Species [Resource Information]

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resouces Audit, 2001.

| Name | Status | Type of Presence |
|-------|--------|------------------|
| Birds | | |

| Name | Status | Type of Presence |
|---|--------|--|
| Acridotheres tristis Common Myna, Indian Myna [387] | | Species or species habitat likely to occur within area |
| Anas platyrhynchos Mallard [974] | | Species or species habitat likely to occur within area |
| Carduelis carduelis European Goldfinch [403] | | Species or species habitat likely to occur within area |
| Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803] | | Species or species habitat likely to occur within area |
| Passer domesticus House Sparrow [405] | | Species or species habitat likely to occur within area |
| Passer montanus Eurasian Tree Sparrow [406] | | Species or species habitat likely to occur within area |
| Streptopelia chinensis Spotted Turtle-Dove [780] | | Species or species habitat likely to occur within area |
| Streptopelia senegalensis Laughing Turtle-dove, Laughing Dove [781] | | Species or species habitat likely to occur within area |
| Sturnus vulgaris Common Starling [389] | | Species or species habitat likely to occur within area |
| | | |
| Mammals | | |
| Mammals Bos taurus Domestic Cattle [16] | | Species or species habitat likely to occur within area |
| Bos taurus | | • |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris | | likely to occur within area Species or species habitat |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris Domestic Dog [82654] Felis catus | | Species or species habitat likely to occur within area Species or species habitat likely to occur within area |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris Domestic Dog [82654] Felis catus Cat, House Cat, Domestic Cat [19] Funambulus pennantii Northern Palm Squirrel, Five-striped Palm Squirrel | | Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris Domestic Dog [82654] Felis catus Cat, House Cat, Domestic Cat [19] Funambulus pennantii Northern Palm Squirrel, Five-striped Palm Squirrel [129] Mus musculus | | Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris Domestic Dog [82654] Felis catus Cat, House Cat, Domestic Cat [19] Funambulus pennantii Northern Palm Squirrel, Five-striped Palm Squirrel [129] Mus musculus House Mouse [120] Oryctolagus cuniculus | | Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area |
| Bos taurus Domestic Cattle [16] Canis lupus familiaris Domestic Dog [82654] Felis catus Cat, House Cat, Domestic Cat [19] Funambulus pennantii Northern Palm Squirrel, Five-striped Palm Squirrel [129] Mus musculus House Mouse [120] Oryctolagus cuniculus Rabbit, European Rabbit [128] | | Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area Species or species habitat likely to occur within area |

| Name | Status | Type of Presence |
|---|-----------------|--|
| Vulpes vulpes | | |
| Red Fox, Fox [18] | | Species or species habitat likely to occur within area |
| Plants | | |
| Asparagus aethiopicus | | |
| Asparagus Fern, Ground Asparagus, Basket Fern, Sprengi's Fern, Bushy Asparagus, Emerald Asparag [62425] Asparagus asparagoides | us | Species or species habitat likely to occur within area |
| Bridal Creeper, Bridal Veil Creeper, Smilax, Florist's Smilax, Smilax Asparagus [22473] | | Species or species habitat likely to occur within area |
| Brachiaria mutica | | |
| Para Grass [5879] | | Species or species habitat may occur within area |
| Cenchrus ciliaris | | |
| Buffel-grass, Black Buffel-grass [20213] | | Species or species habitat may occur within area |
| Chrysanthemoides monilifera | | |
| Bitou Bush, Boneseed [18983] | | Species or species habitat may occur within area |
| Chrysanthemoides monilifera subsp. monilifera | | |
| Boneseed [16905] | | Species or species habitat likely to occur within area |
| Genista sp. X Genista monspessulana | | |
| Broom [67538] | | Species or species habitat may occur within area |
| Lantana camara | | |
| Lantana, Common Lantana, Kamara Lantana, Large leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White Sage, Wild Sag [10892] | | Species or species habitat likely to occur within area |
| Olea europaea | | |
| Olive, Common Olive [9160] | | Species or species habitat may occur within area |
| Pinus radiata | | |
| Radiata Pine Monterey Pine, Insignis Pine, Wilding Pine [20780] | | Species or species habitat may occur within area |
| Rubus fruticosus aggregate | | |
| Blackberry, European Blackberry [68406] | | Species or species habitat likely to occur within area |
| Salix spp. except S.babylonica, S.x calodendron & S | S.x reichardtii | |
| Willows except Weeping Willow, Pussy Willow and Sterile Pussy Willow [68497] | | Species or species habitat likely to occur within area |
| Salvinia molesta Salvinia, Giant Salvinia, Aquarium Watermoss, Karik | oa | Species or species habitat |
| Weed [13665] | | likely to occur within area |
| Tana auto andre Us | | |
| Tamarix aphylla | | |
| Athel Pine, Athel Tree, Tamarisk, Athel Tamarisk, Athel Tamarix, Desert Tamarisk, Flowering Cypress, Salt Cedar [16018] | | Species or species habitat likely to occur within area |
| Reptiles | | |
| Hemidactylus frenatus | | |

Asian House Gecko [1708]

Species or species habitat likely to occur within area

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the gualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

 $-31.46962\ 115.56826, -31.46962\ 115.583, -31.64574\ 115.69925, -31.64574\ 115.68169, -31.46962\ 115.56826, -31.46962\ 115.56826, -31.46962\ 115.583, -31.64574\ 115.69925, -31.64574\ 115.68169, -31.46962\ 115.56826, -31.46962, -31.46962\ 115.56826, -31.46962, -3$

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- -Office of Environment and Heritage, New South Wales
- -Department of Environment and Primary Industries, Victoria
- -Department of Primary Industries, Parks, Water and Environment, Tasmania
- -Department of Environment, Water and Natural Resources, South Australia
- -Department of Land and Resource Management, Northern Territory
- -Department of Environmental and Heritage Protection, Queensland
- -Department of Parks and Wildlife, Western Australia
- -Environment and Planning Directorate, ACT
- -Birdlife Australia
- -Australian Bird and Bat Banding Scheme
- -Australian National Wildlife Collection
- -Natural history museums of Australia
- -Museum Victoria
- -Australian Museum
- -South Australian Museum
- -Queensland Museum
- -Online Zoological Collections of Australian Museums
- -Queensland Herbarium
- -National Herbarium of NSW
- -Royal Botanic Gardens and National Herbarium of Victoria
- -Tasmanian Herbarium
- -State Herbarium of South Australia
- -Northern Territory Herbarium
- -Western Australian Herbarium
- -Australian National Herbarium, Canberra
- -University of New England
- -Ocean Biogeographic Information System
- -Australian Government, Department of Defence
- Forestry Corporation, NSW
- -Geoscience Australia
- -CSIRO
- -Australian Tropical Herbarium, Cairns
- -eBird Australia
- -Australian Government Australian Antarctic Data Centre
- -Museum and Art Gallery of the Northern Territory
- -Australian Government National Environmental Science Program
- -Australian Institute of Marine Science
- -Reef Life Survey Australia
- -American Museum of Natural History
- -Queen Victoria Museum and Art Gallery, Inveresk, Tasmania
- -Tasmanian Museum and Art Gallery, Hobart, Tasmania
- -Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the Contact Us page.

Appendix B Vertebrate Fauna Recorded in Biological Surveys in the Region

Vertebrate Fauna Survey - Yanchep Lagoon, Yanchep

Annandiv R(1) Variabrata fauna accessments

| (=) | . Vertebrate fauna assessme | Surveys | s A | В | (| C | D | E | F | | G | j | | | | | H | | | | I | | J | | | K | | L | | M |
|-------------------|---------------------------------|---------------------------|-----|---|----------|---------|---|---|---------|--------|----------|--------------------|---------|---------|---------|----------|----------|----------|----------|----------|--------|------------|------------|--------|---------------|---------------|-------------------------|--------------|----------|---|
| Family | Species | Common Name | | | Brighton | Unknown | | | Frinity | Acacia | Dryandra | Eucalypt & Banksia | Banksia | Site 01 | Site 02 | Site 10B | Site 02A | Site 10A | Site 01A | Site 14A | Site 4 | Site 4 | Site 2 | Site 1 | Opportunistic | Opportunistic | Site 5 Cassilda Park | Kinsale Park | Brighton | |
| Birds | | | | | _ | | | | | 7 | | | | | 9, 9 | | | | | • | 9, | 9 2 | 9 2 | - | | | | | | X |
| Accipitridae | Accipiter cirrocephalus | Collared Sparrowhawk | | | | | | | | | | | | | | | | | | | | | | T | \top | 1 | _ | 1 | T | T |
| | Accipiter fasciatus | Brown Goshawk | | | | | | | | 1 | 1 | 1 | | | | | | | | | | | | 1 | \top | 2 | \top | 1 | T | |
| | Aquila audax | Wedge-tailed Eagle | | | | | | | | | | | | | | | | | | | | | 1 | 1 | \top | | \top | 1 | T | |
| | Circus approximans | Swamp Harrier | | | | | | | | | | | | | | | | | | | | | | 1 | \top | 3 | \top | 1 | T | |
| | Elanus axillaris | Black-shouldered Kite | | | | | | | | 1 | | | | | | | | | | | | | 1 | 1 | \top | 4 | \top | 1 | T | |
| | Haliastur sphenurus | Whistling Kite | | | | | | | | 1 | | | | | | | | | | | | | | 1 | \top | 2 | \top | 1 | T | |
| | Hieraaetus morphnoides | Little Eagle | 1 | t | | | | | 7 | 1 | | | | 7 | _ | 1 | 1 | 1 | | | | | 1 | 1 | | 3 | 十 | 1 | 1 | |
| | Lophoictinia isura | Square-tailed Kite | | | | | | | | | | | | | | | | | | | | | 1 | 1 | \top | | \top | 1 | T | |
| Anatidae | Tadorna tadornoides | Australian Shelduck | T | 1 | | | | | T | | | | | | | T | Ť | | | | | 5 | | 寸 | 寸 | T | \top | 1 | T | |
| Podargidae | Podargus strigoides | Tawny Frogmouth | | | | | | | | | | | | | | | | | | | | | | 1 | \top | 1 | \top | 1 | T | |
| Casuariidae | Dromaius novaehollandiae | Emu | | | | | | | | | | | | | | | | | | | | 1 | 1 | | 1 | 9 | \top | | | T |
| Laridae | Chroicocephalus novaehollandiae | Silver Gull | | | | | | | | | | | | | | | | | | | | | | 1 | | | \top | | | T |
| Threskiornithidae | Threskiornis spinicollis | Straw-necked Ibis | | | | | | | | | | | | | | | | | | | | | | | - | 92 | + | + | T | |
| Columbidae | Columba livia | Domestic Pigeon | | | | | | | | | | | | | | | | | | | | | 1 | 12 | T | | + | + | T | |
| | Ocyphaps lophotes | Crested Pigeon | | | | | | | | 3 | 7 | | | | | | | | | | | | | | 1 | 15 | + | + | T | |
| | Phaps chalcoptera | Common Bronzewing | | | | | | | | | 2 | 1 | | | | | | | | | 2 | 1 | | | 1 | 14 | + | + | T | |
| | Phaps elegans | Brush Bronzewing | | | | | | | 75 | | | | | | | | | | | | | | | | | | | | | |
| | Spilopelia senegalensis | Laughing Turtle-dove | | | | | | | | 6 | 4 | | | | | | | | | | | | | 6 | | 2 | 1 | 1 | | |
| | Spilpopelia chinensis | Spotted Turtle-dove | | | | | | | | | | | | | | | | | | | | | | 1 | 1 | | \top | 1 | T | |
| Halcyonidae | Dacelo novaeguineae | Laughing Kookaburra | | | | | | | | | | 6 | | | | | | | | | | 1 | | | | 9 | | | | |
| | Todiramphus sanctus | Sacred Kingfisher | | | | | | | | | | 2 | 1 | | | | | | | | 1 | | 1 | 1 | 1 | 1 | 1 | 1 | T | |
| Meropidae | Merops ornatus | Rainbow Bee-eater | | | | | | | | | | | | | | | | | | | 2 | | | 1 | - 1 | 26 | \top | 1 | T | |
| Cuculidae | Cacomantis flabelliformis | Fan-tailed Cuckoo | | | | | | | | | | | | | | | | | | | 1 | 3 | | 1 | \top | | \top | 1 | T | |
| | Chalcites basalis | Horsfield's Bronze-cuckoo | | | | | | | | 1 | 1 | | | | | | | | | | | 2 | 1 | 1 | | 1 | \top | | | T |
| | Chalcites lucidus | Shining Bronze-cuckoo | | | | | | | | | | | | | | | | | | | | 2 | | 1 | \top | | \top | 1 | T | |
| | Heteroscenes pallidus | Pallid Cuckoo | | | | | | | | | | | | | | | | | | | | 2 | | 1 | \top | | \top | 1 | T | |
| Falconidae | Falco berigora | Brown Falcon | | | | | | | | | | | | | | | | | | | | | | | | 2 | | | | |
| Falconidae | Falco cenchroides | Nankeen Kestrel | | | | | | | | | | | 1 | | | | | | | | | | 2 | 1 | | 1 | | 1 | | |
| | Falco longipennis | Australian Hobby | | | | | | | | | | | | | | | | | | | | 1 | | 1 | \top | 1 | \top | 1 | T | |
| | Falco peregrinus | Peregrine Falcon | T | T | | | | | T | | | | | T | | T | 1 | | | | | | 1 | 寸 | 寸 | T | \top | 1 | T | |
| Phasianidae | Coturnix pectoralis | Stubble Quail | T | 1 | | | | | T | | | | | | | T | Ť | | | | | | | 寸 | 寸 | 7 | \top | 1 | T | |
| Otididae | Ardeotis australis | Australian Bustard | | | | | | | T | T | | | | | | | | | 1 | | | | 1 | 寸 | 寸 | T | | 1 | | |
| Rallidae | Porzana tabuensis | Spotless Crake | | T | | 1 | | | T | | | | | 寸 | 1 | 1 | 1 | T | | | | | | 寸 | \dashv | T | \neg | T | 1 | |
| Acanthizidae | Acanthiza apicalis | Inland Thornbill | X | T | | | | | T | | | | | 寸 | 1 | 1 | 1 | T | | | | 1 | | 1 | \dashv | 8 | \neg | T | 1 | |
| | Acanthiza chrysorrhoa | Yellow-rumped Thornbill | X | | | | | | T | T | | | 3 | | | | | | 1 | | 4 | 5 | T | 寸 | _ | 9 | | 1 | | |
| | Acanthiza inornata | Western Thornbill | X | | | | | | T | 7 | | | | T | | | | | 1 | | | 4 | T | 寸 | 1 | 28 | \top | 1 | | |
| | Gerygone fusca | Western Gerygone | X | | | | | | T | 1 | | 10 | 6 | T | | | | | 1 | | 6 | 12 | T | 2 | 1 . | 35 | \top | 1 | | |
| | Sericornis frontalis | White-browed Scrubwren | X | T | | | | | T | | 3 | | | 寸 | 1 | 1 | 1 | T | | | 2 | | | 22 | 1 | 13 | 1 | 1 | 1 | |
| | Smicrornis brevirostris | Weebill | X | T | | | | | T | | | 11 | | 寸 | 1 | 1 | 1 | T | | | | 1 | | 寸 | | 11 | \neg | T | 1 | |
| Acrocephalidae | Acrocephalus australis | Australian Reed-warbler | X | | | | | | | | | | | 1 | | | 1 | T | | | | | - t | 十 | \exists | 7 | 十 | 1 | 1 | |

| | | Survey | A | B | C | 7 | D | E | F | | G | | | | | | Н | | | | I | | J | | | K | | I | , | \mathbf{M} |
|---|-------------------------------|---------------------------|---|---|----------|---------|---|---|----------------|--------|------|--------------------|---------|------------|---------------|----------|-----------|----------|----------|----------|------------|--------|--------|--------|---------------|---------------|---------------|------------------------------|----------|--------------|
| | | Survey | 1 | | Ĭ | | _ | _ | 1 | | Ī | | | | | T | Ť | | | | _ | | Ĭ | | | Î | | Ť | Т | 112 |
| Family | Species | Common Name | | | Brighton | Unknown | | | Frinity | Acacia | | Eucalypt & Banksia | Sanksia | Site 01 | Site 02 | Site 10B | Site 02A | Site 10A | Site 01A | Site 14A | Site 4 | Site 4 | Site 2 | Site 1 | Opportunistic | Opportunistic | Site 5 | Cassuda Park Kinsale Park | Brighton | |
| Artamidae | Artamus cinereus | Black-faced Woodswallow | X | | _ | 5 | | | | 7 | 11 | _ | | 9 2 | 0 2 | 2 0. | | | 0,1 | | 9 2 | | 7 | | | 2 | 6 2 | | | - |
| - II tullilluu | Artamus cyanopterus | Dusky Woodswallow | X | | | | | | | | | | | | | | | | | | | | | | | | | + | + | + |
| | Artamus personatus | Masked Woodswallow | X | | | | | | 1 | | | - | | | | | | | | | | | | | | | | + | + | 1 |
| | Cracticus nigrogularis | Pied Butcherbird | X | | | | | | 1 | | | - | | | | | | | | | | | | | | | | + | + | 1 |
| | Cracticus torquatus | Grey Butcherbird | X | | | | | | 1 | 2 | 6 | 3 | 3 | | | | | | | | | 1 | 3 | 5 | | 4 | | 1 | 1 | 1 |
| | Gymnorhina tibicen | Australian Magpie | X | | | | | | _ | 1 | _ | _ | 6 | | | - | | | | | 4 | 7 | 2 | 3 | _ | 74 | - | 1 1 | 1 | + |
| Campephagidae | Coracina novaehollandiae | Black-faced Cuckoo-shrike | X | | | 5 | | + | _ | _ | _ | _ | 3 | 1 | _ | + | + | | | | - | 3 | 2 | | _ | 12 | - | H | Ť | + |
| Campophagidae | Lalage tricolor | White-winged Triller | X | | | 15 | | 1 | \dashv | | 1 | - | | _ | | + | \dagger | | | | | | - | _ | 7 | | -+ | + | + | + + |
| Corvidae | Corvus coronoides | Australian Raven | X | | | | | + | \dashv | 2 | 1 | \dashv | 1 | \dashv | _ | + | 1 | | 1 | | 4 | 6 | 6 | 6 | 1 | 34 | | 1 | + | \forall |
| Hirundinidae | Cheramoeca leucosterna | White-backed Swallow | X | | | | | | 1 | _ | | - | | | | | | | | | | Ť | 1 | 1 | | 3 | | Ť | + | 1 |
| 1111 01101111000 | Hirundo neoxena | Welcome Swallow | X | | | | | | 1 | 4 | | 1 | | | | | | | | | | | 1 | 4 | _ | 20 | | 1 1 | | \top |
| | Petrochelidon nigricans | Tree Martin | X | | | | | | | Ť | | 1 | | | | | | | | | | | 1 | 18 | | | | + | | \top |
| Maluridae | Malurus lamberti | Variegated Fairy-wren | X | | | 5 | | | | 1 | | 1 | | | | | | | | | | | | | | 24 | | 1 | | \top |
| 111111111111111111111111111111111111111 | Malurus leucopterus | White-winged Fairy-wren | X | | | _ | | | | 1 | | 1 | | | | | | | | | | | | 4 | _ | 23 | | + | | \top |
| | Malurus splendens | Splendid Fairy-wren | X | | | | | | | 12 | 30 2 | 22 | 18 | | | | | | | | 4 | 18 | 8 | 1 | | 204 | | \top | | \top |
| | Stipiturus malachurus | Southern Emu-wren | X | | | | | | | 1 | | 1 | | | | | | | | | | | | | | 1 | | \top | | \top |
| Megaluridae | Cincloramphus mathewsi | Rufous Songlark | X | | | | | | | 1 | | 1 | | | | | | | | | | | | | | 1 | | \top | | \top |
| rregurariane | Acanthorhynchus superciliosus | Western Spinebill | X | | | | | | | 1 | | 1 | | | | | | | | | | 4 | | | | 4 | | \top | | \top |
| | Anthochaera carunculata | Red Wattlebird | X | | | 5 | | | | 1 | 11 | 11 | 15 | | | | | | | | 5 | 6 | 4 | 6 | | 36 | | 1 1 | | \top |
| | Anthochaera chrysoptera | Little Wattlebird | | | | | | | | | | | | | | | | | | | | 14 | 1 | 1 | | 3 | | + | + | + |
| | Anthochaera lunulata | Western Little Wattlebird | X | | | | | | | | | | | | | | | | | | | | | | | | | \top | + | \top |
| | Epthianura albifrons | White-fronted Chat | X | | | | | | | 1 | | 1 | | | | | | | | | | | | | | 15 | | \top | | \top |
| | Gavicalis virescens | Singing Honeyeater | X | | | | | | | 7 | 4 | | 10 | | | | | | | | | 2 | 1 | 12 | _ | 10 | | 1 1 | + | T |
| | Gliciphila melanops | Tawny-crowned Honeyeater | X | | | | | | | | | | | | | | | | | | | | 12 | | | 9 | | \top | + | \top |
| | Lichenostomus ornatus | Yellow-plumed Honeyeater | | | | | | | | 1 | | 1 | | | | | | | | | | 1 | | | | Ť | | \top | | \top |
| | Lichmera indistincta | Brown Honeveater | | | | | | | | 1 | | 1 | | | | | | | | | | | | | | 193 | | \top | | \top |
| | Lichmera indistincta | Brown Honeyeater | X | | | | | | | 3 | 4 | 44 | 11 | | | | | | | | | 21 | 15 | 7 | | | | 1 1 | + | T |
| | Manorina flavigula | Yellow-throated Miner | X | | | | | | | | | | | | | | | | | | | | | | | | | \top | 1 | \top |
| | Melithreptus lunatus | White-naped Honeyeater | Ť | | | | | | | 1 | | 1 | | | | | | | | | | 1 | | | | 1 | | \top | + | \top |
| | Sugomel nigrum | Black Honeyeater | | | | | | | | 1 | | 1 | | | | | | | | | | Ť | | | | 109 | | \top | | \top |
| | Phylidonyris niger | White-cheeked Honeyeater | X | | | 1 | | | 3 | 30 | 230 | 10 | | | | | | | | | | | 5 | 3 | | | | 1 | + | \top |
| Meliphagidae | Phylidonyris novaehollandiae | New Holland Honeyeater | X | | | | | | | | | | | | | | | | | | | 3 | | | | 152 | | 1 1 | + | \top |
| Monarchidae | Grallina cyanoleuca | Magpie-lark | X | | | | | | | 1 | | 1 | | | | | | | | | | 1 | | | _ | 2 | | 1 1 | 1 | \top |
| Motacillidae | Anthus novaeseelandiae | Australasian Pipit | X | | | 15 | | 1 | \dashv | 2 | 1 | | 1 | \dashv | $\neg \vdash$ | \top | † | | | | | | | 1 | | 31 | $\exists f$ | \top | T | \forall |
| Nectariniidae | Dicaeum hirundinaceum | Mistletoe Bird | ╁ | | | 10 | | 1 | \exists | T | 1 | 1 | 7 | 1 | | \vdash | T | | | | | | | | | | | \top | + | \top |
| Pachycephalidae | Colluricincla harmonica | Grey Shrike-thrush | X | | | | | 1 | \dashv | 1 | 5 | 8 | 1 | \dashv | $\neg \vdash$ | \top | † | | | | 2 | 5 | | 1 | 1 | 1 | $\neg \vdash$ | 十 | T | \forall |
| , | Pachycephala pectoralis | Golden Whistler | 1 | | | | | 1 | T | T | _ | _ | 9 | | | 1 | 1 | | | | 1 | | | | T | T | | \top | T | \Box |
| | Pachycephala rufiventris | Rufous Whistler | X | | | | | | T | T | T | T | T | T | | Ť | 1 | | | | 10 | 3 | | | T | 19 | | 十 | T | \top |
| Pardalotidae | Pardalotus punctatus | Spotted Pardalote | X | | | | | 1 | T | T | T | T | T | T | | 1 | 1 | | | | | 4 | | | T | T | | \top | T | \top |
| ** ** * | Pardalotus striatus | Striated Pardalote | X | | | | | 1 | T | T | | 1 | T | | | 1 | 1 | | | | 5 | 8 | | | T | 4 | | 1 | T | \Box |
| Petroicidae | Eopsaltria georgiana | White-breasted Robin | X | | | | | 1 | T | T | T | T | T | 寸 | 1 | 1 | T | | | | | | | | T | 1 | 1 | \top | T | \Box |
| | Petroica boodang | Scarlet Robin | X | | | | | 1 | T | T | | T | T | | | 1 | 1 | | | | | 6 | | | T | T | | 十 | T | \top |

| | | Surveys | A | \mathbf{R} | C | ۲ | D | E | F | | G | 1 | | | | | Η | | | | | T | | J | | I | T | | T. | | \mathbf{M} |
|------------------|------------------------------|----------------------------------|----|--------------|----------|---------|---|---|---------|--------|----------|--------------------|---------|---------|----------------|----------|----------|----------|----------|----------|-------------|--------|--------|--------|---------------|---------------|----------|---------------|--------------|-----------|--------------|
| | | Sur veys | 21 | | Ì | _ | | L | _ | | Ĭ | | | | | | 1 | | | | | _ | | Ť | | _ | | | | Ħ | |
| Family | Species | Common Name | | | Brighton | Unknown | | | Trinity | Acacia | Dryandra | Eucalypt & Banksia | Banksia | Site 01 | Site 02 | Site 10B | Site 02B | Site 02A | Site 10A | Site 01A | Site 14A | Site 4 | Site 2 | Site 1 | Opportunistic | Opportunistic | Site 5 | Cassilda Park | Kinsale Park | Brighton | |
| Rhipiduridae | Rhipidura albiscapa | Grey Fantail | X | | | | | | | | | 4 | 1 | - | - | | | | | | | | 5 | 3 | | 34 | - | 1 | | П | |
| | Rhipidura leucophrys | Willie Wagtail | X | | | 1 | | | | | 1 | | 1 | | | | | | | | | | 1 | 1 | | 11 | | 1 | 1 | | |
| Timaliidae | Zosterops lateralis | Silvereye | X | | | | | | | 62 | 70 | 45 | 11 | | | | | | | | | | 4 | 4 | | 321 | | 1 | 1 | 1 | |
| Cacatuidae | Cacatua sanguinea | Little Corella | | | | | | | | 2 | | | | | | | | | | | | 2 | | | 1 | 14 | | | | | |
| | Calyptorhynchus latirostris | Carnaby's Cockatoo | | | | | | | | | 2 | 1 | 8 | | | | | | | | | 3 4 | 1 1 | | | 240 | | | | | |
| | Eolophus roseicapilla | Galah | | | | | | | | | 2 | 12 | 4 | | | | | | | | | 8 1 | 2 | 5 | 1 | 116 | | 1 | 1 | | |
| Psittacidae | Barnardius zonarius | Australian Ringneck | | | | | | | | | | 14 | | | | | | | | | | 4 1 | 9 | | 1 | 67 | | | 1 | | |
| | Neophema elegans | Elegant Parrot | | | | | | | | | | | | | 1 | + | 1 | 1 | 1 | 1 | 1 | | _ | | | | | | 7 | \exists | \exists |
| | Trichoglossus haematodus | Rainbow Lorikeet | | T | T | T | | | | T | | | | T | 1 | + | 1 | + | 十 | 1 | 7 | 2 | 1 | | 1 | | | | 7 | 寸 | \neg |
| Strigidae | Ninox boobook | Southern Boobook | | | | | | | | | | | | 1 | $\neg \dagger$ | \dashv | 7 | 1 | \dashv | 1 | \dashv | - 1 | 2 | 1 | | | | 1 | 寸 | 寸 | \dashv |
| Fish | | | | T | | | | | | T | | 1 | | T | | 1 | | | 1 | 1 | \dashv | T | 1 | | 1 | | | | 1 | 寸 | \neg |
| Percichthyidae | Bostockia porosa | Nightfish | X | T | | | | | | T | | 1 | | T | | 1 | | | 1 | 1 | \dashv | 1 | 1 | 1 | | | | | | 寸 | \neg |
| Amphibians | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Hylidae | Litoria moorei | Motorbike Frog | X | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Limnodynastidae | Heleioporus eyrei | Moaning Frog | X | 8 | | 40 | | | 1 | | | 1 | 17 | | | 1 (| 6 1 | 2 | | 7 | 13 | 3 | 3 | | | | 3 | | | | |
| | Limnodynastes dorsalis | Western Banjo Frog | X | 15 | 30 | 75 | | | | 4 | 23 | 1 | 2 | | | 2 | 1 . | 5 | 1 | 4 | 2 | | ı | 1 | | | 16 | | | | |
| Myobatrachidae | Crinia insignifera | Squelching Froglet | X | 7 | | 35 | | | | | | | | | | | | | | | | | | | | | | | | | |
| <u></u> | Myobatrachus gouldii | Turtle Frog | | | | | | | | | | | | | | | | | | | | 1 | 2 | | | | | | | | |
| | Pseudophryne guentheri | Gunther's Toadlet | X | 1 | | 1 | | | | | | | | | | | 1 | | 1 | | | | | | | | | | | \neg | |
| Mammals | - zemapmyne garamer | | | | | | | | | | | | | | | | 1 | | 1 | | | | | | | | | | | | = |
| Canidae | Vulpes vulpes | Red Fox | | | | | | | 1 | | | | 1 | | | | | | | | | | 1 | 1 | 1 | | | | | | |
| Felidae | Felis catus | Cat | | | | | | | | | | | | | | | 1 | | 1 | | | | Ť | 1 | 1 | | | | | \neg | |
| Dasvuridae | Dasyurus geoffroii | Chuditch | X | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 usy urrane | Sminthopsis fuliginosus | Grey-bellied Dunnart | | Ť | | | | | | | | | | | | | 1 | | 1 | | | | 1 | | | | | | | \neg | |
| Macropodidae | Macropus fuliginosus | Western Grey Kangaroo | X | 1 | | | | | | 6 | 1 | 3 | | | | | 1 | | 1 | | | 8 | 3 1 | | 1 | | | | | | = |
| | Notamacropus irma | Western Brush Wallaby | | | | | | | | | | - | | | | | | | \dashv | | \dashv | | _ | | Ť | | | | | \dashv | \dashv |
| Tarsipedidae | Tarsipes rostratus | Honey Possum | X | 5 | | 5 | | | | | 16 | | | | | \top | 十 | \dashv | \dashv | \dashv | \dashv | T | 1 | 1 | | | 3 | | 7 | 寸 | \exists |
| Leporidae | Oryctolagus cuniculus | Rabbit | | Ť | | | | | | | | | | | | \top | T | 1 | \dashv | 1 | $\exists t$ | | 1 | 1 | 1 | | | | 1 | 寸 | \exists |
| Peramelidae | Isoodon obesulus | Quenda | Х | | | | | | 1 | | | | | | | \top | T | 1 | \dashv | 1 | \dashv | Ť | Ť | Ť | Ė | t | | | \exists | 寸 | \exists |
| Muridae | Mus musculus | House Mouse | Ť | | 10 | | | | 98 | 13 | 17 | 7 | 4 | 1 | 1 | 2 | 1 | 4 | 5 | 1 | 1 | - 1 | 2 | 6 | | | 5 | | 寸 | 寸 | \dashv |
| | Rattus fuscipes | Bush Rat | X | 6 | | 1 | 4 | 1 | 6 | | 1 | | | 1 | \exists | \dashv | T | _ | 1 | 1 | \dashv | T | \top | 3 | | | Ħ | | 寸 | 寸 | \dashv |
| | Rattus rattus | Black Rat | | Ť | T | T | | | 2 | T | | | | T | 1 | + | 1 | + | 十 | 1 | 7 | 1 | 1 | 2 | 1 | | | | 7 | 寸 | \neg |
| Vespertilionidae | Vespadelus regulus | Southern Forest Bat | X | 1 | | | 1 | | | | | | | 1 | $\neg \dagger$ | \dashv | 7 | 1 | \dashv | 1 | \dashv | | l | 1 | 1 | | | | 寸 | 寸 | \dashv |
| Reptiles | | | | | | | | | | | | | | 1 | $\neg \dagger$ | \dashv | 7 | 1 | \dashv | 1 | \dashv | T | \top | 1 | | | | | 寸 | 寸 | \dashv |
| Agamidae | Ctenophorus adelaidensis | Western Heath Dragon | X | 7 | | 35 | | | | | | | | | | 4 | 5 | 1 | \dashv | 1 | $\exists t$ | \top | \top | 1 | | | | | 1 | 寸 | \exists |
| | Pogona minor | Dwarf Bearded Dragon | X | 8 | | 8 | | | 35 | 2 | | 1 | 3 | 1 | $\neg \dagger$ | | _ | 9 | \dashv | 1 | \dashv | \top | \top | 1 | | | 3 | | 寸 | 寸 | \dashv |
| Diplodactylidae | Crenadactylus ocellatus | Clawless Gecko | X | 3 | | | | | | | | | | | | | T | | T | T | T | 2 | 2 | | | | | | T | 寸 | 1 |
| | Diplodactylus polyophthalmus | Speckled Stone Gecko | X | 2 | | 10 | | | | | | | | | | T | T | T | T | T | T | T | | İ | | | | | T | \exists | 1 |
| | Strophurus elderi | Jewelled Gecko | | | | | | | | 1 | | | | | | T | T | T | T | T | T | T | | | | | | | T | 寸 | 1 |
| | Strophurus spinigerus | South-western Spiny-tailed Gecko | X | 4 | | 4 | | | 81 | 8 | 4 | | | | 1 | 4 | 5 | 7 | 1 | 1 | 1 | - 2 | 2 | | | | | | 7 | \exists | \exists |
| Elapidae | Brachvurophis fasciolata | Narrow-banded Burrowing Snake | X | | | | | | | T | | 1 | | T | | 1 | | | 1 | 1 | \dashv | 1 | 1 | 1 | | | | | | 寸 | \neg |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

| | | Surveys | A | В | (| 3 | D | E | F | | G | | | | |] | H | | | | I | | J | | | K | | I | | M |
|-------------|--|-------------------------------------|----|----|----------|---------|---|----------|---------|----------|---|--------------------|----------|---------|-----------------------|----------|----------|--|----------|----------|----------|----------|----------|------------|---------------|---------------|----------------|-------------------------------|----------|--------|
| Family | Species | Common Name | | | Brighton | Unknown | | | Frinity | Acacia | _ | Eucalypt & Banksia | Banksia | Site 01 | Site 02 Site 10B | Site 02B | Site 02A | Site 10A | Site 01A | Site 14A | Site 4 | Site 4 | Site 2 | Site 1 | Opportunistic | Opportunistic | Site 5 | Cassilda Park Kinsale Park | Brighton | |
| 1 uminy | Demansia psammophis | Yellow-faced Whipsnake | | | | 1 | | | | | | | 2 | 0, | <i>J</i> 2 <i>J</i> . | 2 | | | | | | 02 | | | | - | 1 | | - | + |
| | Echiopsis curta | Bardick | X | 3 | П | 15 | | | 8 | 1 | | | | | | | | | | | | | | | | _ | 2 | \dashv | + | + |
| | Neelaps bimaculatus | Black-naped Burrowing Snake | X | 2 | П | 10 | | | 3 | | | | | | | | | | | | | | | | | _ | 1 | \dashv | + | + |
| | Neelaps calonotus | Black-striped Burrowing Snake | X | 1 | П | | | | | | | | | | | | | | | | | | | | | | \neg | \top | \top | 1 |
| | Parasuta gouldii | Gould's Snake | X | 8 | П | 8 | | | 12 | 1 | | | | | | | 1 | | | | | | | T | 1 | _ | \dashv | + | + | \top |
| | Pseudonaja affinis | Dugite | X | 1 | 2 | 1 | | | 9 | Ť | 1 | 2 | | | | | | | | | | | 1 | 1 | | _ | \dashv | \dashv | + | + |
| | Pseudonaja mengdeni | Western Brown Snake | | | Ħ | | | | 1 | \dashv | - | - | \dashv | | - | | 1 | | | | H | 7 | - | Ť | _ | \dashv | + | + | + | + |
| | Simoselaps bertholdi | Jan's Banded Snake | X | 9 | Н | 9 | | | | 2 | 2 | - | 6 | | | 2 | 1 | | | | | | | - | | \dashv | 1 | + | + | + |
| | Simoselaps littoralis | West Coast Banded Snake | X | | Н | | | | 20 | - | _ | 1 | - | | - | Ť | Ť | | | | | + | + | 7 | | \dashv | - + | + | + | + |
| Gekkonidae | Christinus marmoratus | Marbled Gecko | X | 4 | Н | 20 | | | 10 | - | | \dashv | 1 | - | 1 | + | \vdash | | 4 | 2 | H | \dashv | - | 1 | + | + | + | + | + | + |
| Pygopodidae | Aprasia repens | Southwest Sandplain Worm Lizard | X | 4 | Н | 25 | | | | + | - | 1 | \dashv | | - | | | | Ė | <u> </u> | 2 | + | + | 1 | | \dashv | 1 | + | + | + |
| Тудорошише | Delma concinna | Javelin Lizard | X | | 5 | 20 | | | 6 | | | | | | | | | | | | Ť | | | - | | - | <u> </u> | + | + | + |
| | Delma fraseri | Fraser's Delma | X | 1 | Ť | 5 | | | 7 | + | | | 1 | | | | 1 | | | | | | 1 | - t | _ | \dashv | 1 | + | + | + |
| | Delma grayii | Side-barred Delma | X | 3 | \vdash | 15 | | | 22 | | | | _ | | | 1 | | | | | | | - | t | | -+ | ÷ | + | + | + |
| | Lialis burtonis | Burton's Legless Lizard | X | 4 | H | 20 | | | 57 | | | 1 | 1 | | 1 | 1 | 2 | | | | 6 | 1 | | =t | | + | + | + | + | + |
| | Pletholax gracilis | Keeled Legless Lizard | 21 | • | H | 20 | | | 6 | | | - | _ | | | | 1- | | | | Ü | - | | 1 | | + | + | + | + | + |
| | Pygopus lepidopodus | Common Scaly-foot | X | 2 | 1 | 2 | | | 22 | | 1 | _ | 1 | | | | | | | | | - | | - | | + | + | + | + | + |
| Pythonidae | Morelia spilota | Carpet Python | X | 1 | H | | | | | | - | _ | 1 | | | | | | | | | - | | =t | | + | + | + | + | + |
| Scincidae | Acritoscincus trilineatus | Western Three-lined Skink | X | 2 | H | 10 | | | | | | _ | 1 | | | | | | | 2 | | - | | =t | | + | + | + | + | + |
| Schicidae | Cryptoblepharus buchananii | Buchanan's Snake-eyed Skink | X | 3 | H | 15 | | | 19 | + | 2 | 2 | 5 | + | - | - | 1 | 1 | 8 | 8 | 5 | 2 | + | _ | 1 | + | + | + | + | + |
| | Ctenotus australis | Western Limestone Ctenotus | X | 5 | 10 | 25 | | | | 3 | _ | | 5 | | | 5 | - | 1 | 0 | 0 | 3 | | | =t | 1 | + | 12 | + | + | + |
| | Ctenotus dustratis Ctenotus fallens | West-coast Laterite Ctenotus | X | 5 | 20 | 25 | | | _ | _ | _ | _ | 16 | | | 2 | | | 3 | 2 | 2 | | | 1 | | | 22 | + | + | + |
| | Cyclodomorphus celatus | Western Slender Bluetongue | X | 6 | 20 | 30 | | | 31 | _ | _ | 1 | 10 | | | | 3 | | 3 | | 2 | - | | 1 | | | 2 | + | + | + |
| | Egernia kingii | King's Skink | 71 | U | H | 50 | | | | | | 1 | 1 | | | | | | | | | - | | 1 | | + | | + | + | + |
| | Egernia napoleonis | Southwestern Crevice Skink | X | 2 | 5 | 10 | | | 20 | | | 3 | | | | | | | | | | | | - | | + | + | + | + | + |
| | Hemiergis quadrilineatum | Two-toed Earless Skink | X | 8 | | | | | 123 | 18 | | 32 | 13 | + | 7 | 3 | 11 | 5 | 20 | 3 | 63 | 3 | 1 | 1 | - | + | 2 | + | + | + |
| | Lerista distinguenda | South-western Orange-tailed Slider | 71 | U | 7.5 | 70 | | | 1 | 10 | | 32 | 7.5 | | | | 11 | 5 | 20 | 3 | 4 | 5 | 1 | 1 | | + | | + | + | + |
| | Lerista distinguenda Lerista elegans | West Coast Four-toed Lerista | v | 11 | 20 | 55 | | | 1 | + | + | - | + | + | - | 9 | 18 | | 4 | 48 | | 2 | 2 | 2 | - | + | 1 | + | + | + |
| | Lerista lineopunctulata | Dotted-line Robust Slider | X | 4 | 20 | 20 | | | 2 | + | + | - | + | + | - | 1 | 10 | | 7 | 40 | 2 | | | _ | - | + | + | + | + | + |
| | Lerista uneopunctutata Lerista praepedita | Blunt-tailed West-coast Slider | X | 13 | Н | 65 | | H | 11 | \dashv | - | \dashv | \dashv | - | + | 1 | 1 | | | | 2 | 1 | \dashv | ┪ | \dashv | + | + | + | + | 十 |
| | Menetia greyii | Common Dwarf Skink | X | 5 | 5 | 25 | | | | 2 | 2 | 4 | 3 | - | 2 | 1 | 3 | 2 | 24 | 57 | 5 | - | 6 | \dashv | - | + | 2 | + | + | + |
| | Morethia lineoocellata | Pale-flecked Morethia | Λ | J | ر | 23 | | | 8 | _ | _ | + | 3 | - | + 2 | 1 | 3 | | 24 | 31 | 1 | 1 | 1 | \dashv | - | + | | + | + | + |
| | Morethia imeooceiiata Morethia obscura | Shrubland Pale-flecked Morethia | X | 2 | 20 | 10 | | | 40 | + | 3 | 3 | 4 | | - | 2 | 4 | 2 | 6 | 21 | | 1 | 1 | \dashv | \dashv | + | + | + | + | + |
| | Tiliqua occipitalis | Western Blue-tongued Lizard | Λ | | 20 | 10 | | | 7 | - | 3 | _ | 2 | | 2 | - 2 | 1 | | 0 | 21 | 0 | - | 4 | 2 | 1 | + | 9 | + | + | + |
| | Tiliqua occipitatis Tiliqua rugosa | Bobtail | X | 1 | 2 | | | \vdash | | 2 | 3 | | _ | | 16 1 | 1 | <u> </u> | - | | - | 1 | _ | 4 | _ | 1 | | 4 | + | + | + |
| Typhlopidae | Anilios australis | Austral Blind Snake | X | 1 | \vdash | 2 | | \vdash | 46 | _ | J | 3 | 1 | 7 | 10 1 | 1 | 4 | | 1 | | 1 | \dashv | \dashv | \dashv | \dashv | \dashv | + | + | + | + |
| туршориае | Anilios austraiis Anilios pinguis | Rotund Blind Snake | Λ | 1 | Н | | | | 2 | + | - | + | 1 | | - | - | 1 | | 1 | - | H | + | + | \dashv | \dashv | + | + | + | + | + |
| Varanidas | Varanus gouldii | | | | Н | | | | 1 | + | + | + | + | - | - | | 1 | _ | 1 | | \vdash | \dashv | + | \dashv | + | + | + | + | + | + |
| Varanidae | Varanus gouiau Varanus tristis | Gould's Goanna Black-headed Monitor | X | 1 | Н | 1 | | | 1 | + | - | + | + | | | | 1 | _ | 1 | | | + | + | - | + | + | + | + | + | + |
| Chalidaa | | | Λ | 1 | Н | 5 | | \vdash | | - | - | \dashv | + | + | - | - | ╁ | - | 1 | | \vdash | + | - | - | + | + | + | + | + | + |
| Chelidae | Chelodina colliei | Collie's Snake-necked Turtle | | 1 | ш | J | | | | | | | | | | <u> </u> | 1 | <u> </u> | <u> </u> | <u> </u> | Ш | | | | | \perp | L | | — | Щ |

Atlas of Living Australia Western Australian Museum A

В

- C NatureMap Brighton
- D NatureMap Gnangara
- E NatureMap Mitchell Freeway
- F Terrestrial Ecosystems (2012) Trinity
- G ATA Environmental (2008) Vertebrate Fauna Assessment Lot 3 Romeo Road, Alkimos, Unpublished report for Northern Corridor Developments Limited.
- H Valentine, I.E., Wilson, B.A., Reaveley, A., Huang, N., Johnson, B. and Brown, P.R. (2009) *Patterns of Ground-dwelling Vertebrate Biodiversity in the Gnangara Sustainability Strategy Study Area*, Unpublished report for the Department of Environment and Conservation, Perth.
- I GHD (2014b) Neerabup Road Extension Level 2 Fauna Survey, Unpublished report for Main Roads Western Australia, Perth
- J ATA Environmental (1991) Yanchep Structure Plan Vertebrate Fauna Survey, Unpublished report for Tokyu Corporation, Perth.
- K ATA Environmental (2007) Vertebrate Fauna Assessment St Andrews Estate (Southern Precinct), Yanchep, Unpublished report for Yanchep Sun City Pty Ltd, Perth.
- L Gole, C.A. (2003) Bird Survey in selected Perth Metropolitan Reserves. A Joint Biodiversity Conservation Project between Birds Australia WA and Perth Biodiversity Project, Unpublished report Birds Australia and Perth Biodiversity Project, Perth.
- M Bamford Consulting Ecologists (2005) Alkimos Proposed Wastewater Treatment Plant: Fauna Assessment, Unpublished report for Water Corporation, Perth.

Appendix B(2). Vertebrate fauna assessments

| rippendix D(2) | . Vertebrate fauna asses | Surve | v | | | A | | | |
|----------------------|--|---------------------------|--------|--------|--------|--------|--------|--------|-------------|
| Family Amphibians | Species | Common Name | Site 3 | Site 6 | Site 5 | Site 4 | Site 2 | Site 1 | Opportunist |
| Limnodynastidae | Heleioporus eyrei | Moaning Frog | 24 | 12 | 13 | 27 | 9 | 5 | |
| Lillillodyllastidae | Limnodynastes dorsalis | Western Banjo Frog | 24 | 12 | 13 | 4 | 9 | 3 | |
| Mysobotusobidos | | Ouacking Frog | - | | | 4 | | | |
| Myobatrachidae | Crinia georgiana Crinia insignifera | Squelching Froglet | - | | | 1 | | | |
| | 0 2 | Gunther's Toadlet | - | | | 1 | | | |
| Birds | Pseudophryne guentheri | Guittier's Toadlet | - | | | 1 | | | |
| Accipitridae | A a simit on faces interes | Brown Goshawk | 1 | | | | | | |
| Accipitridae | Accipiter fasciatus | | 1 | | | | | | |
| | Elanus axillaris | Black-shouldered Kite | 1 | | | | 1 | | |
| A .: 1 | Hieraaetus morphnoides | Little Eagle | | | | 2 | 1 | | |
| Anatidae | Anas superciliosa | Pacific Black Duck | - | | | 2 | | | _ |
| Ardeidae | Egretta novaehollandiae | White-faced Heron | 1 | | | | | | 1 |
| Threskiornithidae | Threskiornis molucca | Australian White Ibis | 6 | | | 1 | | | |
| Columbidae | Phaps chalcoptera | Common Bronzewing | 1 | | | 1 | | | |
| | Spilopelia senegalensis | Laughing Turtle-dove | | | | 2 | | | |
| Halcyonidae | Dacelo novaeguineae | Laughing Kookaburra | | 1 | _ | 7 | 2 | _ | |
| | Todiramphus sanctus | Sacred Kingfisher | | 1 | 1 | 6 | 3 | 1 | |
| Meropidae | Merops ornatus | Rainbow Bee-eater | | | | 1 | 8 | 1 | 1 |
| Cuculidae | Chalcites basalis | Horsfield's Bronze-cuckoo | | | | | 1 | | |
| | Chalcites lucidus | Shining Bronze-cuckoo | | 3 | | | | | |
| | Heteroscenes pallidus | Pallid Cuckoo | | | | | | 1 | |
| Falconidae | Falco berigora | Brown Falcon | | 1 | | | | | |
| Acanthizidae | Acanthiza apicalis | Inland Thornbill | | | | 4 | | | |
| | Acanthiza chrysorrhoa | Yellow-rumped Thornbill | | 10 | | 25 | | 9 | |
| | Acanthiza inornata | Western Thornbill | | 9 | | | 17 | | |
| | Gerygone fusca | Western Gerygone | 20 | 24 | 10 | 6 | 47 | 25 | 2 |
| | Smicrornis brevirostris | Weebill | 8 | 14 | 1 | | 12 | 3 | |
| Artamidae | Cracticus torquatus | Grey Butcherbird | 2 | 8 | 8 | 7 | 4 | 2 | 1 |
| | Gymnorhina tibicen | Australian Magpie | 9 | | 6 | 21 | 6 | 11 | 3 |
| Campephagidae | Coracina novaehollandiae | Black-faced Cuckoo-shrike | | 7 | 2 | 1 | 5 | 11 | 1 |
| Corvidae | Corvus coronoides | Australian Raven | 3 | 5 | 8 | 6 | 14 | 22 | 1 |
| Maluridae | Malurus splendens | Splendid Fairy-wren | | | | 5 | 14 | | |
| Meliphagidae | Anthochaera carunculata | Red Wattlebird | 2 | 4 | 10 | 14 | 13 | 12 | 1 |
| | Gavicalis virescens | Singing Honeyeater | | 6 | 7 | 4 | 7 | 6 | |
| | Lichmera indistincta | Brown Honeyeater | | | | 3 | 5 | 9 | |
| | Phylidonyris niger | White-cheeked Honeyeater | | | 6 | 3 | 4 | 1 | |
| Monarchidae | Grallina cyanoleuca | Magpie-lark | | 2 | 2 | 3 | 1 | 1 | |
| Pachycephalidae | Colluricincla harmonica | Grey Shrike-thrush | | | | | 2 | | |
| | Pachycephala pectoralis | Golden Whistler | 2 | 1 | | | | | |
| | Pachycephala rufiventris | Rufous Whistler | 2 | 6 | 1 | 3 | 5 | 1 | 1 |
| Pardalotidae | Pardalotus striatus | Striated Pardalote | 1 | | 4 | | 3 | 4 | |
| Petroicidae | Petroica boodang | Scarlet Robin | | | | | 1 | | |
| Rhipiduridae | Rhipidura albiscapa | Grey Fantail | 2 | 2 | 1 | 50 | 11 | 3 | 1 |
| Timaliidae | Zosterops lateralis | Silvereye | 4 | 8 | 15 | 34 | 90 | 26 | |

| | | Survey | | | | A | | | |
|------------------|-----------------------------|---------------------------------|--------|--------|--------|----|--------|--------|-------------|
| Family | Species | Common Name | Site 3 | Site 6 | Site 5 | 4 | Site 2 | Site 1 | Opportunist |
| Cacatuidae | Cacatua sanguinea | Little Corella | | 91 | | 9, | 5 | | |
| | Calyptorhynchus latirostris | Carnaby's Cockatoo | 3 | | 2 | | 6 | | |
| | Eolophus roseicapilla | Galah | | 5 | 6 | 2 | 13 | 15 | |
| Psittacidae | Barnardius zonarius | Australian Ringneck | 3 | 7 | 26 | 33 | 33 | 36 | 1 |
| | Purpureicephalus spurius | Red-capped Parrot | | 1 | | 1 | | | |
| | Trichoglossus haematodus | Rainbow Lorikeet | | | | | 4 | 10 | |
| Strigidae | Ninox boobook | Southern Boobook | | | | | 1 | | |
| Molossidae | Austronomus australis | White-striped Freetail Bat | | | 1 | | | | |
| Vespertilionidae | Chalinolobus gouldii | Gould's Wattled Bat | | 1 | | | | | |
| Mammals | | | | | | | | | |
| Macropodidae | Macropus fuliginosus | Western Grey Kangaroo | 2 | 1 | 1 | 11 | 5 | 2 | |
| Phalangeridae | Trichosurus vulpecula | Common Brushtail Possum | 4 | 1 | | | 4 | | |
| Leporidae | Oryctolagus cuniculus | Rabbit | | | | 1 | 1 | | |
| Peramelidae | Isoodon obesulus | Southern Brown Bandicoot | | | 1 | | | | |
| Equidae | Equus caballus | Horse | | | | | | 1 | |
| Muridae | Mus musculus | House Mouse | | | 3 | | 1 | 8 | |
| Reptiles | | | | | | | | | |
| Agamidae | Pogona minor | Western Bearded Dragon | | 1 | 2 | | 1 | 1 | |
| Diplodactylidae | Oedura marmorata | Marbled Velvet Gecko | | | | | 1 | 1 | |
| Elapidae | Neelaps bimaculatus | Black-naped Burrowing Snake | 1 | | | | | | |
| | Pseudonaja affinis | Dugite | 1 | | | | 1 | | |
| | Simoselaps bertholdi | Jan's Banded Snake | | | | | 1 | | |
| Pygopodidae | Aprasia repens | Southwest Sandplain Worm Lizard | 1 | 4 | | 1 | | 2 | |
| | Lialis burtonis | Burton's Legless Lizard | 4 | 2 | 9 | | 10 | 13 | |
| | Pletholax gracilis | Keeled Legless Lizard | | | | | | 1 | |
| Pythonidae | Morelia spilota | Carpet Python | | | | | 2 | | |
| Scincidae | Acritoscincus trilineatus | Western Three-lined Skink | | 1 | | 3 | | | |
| | Cryptoblepharus buchananii | Buchanan's Snake-eyed Skink | 4 | 8 | 5 | | 7 | 3 | |
| | Ctenotus australis | Western Limestone Ctenotus | | | 2 | | 1 | 2 | |
| | Ctenotus fallens | West-coast Laterite Ctenotus | | | 1 | | 7 | 8 | |
| | Cyclodomorphus celatus | Western Slender Bluetongue | | | | | 1 | | |
| | Hemiergis quadrilineatum | Two-toed Earless Skink | 14 | 15 | 9 | 4 | 37 | 28 | |
| | Lerista elegans | West Coast Four-toed Lerista | 18 | 7 | 6 | | 5 | 1 | |
| | Lerista praepedita | Blunt-tailed West-coast Slider | | | 1 | | | 1 | |
| | Menetia greyii | Common Dwarf Skink | 18 | 6 | 2 | 20 | 1 | 3 | |
| | Morethia lineoocellata | Pale-flecked Morethia | | 3 | 1 | | 4 | 2 | |
| | Morethia obscura | Shrubland Pale-flecked Morethia | 5 | 3 | 3 | | 2 | 1 | |
| | Tiliqua rugosa | Bobtail | 3 | 3 | 3 | 4 | 7 | 7 | |
| Typhlopidae | Anilios australis | Austral Blind Snake | 1 | | | | | | |
| Varanidae | Varanus tristis | Black-headed Monitor | 1 | | | | | | |

A Biota Environmental Sciences (2000) Lot 52 Burns Beach Road Fauna Survey, Perth.

Appendix C Definitions of Significant Fauna under the WA *Biodiversity Conservation Act 2016* and Priority Species

Vertebrate Fauna Survey - Yanchep Lagoon, Yanchep

ATTACHMENT C DEFINITIONS OF SIGNIFICANT FAUNA UNDER THE WA BIODIVERSITY CONSERVATION ACT 2016

Threatened, Extinct and Specially Protected fauna or flora¹ are species² which have been adequately searched for and are deemed to be, in the wild, threatened, extinct or in need of special protection, and have been gazetted as such. The *Wildlife Conservation (Specially Protected Fauna) Notice 2018* and the *Wildlife Conservation (Rare Flora) Notice 2018* have been transitioned under regulations 170, 171 and 172 of the *Biodiversity Conservation Regulations 2018* to be the lists of Threatened, Extinct and Specially Protected species under Part 2 of the *Biodiversity Conservation Act 2016*. Categories of Threatened, Extinct and Specially Protected fauna and flora are:

T Threatened Species

Listed by order of the Minister as Threatened in the category of critically endangered, endangered or vulnerable under section 19(1), or is a rediscovered species to be regarded as threatened species under section 26(2) of the *Biodiversity Conservation Act 2016* (BC Act).

Threatened fauna is that subset of 'Specially Protected Fauna' listed under schedules 1 to 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for Threatened Fauna.

Threatened flora is that subset of 'Rare Flora' listed under schedules 1 to 3 of the *Wildlife Conservation (Rare Flora) Notice 2018* for Threatened Flora.

The assessment of the conservation status of these species is based on their national extent and ranked according to their level of threat using IUCN Red List categories and criteria as detailed below.

CR Critically endangered species

Threatened species considered to be "facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with criteria set out in the ministerial guidelines".

Listed as critically endangered under section 19(1)(a) of the BC Act in accordance with the criteria set out in section 20 and the ministerial guidelines. Published under schedule 1 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for critically endangered fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for critically endangered flora.

EN Endangered species

Threatened species considered to be "facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines".

Listed as endangered under section 19(1)(b) of the BC Act in accordance with the criteria set out in section 21 and the ministerial guidelines. Published under schedule 2 of the *Wildlife Conservation (Specially Protected Fauna) Notice* 2018 for endangered fauna or the *Wildlife Conservation (Rare Flora) Notice* 2018 for endangered flora.

VU Vulnerable species

Threatened species considered to be "facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with criteria set out in the ministerial guidelines".

Listed as vulnerable under section 19(1)(c) of the BC Act in accordance with the criteria set out in section 22 and the ministerial guidelines. Published under schedule 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice* 2018 for vulnerable fluora.

Extinct Species

¹ The definition of flora includes algae, fungi and lichens

² Species includes all taxa (plural of taxon - a classificatory group of any taxonomic rank, e.g. a family, genus, species or any infraspecific category i.e. subspecies or variety, or a distinct population).

Listed by order of the Minister as extinct under section 23(1) of the BC Act as extinct or extinct in the wild.

EX Extinct species

Species where "there is no reasonable doubt that the last member of the species has died", and listing is otherwise in accordance with the ministerial guidelines (section 24 of the BC Act).

Published as presumed extinct under schedule 4 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018 for extinct fauna or the Wildlife Conservation (Rare Flora) Notice 2018 for extinct flora.

EW Extinct in the wild species

Species that "is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; and it has not been recorded in its known habitat or expected habitat, at appropriate seasons, anywhere in its past range, despite surveys over a time frame appropriate to its life cycle and form", and listing is otherwise in accordance with the ministerial guidelines (section 25 of the BC Act).

Currently there are no threatened fauna or threatened flora species listed as extinct in the wild. If listing of a species as extinct in the wild occurs, then a schedule will be added to the applicable notice.

Specially Protected Species

Listed by order of the Minister as specially protected under section 13(1) of the BC Act. Meeting one or more of the following categories: species of special conservation interest; migratory species; cetaceans; species subject to international agreement; or species otherwise in need of special protection.

Species that are listed as threatened species (critically endangered, endangered or vulnerable) or extinct species under the BC Act cannot also be listed as Specially Protected species.

MI Migratory birds protected under an international agreement

Fauna that periodically or occasionally visit Australia or an external Territory or the exclusive economic zone; or the species is subject of an international agreement that relates to the protection of migratory species and that binds the Commonwealth; and listing is otherwise in accordance with the ministerial guidelines (section 15 of the BC Act).

Includes birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and fauna subject to the *Convention on the Conservation of Migratory Species of Wild Animals* (Bonn Convention), an environmental treaty under the United Nations Environment Program. Migratory species listed under the BC Act are a subset of the migratory animals, that are known to visit Western Australia, protected under the international agreements or treaties, excluding species that are listed as Threatened species.

Published as migratory birds protected under an international agreement under schedule 5 of the *Wildlife Conservation* (Specially Protected Fauna) Notice 2018.

CD Species of special conservation interest (conservation dependant fauna)

Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened, and listing is otherwise in accordance with the ministerial guidelines (section 14 of the BC Act).

Published as conservation dependent fauna under schedule 6 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018.

OS Other specially protected species

Fauna otherwise in need of special protection to ensure their conservation, and listing is otherwise in accordance with the ministerial guidelines (section 18 of the BC Act).

Published as other specially protected fauna under schedule 7 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018.*

Priority species

Possibly threatened species that do not meet survey criteria, or are otherwise data deficient, are added to the Priority Fauna or Priority Flora Lists under Priorities 1, 2 or 3. These three categories are ranked in order of priority for survey and evaluation of conservation status so that consideration can be given to their declaration as threatened fauna or flora.

Species that are adequately known, are rare but not threatened, or meet criteria for near threatened, or that have been recently removed from the threatened species or other specially protected fauna lists for other than taxonomic reasons, are placed in Priority 4. These species require regular monitoring.

Assessment of Priority codes is based on the Western Australian distribution of the species, unless the distribution in WA is part of a contiguous population extending into adjacent States, as defined by the known spread of locations

P1 Priority 1: Poorly-known species

Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.

P2 Priority 2: Poorly-known species

Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.

P3 Priority 3: Poorly-known species

Species that are known from several locations, and the species does not appear to be under imminent threat, or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. Such species are in need of further survey.

P4 Priority 4: Rare, Near Threatened and other species in need of monitoring

- (a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection but could be if present circumstances change. These species are usually represented on conservation lands.
- (b) Near Threatened. Species that are considered to have been adequately surveyed and that are close to qualifying for vulnerable but are not listed as Conservation Dependent.
- (c) Species that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.

Appendix D Fauna habitat assessment results

Vertebrate Fauna Survey – Yanchep Lagoon, Yanchep

Date: 29-Aug-19 Habitat Assessment #: Y1 Observers: Dr Scott Thompson

Zone: 50 Easting: 369503 mE Northing: 6508473 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y2 Observers: Dr Scott Thompson

Zone: 50 Easting: 369480 mE Northing: 6508482 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y3 Observers: Dr Scott Thompson

Zone: 50 Easting: 369530 mE Northing: 6508434 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone






Date: 29-Aug-19 Habitat Assessment #: Y4 Observers: Dr Scott Thompson

Zone: 50 Easting: 369555 mE Northing: 6508417 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y5 Observers: Dr Scott Thompson

Zone: 50 Easting: 369580 mE Northing: 6508380 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y6 Observers: Dr Scott Thompson

Zone: 50 Easting: 369612 mE Northing: 6508407 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y7 Observers: Dr Scott Thompson

Zone: 50 Easting: 369631 mE Northing: 6508411 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y8 Observers: Dr Scott Thompson

Zone: 50 Easting: 369662 mE Northing: 6508429 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone

Habitat Structure: Mixed open and closed shrubland and heath on sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y9 Observers: Dr Scott Thompson

Zone: 50 Easting: 369698 mE Northing: 6508400 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone

Habitat Structure: Mixed open and closed shrubland and heath on sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y10 Observers: Dr Scott Thompson

Zone: 50 Easting: 369729 mE Northing: 6508385 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y11 Observers: Dr Scott Thompson

Zone: 50 Easting: 369712 mE Northing: 6508438 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y12 Observers: Dr Scott Thompson

Zone: 50 Easting: 369711 mE Northing: 6508475 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y13 Observers: Dr Scott Thompson

Zone: 50 Easting: 369676 mE Northing: 6508506 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y14 Observers: Dr Scott Thompson

Zone: 50 Easting: 369650 mE Northing: 6508521 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y15 Observers: Dr Scott Thompson

Zone: 50 Easting: 369619 mE Northing: 6508489 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y16 Observers: Dr Scott Thompson

Zone: 50 Easting: 369689 mE Northing: 6508555 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y17 Observers: Dr Scott Thompson

Zone: 50 Easting: 369719 mE Northing: 6508578 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y18 Observers: Dr Scott Thompson

Zone: 50 Easting: 369752 mE Northing: 6508607 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y19 Observers: Dr Scott Thompson

Zone: 50 Easting: 369747 mE Northing: 6508638 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y20 Observers: Dr Scott Thompson

Zone: 50 Easting: 369710 mE Northing: 6508626 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y21 Observers: Dr Scott Thompson

Zone: 50 Easting: 369685 mE Northing: 6508594 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y22 Observers: Dr Scott Thompson

Zone: 50 Easting: 369650 mE Northing: 6508563 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y23 Observers: Dr Scott Thompson

Zone: 50 Easting: 369625 mE Northing: 6508561 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Grass and garden

Habitat Structure: Highly disturbed, cleared habitat or planted vegetation





Date: 29-Aug-19 Habitat Assessment #: Y24 Observers: Dr Scott Thompson

Zone: 50 Easting: 369610 mE Northing: 6508591 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y25 Observers: Dr Scott Thompson

Zone: 50 Easting: 369631 mE Northing: 6508648 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y26 Observers: Dr Scott Thompson

Zone: 50 Easting: 369663 mE Northing: 6508633 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y27 Observers: Dr Scott Thompson

Zone: 50 Easting: 369694 mE Northing: 6508628 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y28 Observers: Dr Scott Thompson

Zone: 50 Easting: 369707 mE Northing: 6508670 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y29 Observers: Dr Scott Thompson

Zone: 50 Easting: 369687 mE Northing: 6508694 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y30 Observers: Dr Scott Thompson

Zone: 50 Easting: 369660 mE Northing: 6508679 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y31 Observers: Dr Scott Thompson

Zone: 50 Easting: 369621 mE Northing: 6508684 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y32 Observers: Dr Scott Thompson

Zone: 50 Easting: 369553 mE Northing: 6508677 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y33 Observers: Dr Scott Thompson

Zone: 50 Easting: 369562 mE Northing: 6508647 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y34 Observers: Dr Scott Thompson

Zone: 50 Easting: 369535 mE Northing: 6508620 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y35 Observers: Dr Scott Thompson

Zone: 50 Easting: 369508 mE Northing: 6508604 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y36 Observers: Dr Scott Thompson

Zone: 50 Easting: 369489 mE Northing: 6508589 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y37 Observers: Dr Scott Thompson

Zone: 50 Easting: 369500 mE Northing: 6508641 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y38 Observers: Dr Scott Thompson

Zone: 50 Easting: 369523 mE Northing: 6508682 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y39 Observers: Dr Scott Thompson

Zone: 50 Easting: 369505 mE Northing: 6508707 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y40 Observers: Dr Scott Thompson

Zone: 50 Easting: 369541 mE Northing: 6508720 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y41 Observers: Dr Scott Thompson

Zone: 50 Easting: 369608 mE Northing: 6508717 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y42 Observers: Dr Scott Thompson

Zone: 50 Easting: 369645 mE Northing: 6508723 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y43 Observers: Dr Scott Thompson

Zone: 50 Easting: 369682 mE Northing: 6508726 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y44 Observers: Dr Scott Thompson

Zone: 50 Easting: 369708 mE Northing: 6508739 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y45 Observers: Dr Scott Thompson

Zone: 50 Easting: 369709 mE Northing: 6508769 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y46 Observers: Dr Scott Thompson

Zone: 50 Easting: 369688 mE Northing: 6508775 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y47 Observers: Dr Scott Thompson

Zone: 50 Easting: 369657 mE Northing: 6508767 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y48 Observers: Dr Scott Thompson

Zone: 50 Easting: 369600 mE Northing: 6508737 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y49 Observers: Dr Scott Thompson

Zone: 50 Easting: 369551 mE Northing: 6508744 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y50 Observers: Dr Scott Thompson

Zone: 50 Easting: 369527 mE Northing: 6508766 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y51 Observers: Dr Scott Thompson

Zone: 50 Easting: 369505 mE Northing: 6508811 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y52 Observers: Dr Scott Thompson

Zone: 50 Easting: 369546 mE Northing: 6508795 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y53 Observers: Dr Scott Thompson

Zone: 50 Easting: 369607 mE Northing: 6508789 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y54 Observers: Dr Scott Thompson

Zone: 50 Easting: 369650 mE Northing: 6508794 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y55 Observers: Dr Scott Thompson

Zone: 50 Easting: 369692 mE Northing: 6508818 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y56 Observers: Dr Scott Thompson

Zone: 50 Easting: 369708 mE Northing: 6508858 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y57 Observers: Dr Scott Thompson

Zone: 50 Easting: 369707 mE Northing: 6508933 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y58 Observers: Dr Scott Thompson

Zone: 50 Easting: 369682 mE Northing: 6508881 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y59 Observers: Dr Scott Thompson

Zone: 50 Easting: 369673 mE Northing: 6508859 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y60 Observers: Dr Scott Thompson

Zone: 50 Easting: 369651 mE Northing: 6508848 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y61 Observers: Dr Scott Thompson

Zone: 50 Easting: 369600 mE Northing: 6508864 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y62 Observers: Dr Scott Thompson

Zone: 50 Easting: 369563 mE Northing: 6508872 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y63 Observers: Dr Scott Thompson

Zone: 50 Easting: 369523 mE Northing: 6508888 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y64 Observers: Dr Scott Thompson

Zone: 50 Easting: 369473 mE Northing: 6508882 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Garden

Habitat Structure: Highly disturbed, cleared habitat or planted vegetation





Date: 29-Aug-19 Habitat Assessment #: Y65 Observers: Dr Scott Thompson

Zone: 50 Easting: 369426 mE Northing: 6508912 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y66 Observers: Dr Scott Thompson

Zone: 50 Easting: 369402 mE Northing: 6508893 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y67 Observers: Dr Scott Thompson

Zone: 50 Easting: 369385 mE Northing: 6508871 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y68 Observers: Dr Scott Thompson

Zone: 50 Easting: 369350 mE Northing: 6508864 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y69 Observers: Dr Scott Thompson

Zone: 50 Easting: 369377 mE Northing: 6508830 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y70 Observers: Dr Scott Thompson

Zone: 50 Easting: 369351 mE Northing: 6508826 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y71 Observers: Dr Scott Thompson

Zone: 50 Easting: 369368 mE Northing: 6508786 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y72 Observers: Dr Scott Thompson

Zone: 50 Easting: 369391 mE Northing: 6508724 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y73 Observers: Dr Scott Thompson

Zone: 50 Easting: 369412 mE Northing: 6508696 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y74 Observers: Dr Scott Thompson

Zone: 50 Easting: 369412 mE Northing: 6508674 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y75 Observers: Dr Scott Thompson

Zone: 50 Easting: 369419 mE Northing: 6508644 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone






Date: 29-Aug-19 Habitat Assessment #: Y76 Observers: Dr Scott Thompson

Zone: 50 Easting: 369431 mE Northing: 6508723 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Garden

Habitat Structure: Highly disturbed, cleared habitat or planted vegetation





Date: 29-Aug-19 Habitat Assessment #: Y77 Observers: Dr Scott Thompson

Zone: 50 Easting: 369429 mE Northing: 6508638 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y78 Observers: Dr Scott Thompson

Zone: 50 Easting: 369436 mE Northing: 6508593 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y79 Observers: Dr Scott Thompson

Zone: 50 Easting: 369447 mE Northing: 6508544 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y80 Observers: Dr Scott Thompson

Zone: 50 Easting: 369485 mE Northing: 6508510 mN

Landform: Flat Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Garden

Habitat Structure: Highly disturbed, cleared habitat or planted vegetation





Date: 29-Aug-19 Habitat Assessment #: Y81 Observers: Dr Scott Thompson

Zone: 50 Easting: 369270 mE Northing: 6509252 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand

Habitat Structure: Highly disturbed, cleared habitat or planted vegetation





Date: 29-Aug-19 Habitat Assessment #: Y82 Observers: Dr Scott Thompson

Zone: 50 Easting: 369256 mE Northing: 6509268 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y83 Observers: Dr Scott Thompson

Zone: 50 Easting: 369222 mE Northing: 6509255 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y84 Observers: Dr Scott Thompson

Zone: 50 Easting: 369218 mE Northing: 6509203 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y85 Observers: Dr Scott Thompson

Zone: 50 Easting: 369185 mE Northing: 6509193 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y86 Observers: Dr Scott Thompson

Zone: 50 Easting: 369164 mE Northing: 6509181 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y87 Observers: Dr Scott Thompson

Zone: 50 Easting: 369110 mE Northing: 6509171 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y88 Observers: Dr Scott Thompson

Zone: 50 Easting: 369153 mE Northing: 6509133 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y89 Observers: Dr Scott Thompson

Zone: 50 Easting: 369202 mE Northing: 6509132 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y90 Observers: Dr Scott Thompson

Zone: 50 Easting: 369240 mE Northing: 6509161 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y91 Observers: Dr Scott Thompson

Zone: 50 Easting: 369230 mE Northing: 6509098 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y92 Observers: Dr Scott Thompson

Zone: 50 Easting: 369208 mE Northing: 6509064 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y93 Observers: Dr Scott Thompson

Zone: 50 Easting: 369264 mE Northing: 6509060 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y94 Observers: Dr Scott Thompson

Zone: 50 Easting: 369268 mE Northing: 6509027 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y95 Observers: Dr Scott Thompson

Zone: 50 Easting: 369296 mE Northing: 6509002 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y96 Observers: Dr Scott Thompson

Zone: 50 Easting: 369282 mE Northing: 6508984 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y97 Observers: Dr Scott Thompson

Zone: 50 Easting: 369297 mE Northing: 6508945 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y98 Observers: Dr Scott Thompson

Zone: 50 Easting: 369305 mE Northing: 6508952 mN

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y99 Observers: Dr Scott Thompson

Zone: 50 Easting: 369334 mE Northing: 6508925 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Poor

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y100 Observers: Dr Scott Thompson

Zone: 50 Easting: 369362 mE Northing: 6508893 mN

Landform: Coastal undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y101 Observers: Dr Scott Thompson

Zone: 50 Easting: 369373 mE Northing: 6508956 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y102 Observers: Dr Scott Thompson

Zone: 50 Easting: 369434 mE Northing: 6508952 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y103 Observers: Dr Scott Thompson

Zone: 50 Easting: 369462 mE Northing: 6508967 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y104 Observers: Dr Scott Thompson

Zone: 50 Easting: 369504 mE Northing: 6508930 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y105 Observers: Dr Scott Thompson

Zone: 50 Easting: 369549 mE Northing: 6508913 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y106 Observers: Dr Scott Thompson

Zone: 50 Easting: 369583 mE Northing: 6508914 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y107 Observers: Dr Scott Thompson

Zone: 50 Easting: 369616 mE Northing: 6508938 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y108 Observers: Dr Scott Thompson

Zone: 50 Easting: 369650 mE Northing: 6508947 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y109 Observers: Dr Scott Thompson

Zone: 50 Easting: 369678 mE Northing: 6508980 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y110 Observers: Dr Scott Thompson

Zone: 50 Easting: 369706 mE Northing: 6509022 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y111 Observers: Dr Scott Thompson

Zone: 50 Easting: 369683 mE Northing: 6509056 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y112 Observers: Dr Scott Thompson

Zone: 50 Easting: 369626 mE Northing: 6509065 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y113 Observers: Dr Scott Thompson

Zone: 50 Easting: 369601 mE Northing: 6509030 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y114 Observers: Dr Scott Thompson

Zone: 50 Easting: 369601 mE Northing: 6508988 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y115 Observers: Dr Scott Thompson

Zone: 50 Easting: 369565 mE Northing: 6508970 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y116 Observers: Dr Scott Thompson

Zone: 50 Easting: 369513 mE Northing: 6508963 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y117 Observers: Dr Scott Thompson

Zone: 50 Easting: 369513 mE Northing: 6509019 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y118 Observers: Dr Scott Thompson

Zone: 50 Easting: 369505 mE Northing: 6509062 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y119 Observers: Dr Scott Thompson

Zone: 50 Easting: 369486 mE Northing: 6509095 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good to very good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y120 Observers: Dr Scott Thompson

Zone: 50 Easting: 369485 mE Northing: 6509137 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y121 Observers: Dr Scott Thompson

Zone: 50 Easting: 369443 mE Northing: 6509100 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y122 Observers: Dr Scott Thompson

Zone: 50 Easting: 369450 mE Northing: 6509149 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y123 Observers: Dr Scott Thompson

Zone: 50 Easting: 369401 mE Northing: 6509131 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Good

Soil Type: Sand Surface: Sand and limestone





Date: 29-Aug-19 Habitat Assessment #: Y124 Observers: Dr Scott Thompson

Zone: 50 Easting: 369344 mE Northing: 6509136 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand





Date: 29-Aug-19 Habitat Assessment #: Y125 Observers: Dr Scott Thompson

Zone: 50 Easting: 369344 mE Northing: 6509173 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand





Date: 29-Aug-19 Habitat Assessment #: Y126 Observers: Dr Scott Thompson

Zone: 50 Easting: 369351 mE Northing: 6509229 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand





Date: 29-Aug-19 Habitat Assessment #: Y127 Observers: Dr Scott Thompson

Zone: 50 Easting: 369339 mE Northing: 6509305 mN

Landform: Undulating Fire History: > 5 years Habitat Quality: Disturbed

Soil Type: Sand Surface: Sand



