



Pilbara Green Link: Preliminary Investigation Activities

Supporting Document: NVCP 2 Clearing Application

Horizon Power

5 June 2025

→ **The Power of Commitment**



Project name		Pilbara Green Link					
Document title		Pilbara Green Link: Preliminary Investigation Activities - Supporting Document: NVCP 2 Clearing Application					
Project number		12657458					
GHD Doc No		12657458-GHD-00-00-RPT-EN-00002					
HP Doc No		P1001373-P91210-PIL-EN-RPT-0006					
File name		12657458-GHD-00-00-RPT-EN-00002_NVCP2_SupportingDocument_Rev0.docx					
Status Code	Revision	Author	Reviewer		Approved for issue		
			Name	Signature	Name	Signature	Date
S0	A			On file		On file	29/05/2025
S4	0			On file		On file	05/06/2025

GHD Pty Ltd | ABN 39 008 488 373
Contact: , Project Manager | GHD
999 Hay Street, Level 10
Perth, Western Australia 6000, Australia
T +61 8 6222 8222 | **F** +61 8 6222 8555 | **E** permail@ghd.com | **ghd.com**

© GHD 2025
This document is and shall remain the property of Horizon Power. The document may only be used for the purpose for which it was commissioned and in accordance with the Terms of Engagement for the commission. Unauthorised use of this document in any form whatsoever is prohibited.

Contents

1. Introduction	5
1.1 Proposal background	5
1.2 Purpose of this report	8
2. Description of clearing activities	10
2.1 Access tracks	10
2.2 Geotechnical Investigations	11
2.3 Groundwater Investigations	11
2.4 Environmental Management Plan (EMP)	12
3. Existing environment	17
3.1 Climate	17
3.2 Flora and vegetation	18
3.3 Fauna	36
3.4 Conservation Areas	45
3.5 Environmentally Sensitive Areas	45
3.6 Hydrology	45
3.7 Land use	48
4. Assessment of impacts	50
4.1 Potential impacts to vegetation and flora	50
4.2 Potential impacts to fauna and fauna habitat	50
5. Assessment against 10 clearing principles	53
6. Environmental Avoidance and Management	68
6.1 Impact avoidance and minimisation through design and planning	68
6.2 Impact avoidance and minimisation measures applied on site	68
7. Other approvals	71
7.1 Referral to EPA	71
7.2 Approvals under the RiWI Act	71
7.3 Aboriginal Heritage	71
8. Offsets	72
9. References	73

Table index

Table 1	Project definitions	6
Table 2	Clearing type for each Investigation Activity	11
Table 3	Landscape Zones intersecting the DE	19
Table 4	Soil-Landscape Systems intersecting the DE	19
Table 5	Mapped pre-European vegetation associations	20

Table 6	Representation of Pre-European Vegetation Associations recorded within the DE.	20
Table 7	Vegetation condition categories in the DE	22
Table 8	Vegetation types recorded in the PGL Surveyed Area	23
Table 9	Conservation significant flora - Likelihood of occurrence assessment (Biota 2025)	34
Table 10	Flora likelihood of occurrence assessment against relevant guidelines (Biota 2025)	35
Table 11	Fauna Habitats within the PGL Surveyed Area	37
Table 12	Conservation Significant Fauna Species in the PGL Surveyed Area	43
Table 13	Microhabitats likely to support SRE species	44
Table 14	Water Resources within the DE	45
Table 15	Catchments that intersect the DE	46
Table 16	Localities surrounding the PGL Surveyed Area	48
Table 17	Vegetation types and condition within the Indicative Impact Area	50
Table 18	Fauna habitat types within the Indicative Impact Area	51
Table 19	Disruption of potential habitat clearing on Conservation Significant fauna species	51
Table 20	Impact of DE clearing on Conservation Significant Fauna Species	56
Table 21	Definition of acronyms	76
Table 22	Investigation Pads and activities	76
Table 23	Impacts of proposed clearing on Land Parcels	80

Figure index

Figure 1	Project Location	9
Figure 2	Investigation Site locations	13
Figure 3	Port Hedland Airport monthly climate statistics (BoM weather station: 004032)	17
Figure 4	Mandora monthly climate statistics (BoM weather station: 004019)	17
Figure 5	Environmental Constraint: Vegetation Type	26
Figure 6	Environmental Constraint: Vegetation Condition	30
Figure 7	Environmental Constraint: Fauna Habitats	39
Figure 8	Environmental Constraint: Hydrological	47
Figure 9	Environmental Constraint: Acid Sulfate Soils	49

Appendices

Appendix A	Investigation sites and activities
Appendix B	Land parcel clearing
Appendix C	Dandjoo Database Search Results

Limitations

This report has been prepared by GHD for Horizon Power and may only be used and relied on by Horizon Power for the purpose agreed between GHD and Horizon Power as set out in section 1.2 this report.

GHD otherwise disclaims responsibility to any person other than Horizon Power arising in connection with this report. GHD also excludes implied warranties and conditions, to the extent legally permissible.

The services undertaken by GHD in connection with preparing this report were limited to those specifically detailed in the report and are subject to the scope limitations set out in the report.

The opinions, conclusions and any recommendations in this report are based on conditions encountered and information reviewed at the date of preparation of the report. GHD has no responsibility or obligation to update this report to account for events or changes occurring subsequent to the date that the report was prepared.

GHD has prepared this report on the basis of information provided by Horizon Power and others who provided information to GHD (including Government authorities), which GHD has not independently verified or checked beyond the agreed scope of work. GHD does not accept liability in connection with such unverified information, including errors and omissions in the report which were caused by errors or omissions in that information.

Abbreviations

Abbreviation	Definition
AREH	Australian Renewable Energy Hub
BoM	Bureau of Meteorology
DBCA	Department of Biodiversity, Conservation and Attractions
DCCEEW	Department of Climate Change, Energy, the Environment and Water
DPIRD	Department of Primary Industries and Regional Development
DPLH	Department of Planning, Lands and Heritage
DWER	Department of Water and Environmental Regulation
DE	Development Envelope
EP Act	<i>Environmental Protection Act 1986</i>
EPA	Environmental Protection Authority
EPBC Act	<i>Environmental Protection, Biodiversity and Conservations Act 1999</i>
ERD	Environmental Review Document
ESA	Environmentally Sensitive Area
GDE	Groundwater Dependent Ecosystem
GDV	Groundwater Dependent Vegetation
HP	Horizon Power
IBRA	Interim Biogeographic Regionalisation for Australia
MNES	Matters of National Environmental Significance
NVCP	Native Vegetation Clearing Permit
PEC	Priority Ecological Community
PGL	Pilbara Green Link
PMST	Protected Matters Search Tool
RiWI Act	<i>Rights in Water and Irrigation Act 1914</i>
TEC	Threatened Ecological Community

1. Introduction

1.1 Proposal background

Horizon Power (HP) is a Western Australian (WA) Government Trading Enterprise (GTE) and the state's regional and remote energy utility. Horizon power operates under the *Electricity Corporations Act 2005* and is governed by a Board of Directors accountable to the Minister for Energy. Horizon Power is an experienced asset manager undertaking active management of vast electricity networks and generation assets across regional WA, utilising mature and robust operational, health and safety, and environmental management systems.

HP anticipates significant transmission developments in the Pilbara as industry prepares to further electrify and decarbonise its operations. The electricity requirements for committed and planned projects are substantial, and the need for all parties to transition to renewable energy solutions provides unique challenges for the State of Western Australia. The Australian Renewable Energy Hub (AREH) renewable project is one of many private developments that the State can support to transition the Pilbara region to a low-carbon energy future.

Working with the AREH Project, HP is proposing to construct a 330 kV transmission line as part of the Pilbara Green Link (PGL) Project. The PGL Project will interconnect with HP's existing network at Port Hedland and extends approximately 250 km east towards the AREH site (Figure 1). PGL was selected as a priority project by WA Government, forming the Great Sandy Desert Corridor, one of the 4 Priority (transmission line) Corridors identified as part of the Pilbara Energy Transition (PET) Plan.

As part of Project feasibility stage, PGL requires preliminary geotechnical and groundwater investigations (Investigation Activities) to be undertaken to assist in determining the alignment of the proposed transmission line, and substations, which ultimately will inform the detailed design.

Horizon Power engaged GHD Pty Ltd (GHD) to prepare a Native Vegetation Clearing Permit (NVCP) application for the Investigation Activities. Given the need to assess two distinct areas, in consultation with DWER in November 2024, it was determined that submitting separate NVCPs would be the most appropriate approach.

NVCP 1 was lodged on 6 February 2025 and is currently awaiting determination. This document serves as the supporting submission for the second application, referred to as NVCP 2. The clearing proposed under NVCP 2 is intended for a specific purpose and will occur on land not owned by Horizon Power. However, Horizon Power will have the necessary legal access to conduct the investigations. Given these circumstances, a purpose permit is considered the appropriate mechanism.

The Investigation Activities are proposed to occur between September 2025 and August 2027 as outlined in the NVCP 2 application form. As an 'energy operator', Horizon Power has certain rights under Sections 46 and 49 of the *Energy Operators (Powers) Act 1979* which allow it to access and use land for the purpose of constructing, maintaining and operating electricity infrastructure. Horizon Power will utilise these access powers for the Investigation Activities. A Notice of Entry as required under the Act, will be issued to all relevant landowners or occupiers.

1.1.1 Stakeholder engagement

Ecological surveys of the proposed Sites associated with the Investigation Activities (see below) has involved engagement with Pastoral station owners and Traditional Owners. Engagement with these, and other stakeholders, will continue through the development of the Investigation Activities and the PGL project more broadly.

1.1.2 PGL Surveyed Area and the Development Envelope

The PGL Surveyed Area extends from Port Hedland and runs approximately 250 km east towards the AREH site. The early investigative stage of the PGL Project associated with NVCP 2 will consist of *up to*:

- 16 groundwater bores
- 28 test pits and 13 shallow bores for geotechnical investigations

It should be noted that several of the groundwater bores are contingency sites and are unlikely to be developed. Where required, access tracks to geotechnical and groundwater bore locations will be developed to enable the investigation program. For the purpose of this NVCP assessment, the following areas are defined in Table 1. The Investigation Activities described in this NVCP propose to clear up to 93.3 ha of native vegetation, within the 232.6 ha DE, for the purposes of geotechnical and groundwater investigations (Figure 2).

Table 1 Project definitions

Project definition	Purpose	Area (hectares (ha))
Investigation Activities	The preliminary geotechnical and groundwater investigations to be undertaken to assist in determining the alignment of the proposed transmission line and substations which ultimately will assist in informing detailed design	Described in the rows below
PGL Surveyed Area	<p>This is an area consisting of both the PGL Alignment Area (50,057.8 ha) and Additional Targeted Survey Area (145.5 ha). The PGL Alignment Area is an area along the 250 km proposed transmission line route, approximately 2 km in width, which was surveyed by Biota (2024) to inform the ecological constraints.</p> <p>The Additional Targeted Survey Area included areas outside the PGL Alignment Area which were surveyed by Biota (2025) to further inform the ecological constraints associated with the Investigation Activities.</p>	50,203.3 ha
Development Envelope (DE)	<p>The maximum extent within which limited clearing will occur to enable Investigation Activities, consisting of access track corridors and Investigation Pads (38 total).</p> <p>A full outline of Investigation Pads and the associated Investigation Activities is found in Table 22.</p>	<p>232.6ha</p> <p>Consisting of:</p> <ul style="list-style-type: none"> – 38.0 ha of Investigation Pads – 194.6 ha of Access track corridors
Investigation Pad	<p>The pads consist of the following combinations of Investigation Activities:</p> <ul style="list-style-type: none"> – 22 pads for geotechnical investigations only – 10 pads for groundwater investigations only – 6 pads for co-located groundwater and geotechnical investigations <p>Each pad allocated a maximum area of 100m x 100m to allow flexibility in micro-siting to account for local constraints and minimise the impacts of clearing.</p> <p>Of note, not all Investigation Pads may be required, and therefore the total number is considered a theoretical maximum.</p>	
Access track corridor	<ul style="list-style-type: none"> – 21 access track corridors <p>Track corridors were allocated a buffer width of 25m to allow flexibility in micro-siting to account for local constraints and minimise the impacts of clearing</p>	
Indicative Impact Area:	<p>The maximum extent of impact to vegetation within the DE. Consisting of Investigation Activity Pads and access tracks.</p> <ul style="list-style-type: none"> – The total number of Investigation Pads is considered a maximum that may be required for the Investigation Activities. There is not a guaranteed need for all Sites with contingency Sites included in case successful boreholes cannot be established – Where a geotechnical Investigation Site is co-located with a groundwater Investigation Site, the clearing required within the Investigation Pad will be approximately 70 m by 70 m to account for all Investigation Activities. – Clearing totals for Investigation Activities below are based on whether they are co-located, as described above. 	<p>93.3 ha</p> <p>Consisting of:</p> <ul style="list-style-type: none"> – 0.5 ha of geotechnical test pits – 0.8 ha of geotechnical boreholes – 7.8 ha of groundwater boreholes – 84.1 ha of access tracks

Project definition	Purpose	Area (hectares (ha))
Geotechnical test pit	<p>28 test pits:</p> <ul style="list-style-type: none"> – 12 containing geotechnical test pit sites only – 10 co-located with geotechnical boreholes – 3 co-located with groundwater boreholes – 3 co-located with groundwater boreholes and geotechnical boreholes – A test pit will require approximately 15 m by 15 m clearing within an Investigation Pad. This allows for the movement of excavator equipment, placement of any stripped vegetation and topsoil, spoil heap and appropriate area to operate this investigation in a safe manner – A 5% clearing contingency is applied 	
Geotechnical boreholes	<p>13 geotechnical boreholes:</p> <ul style="list-style-type: none"> – 10 co-located with geotechnical test-pits only – 3 co-located with both groundwater boreholes and geotechnical test-pits – A Geotechnical borehole will require approximately 30 m by 25 m clearing within an Investigation Pad. This is required to accommodate the drilling of a borehole using a truck mounted auger, a laydown area for equipment, and drill spoil – A 5% clearing contingency is applied 	
Groundwater boreholes	<p>16 groundwater boreholes:</p> <ul style="list-style-type: none"> – 10 groundwater borehole sites only – 6 co-located with geotechnical investigations activities (i.e. geotechnical test pit and/or borehole) – Groundwater borehole establishment will require approximately 70 m by 70 m clearing within an Investigation Pad to accommodate the placement of stripped vegetation and topsoil, drilling of a borehole using a truck mounted auger, laydown area for equipment and materials, sump for water, and drill spoil – This total includes contingency sites. Total clearing is expected to be lower if not all sites are required 	
Access tracks	<p>Approximately 79.8 km</p> <ul style="list-style-type: none"> – A maximum width of 10 m of clearing has been allocated to access tracks. This is an upper limit value and is unlikely to be needed for most of the proposed access tracks required, with a typical access track estimated to be 3m to 4m wide if active vegetation clearing is required. – This area has been allocated for access tracks, based on desktop assessment of clearing that may be required at specific locations (e.g. for occasional turning areas or bends in the road), clearing vegetation associated with existing tracks, or in the development of new, short-distance access tracks. Of note, the total clearing area includes some alternative or contingency tracks where more than one access to a pad has been identified (e.g. TP44 and TP45; TP46 and TP47). It is not intended to clear vegetation for more than one access track to each pad where a potential alternative track has been identified. – For new access tracks to pads containing geotechnical test pits and/or boreholes, where practicable, low vegetation will be driven over rather than cleared by blade to minimise vegetation disturbance / clearing. – A 5% clearing contingency has been applied 	

1.2 Purpose of this report

The purpose of this document is to support the second NVCP application by describing the ten clearing principles assessment and outcome for the Investigation Activities, assess the potential impacts that may result from the proposed Investigation Activities, together with broad management measures to avoid and minimise potential impacts.

This document has been prepared in support of an application for a NVCP (purpose) under Section 51E of Part V of the *Environmental Protection Act 1986* (WA) (EP Act).

This document includes:


- An overview of works required and description of clearing activities to be undertaken (Section 2)
- An overview of existing environment (Section 3)
- An assessment of potential impacts identified (Section 4)
- An assessment against the Ten Clearing Principles, as defined in Schedule 5 of the EP Act (Section 5)
- Environmental management measures to be implemented to minimise clearing impacts (Section 6)
- Identification of other environmental and heritage approvals applicable to the proposed works (Section 7)
- Offsets applicable to the proposed works (Section 8)



Paper Size ISO A3
at Scale: 1:700,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020



Pilbara Green Link
Investigation Area - Permit Activities

PGL Alignment Area

Project No. 12657458
Revision No. B
Date 4/06/2025

FIGURE 1

2. Description of clearing activities

The proposed geotechnical test pit and borehole locations and the groundwater borehole locations are shown in Figure 2. Investigation Sites were informed by the requirements of geotechnical and groundwater studies as described below, together with the environmental constraints identified through the desktop and field survey assessments. Positioning of the Investigation Sites followed the PGL Preliminary Design Principles and application of DWER's clearance principles. The investigation will avoid and minimise impacts to significant flora, fauna and water courses where possible, through:

- Applying the native vegetation disturbance hierarchy throughout the design process, to avoid and minimise disturbance
- Avoiding placements of pads 50m either side of defined waterways where practicable
- Locating the investigation pads adjacent to, or near, existing tracks where practicable
- Avoiding disturbance to boulder outcrops / boulder fields and breakaways; caves and their immediate (200m radius) surrounds, gorges and deep gullies and clay pans
- Avoiding disturbance within 300m radius, and preferably 500m radius of identified Bilby burrows
- Avoiding disturbance to known Aboriginal sites.

The proposed locations are subject to change if there is an unforeseen constraining factor (such as previously unidentified Aboriginal heritage) to allow flexibility in the placement of geotechnical and groundwater investigations. The larger PGL Surveyed Area identifies the area within which access tracks will be constrained. A summary of the clearing required for each Investigation Activity is described in Table 1 above. The full breakdown of Investigation Pads, Investigation Activities and the clearing required can be found in Appendix A.

The type of clearing and actions for the cleared Indicative Impact Area following the completion of Investigation Activities is summarised in Table 2 below. Where geotechnical investigations occur alongside groundwater bores, rehabilitation of the pad will occur after the groundwater bore is established. Where a groundwater borehole is developed into a monitoring bore, the access track will remain in place to enable periodic monitoring. Where a groundwater borehole will not be developed into a monitoring bore (e.g. no water was located), the borehole, pad and access track (if new) will be rehabilitated.

2.1 Access tracks

To conduct the investigative works, some native vegetation is proposed to be cleared along existing and new access tracks within the DE. A truck mounted drill rig and light vehicles will need to drive between investigation test locations, targeting identified locations that will inform the placement of infrastructure. There will be some clearing required to increase the accessibility of existing tracks by removing regrowth or to create new tracks. Where feasible, track development will avoid clearing vegetation with a blade and instead low vegetation may be driven over.

Although new access tracks would typically be 3 to 4 m wide, a maximum width of 10 m has been allowed for to enable larger groundwater drilling rigs to turn around intersections. This is within a larger 25 m wide DE, which enables contractors to avoid constraints when accessing each Investigation Pad. Further, the total clearing area (i.e. Indicative Impact Area) includes some alternative or contingency tracks where more than one access to a pad has been identified (e.g. TP44 and TP45; TP46 and TP47). It is not intended to clear vegetation for more than one access track to each pad where a potential alternative track has been identified which will result in a reduced Indicative Impact Area.

An approach which avoids and minimises environmental impact will apply during the investigation program meaning the final clearing is expected to be less than the total outlined below. For the Indicative Impact Area of each access track, a 5% clearing contingency is applied to allow for flexibility when undertaking the works while also accounting for typical GPS inaccuracies. With a maximum width requirement of 10 m and 5% contingency applied, access tracks account for 84.1 ha (90.2%) of the maximum 93.3 ha of clearing proposed, with a total length of 79.8 km.

Following the completion of Investigation Activities any clearing associated with existing tracks will be retained. New access tracks developed to pads involving groundwater bores developed into monitoring bores will remain cleared. Where access to a pad is no longer required, and the track was established specifically to access the pad, the track will be rehabilitated.

2.2 Geotechnical Investigations

For the geotechnical investigations, an indicative test spacing of approximately 20 km between boreholes and approximately 10 km between test pits along the proposed PGL Surveyed Area was used. Proposed test locations were then reviewed in terms of accessibility, aerial imagery, topography, and underlying geology to capture relevant geotechnical information along the PGL Surveyed Area.

Up to 28 geotechnical test pits and 13 boreholes are proposed within the PGL Surveyed Area as part of this NVCP (Figure 2). For the geotechnical boreholes a clearing area of approximately 30 m by 25 m is required while for test pits, a clearing area of approximately 15 m by 15 m is required. For each geotechnical Investigation Activity, a 5% clearing contingency is applied to allow for flexibility when undertaking the works while also accounting for typical GPS inaccuracies. Of the 93.3 ha of clearing proposed, 1.3 ha (1.4%) is for geotechnical boreholes and test pits (inclusive of maximum clearing for the activity and a 5% contingency).

Where feasible and safe to do so, the location of boreholes and test pits will minimise clearing vegetation by traversing over low vegetation rather than using a blade.

2.3 Groundwater Investigations

Groundwater drilling sites were selected after an extensive assessment of the known hydrogeological resources present within or adjacent to the Alignment Area. Estimated yield, hydraulic conductivity, water quality, aquifer homogeneity and isotropy, local prospectivity (topography and hydrology) were the technical aspects assessed, along with existing access and optimising any existing bore locations. Spatial distribution of target sites along the PGL Surveyed Area was determined by estimated Project water demands at specific locations, with a view to co-locate target sites with geotechnical sites where practicable to minimise disturbance.

Up to 16 groundwater boreholes are proposed within the PGL Surveyed Area (Figure 2) as part of this NVCP. For the groundwater boreholes an area of approximately 70 m by 70 m is required to accommodate the placement of stripped vegetation and topsoil, drilling of a borehole using a truck mounted auger, a laydown area for equipment and materials, sump for water and drill spoil. Of the 93.3 ha of clearing proposed, 7.8 ha (8.4%) is for groundwater boreholes.


Table 2 Clearing type for each Investigation Activity

Investigation Activity	Clearing type	Action following completion of Investigation Activity
Access track – Geotechnical	Temporary	<ul style="list-style-type: none"> Tracks only cleared for the Investigation Activity will be rehabilitated Clearing associated with existing tracks will be retained
Geotechnical test pits and boreholes	Temporary	<ul style="list-style-type: none"> Sites will be left to be revegetated through active rehabilitation. Sites will be restored and reprofiled to natural levels and contours Sites co-located with groundwater borehole will be rehabilitated after the groundwater bore is established
Access track - Groundwater	Permanent	<ul style="list-style-type: none"> New access tracks developed and existing tracks involving groundwater bores will remain cleared
Groundwater boreholes	Permanent	<ul style="list-style-type: none"> If a groundwater bore is successful in locating a suitable water supply, the bore will be a monitoring bore as part of the Investigation Activities. Any part of the pad area no longer required will be rehabilitated. The access track will be maintained for monitoring access. Monitoring bores deemed suitable for water supply will be established as operational areas during construction phase of the PGL Project with clearing maintained. Post-construction it is likely groundwater bore ownership will be passed on to an interested party (i.e. pastoralist) or otherwise decommissioned and rehabilitated.

Investigation Activity	Clearing type	Action following completion of Investigation Activity
Groundwater borehole – unsuccessful establishment	Temporary	<ul style="list-style-type: none"> – If a groundwater bore is not successful in locating water, the bore will be capped and the Investigation Pad area rehabilitated – If a new access track was developed to the site, it will be rehabilitated

2.4 Environmental Management Plan (EMP)


Prior to the commencement of the Investigation Activities, the Contractor for each investigation type will, as a minimum, be required to prepare an Environmental Management Plan (EMP) incorporating the requirements of this NVCP and any conditions. The avoidance and mitigation measures outlined in Section 6.2 demonstrate the minimum requirements (commitments) expected to be included in each Contractor developed EMP.



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres


Horizontal Datum: GDA2020
Grid: GDA2020



Pilbara Green Link
Investigation Area – Permit Activities

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 2.1



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area - Permit Activities

Investigation Site Locations

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 2.2



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area – Permit Activities

Investigation Site Locations

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 2.3



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020



Horizon Power
Investigation Area - Permit Activities

Investigation Site Locations

Project No. 12657458
Revision No. B
Date 4/06/2025

FIGURE 2.4

3. Existing environment

3.1 Climate

The Pilbara region that the Project is located within has a climate of very hot summers, mild winters, and generally low rainfall year-round. The climate is characteristic of hot grassland in the north-west part of the region, where the Project is located.

The western portion of the PGL Surveyed Area is near Port Hedland and the eastern portion extends towards Mandora Station, hence the typical weather across the PGL Surveyed Area is represented by the weather at these locations.

At the Port Hedland Airport weather station (site number: 004032, BoM, Figure 3) the mean maximum temperature is 36.8°C, occurring in the months of March and December. The mean minimum temperature is in July at 12.5°C. February receives the highest mean rainfall of 88.3 mm, and October the lowest of 0.9 mm. The mean annual rainfall is 314.1 mm.




Figure 3 Port Hedland Airport monthly climate statistics (BoM weather station: 004032)

At the Mandora weather station (site number: 004019, BoM, Figure 4) the mean maximum temperature is 37.0°C, occurring in March. The mean minimum temperature is in July at 12.5°C. February receives the highest mean rainfall of 100.9 mm, and September the lowest of 1.1 mm. The mean annual rainfall is 375.0 mm.




Figure 4 Mandora monthly climate statistics (BoM weather station: 004019)

3.2 Flora and vegetation

In May 2024, Biota conducted reconnaissance and targeted level field surveys of the PGL Surveyed Area to inform the design of the PGL Project. Following the initial biological survey, additional development areas were identified outside of the previous survey area. These areas are part of early investigation works to assist in determining the alignment of the proposed transmission line. In March 2025, Biota conducted reconnaissance and targeted level field surveys on the additional development areas (termed in this NVCP as the Additional Targeted Survey Area).

These activities were completed in line with the following policy documents and technical guidance:

- Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment (EPA 2016a)
- Technical Guidance – Sampling of Short-range Endemic Invertebrate Fauna (EPA 2016b)
- Technical Guidance – Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment (EPA 2020)
- Survey guidelines for Australia's threatened mammals: Guidelines for detecting mammals listed as threatened under the EPBC Act (DSEWPaC 2011a)
- Survey guidelines for Australia's threatened birds: Guidelines for detecting birds listed as threatened under the EPBC Act (DEWHA 2010a)
- Survey guidelines for Australia's threatened reptiles: Guidelines for detecting reptiles listed as threatened under the EPBC Act (DSEWPaC 2011b)
- Survey guidelines for Australia's threatened bats: Guidelines for detecting bats listed as threatened under the EPBC Act (DEWHA 2010b)
- A review of ghost bat ecology, threats and survey requirements (Bat Call WA 2021a)
- A review of Pilbara leaf-nosed bat ecology, threats and survey requirements (Bat Call WA 2021b).

In line with the Environmental Protection Authority (EPA) technical guidance (EPA 2016a, 2020) potential limitations of the surveys were identified by Biota (2024 & 2025). The following limitations were identified:

- May 2024 survey - Rainfall: Conditions were drier than expected. This resulted in the collection of poor material for some species, hindering identification, and some annual and cryptic perennial taxa are unlikely to have been present, particularly in the clay plains habitats. This would include some Priority flora taxa but no Threatened flora taxa likely for the survey area.
- May 2024 and March 2025 surveys - Fire: Considered a limitation for the flora survey, but one that is typically encountered in the Pilbara region and unlikely to have affected overall survey adequacy.

3.2.1 Regional biogeography

The PGL Surveyed Area lies within the McLarty, Chichester and Roebourne subregions of the Great Sandy Desert and Pilbara bioregions (Department of Conservation and Land Management (DCLM) 2002).

- The McLarty subregion (GSD1) is described as “mainly tree steppe grading to shrub steppe in south, comprising open hummock grassland of *Triodia pungens* and *Triodia schinzii* with scattered trees of *Owenia reticulata* and Bloodwoods, and shrubs of *Acacia spp*, *Grevillea wickhamii* and *G. refracta*, on Quaternary red longitudinal sand dune fields overlying Jurassic and Cretaceous sandstones of the Canning and Armadeus Basins. *Casuarina decaisneana* (Desert Oak) occurs in the far east of the region. Gently undulating lateritised uplands support shrub steppe such as *Acacia pachycarpa* shrublands over *Triodia pungens* hummock grass. Calcrete and evaporite surfaces are associated with occluded palaeo-drainage systems that traverse the desert; these include extensive salt-lake chains with samphire low shrublands, and *Melaleuca glomerata* - *M. lasiandra* shrublands. It includes the Mandora Paleoriver System. Red-brown dunefields with finer texture than further south. Includes gravelly surfaces of Anketell Ridge along its northern margin. The subregion is arid tropical with summer rain and is influenced by monsoonal activity. Morning fogs are recorded during the dry season. The total subregional area is 13,173, 266 ha.” (DCLM 2002)
- The Chichester subregion (PIL1) comprises of the northern section of the Pilbara Craton. Undulating Archaean granite and basalt plains include significant areas of basaltic ranges. Plains support a shrub steppe characterised by *Acacia inaequilatera* over *Triodia wiseana* (formerly *Triodia pungens*) hummock grasslands, while *Eucalyptus leucophloia* tree steppes occur on ranges. The climate is Semi-desert-tropical

and receives 300 mm of rainfall annually. Drainage occurs to the north via numerous rivers (e.g. De Grey, Oakover, Nullagine, Shaw, Yule, Sherlock). Subregional area is 9,044,560 ha" (DCLM 2002).

- The Roebourne subregion (PIL4) comprises of quaternary alluvial and older colluvial coastal and sub-coastal plains with a grass savannah of mixed bunch and hummock grasses, and dwarf shrub steppe of *Acacia stellaticeps* or *A. pyrifolia* and *A. inaequilatera*. Uplands are dominated by *Triodia* hummock grasslands. Ephemeral drainage lines support *Eucalyptus victrix* or *Corymbia hamersleyana* woodlands. Samphire, *Sporobolus* and mangal occur on marine alluvial flats and river deltas. Resistant linear ranges of basalts occur across the coastal plains, with minor exposures of granite. Islands are either Quaternary sand accumulations, or composed of basalt or limestone, or combinations of any of these three. Climate is arid (semi-desert) tropical with highly variable rainfall, falling mainly in summer. Cyclonic activity is significant, with several systems affecting the coast and hinterland annually. Subregional area is 2,008,983 ha" (DCLM 2002).

3.2.2 Land Systems

The DE intersects with three unique soil landscape zones (Department of Primary Industries and Regional Development (DPIRD) 2022a) described in the table below (Table 3).

Table 3 Landscape Zones intersecting the DE

Zone	Description
Zone 117 (Nita Sandplain Zone)	Sandplains and dunes on cretaceous canning basin sedimentary rocks with red deep sands and some red sandy earths.
Zone 280 (Nullagine Hills Zone)	Hills and ranges (with some stony plains) on volcanic and sedimentary rocks of the Pilbara craton (including the Hamersley Basin) with stony soils and red shallow loams and sands.
Zone 281 (De Grey-Roebourne Lowlands Zone)	Alluvial plains and sandplains on alluvial and marine deposits over the northern Pilbara craton with red deep sandy duplexes, red loamy earths, red/brown non-cracking clays, cracking clays, red sandy earths and red deep loamy duplexes.

The landscape zones describe above have been further classified into soil-landscape systems (DPIRD 2022b). The DE intersects with seven soil-landscape systems described below (Table 4).

Table 4 Soil-Landscape Systems intersecting the DE

System	Description
Boolgeeda System (281Bg)	Stony lower slopes and plains below hill systems supporting hard and soft spinifex grasslands or mulga shrublands.
Callawa System (117Cl)	Highly dissected low hills, mesas and gravelly plains on sandstone and conglomerate supporting soft and hard spinifex grasslands.
Capricorn System (280Cp & 281Cp)	Rugged sandstone hills, ridges, stony foot slopes and interfluvies supporting low acacia shrublands or hard spinifex grasslands with scattered shrubs.
Macroy System (281Mc)	Stony plains and occasional tor fields based on granite supporting hard and soft spinifex shrubby grasslands.
Nita System (117Nt & 281Nt)	Sandplains supporting shrubby spinifex grasslands with occasional trees.
River System (281Ri)	Narrow, seasonally active flood plains and major river channels supporting moderately close, tall shrublands or woodlands of acacias and fringing communities of eucalypts sometimes with tussock grasses or spinifex
Uaroo System (281Ua)	Broad sandy plains, pebbly plains and drainage tracts supporting hard and soft spinifex hummock grasslands with scattered acacia shrubs.

3.2.3 Broad vegetation mapping and extents

Broad-scale (1:250,000) pre-European vegetation mapping identified five Vegetation Associations present in the DE (DPIRD 2019). These are shown in Table 5 below.

Table 5 Mapped pre-European vegetation associations

Pre-European vegetation association	Area within DE (ha)
93: Hummock grasslands, shrub steppe; kanji over soft spinifex	10.5
101: Hummock grasslands, shrub steppe; <i>Acacia pachycarpa</i> over soft spinifex	72.3
117: Hummock grasslands, grass steppe; soft spinifex	147.8
589: Mosaic: Short bunch grassland – savanna / grass plain (Pilbara) / Hummock grasslands, grass steppe; soft spinifex	1.0
647: Hummock grasslands, dwarf-shrub steppe; <i>Acacia translucens</i> over soft spinifex	1.0

All Vegetation Associations are well represented across State, Interim Biogeographic Regionalisation for Australia (IBRA) Region, IBRA Subregion, and Local Government Area (LGA) extents. The majority of pre-European vegetation associations within the PGL Surveyed Area have 99% or more of their pre-European extent remaining (Table 6). Vegetation Association (Veg Assoc) No. 117, described as “*hummock grasslands, grass steppe; soft spinifex*”, has historically been subject to the highest proportional amount of clearing, seen at a subregional scale (Pindanland). Retention of pre-European vegetation at this scale is approximately 76%. However, a relatively high proportion of this remaining vegetation now exists in Department of Biodiversity, Conservation and Attraction (DBCA) reserves where it is protected from future clearing.

Table 6 Representation of Pre-European Vegetation Associations recorded within the DE.

Pre-European Vegetation Association	Scale		Pre-European (ha)	Current Extent (ha)	% Remaining	% Remaining in DBCA Reserves
Veg Assoc No. 93	Statewide WA		3,044,310	3,040,641	99.88	1.96
	IBRA Bioregion	Pilbara	3,041,114	3,038,472	99.88	1.96
		Great Sandy Desert	1,107	1,096	98.97	0
	IBRA Sub-region	Roebourne	46,361	46,334	99.94	0
		Chichester	2,940,348	2,936,732	99.88	2.02
		McLarty	1,107	1,096	98.97	0
	Local Government Authority	Town of Port Hedland	1,015,339	1,014,600	99.93	0
		Shire of East Pilbara	1,709,522	1,706,781	99.84	2.70
Veg Assoc No. 101	Statewide WA		1,191,084	1,191,038	100.00	1.67
	IBRA Bioregion	Pilbara	341	341	100.00	0
		Great Sandy Desert	961,170	961,124	100.00	2.07
	IBRA Sub-region	Roebourne	56	56	100.00	0
		Chichester	285	285	100.00	0
		McLarty	625,007	624,961	99.99	3.18
	Local Government Authority	Town of Port Hedland	6,958	6,928	99.57	0
		Shire of East Pilbara	612,920	612,904	100.00	3.25
Veg Assoc No. 117	Statewide WA		919,517	886,006	96.36	15.47
	IBRA Bioregion	Pilbara	82,706	78,097	94.43	21.69
		Great Sandy Desert	467,579	467,122	99.90	0.19
		Dampierland	28,895	22,074	76.39	30.54
	IBRA Sub-region	Roebourne	50,963	46,902	92.03	35.19
		Chichester	31,743	31,195	98.27	0
		Pindanland	28,895	22,074	76.39	30.54

Pre-European Vegetation Association	Scale		Pre-European (ha)	Current Extent (ha)	% Remaining	% Remaining in DBCA Reserves
		McLarty	247,331	246,923	99.84	0
	Local Government Authority	Town of Port Hedland	26,426	22,987	86.99	0
		Shire of East Pilbara	655,721	654,571	99.82	16.04
Veg Assoc No. 589	Statewide WA		807,699	802,713	99.38	1.9
	IBRA Bioregion	Pilbara	728,768	724,696	99.44	2.10
		Great Sandy Desert	26	26	100	0
		Dampierland	90	90	100	0
	IBRA Sub-region	Roebourne	675,392	671,327	99.4	2.13
		Chichester	53,376	53,368	99.98	1.78
		McLarty	26	26	100	0
		Pindanland	90	90	100	0
	Local Government Authority	Town of Port Hedland	338,269	335,921	99.31	0
		Shire of East Pilbara	64,182	64,112	99.89	0.07
Veg Assoc No. 647	Statewide WA		195,861	191,711	97.88	0
	IBRA Bioregion Pilbara		195,860	191,711	97.88	0
	IBRA Sub-region	Roebourne	188,901	184,775	97.82	0
		Chichester	6,959	6,936	99.68	0
	Local Government Authority	Town of Port Hedland	180,908	176,759	97.71	0

3.2.4 Vegetation types and condition

3.2.4.1 Vegetation types

Biota (2024 & 2025) completed reconnaissance and targeted level field surveys of the PGL Surveyed Area. Twenty-four vegetation types were identified from the PGL Surveyed Area (50,203.3 ha), associated with the following six broad landforms, with the remaining 603.5 ha consisting of cleared areas:

- Plains (P) – 33,961.4 ha
- Low Stony Rises (H) – 5,934.9 ha
- Rocky Outcrops and Breakaways (R) – 4,401.0 ha
- Clay Plains (C) – 4,274.9 ha
- Drainage Areas (D) – 1,006.7 ha
- Sand Dunes (S) – 21.0 ha

Each vegetation type is coded based on the broad landform to which it belongs. Vegetation types within each broad landform are shown in Figure 5 and Table 8. The following two vegetation types dominate the PGL Surveyed Area, both belonging to the Plains broad landform (Biota 2025):

- P3: *Corymbia zygophylla* scattered low trees to low open woodland over *Grevillea refracta* subsp. *refracta*, *Acacia tumida*, *Acacia ancistrocarpa*, *Acacia eriopoda*, *Acacia monticola* tall open shrubland over *Jacksonia aculeata*, *Croton aridus* low open shrubland – 15,533.3 ha (30.9% of the PGL Surveyed Area)
- P1: *Acacia inaequilatera* scattered tall shrubs over *Acacia ancistrocarpa* scattered shrubs over *Acacia stellaticeps* scattered to low open shrubland over *Triodia epactia* scattered tussocks to open hummock grassland - 13,174.9 ha (26.2% of the PGL Surveyed Area)

The third most abundant vegetation type, H2: *Acacia robeorum* scattered tall shrubs over *Acacia stellaticeps* low open shrubland over *Triodia angusta* (*Triodia wiseana*, *T. epactia*) hummock grassland to very open hummock grassland, represents comparatively less vegetation within the PGL Surveyed Area, with 3,790.9 ha (7.6% of the PGL Surveyed Area).

3.2.4.2 Condition

Biota (2024 & 2025) mapped vegetation condition over the PGL Surveyed Area using the condition categories from EPA (2016). The vegetation condition is presented in Table 7 and Figure 6. The mapping indicates that the PGL Surveyed Area is predominantly in Excellent and Very Good to Excellent condition, with each condition representing 34.5% and 37.6% respectively.

Areas completely devoid of native vegetation were mapped as Cleared and were not assigned a condition rating. In total, 1.2% of the PGL Surveyed Area has been completely cleared of vegetation (e.g. roads, rails, mining area).

As observed by Biota (2025), the main disturbance factors in the PGL Surveyed Area comprised weed invasion and cattle grazing and/or trampling. The areas in the worst condition were associated with drainage lines and floodplains and usually supported high densities of weed species.

Table 7 Vegetation condition categories in the DE

Condition Rating	Area (ha) in the DE	Proportion of DE (%)	Proportion of PGL Surveyed Area (%)
Excellent	81.3	87	0.2
Very Good to Excellent	2.7	2.9	<0.1
Very Good	8.5	9.1	<0.1
Good to Very Good	0	0	<0.1
Good	0	0	<0.1
Poor to Good	0.02	<0.1	<0.1
Cleared/NA	0.8	0.9	<0.1

PROTECTED

Table 8 Vegetation types recorded in the PGL Surveyed Area


Landform	Vegetation type	Description	Extent (ha)	% of PGL Surveyed Area
Drainage Areas	D1	<i>Melaleuca argentea</i> , <i>Eucalyptus camaldulensis</i> subsp. <i>refulgens</i> open forest over <i>Ficus aculeata</i> var. <i>indecora</i> scattered low trees over <i>*Calotropis procera</i> tall open scrub over <i>Indigofera oblongifolia</i> tall shrubland over <i>*Cenchrus setiger</i> , <i>*Cynodon dactylon</i> open tussock grassland over <i>*Argemone ochroleuca</i> subsp. <i>ochroleuca</i> scattered herbs.	342.4	0.7
	D2	<i>Eucalyptus camaldulensis</i> subsp. <i>refulgens</i> and/or <i>Eucalyptus victrix</i> open to low open woodland over <i>Melaleuca argentea</i> tall open scrub to open woodland over <i>Acacia trachycarpa</i> , <i>Acacia colei</i> tall shrubland over <i>Corchorus incanus</i> subsp. <i>incanus</i> scattered low shrubs over <i>Triodia epactia</i> open to very open hummock grassland over <i>*Cenchrus</i> spp. scattered to very open tussock grassland.	577.2	1.1
	D3	<i>Corymbia flavescens</i> low open woodland over <i>Acacia tumida</i> var. <i>pilbarensis</i> over <i>Triodia epactia</i> scattered to very open hummock grassland over mixed native tussock grasses.	64.9	0.1
	D4	<i>Eucalyptus victrix</i> low woodland over <i>Lysiphyllum cunninghamii</i> low open woodland over <i>Acacia trachycarpa</i> , <i>Acacia colei</i> var. <i>colei</i> , <i>Acacia ancistrocarpa</i> , <i>Atalaya hemiglauc</i> tall shrubland over <i>Carissa lanceolata</i> open shrubland over <i>Chrysopogon fallax</i> open tussock grassland.	22.1	<0.1
	Total		1,006.7	2.0
Clay Plains	C1	<i>Eriachne</i> spp. open to closed tussock grassland.	1,330.3	2.6
	C2	<i>Acacia</i> spp., <i>Indigofera</i> spp., <i>Senna</i> spp. scattered low shrubs over <i>Triodia epactia</i> very open to hummock grassland.	1,252.4	2.5
	C3	<i>Triodia epactia</i> open hummock to hummock grassland.	1,572.0	3.1
	C4	<i>Eucalyptus victrix</i> , <i>Lysiphyllum cunninghamii</i> low open woodland over <i>Acacia tumida</i> var. <i>pilbarensis</i> scattered tall shrubs over <i>Atalaya hemiglauc</i> , <i>Acacia stellaticeps</i> low open shrubland over <i>Corchorus incanus</i> subsp. <i>incanus</i> , <i>Bonamia alatisemina</i> scattered low shrubs over <i>Triodia epactia</i> open hummock grassland.	120.2	0.2
	Total		4,274.9	8.5
Sand Dunes	S1	<i>Corymbia flavescens</i> low open woodland over <i>Acacia tumida</i> var. <i>pilbarensis</i> , <i>Acacia ancistrocarpa</i> tall shrubland over <i>Grevillea refracta</i> subsp. <i>refracta</i> , <i>Petalostylis labicheoides</i> open shrubland over <i>Acacia adoxa</i> var. <i>adoxa</i> , <i>Acacia hilliana</i> low shrubland over <i>Triodia epactia</i> open hummock grassland over <i>Grevillea wickhamii</i> , <i>Bonamia erecta</i> low shrubland.	10.7	<0.1
	S2	<i>Corymbia zygophylla</i> scattered low trees over <i>Acacia</i> spp. scattered tall shrubs over <i>Dicrastylis doranii</i> low open shrubland over <i>Triodia schinzii</i> very open hummock grassland over <i>Aristida holathera</i> var. <i>holathera</i> scattered tussock grasses.	10.3	<0.1
	Total		21.0	<0.1
Plains	P1	<i>Acacia inaequilatera</i> scattered tall shrubs over <i>Acacia ancistrocarpa</i> scattered shrubs over <i>Acacia stellaticeps</i> scattered to low open shrubland over <i>Triodia epactia</i> scattered tussocks to open hummock grassland.	13,174.9	26.2
		<i>Eucalyptus victrix</i> low woodland over <i>Corymbia hamersleyana</i> scattered low trees over <i>Acacia colei</i> var. <i>colei</i> tall open shrubland over <i>Triodia epactia</i> open hummock grassland.		
	P2	<i>Lysiphyllum cunninghamii</i> scattered low trees over <i>Acacia inaequilatera</i> tall shrubland over <i>Triodia longiceps</i> very open hummock grassland.	1,172.1	2.3

PROTECTED

Landform	Vegetation type	Description	Extent (ha)	% of PGL Surveyed Area
	P3	<i>Corymbia zygophylla</i> scattered low trees to low open woodland over <i>Grevillea refracta</i> subsp. <i>refracta</i> , <i>Acacia tumida</i> , <i>Acacia ancistrocarpa</i> , <i>Acacia eriopoda</i> , <i>Acacia monticola</i> tall open shrubland over <i>Jacksonia aculeata</i> , <i>Croton aridus</i> low open shrubland over <i>Triodia schinzii</i> (<i>T. epactia</i>) open to very open hummock grassland	15,533.3	30.9
	P4	<i>Acacia inaequilatera</i> , <i>Acacia ancistrocarpa</i> , <i>Grevillea wickhamii</i> tall open shrubland over <i>Acacia stellaticeps</i> low open shrubland over <i>Triodia epactia</i> (<i>T. wiseana</i> , <i>T. angusta</i>) open to very open hummock grassland.	2,504.5	5.0
	P5	<i>Corymbia hamersleyana</i> scattered low trees over <i>Acacia stellaticeps</i> , <i>Acacia adoxa</i> var. <i>adoxa</i> low open shrubland over <i>Triodia epactia</i> hummock grassland.	1,576.7	3.1
	Total		33,961.4	67.6
Low Stony Rises	H1a	<i>Grevillea wickhamii</i> scattered tall shrubs <i>Acacia hilliana</i> , <i>Acacia stellaticeps</i> low open shrubland over <i>Triodia scintillans</i> hummock grassland to open hummock grassland.	1,933.7	3.9
	H1b	<i>Corymbia hamersleyana</i> scattered low trees over <i>Grevillea wickhamii</i> , <i>Acacia</i> spp. scattered tall shrubs over <i>Acacia hilliana</i> , <i>Acacia stellaticeps</i> scattered low shrubs over <i>Triodia epactia</i> open to very open hummock grassland.		
	H1c	<i>Acacia bivenosa</i> , <i>Carissa lanceolata</i> scattered shrubs over <i>Acacia adoxa</i> var. <i>adoxa</i> scattered low shrubs over <i>Triodia wiseana</i> very open hummock grassland over scattered native tussock grasses.		
	H1d	<i>Corymbia flavescens</i> scattered low trees to low open woodland over <i>Acacia elachantha</i> , <i>Acacia ancistrocarpa</i> , <i>Acacia colei</i> tall open shrubland over <i>Triodia epactia</i> open to very open hummock grassland.		
	H2a	<i>Acacia robeorum</i> scattered tall shrubs over <i>Acacia stellaticeps</i> low open shrubland over <i>Triodia angusta</i> (<i>Triodia wiseana</i> , <i>T. epactia</i>) hummock grassland to very open hummock grassland.	3,790.9	7.6
	H2b	<i>Corymbia hamersleyana</i> scattered low trees over <i>Acacia stellaticeps</i> , <i>Acacia bivenosa</i> low shrubland to open shrubland over <i>Triodia epactia</i> open to very open hummock grassland.		
	H2c	<i>Gyrocarpus americanus</i> subsp. <i>pachyphyllus</i> scattered low trees over <i>Carissa lanceolatum</i> , <i>Flueggea virosa</i> subsp. <i>melanthesoides</i> scattered tall shrubs over * <i>Cenchrus ciliaris</i> very open tussock grassland and <i>Triodia epactia</i> scattered hummock grasses.		
	H2d	<i>Acacia inaequilatera</i> scattered tall shrubs over <i>Triodia wiseana</i> (<i>T. epactia</i>) open hummock grassland.		
	H2e	<i>Corymbia flavescens</i> scattered low trees over <i>Acacia colei</i> var. <i>colei</i> scattered tall shrubs over <i>Acacia stellaticeps</i> low open shrubland over <i>Triodia epactia</i> open hummock grassland over * <i>Cenchrus ciliaris</i> scattered tussock grasses.		
	H3	<i>Acacia inaequilatera</i> scattered shrubs over <i>Triodia epactia</i> scattered to open hummock grassland.	210.3	0.4
	Total		5,934.9	11.8
Rocky Outcrops and Breakaways	R1	<i>Atalaya hemiglauca</i> scattered low trees over <i>Acacia colei</i> scattered shrubs over <i>Triodia epactia</i> very open hummock grassland.	52.0	0.1
	R2a	<i>Corymbia hamersleyana</i> low open woodland over <i>Acacia tumida</i> var. <i>pilbarensis</i> , <i>Acacia acradenia</i> tall open scrub over <i>Triodia epactia</i> (<i>T. wiseana</i>) open to very open hummock grassland over * <i>Cenchrus ciliaris</i> scattered tussock grasses.	3,002.8	6.0

PROTECTED


Landform	Vegetation type	Description	Extent (ha)	% of PGL Surveyed Area
	R2b	<i>Triodia epactia</i> hummock grassland.		
	R2c	<i>Terminalia circumalata</i> scattered tall shrubs to tall shrubland over <i>Triodia epactia</i> hummock grassland to very open hummock grassland over mixed scattered tussock grasses.		
	R2d	<i>Acacia inaequilatera</i> , <i>Grevillea</i> spp. scattered tall shrubs over <i>Triodia epactia</i> open hummock grassland.		
	R3	<i>Acacia orthocarpa</i> open shrubland over <i>Triodia epactia</i> very open hummock grassland over <i>Eriachne mucronata</i> (Typical Form) scattered tussock grasses over <i>Goodenia scaevolina</i> scattered herbs.	17.0	<0.1
	R4	<i>Acacia colei</i> scattered to tall open shrubland over <i>Triodia epactia</i> very open hummock grassland.	7.4	<0.1
	R5a	<i>Acacia inaequilatera</i> scattered tall shrubs over <i>Triodia epactia</i> open to very open hummock grassland.	352.7	0.7
	R5b	<i>Corymbia hamersleyana</i> scattered low trees over <i>Acacia tumida</i> var. <i>pilbarensis</i> tall shrubland over <i>Triodia epactia</i> open hummock grassland.		
	R6	<i>Acacia ancistrocarpa</i> , <i>Acacia colei</i> tall open shrubland over <i>Acacia adoxa</i> var. <i>adoxa</i> low open shrubland over <i>Triodia epactia</i> open hummock grassland.	969.1	1.9
	Total		4,401.0	8.8
Cleared/ N/A	NA	Cleared vegetation for roads, mining, rails and existing tracks.	603.5	1.2



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area - Permit Activities

**Environmental Constraints
Vegetation Type**

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 5.1



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area- Permit Activities

Environmental Constraints
Vegetation Type

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 5.2



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020





Pilbara Green Link
Investigation Area - Permit Activities

**Environmental Constraints
Vegetation Type**

Project No. 12657458
Revision No. B
Date 4/06/2025

FIGURE 5.3






Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area - Permit Activities

Environmental Constraints
Vegetation Condition

Project No. 12657458
Revision No. B
Date 06/01/2025


FIGURE 6.1



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres


Horizontal Datum: GDA2020
Grid: GDA2020



Pilbara Green Link
Investigation Area - Permit Activities
**Environmental Constraint
Vegetation Condition**

Project No. 12657458
Revision No. B
Date 06/01/2025


FIGURE 6.2



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020



Pilbara Green Link
Investigation Area - Permit Activities

**Environmental Constraints
Vegetation Condition**

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 6.3



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020



Horizon Power
Pilbara Green Link - Permit Activities

**Environmental Constraints -
Vegetation Condition**

Project No. 12657458
Revision No. B
Date 5/06/2025

FIGURE 6.4

3.2.5 Significant ecological communities

The Protected Matters Search Tool (PMST) search did not identify any Threatened Ecological Communities (TECs) previously recorded within the PGL Surveyed Area. The database search indicated one Priority Ecological Community (PEC) intersecting the PGL Surveyed Area, the Gregory Land System (DBCA 2022). DBCA listed this PEC as Priority 3.

Results of the Biota (2024) survey indicated no vegetation types representative of TECs or PECs were recorded. Post-field survey, the Gregory Land System classification was identified as a misclassification by DBCA. The area is instead classified as a non-listed Granitic Land System, therefore the PGL Alignment Area does not intersect a significant ecological community.

Similarly, the results of the Biota (2025) survey indicated no vegetation types representative of TECs or PECs were recorded.

The riparian vegetation type D2 (the De Grey River and its tributaries) is considered as having a high potential to be a Groundwater Dependent Ecosystem (GDE) or Groundwater Dependent Vegetation (GDV) due to the phreatophytic species *Eucalyptus camaldulensis*, *Eucalyptus victrix* and *Melaleuca argentea* being dominant taxa. In total, vegetation type D2 represents 577.2 ha of the PGL Surveyed Area (1.1% of the PGL Surveyed Area).

3.2.6 Flora diversity

The Dandjoo search identified 871 vascular flora species previously recorded within the PGL Surveyed Area (Appendix C). During the field survey, Biota (2024 & 2025) recorded a total of 426 native flora species from 148 genera and 51 families. Fabaceae and Poaceae represented the two most dominant native families. These families and genera are typical of species lists from the region.

3.2.7 Conservation significant flora

No threatened species were recorded during the Biota field surveys (2024 & 2025).

The Biota (2024 & 2025) field surveys identified eleven Priority species, comprising of one P2 species, ten P3 species across the PGL Surveyed Area (Table 9 and Figure 5). Records of all Priority species represented new populations of the species. For two of the recorded Priority species, these new populations are considered significant range extensions:

- *Bonamia oblongifolia* (P3)
- *Euphorbia inappendiculata* var. *queenslandica* (P3)

Biota (2024) concluded an additional 7 Priority species, which were not detected during the field survey, as *likely to occur* and an additional 8 species as *may occur* in the PGL Surveyed Area (Table 9). No additional species were identified during the 2025 Biota field survey. The guidelines applied to determine the flora likelihood of occurrence are detailed in Table 10.

Table 9 Conservation significant flora - Likelihood of occurrence assessment (Biota 2025)

Species	EP Act/DBCA Status
Known	
<i>Goodenia hartiana</i>	P2
<i>Abutilon</i> sp. <i>Pritzelianum</i> (S. van Leeuwen 5095)	P3
<i>Bonamia oblongifolia</i>	P3
<i>Bulbostylis burbridgeae</i>	P3
<i>Croton aridus</i>	P3
<i>Euphorbia clementii</i>	P3
<i>Euphorbia inappendiculata</i> var. <i>inappendiculata</i>	P3
<i>Euphorbia inappendiculata</i> var. <i>queenslandica</i>	P3

Species	EP Act/DBCA Status
<i>Indigofera ammobia</i>	P3
<i>Polymeria</i> sp. Broome (K.F. Kenneally 9759)	P3
<i>Tribulopsis marliesiae</i>	P3
Likely	
<i>Triodia degreyensis</i>	P1
<i>Euploca mutica</i>	P3
<i>Hibiscus</i> aff. <i>krichauffianus</i>	P3
<i>Nicotiana umbratica</i>	P3
<i>Phyllanthus</i> sp. aff. <i>herbecarpus</i>	P3
<i>Rothia indica</i> subsp. <i>australis</i>	P3
<i>Terminalia kumpaja</i>	P3
May	
<i>Acacia cyperophylla</i> var. <i>omearana</i>	P1
<i>Euploca parviantrum</i>	P1
<i>Tephrosia rosea</i> var. Port Hedland	P1
<i>Gomphrena leptophylla</i>	P3
<i>Gymnanthera cunninghamii</i>	P3
<i>Heliotropium murinum</i>	P3
<i>Triodia chichesterensis</i>	P3
<i>Ptilotus mollis</i>	P4

Table 10 Flora likelihood of occurrence assessment against relevant guidelines (Biota 2025)

Rank/ Likelihood	Criteria
Recorded	1. The species has been recorded in the PGL Surveyed Area
Likely to occur/ High	1. There are existing records of the species in proximity to the PGL Surveyed Area; and <ul style="list-style-type: none"> The species is strongly linked to a specific habitat, which is present in the PGL Surveyed Area; or The species has more general habitat preferences, and suitable habitat is present.
May occur / Moderate	1. There are existing records of the species from the locality, however <ul style="list-style-type: none"> The species is strongly linked to a specific habitat, of which only a small amount is present in the PGL Surveyed Area; or The species has more general habitat preferences, but only some suitable habitat is present 2. There is suitable habitat in the PGL Surveyed Area, but the species is recorded infrequently in the locality.
Unlikely to occur/ Low	1. The species is linked to a specific habitat, which is absent from the PGL Surveyed Area; or 2. Suitable habitat is present, however there are no existing records of the species from the locality despite reasonable previous search effort in suitable habitat; or 3. There is some suitable habitat in the PGL Surveyed Area, however the species is very infrequently recorded in the locality, or the only records are historical (>40 years old).
Would not occur/ Negligible	1. The species is strongly linked to a specific habitat, which is absent from the PGL Surveyed Area; or 2. The species' range is very restricted and does not include the PGL Surveyed Area; or 3. The species is not considered extant in the locality

3.2.8 Weeds

Biota (2024 & 2025) recorded the presence of 15 introduced species within the PGL Surveyed Area (Figure 5). Two of these weed species represented Declared Pests; *Calotropis procera* (Calotrope) and *Parkinsonia aculeata* (Jerusalem thorn). The latter species is also considered a weed of national significance (Department of Agriculture and Water Resources 2017).

3.3 Fauna

3.3.1 Vertebrate fauna habitat

From the field surveys (Biota 2024 & 2025), nine fauna habitats were identified within PGL Surveyed Area (Figure 7):

- Acacia shrubland on spinifex sandplain – 32,790.2 ha
- Granite boulders – 3,836.2 ha
- Gorges and gullies – 3,000.6 ha
- Claypan – 2,902.3 ha
- Minor/moderate drainage lines – 2,631.6 ha
- Low stony rises – 2,505.3 ha
- Rocky outcrops – 993.5 ha
- Major drainage lines – 919.0 ha
- Sand dunes – 21.0 ha
- Cleared areas – 603.5 ha

Acacia shrubland on spinifex sandplain was the dominant habitat type within the PGL Surveyed Area, representing 65.3%. The other nine habitat types represent between 7.6% and <0.1% of the PGL Surveyed Area. These habitats are described in further detail below in Table 11 and shown Figure 7.

Significant fauna species likely to inhabit each habitat have been identified by the Biota (2024 & 2025) and listed in Table 11. Migratory species were not included in the table as most habitat types were deemed as suitable and/or the habitats were not breeding dependent (Biota 2025). These species will be reviewed in the assessment against the 10 clearing principles.


PROTECTED

Table 11 *Fauna Habitats within the PGL Surveyed Area*

Fauna Habitat	Habitat description	Area (ha)	Proportion of PGL Surveyed Area (%)	Fauna species likely to inhabit
Acacia shrubland on spinifex sandplain	Open Acacia shrubland with some denser patches and scattered <i>Corymbia</i> and <i>Eucalyptus victrix</i> , including <i>A. stellaticeps</i> , <i>A. ancistrocarpa</i> , <i>A. monticola</i> , over hummock grasslands of spinifex (<i>Triodia</i> spp.) on sandplain	32,790.2	65.3	<ul style="list-style-type: none"> – Bilby – Brush-tailed Mulgara – Northern Quoll – Spectacled Hare-wallaby – Western Pebble-mound Mouse – Pilbara Leaf-nosed Bat – Grey Falcon
Granite boulders	Granite boulders with scattered <i>Corymbia</i> spp. and <i>Acacia</i> spp. over open <i>Triodia</i> hummock grassland.	3,836.2	7.6	<ul style="list-style-type: none"> – Northern Quoll – Ghost Bat – Pilbara Leaf-nosed Bat – Grey Falcon – Black-footed Rock-wallaby
Gorges and gullies	Gorges, gullies, large breakaways and associated foot slopes. <i>Corymbia hamersleyana</i> low open woodland over <i>Acacia</i> spp., over <i>Triodia</i> grassland.	3000.6	6.0	<ul style="list-style-type: none"> – Northern Quoll – Western Pebble-mound Mouse – Pilbara Olive Python – Ghost Bat – Pilbara Leaf-nosed Bat – Grey Falcon – Black-footed Rock-wallaby
Claypan	<i>Triodia</i> hummock and tussock grasslands on clay.	2,902.3	5.8	<ul style="list-style-type: none"> – Short-tailed Mouse
Minor/moderate drainage line	Minor to moderate drainage lines and floodplains fringed with low eucalypts (<i>Eucalyptus victrix</i> , <i>Corymbia flavescens</i>) and <i>Lysiphyllum cunninghamii</i> over sparse shrubland and tussock grassland	2,631.6	5.2	<ul style="list-style-type: none"> – Brush-tailed Mulgara – Ghost Bat – Northern Quoll – Pilbara Olive Python – Pilbara Leaf-nosed Bat – Grey Falcon
Low stony rises	Rocky undulating plains, rises and slopes with small outcrops. Mixed <i>Acacia</i> tall shrubland with some scattered <i>Corymbia</i> ; <i>Grevillea wickhamii</i> , <i>Triodia epactia</i> , and open tussock grasses.	2,505.3	5.0	<ul style="list-style-type: none"> – Northern Quoll – Western-Pebble-mound Mouse
Rocky outcrops	Breakaways and large complex rocky outcropping and, vegetated with <i>Corymbia</i> spp., mixed <i>Acacia</i> spp., <i>Triodia epactia</i> and scattered tussock grasses.	993.5	2.0	<ul style="list-style-type: none"> – Western Pebble-mound Mouse – Northern Quoll – Ghost Bat – Pilbara Leaf-nosed Bat – Black-footed Rock-wallaby

PROTECTED


Fauna Habitat	Habitat description	Area (ha)	Proportion of PGL Surveyed Area (%)	Fauna species likely to inhabit
Major drainage lines	Major drainage lines fringed by eucalypts (<i>Eucalyptus camaldulensis</i> , <i>E. victrix</i>), <i>Melaleuca argentea</i> over tall shrubland and open <i>Triodia</i> grassland.	919.0	1.8	<ul style="list-style-type: none"> – Northern Quoll – Ghost Bat – Pilbara Leaf-nosed Bat – Pilbara Olive Pythons – Brush-tailed Mulgara – Short-tailed Mouse – Grey Falcon
Sand dunes	<i>Corymbia</i> low open woodland over <i>Acacia</i> tall shrubland and open grassland on sand dunes.	21.0	<0.1	<ul style="list-style-type: none"> – Bilby – Brush-tailed Mulgara – Dampierland Plain Slider
Cleared areas		603.5	1.2	



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area - Permit Activities

Environmental Constraints
Fauna Habitat

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 7.1



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020




Pilbara Green Link
Investigation Area- Permit Activities

**Environmental Constraints
Fauna Habitat**

Project No. 12657458
Revision No. B
Date 4/06/2025


FIGURE 7.2



Paper Size ISO A3
at Scale: 1:200,000

0 0.5 1
Kilometres

Horizontal Datum: GDA2020
Grid: GDA2020



Pilbara Green Link
Investigation Area - Permit Activities

Environmental Constraints
Fauna Habitat

Project No. 12657458
Revision No. B
Date 5/06/2025

FIGURE 7.3

3.3.2 Vertebrate fauna diversity

The Dandjoo database search (Appendix C) identified 642 vertebrate fauna species previously recorded within 20 km of the PGL Surveyed Area. This included:

- 309 Birds
- 170 Reptiles
- 75 Fish
- 71 Mammals
- 17 Amphibians

Biota (2024 & 2025) identified 130 vertebrate species comprising of 20 mammals (including 11 bat species), 81 bird species, 24 reptile species and one amphibian species within the PGL Surveyed Area.

Of the four species, four introduced fauna species were recorded: Dingo (*Canis familiaris*), Feral Domestic Cat (*Felis catus*), Camel (*Camelus dromedarius*) and European Cattle (*Bos primigenius taurus*).

3.3.3 Conservation significant vertebrate fauna

The Biota field surveys (2024 & 2025) confirmed the presence of eight Conservation Significant fauna species within the PGL Surveyed Area (Table 12). A further five species have been described as likely to occur due based on previous records within the PGL Surveyed Area and/or the availability of suitable habitat. Ten species were identified as may occur in the PGL Surveyed Area, by Biota (2024 & 2025).

Six of the known species are listed under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) as Matters of National Environmental Significance (MNES) and under the Western Australian *Biodiversity Conservation Act 2016* (BC Act). These species are the Northern Quoll, Bilby and Pilbara Leaf-nosed Bat, Grey Falcon, Pacific Swift, Oriental Pratincole (Department of Climate Change, Energy, the Environment and Water (DCCEEW) 2025). The two other known significant species, the Brush-tailed Mulgara and Western Pebble-mound Mouse, are listed as Priority 4 by DBCA under the BC Act.

Table 12 Conservation Significant Fauna Species in the PGL Surveyed Area

Taxon	Common Name	Status	
		BC Act/DBCA	EPBC Act
Known			
<i>Dasyurus hallucatus</i>	Northern Quoll	Endangered	Endangered
<i>Macrotis lagotis</i>	Bilby	Vulnerable	Vulnerable
<i>Rhinonicteris aurantia</i> (Pilbara form)	Pilbara Leaf-nosed Bat	Vulnerable	Vulnerable
<i>Falco hypoleucos</i>	Grey Falcon	Vulnerable	Vulnerable
<i>Apus pacificus</i>	Pacific Swift	Migratory	Migratory
<i>Glareola maldivarum</i>	Oriental Pratincole	Migratory	Migratory
<i>Dasycercus blythi</i>	Brush-tailed Mulgara	Priority 4	-
<i>Pseudomys chapmani</i>	Western Pebble-mound Mouse	Priority 4	-
Likely			
<i>Liasis olivaceus barroni</i>	Pilbara Olive Python	Vulnerable	Vulnerable
<i>Macroderma gigas</i>	Ghost Bat	Vulnerable	Vulnerable
<i>Gelochelidon [nilotica] macrotarsa</i>	Australian [Gull-billed] Tern	Migratory	Migratory
<i>Hydroprogne caspia</i>	Caspian Tern	Migratory	Migratory

Taxon	Common Name	Status	
		BC Act/DBCA	EPBC Act
<i>Falco peregrinus</i>	Peregrine Falcon	Other specially protected	-
May			
<i>Petrogale lateralis lateralis</i>	Black-footed Rock-wallaby	Endangered	Endangered
<i>Actitis hypoleucos</i>	Common Sandpiper	Migratory	Migratory
<i>Anarhynchus veredus</i>	Oriental Plover	Migratory	Migratory
<i>Chlidonias leucopterus</i>	White-winged Tern	Migratory	Migratory
<i>Plegadis falcinellus</i>	Glossy Ibis	Migratory	Migratory
<i>Numenius minutus</i>	Little Curlew	Migratory	Migratory
<i>Tringa glareola</i>	Wood Sandpiper	Migratory	Migratory
<i>Lerista separanda</i>	Dampierland Plain Slider	Priority 2	-
<i>Lagorchestes conspicillatus</i>	Spectacled Hare-Wallaby	Priority 4	-
<i>Leggadina lakedownesis</i>	Short-tailed Mouse	Priority 4	-

3.3.4 Invertebrate fauna habitats

Ten fauna habitats were identified and mapped through field habitat assessment and consideration of the ecological niches relevant to SRE fauna (Biota 2024). Table 13 describes the microhabitats likely to support SRE species within each of the broader fauna habitat units of the PGL Surveyed Area.

Based on the review of spatial data and ground-truthing in contextual areas, the SRE fauna habitats present are not confined to the PGL Surveyed Area and extend contiguously beyond, reducing the likelihood that any species would be restricted in distribution to the PGL Surveyed Area.

Table 13 Microhabitats likely to support SRE species

Fauna Habitat	SRE Microhabitat Description
Acacia shrubland on spinifex sandplain	Accumulations of leaf litter, areas of clayey soils, mature <i>Triodia</i> hummocks.
Granite boulders	Mature <i>Triodia</i> hummocks, leaf litter accumulations, granite outcrops.
Gorges and gullies	Accumulations of leaf litter, areas of clayey soils, mature <i>Triodia</i> hummocks.
Claypan	Extensive areas of high clay content soils, mature <i>Triodia</i> hummocks
Minor/moderate drainage line	Leaf litter accumulations, areas of high clay-content soils on floodplains.
Low stony rises	Sandy/stony soils, accumulations of leaf litter, patches of mature <i>Triodia</i> hummocks.
Rocky outcrops	Accumulations of leaf litter.
Major drainage lines	Floodplains of high clay content soils, permanent pools, leaf litter accumulations
Sand dunes	Nil
Cleared areas	Nil

3.3.5 Invertebrate diversity

The Biota 2024 field survey identified the following invertebrates across the PGL Alignment Area:

- 11 Mygalomorph spiders from 11 sites
- 16 Camaenid Snails from six sites
- Two Scorpions from two sites
- Two Pseudoscorpions from one site

- 11 Isopods from six sites
- Three Centipedes from three sites
- One Selenopid spider from one site

The Biota 2025 field survey did not survey for invertebrates.

3.4 Conservation Areas

There are no DBCA Legislated Lands or Waters occurring within the DE (DWER 2024).

3.5 Environmentally Sensitive Areas

There is one Environmentally Sensitive Area (ESA) that intersects the PGL Surveyed Area, the De Grey River (DWER 2021b). The PGL Surveyed Area overlaps 300 ha of the De Grey River ESA, which is listed on the Directory of Important Wetlands of Australia (DCCEEW 2018), and is also listed as an ESA under the EP Act (DWER 2021b)(Figure 8). This wetland is known to support some significant fauna species such as the Grey Falcon (Vulnerable), the Far Eastern Curlew (Vulnerable) and the Pilbara Dragonfly (Priority 2). The De Grey River also has high potential to support vegetation representing GDEs (Bureau of Meteorology 2024).

In line with the PGL Preliminary Design Principles, wherever possible, the Investigation pads are not within 50 m of the De Grey ESA. There is however, one Investigation Pad (Investigation Pad 6 containing a single geotechnical test pit) which is located approximately 10 m from the ESA boundary. All other Investigation Pads identified in this NVCP are not located within the De Grey River.

3.6 Hydrology

Desktop review of Government of Western Australia's hydrology related data layers identified the water resources present in the DE. There are detailed below in Table 14.

Table 14 Water Resources within the DE

Aspect	Details	Results
Groundwater Areas (Department of Water and Environmental Regulation (DWER), 2018a)	Groundwater areas proclaimed under the <i>Rights in Water and Irrigation Act 1914</i> (RiWI Act)	Pilbara Groundwater Area <ul style="list-style-type: none"> – Ashburton Subarea – East Pilbara Subarea Canning-Kimberly Groundwater Area <ul style="list-style-type: none"> – West Canning-Yarrie Subarea
Surface Water Areas	Surface water areas proclaimed under the RiWI Act	Pilbara Surface water Area
Irrigation District	Irrigation Districts proclaimed under the RiWI Act	None present
Rivers	Rivers proclaimed under the Rights in RiWI Act	None present
Public Drinking Water Source Areas (DWER 2024b)	PDWSA is a collective term used for the description of Water Reserves, Catchment Areas and Underground Pollution Control Areas declared (gazetted) under the provisions of the <i>Metropolitan Water Supply, Sewage and Drainage Act 1909</i> (WA) or the <i>Country Areas Water Supply Act 1947</i> (WA)	De Grey River Water Reserve (shown in Figure 8)
Waterways Management Area	Areas proclaimed under the Waterway Conservation Act 1976 (WA). These are Albany waterways, Avon River, Wilson Inlet, Peel– Harvey estuaries and Leschenault Inlet.	None present

3.6.1 Groundwater

The DE intersects one Public Drinking Water Source Area: De Grey River Water Reserve (DWER 2024) (Figure 8). Within this water reserve, the De Grey River wellfield is used to supply water for Port Hedland (DWER 2018b). The De Grey River wellfield is operated by the Water Corporation and draws groundwater from the alluvial deposits of the De Grey River.

The DE falls within the Pilbara Groundwater Area and Canning-Kimberly Groundwater Area. Groundwater in the Pilbara originates from direct infiltration by rainfall and surface water flows, such as rivers. The most significant aquifers in the region can be grouped into three types:

- Alluvial aquifers that are either unconsolidated sedimentary aquifers or chemically deposited aquifers
- Consolidated sedimentary (or sedimentary rock) aquifers
- Fractured rock aquifers

Groundwater in the Pilbara region is typically fresh, with low salinity levels (200-1500 mg/L). Salinity is higher in well sites than bores and mean Total Dissolved Solids (TDS) was found to increase toward the coast (Reeves et al. 2007).

3.6.2 Surface water and drainage

De Grey River is intersected by the PGL Surveyed Area, crossing through the central portion of the area. This system of permanent river pools constitutes a significant drought refuge area for freshwater fishes and waterbirds in the bioregion (DWER 2018c). Drainage through the PGL Surveyed Area is shown in Figure 8.

The DE is within the Indian Ocean and Western Plateau catchment divisions of Western Australia. Within these divisions, the DE falls within the De Grey River, Port Hedland Coast, and Sandy Desert basins. The DE is spread out between three different catchments, and three different sub catchment management areas within these catchments, as detailed in Table 15.


Table 15 *Catchments that intersect the DE*

Catchment Order				Area (ha)
Division	Basin	Catchment	Sub catchment	
Indian Ocean	De Grey River	De Grey River	De Grey	38.4
		Coastal	De Grey	28.8
	Port Hedland Coast	Coastal	Turner	7.5
Western Plateau	Sandy Desert	Sandy Desert Lake Dora	Sandy Desert Basin	158.0

3.6.3 Wetlands

The PMST search identified one Wetland of International Importance (Ramsar) within the 30-40 km of the closest point of the DE, Eighty Mile Beach (DBCA 2017). No indirect impacts to the Ramsar site are anticipated.

The De Grey River, which is also classified as a Nationally Important Wetland, is in close proximity to the DE (Figure 8) (DBCA 2018). In line with the PGL Preliminary Design Principles, wherever possible, the Investigation pads are not within 50 m of the riverbed. There is however, one Investigation Pad (Investigation Pad 6 containing a single geotechnical test pit) which is located approximately 10 m from the wetland boundary. All other Investigation Pads identified in this NVCP are not located within the De Grey River.



3.7 Land use

3.7.1 Land use and sensitive receptors

The Pilbara area and landscape surrounding the Investigation Activities are remote and mainly used for iron ore mining and pastoral activities (cattle grazing). Within this broader region, notable sensitive receptors are conservation areas and reserves. No conservation areas/reserves are intersected by the DE or the broader PGL Alignment Area (Landgate 2024).

The nearest localities to the PGL Surveyed Area are listed in Table 16 below. Table 23 identifies the Land Parcels which will be impacted by the clearing required for Investigation Activities to occur. Ten Investigation Pads will be located within the former Goldsworthy townsite.

Table 16 Localities surrounding the PGL Surveyed Area


Locality	Distance from the PGL Surveyed Area
Goldsworthy (historic locality)	Intersected by the PGL Surveyed Area
Shay Gap (historic locality)	Intersected by the PGL Surveyed Area
Port Hedland	1.8 km
Finucane Island	12.6 km
Shellborough (historic locality)	31.7 km

3.7.2 Contaminated Sites

There are no contaminated sites mapped within the DE (DWER 2024a).

3.7.3 Acid Sulfate Soils

Within the PGL Surveyed Area, the area surrounding the De Grey River and its tributaries is classified as Moderate to low risk of Acid Sulfate Soils (ASS) (DWER 2024) (Figure 9). The DE associated with TP18 intersects the Moderate to low-risk ASS area.



\\ghdnet\ghd\AU\Perth\Projects\12657458\GIS\Maps\Working\12657458_NVCP2_Figures\12657458_NVCP2_Figures.aprx\12657458_01_A_05AcidSulfateSoils_A3L1
Print date: 04 Jun 2025 - 16:08

Data source: World Imagery; Earthstar Geographics
World Imagery: Maxar. Created by: nstjepanovic

4. Assessment of impacts

4.1 Potential impacts to vegetation and flora

4.1.1 Vegetation

Potential impacts to vegetation include:

- Direct impact to native vegetation as a result of the mechanical clearing of a maximum of 84.1 ha (within the 232.6 ha DE) to allow for the development of access tracks to enable light and heavy vehicle access to the investigation locations.
- Direct impact to native vegetation as a result of the mechanical clearing of a maximum of 9.1 ha (within the 232.6 ha DE) for geotechnical boreholes and test pits and groundwater boreholes.
- Indirect impact to native vegetation as a result of the potential introduction or spread of weeds during Investigation Activities.
- Indirect impact to native vegetation as a result of fugitive dust generated during Investigation Activities

The vegetation types and associated condition for each vegetation type proposed to be cleared for the Investigation Activities is detailed in Table 17.

Table 17 Vegetation types and condition within the Indicative Impact Area

Vegetation Type (VT)	Vegetation Condition (ha)					Total Indicative Impact Area (ha)
	Excellent	Very Good - Excellent	Very Good	Poor - good	Cleared/ Degraded	
P1	1.5	0.1	1.8			3.1
P2				<0.1		<0.1
P3	64.7	1.7				66.3
P4			5.8			5.8
H1	12.3	1.0				13.2
H2	2.5					2.5
R2			0.6			0.6
R5	0.1					0.1
D2			0.3			0.3
S2	0.3					0.3
Cleared					0.7	0.7
Total						93.3

4.1.2 Flora

No Threatened flora species are proposed to be cleared within the DE.

Priority 3 flora species' *Croton aridus* and *Triobulopsis marliesiae* were identified along areas of the DE designated to potential access tracks. There is potential for each species to be impacted by the maximum proposed clearing of 84.1 ha for access tracks. However, as detailed in Section 2.1, the extent of clearing for tracks is likely to be substantially less than the maximum proposed, and the distribution of both species is widespread across the north-east of Western Australia and Northern Territory (Atlas of Living Australia, 2025 & Biota, 2025).

4.2 Potential impacts to fauna and fauna habitat

Potential impacts to fauna and fauna habitat include:

- Direct impact to fauna habitat as a result of approximately 93.3 ha of largely temporary clearing (total) (Table 18)
- Direct impact to Conservation Significant fauna habitat as a result of approximately 93.3 ha of largely temporary clearing (total) (Table 19).
- Direct impact to Conservation Significant fauna species as a results of increased vehicle movements and potential vehicle strike
- Indirect impact to fauna habitat through the potential introduction and/or spread of weeds as a result of clearing, increased vehicle movement and topsoiling during the Investigation Activities.

There is potential for Conservation Significant fauna and fauna habitat to be impacted by the Investigation Activities. However, clearing of the maximum 93.3 ha in the Indicative Impact Area amounts to 0.2% of the 50,203.3 ha of native vegetation within the PGL Surveyed Area.

Table 18 Fauna habitat types within the Indicative Impact Area

Fauna Habitat type	Description	Area within PGL Surveyed Area (ha)	Indicative Impact Area – maximum proposed clearing (ha)	Proportion of potential habitat in PGL Surveyed Area (%)
Acacia shrubland on spinifex sandplain	Open Acacia shrubland with some denser patches and scattered <i>Corymbia</i> and <i>Eucalyptus victrix</i> , including <i>A. stellaticeps</i> , <i>A. ancistrocarpa</i> , <i>A. monticola</i> , over hummock grasslands of spinifex (<i>Triodia</i> spp.) on sandplain	32,790.2	74.6	0.2%
Low stony rises	Rocky undulating plains, rises and slopes with small outcrops. Mixed <i>Acacia</i> tall shrubland with some scattered <i>Corymbia</i> ; <i>Grevillea wickhamii</i> , <i>Triodia epactia</i> , and open tussock grasses.	2505.3	17.1	0.7%
Minor/moderate drainage lines	Minor to moderate drainage lines and floodplains fringed with low eucalypts (<i>Eucalyptus victrix</i> , <i>Corymbia flavescentes</i>) and <i>Lysiphyllum cunninghamii</i> over sparse shrubland and tussock grassland	2,631.6	0.4	<0.1%
Sand dunes	<i>Corymbia</i> low open woodland over <i>Acacia</i> tall shrubland and open grassland on sand dunes.	21.0	0.3	1.6%
Granite Boulders	Granite boulders with scattered <i>Corymbia</i> spp. and <i>Acacia</i> spp. over open <i>Triodia</i> hummock grassland.	3,836.2	<0.1	<0.1%
Cleared areas	Mining, disturbances or cleared areas.	603.5	0.7	0.1%

The impact to suitable habitat for each Conservation Significant fauna species (Table 19) was calculated from the known fauna habitat used by the species (Table 11) and the clearing proposed within each fauna habitat (Table 18).

Table 19 Disruption of potential habitat clearing on Conservation Significant fauna species

Fauna Species	BC Act/ DBCA	Maximum potential area of habitat to be cleared (ha)	Total area of potentially suitable habitat within PGL Surveyed Area (ha)	Proportion of potential habitat impacted (%)
<i>Dasyurus hallucatus</i> - Northern Quoll	EN	92.2	41,763.3	0.2%
<i>Macrotis lagotis</i> - Bilby	V	74.6	32811.2	0.2%
<i>Rhinonicteris aurantia</i> (Pilbara form) - Pilbara Leaf-nosed Bat	V	75.1	39258.0	0.2%

PROTECTED

Fauna Species	BC Act/ DBCA	Maximum potential area of habitat to be cleared (ha)	Total area of potentially suitable habitat within PGL Surveyed Area (ha)	Proportion of potential habitat impacted (%)
<i>Dasycercus blythi</i> - Brush-tailed Mulgara	P4	75.4	35442.8	0.2%
<i>Pseudomys chapmani</i> - Western Pebble-mound Mouse	P4	91.4	35295.5	0.3%
<i>Macroderma gigas</i> - Ghost Bat	V	0.8	6467.8	<0.1%
<i>Falco hypoleucos</i> - Grey Falcon	V	75.1	39258.0	0.2%
<i>Liasis olivaceus barroni</i> - Pilbara Olive Python	V	0.7	2631.6	<0.1%
<i>Petrogale lateralis lateralis</i> - Black-footed Rock-Wallaby	EN	<0.1	3836.2	<0.1%
<i>Lagorchestes conspicillatus</i> - Spectacled Hare-Wallaby	P4	74.3	32790.2	0.2%
<i>Lerista separanda</i> – Dampierland Plain Slider	P2	0.3	21.0	<0.1%

5. Assessment against 10 clearing principles

The clearing of vegetation in Western Australia is regulated by DWER and requires a permit under Part V of the EP Act, except when a Project is assessed under Schedule 6 of the Act or is prescribed by regulation in the *Environmental Protection (Clearing Native Vegetation) Regulations 2004*.

In making a decision about a clearing permit application under section 51O of the EP Act, the CEO of DWER must consider the clearing principles contained in Schedule 5 of the EP Act so far as they are relevant to the matter under consideration. The ten clearing principles aims to ensure that potential impacts resulting from removal of native vegetation can be assessed holistically.

To support the NVCP application for the Investigation Activities associated with the prefeasibility phase of the PGL Project, an assessment of the proposed clearing against the ten clearing principles outlined in Schedule 5 of the EP Act has been undertaken.

The assessment was undertaken with reference to DWER guideline “*A guide to the assessment of applications to clear native vegetation under Part V Division 2 of the Environmental Protection Act 1986*” (DWER 2014).

This assessment concluded the proposed clearing associated with the Investigation Activities is unlikely to be or not at variance to the clearing principles.

(a) Native vegetation should not be cleared if it comprises a high level of biological diversity.

Native vegetation should not be cleared if it comprises a high level of biological diversity.

Proposed clearing is not at variance to this Principle.

Assessment

Vegetation

The vegetation within the DE, is consistent and contiguous with the native vegetation in the surrounding area. As discussed in 3.2.2, all broad-scale Vegetation Associations are well represented across State, IBRA Region, IBRA Subregion, and LGA extents. Most pre-European vegetation associations have 99% or more of their pre-European extent remaining.

24 vegetation types were identified and mapped, stemming from six broad landforms, within the PGL Surveyed Area. Four units from drainage areas, four from clay plains, five from plains, three from low stony rises, six from rocky outcrops and breakaways, and two units from sand dunes. The DE will involve the clearing of vegetation within nine of these vegetation types, described in Table 17.

Most clearing will occur in vegetation types P3 and H1, with a loss of 0.4% (66.3 ha) and 0.7% (13.2 ha) of available vegetation type in the PGL Surveyed Area, respectively. All vegetation types are typical of the wider region. The proportion of clearing proposed in each vegetation type is unlikely to significantly impact the biological diversity of the region.

The condition of the mapped vegetation types ranged from Cleared to Excellent. The mapping indicates that the PGL Surveyed Area is predominantly in Excellent and Very Good to Excellent condition, with these conditions representing 34.5% and 37.6% respectively. Areas completely devoid of native vegetation were mapped as Cleared ('NA') and were not assigned a condition rating. In total, 1.20% of the PGL Surveyed Area has been completely cleared of vegetation (e.g. roads, rails, mining area). As it is a high proportion of the PGL Surveyed Area, clearing of vegetation is proposed to occur in areas that are considered Excellent and Very Good to Excellent condition. Where possible, clearing for the DE has been avoided by using previously cleared areas (e.g. use of existing tracks) or prioritised in areas of lesser condition value.

The condition of vegetation proposed to be cleared within the DE is described below

Vegetation condition	Proposed Indicative Impact Area (ha)	Proportion of PGL Surveyed Area (%)
Excellent	81.3	0.2
Very Good to Excellent	2.7	<0.1
Very Good	8.5	<0.1
Good to Very Good	0	0
Good	0	0
Poor to Good	0.7	<0.1

Threatened and Priority Ecological Communities

The desktop assessment and Biota field survey (2025) identified no potential TECs or PECs within the PGL Surveyed Area.

The riparian vegetation types D1 and D2 (the De Grey River and its tributaries) are considered as having a high potential to be GDEs or GDVs due to the phreatophytic species *Eucalyptus camaldulensis*, *Eucalyptus victrix* and *Melaleuca argentea* being dominant taxa. The DE intersects 0.3 ha of D2 with 1.0 ha associated with DS_01b and the remaining 0.3 ha associated with a section of the proposed access track to TP50. Micro-siting of the Indicative Impact Area within Investigation Pad 3 (DS_01b) and clearing for access track to TP50 will aim to minimise potential clearing of vegetation type D2. The DE of 0.6 ha of this vegetation type represents <0.1% of the available vegetation type within the PGL Surveyed Area.

Flora Diversity

The Dandjoo search identified 871 vascular flora species previously recorded within the PGL Surveyed Area. During the field survey, Biota (2024 & 2025) recorded a total of 426 native flora species from 148 genera and 51 families. Fabaceae and Poaceae represented the two most dominant native families. These families and genera recorded within, are typical of species lists from the region.

Significant flora

No threatened flora taxa pursuant to the EPBC Act and/or gazetted as Threatened pursuant to the Biodiversity and Conservation Act 2016 (BC Act) were recorded within the PGL Surveyed Area. No Threatened flora species are proposed to be cleared within the DE

The Biota (2024 & 2025) field survey has identified eleven DBCA Priority species occurring with the PGL Surveyed Area comprising:

- One Priority 2 species
- Nine Priority 3 species; and
- One Priority 4 species

Native vegetation should not be cleared if it comprises a high level of biological diversity.

Following a post-survey analysis an additional 15 conservation significant flora species were included in the likelihood of occurrence assessment. Of the 26 significant flora species, 18 were considered to have a high likelihood of occurrence (Table 9).

Priority 3 flora species' *Croton aridus* and *Triobulopsis marliesiae* were identified along areas of the DE designated to potential access tracks. There is potential for each species to be impacted by the maximum proposed clearing of 84.1 ha for access tracks. However, as detailed in Section 2.1, the extent of clearing for tracks is likely to be substantially less than the maximum proposed, and the distribution of both species is widespread across the north-east of Western Australia and Northern Territory (Atlas of Living Australia, 2025 & Biota, 2025).

With Investigation Activities able to avoid environmental constraints within each Investigation Pad, an approach of avoid and minimise during clearing will reduce the impact on significant flora species.

Fauna Habitat

From the field survey (Biota, 2025), nine fauna habitats were identified within PGL Surveyed Area. Five of these habitats will be impacted by clearing within the DE (shown Figure 7). The proportion of each fauna habitat impacted is as below:

- Acacia shrubland on spinifex sandplain = 0.2% (74.6 ha)
- Low stony rises = 0.7% (17.1 ha)
- Minor/moderate drainage line = <0.1% (0.4 ha)
- Granite boulders = <0.1% (<0.1 ha)
- Sand dunes = 1.6% (0.3 ha)

The impact to each fauna habitat type within the PGL Surveyed Area is not expected to impact the availability of significant habitat for fauna indigenous to Western Australia. The habitat proposed to be cleared as part of the DE is available consistently and in comparable condition beyond the PGL Surveyed Area. Given the temporary nature of most clearing proposed and distribution of the DE across 250km, it is unlikely there will be a significant impact the availability of habitat for conservation significant fauna

Methodology

- Biological Survey (Biota 2024 & 2025)
- DCCEEW Protected Matters Search Tool Report (DCCEEW 2024b)
- Dandjoo Database Search (DBCA 2024)
- Government GIS Shapefiles:
- DBCA Threatened and Priority Ecological Community database search (Accessed November 2024)
- DBCA Threatened and Priority flora database search (Accessed November 2024)
- DBCA Threatened and Priority fauna database search (Accessed November 2025)
- WA Herbarium Threatened and Priority flora (Accessed November 2024)
- Statewide Vegetation Statistics (Government of Western Australia)

(b) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

Proposed clearing is unlikely to be at variance to this Principle.

Assessment

Fauna Habitat

From the field survey (Biota, 2025), nine fauna habitats were identified within PGL Surveyed Area. Five of these habitats will be impacted by potential clearing within the DE (shown in Table 18). The proportion of each fauna habitat impacted is as below:

- Acacia shrubland on spinifex sandplain = 0.2% (74.6 ha)
- Low stony rises = 0.7% (17.1 ha)
- Minor/moderate drainage line = <0.1% (0.4 ha)
- Granite boulders = <0.1% (<0.1 ha)
- Sand dunes = 1.6% (0.3 ha)

The impact to each fauna habitat type within the PGL Surveyed Area is not expected to impact the availability of significant habitat for fauna indigenous to Western Australia. The habitat proposed to be cleared as part of the DE is available consistently and in comparable condition beyond the PGL Surveyed Area.

Fauna

The field survey (Biota 2025) recorded the presence of five conservation significant vertebrate fauna species. An additional eight species are considered likely to occur or have been previously recorded in the PGL Surveyed Area and a further ten species which may occur within the PGL Surveyed Area.

The impact of the maximum proposed clearing was assessed based on the proportion of habitat the activities would disturb for each species within the PGL Surveyed Area. This is described in the table below (Table 20) for all species considered to potentially occur.

Table 20 Impact of DE clearing on Conservation Significant Fauna Species

Scientific name / Common Name	BC Act / DBCA	Likelihood of occurrence	Impact
<i>Dasyurus hallucatus</i> Northern Quoll	EN	Known	There is 46,676.4 ha of habitat for the Northern Quoll in the PGL Surveyed Area. The proposed clearing will result in impact to 0.2% (92.2 ha) of suitable habitat for the Northern Quoll within the PGL Surveyed Area. The habitats available within the PGL Surveyed Area are representative of the greater Pilbara region. Placement of the proposed DE has avoided areas where the Northern Quoll was recorded during field surveys (Biota 2024 & 2025). A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the local population.
<i>Macrotis lagotis</i> Bilby	V	Known	There is 32,811.2 ha of suitable habitat for Bilby's available within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (75.0 ha) of suitable habitat for the Bilby within the PGL Surveyed Area. Placement of the proposed DE has avoided areas where the Bilby was recorded during field surveys (Biota 2024 & 2025). A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the local population.
<i>Rhinonicteris aurantia</i> (Pilbara form) Pilbara Leaf-nosed Bat	V	Known	There is 44,171.1 ha of suitable habitat for Pilbara Leaf-nosed Bat within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (75.1 ha) of suitable habitat for the Pilbara Leaf-nosed Bat within the PGL Surveyed Area. No Suitable roosting and breeding sites were detected during the field survey (Biota 2024 & 2025). The observed foraging habitat for the species is found consistently within the PGL Surveyed Area and the greater region. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the local population.

Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

<i>Falco hypoleucos</i> Grey Falcon	V	Likely	There is 44,171.1 ha of suitable habitat for Grey Falcon within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (75.1 ha) of suitable habitat for the Grey Falcon within the PGL Surveyed Area. The species has a widespread distribution across northern and arid inland Australia. The observed foraging habitat for the species is found consistently within the PGL Surveyed Area and the greater region (Biota 2024 & 2025). A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Dasycercus blythi</i> Brush-tailed Mulgara	P4	Known	There is 36,361.8 ha of suitable habitat for Brush-tailed Mulgara within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (75.4 ha) of suitable habitat for the Brush-tailed Mulgara within the PGL Surveyed Area. The species has a widespread distribution across WA, typically occurring in spinifex grasslands on sandplains from the Pilbara down to the Gascoyne. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Pseudomys chapmani</i> Western Pebble-mound Mouse	P4	Known	There is 39,289.6 ha of suitable habitat for Western Pebble-mound Mouse within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (91.7 ha) of suitable habitat for the Western Pebble-mound Mouse within the PGL Surveyed Area. The species has a widespread distribution across the Pilbara and Gascoyne regions. Placement of the proposed DE has avoided known Western Pebble-mound Mouse mounds. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Macroderma gigas</i> Ghost Bat	V	Likely	There is 11,380.9 ha of suitable habitat for Ghost Bats within the PGL Surveyed Area. The proposed clearing will result in an impact to <0.01% (0.4 ha) of suitable habitat for the Ghost Bat within the PGL Surveyed Area. No Suitable roosting and breeding sites were detected during the field survey (Biota 2024 & 2025). Foraging habitat for the species is readily available across most habitat types found within the PGL Surveyed Area. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the local population.
<i>Liasis olivaceus barroni</i> Pilbara Olive Python	V	Likely	There is 6,551.1 ha of suitable habitat for Pilbara Olive Python within the PGL Surveyed Area. The proposed clearing will result in an impact 0.01% (0.4 ha) of suitable habitat for the Pilbara Olive Python within the PGL Surveyed Area. The species is distributed across the Pilbara region with large home ranges (between 88 – 449 ha) (Biota 2025). The observed suitable habitat for the species is found consistently within the PGL Surveyed Area and the greater region. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Petrogale lateralis lateralis</i> Black-footed Rock-Wallaby	EN	May	There is 7,830.3 ha of suitable habitat for Black-footed Rock-wallaby within the PGL Surveyed Area. The proposed clearing will result in an impact to <0.01% (0.02 ha) of suitable habitat for the Black-footed Rock-wallaby within the PGL Surveyed Area. Rocky outcrops and boulder fields were avoided in the development of the DE. The observed suitable habitat for the species is found consistently within the PGL Surveyed Area and the greater region. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Lerista separanda</i> Dampierland Plain Slider	P2	May	There is 21 ha of suitable habitat for Dampierland Plain Slider within the PGL Surveyed Area. The proposed clearing will result in an impact to <0.01% (0.3 ha) of suitable habitat for the Dampierland Plain Slider within the PGL Surveyed Area. The observed suitable habitat for the species is found consistently within the PGL Surveyed Area and the greater region. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
<i>Leggadina lakedownesis</i> Short-tailed Mouse	P4	May	There is 3,821.3 ha of suitable habitat for Short-tailed Mouse within the PGL Surveyed Area. The proposed Investigation Project Area will not impact suitable habitat for the Short-tailed Mouse and therefore is unlikely to pose a significant risk to the survival of the species.

Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

<i>Lagorchestes conspicillatus</i> Spectacled Hare-Wallaby	P4	May	There is 32,790.2 ha of suitable habitat for Spectacled Hare-wallaby within the PGL Surveyed Area. The proposed clearing will result in an impact to 0.2% (74.6 ha) of suitable habitat for the Spectacled Hare-wallaby within the PGL Surveyed Area. The observed suitable habitat for the species is found consistently within the PGL Surveyed Area and the greater region. A temporary loss of this proportion of habitat is unlikely to pose a significant risk to the survival of the species.
---	----	-----	--

The following migratory species were excluded from the assessment:

- *Actitis hypoleucos* – Common Sandpiper
- *Anarhynchus veredus* – Oriental Plover
- *Apus pacificus* - Pacific Swift
- *Chlidonias leucopterus* – White-winged Tern
- *Gelochelidon [nilotica] macrotarsa* - Australian [Gull-billed] Tern
- *Glareola maldivarum* - Oriental Pratincole
- *Hydroprogne caspia* – Caspian Tern
- *Numerius minutus* – Little Curlew
- *Plegadis falcinellus* – Glossy Ibis
- *Tringa glareola* – Wood Sandpiper

Foraging for these species either occurs aerially or across most habitats available within the PGL Surveyed Area. All species attend the PGL Surveyed Area as non-breeding migrants and therefore no breeding sites will be impacted by the Investigation Activities. These species are likely to predominately use the PGL Surveyed Area for gathering around drainage areas. A maximum clearing of 0.3 ha in minor/moderate drainage lines associated with access tracks is proposed within the DE. A clearing of this proportion represents <0.1% of this habitat with the PGL Surveyed Area and is unlikely to significantly impact the availability of this habitat for migratory species.

The clearing of 93.3 ha amounts to 0.2% of the 50,203.2 ha of native vegetation within the PGL Surveyed Area. The potential fauna habitat types within the Indicative Impact Area will remain well connected and part of a larger contiguous landscape of similar habitats within the local area and surrounding region. The Indicative Impact Area is unlikely to support fauna habitat that is in better condition than the surrounding available habitat. Furthermore, the Indicative Impact Area is not likely to comprise significant locally or regionally unique habitat for indigenous fauna dependent on the habitats present in within the PGL Surveyed Area. It is likely these habitats are well represented in the local and regional area given the extent of native vegetation adjacent to the PGL Surveyed Area footprint, and in nearby conservation areas.

Given the temporary nature of most clearing proposed and distribution of the DE across 250km, it is unlikely there will be a significant impact the availability of habitat for conservation significant fauna

Methodology

- Biological Survey (Biota 2024 & 2025)
- DCCEEW Protected Matters Search Tool Report (DCCEEW 2024b)
- Dandjoo Database Search (DBCA 2024)
- Government GIS Shapefiles:
 - DBCA Threatened and Priority fauna database search (Accessed November 2024)
- Species specific conservation listing advice and recovery plans

(c) Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, threatened flora.

Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, threatened flora.

Proposed clearing is unlikely to be at variance to this Principle.

Assessment

No EPBC Act and/or BC Act listed Threatened flora species have been identified within the PGL Surveyed Area (Biota 2024 & 2025). Further, no threatened flora is considered likely to occur within the PGL Surveyed Area based on the likelihood of occurrence assessment (Biota 2024 & 2025).

Impacts to vegetation as a result of the Investigation Activities is not likely to include or be necessary for the continued existence of rare/threatened flora.

The PGL Surveyed Area, and subsequent Indicative Impact Areas are part of contiguous vegetation that is not considered more diverse or in better condition than the surrounding vegetation.

It is considered that the proposed clearing is unlikely to be at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- DCCEEW Protected Matters Search Tool Report (DCCEEW 2025)
- Government GIS shapefiles:
 - DBCA Threatened flora database search (Accessed March 2025)
 - WA Herbarium Threatened and Priority flora (Accessed March 2024)

(d) Native vegetation should not be cleared if it comprises the whole or a part of or is necessary for the maintenance of a threatened ecological community.

Native vegetation should not be cleared if it comprises the whole or a part of or is necessary for the maintenance of a threatened ecological community.

Proposed clearing is not at variance to this Principle.

Assessment

The EPBC Act PMST and DBCA TEC databases did not identify any TECs within 20 km of the PGL Surveyed Area.

The Biota (2024 & 2025) field survey did not identify any vegetation within the PGL Surveyed Area that is representative of TECs.

The proposed clearing is not at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- DCCEEW Protected Matters Search Tool Report (DCCEEW 2025b)
- Government GIS shapefiles:
 - DBCA Threatened Ecological Community database search

(e) Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.

Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.

Proposed clearing is not at variance to this Principle.

Assessment

Vegetation Association

Broad-scale (1:250,000) pre-European vegetation mapping identified seven Vegetation Associations present in the PGL Surveyed Area (DPIRD 2019). These are shown below.

Pre-European Vegetation Association	Area (ha) in PGL Surveyed Area	Indicative Impact Area (ha)	Prop. of Veg Association clearing (%)
93: Hummock grasslands, shrub steppe; kanji over soft spinifex	15,536.0	3.7	<0.1
101: Hummock grasslands, shrub steppe; <i>Acacia pachycarpa</i> over soft spinifex	5,359.5	30.6	0.6
117: Hummock grasslands, grass steppe; soft spinifex	18,482.0	51.4	0.3
171: Hummock grasslands, low tree steppe; snappy gum over soft spinifex & <i>Triodia brizoides</i>	544.3	0	0
589: Mosaic: Short bunch grassland – savanna / grass plain (Pilbara) / Hummock grasslands, grass steppe; soft spinifex	7,722.3	<0.1	<0.1
619: Medium woodland; river gum (<i>Eucalyptus camaldulensis</i>)	723.2	0	0
647: Hummock grasslands, dwarf-shrub steppe; <i>Acacia translucens</i> over soft spinifex	1,685.9	0.5	<0.1

All Vegetation Associations are well represented across State, IBRA Region, IBRA Subregion, and Local Government Area (LGA) extents. Most pre-European vegetation associations have 99% or more of their pre-European extent remaining (Table 6). Veg Assoc No. 117, described as “hummock grasslands, grass steppe; soft spinifex”, has historically been subject to the highest proportional amount of clearing, seen at a subregional scale (Pindanland) and LGA scale (Shire of Broome). Retention of pre-European vegetation at this scale is approximately 76% and 63%, respectively. However, a relatively high proportion of this remaining vegetation now exists in DBCA reserves where it is protected from future clearing.

The National Objectives and Targets for Biodiversity Conservation Australia have been set to prevent clearance of ecological communities with less than 30% of their pre-European extent, below which species loss appears to accelerate exponentially (Commonwealth of Australia 2001). The current extent of all Vegetation Associations is higher than 30% for pre-European extent at all scales (Statewide, IBRA Bioregion, IBRA Subregion, LGA). The proposed clearing of 51.4 ha of Veg Assoc. No. 117 will reduce the total remain of the pre-European extent by <0.01%.

Approximately 50,203.3 ha of native vegetation is present within the PGL Surveyed Area. Potential clearing of 93.3 ha amounts to 0.2% of the 50,203.3 ha of native vegetation within the PGL Surveyed Area. After native vegetation clearing has been undertaken, sufficient pre-European vegetation remains in the region. The proposed clearing will not impact the maintenance of ecological values in the landscape. The proposed clearing is not at variance to this principle.

Vegetation proposed to be cleared is not significant as a remnant of native vegetation.

Methodology

- Biological Survey (Biota 2024 & 2025)
- Government GIS shapefiles:
 - Pre-European vegetation (Accessed September 2024)

(f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.

Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.

Proposed clearing is unlikely at variance to this principle

Assessment

The DE is not located within and does not intersect an internationally Important Wetland (RAMSAR). The nearest RAMSAR wetland is the Eighty Mile Beach System, located approx. 19 km north-east of the PGL Surveyed Area at its closest point. The DE will not have direct impacts to this RAMSAR wetland, nor will it have indirect impacts based on current understanding.

The PGL Surveyed Area does intersect a Nationally Important Wetland, the De Grey River. The De Grey River is also identified as an ESA. However, the location of the Investigation Activities has considered and avoided the De Grey River (Figure 9.2).

Biota (2025) mapped two vegetation types as potentially containing GDV, D1 and D2. Based on this mapping, there are seven occurrences of GDVs within the PGL Surveyed Area, totalling 919.6 ha, or 1.8%. The D1 vegetation type is associated with the De Grey River, while D2 is associated with other drainage lines found between the De Grey River and Boodarie. The riparian vegetation types D1 and D2 are considered as having a high potential to be GDEs or GDV due to the phreatophytic species *Eucalyptus camaldulensis*, *Eucalyptus victrix* and *Melaleuca argentea* being dominant taxa. The DE intersects 0.3 ha of D2 with 1.0 ha associated with DS_01b and the remaining 0.3 ha associated with a section of the proposed access track to TP50.

In line with the PGL Preliminary Design Principles, the proposed investigation locations were selected to avoid minor intermittent surface water drainage lines where riparian vegetation is present, with a buffer of 50m applied to these riparian areas. However, the DE intersects the borders of minor/broad ephemeral drainage habitats. These drainage habitats are likely to be associated with infrequent surface water caused by sporadic weather events like seasonal cyclones.

As the Important Wetlands are not intersected by the proposed DE it is considered unlikely that significant impacts will occur as a result of the Investigation Activities.

Micro-siting of the Indicative Impact Area within Investigation Pad 3 and clearing for access track to TP50 will aim to minimise potential clearing of vegetation type D2. The DE of 0.6 ha of this vegetation type represents <0.1% of the available vegetation type within the PGL Surveyed Area.

The proposed clearing is unlikely to be at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- Government GIS shapefiles:
 - Geomorphic Wetlands (Accessed September 2024)
 - Ramsar Wetlands (Accessed September 2024)
 - Important Wetlands (Accessed September 2024)
 - Watercourses (Accessed September 2024)
 - RiWI Act Rivers (Accessed September 2024)

(g) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

Proposed clearing is unlikely to be at variance to this Principle.

Assessment

According to DPIRD (2022a) the PGL Surveyed Area is located across four unique soil landscape zones; The Nita Sandplain (Zone 117), De Grey-Roebourne Lowlands (Zone 281), Great Sandy Desert (Zone 112) and the Nullagine Hills (Zone 280).

The table below shows the effects to the three soil landscape zones impacted by clearing within the Indicative Impact Area.

Zone	Description	Indicative Impact Area (ha)
Zone 117 (Nita Sandplain Zone)	Sandplains and dunes on cretaceous canning basin sedimentary rocks with red deep sands and some red sandy earths.	80.5
Zone 281 (De Grey-Roebourne Lowlands Zone)	Alluvial plains and sandplains on alluvial and marine deposits over the northern Pilbara craton with red deep sandy duplexes, red loamy earths, red/brown non-cracking clays, cracking clays, red sandy earths and red deep loamy duplexes.	12.7
Zone 280 (Nullagine Hills Zone)	Hills and ranges (with some stony plains) on volcanic and sedimentary rocks of the Pilbara craton (including the Hamersley Basin) with stony soils and red shallow loams and sands.	0.1

Land zones have been further classified into soil-landscape systems (DPIRD 2022b). The PGL Surveyed Area is located across 16 soil-landscape systems. The table below shows the impacts to the nine soil landscape systems impacted by clearing within the Indicative Impact Area.

System	Description	Indicative Impact Area (ha)
Boolgeeda System (281Bg)	Stony lower slopes and plains below hill systems supporting hard and soft spinifex grasslands or mulga shrublands.	4.2
Callawa System (117CI)	Highly dissected low hills, mesas and gravelly plains on sandstone and conglomerate supporting soft and hard spinifex grasslands.	15.6
Capricorn System (280Cp & 281Cp)	Rugged sandstone hills, ridges, stony foot slopes and interfluvies supporting low acacia shrublands or hard spinifex grasslands with scattered shrubs.	0.1
Macroy System (281Mc)	Stony plains and occasional tor fields based on granite supporting hard and soft spinifex shrubby grasslands.	2.6
Nita System (117Nt & 281Nt)	Sandplains supporting shrubby spinifex grasslands with occasional trees.	67.7
River System (281Ri)	Narrow, seasonally active flood plains and major river channels supporting moderately close, tall shrublands or woodlands of acacias and fringing communities of eucalypts sometimes with tussock grasses or spinifex	0.5
Uaroo System (281Ua)	Broad sandy plains, pebbly plains and drainage tracts supporting hard and soft spinifex hummock grasslands with scattered acacia shrubs.	2.6

An inventory and condition survey of the Pilbara region, undertaken by van Vreeswyk et al. (2004), for the Western Australian Department of Agriculture (now DPIRD), profiled each soil-landscape system, the current condition and the management measures required to maintain the system. Most soil-landscape systems described above are known to be resistant to and have experience nil to minimal soil erosion while being largely resistant to degradation. The River system was identified as being potentially susceptible to soil erosion and degradation when vegetation is removed.

Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

The DE intersects only a small portion of the river soil-landscape system (0.5 ha). By extension, clearing of vegetation within this system will be minimal and not pose significant impacts to the entirety of the system.

The SLIP/ASRIS database indicates that the area surrounding the De Grey River and its tributaries is classified as Moderate to low risk of Acid Sulfate Soils (ASS). The proposed vegetation clearing is not expected to disturb ASS, however, ASS management measures will be incorporated into an EMP to appropriately manage the associated Investigation Activities and will ensure that any potential disturbance of ASS will be appropriately managed and adverse ASS related impacts are avoided.

Given that risk to soil systems is likely to be minimal and that risk of ASS is likely low, it is unlikely that clearing is at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- CSIRO Atlas of Acid Sulfate Soils (Accessed November 2025)

(h) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.

Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.

Proposed clearing is not at variance to this Principle.

Assessment

No conservation areas are located within the PGL Surveyed Area and therefore will not be impacted by the proposed clearing within the DE.

The proposed clearing will not impact the environmental values of conservation areas and is therefore not at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- DCCEEW Protected Matters Search Tool Report (DCCEEW 2025b)
- Government GIS Shapefiles:
 - DBCA Legislated Lands and Waters & Lands of Interest (Accessed May 2025)

(i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.

Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.

Proposed clearing is not likely to be at variance to this Principle.

Assessment

The climate of the Pilbara is described as arid-tropical with two distinct seasons. Rainfall in the Pilbara is highly variable and may occur during both seasons. Average long-term annual rainfall for the area is between 314.1 mm and 375.0 mm (Port Hedland Airport, station number 4032 and Madora, station number 4019) which can occur in heavy localised falls. Based on very high annual evaporation rates, any surface runoff resulting from rainfall events is likely to be relatively short lived. In addition, the PGL Surveyed Area is largely surrounded by native vegetation, and it is likely that a large proportion of runoff will be absorbed by this natural environment.

The PGL Surveyed Area lies within the Pilbara Groundwater Area and Canning-Kimberly Groundwater Area which are proclaimed under the RiWI Act 1914 (DWER 2018). Within this groundwater area, the PGL Surveyed Area intersects three groundwater subareas: Ashburton, West Canning - Yarrie and East Pilbara. The PGL Surveyed Area intersects one Public Drinking Water Source Area, the De Grey River Water Reserve (DWER 2024b). Within this water reserve, the De Grey River wellfield is used to supply water for Port Hedland (Water and Rivers Commission 1999). The De Grey River wellfield is operated by the Water Corporation and draws groundwater from the alluvial deposits of the De Grey River.

It is considered unlikely any clearing will significantly disturb or interrupt natural drainage and surface run-off patterns. However, during heavy localised rainfall events, erosion may occur in cleared areas resulting in localised, short-term soil erosion and/or sedimentation. It is unlikely clearing will have an impact on groundwater levels or quality. The proposed Investigation Activities associated with the clearing will not alter the current hydrological regime. Due to small proportion of clearing proposed within each area, the surrounding native vegetation and natural ground will minimise sheet flow over the duration of each Investigation Activity

Standard construction measures regarding erosion and sediment control (include topsoil management) will be included as part of an EMP and implemented during construction works to manage the risks of erosion within cleared areas.

The proposed clearing is unlikely to cause deterioration in surface water and ground water quality

Methodology

- Biological Survey (Biota 2024 & 2025)
- Government GIS Shapefiles:
 - RiWI Act, Surface Water Areas and Irrigation Districts (Accessed May 2025)
 - RiWI Act, Groundwater Areas (Accessed May 2025)
 - Public Drinking Water Source Areas (Accessed May 2025)

(j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence or intensity of flooding.

Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence or intensity of flooding.

Proposed clearing is not at variance to this Principle.

Assessment

The Pilbara region receives an average rainfall between 314.1 – 375.0 mm per year (Port Hedland Airport weather station (site number: 4032) & Madora weather station (site number: 4019)) (BoM 2024). Seasonal drainage lines are common in the region and temporary localised flooding may occur briefly following heavy rainfall events.

Removal of vegetation in areas with clay soils can exacerbate the incidence or intensity of flooding or localised waterlogging. However, the total DE and Indicative Impact Areas proposed is relatively small considering the amount of remaining native vegetation in the surrounding area. The surrounding areas are predominantly undeveloped. Therefore, it is unlikely clearing will increase the incidence of flooding and compaction of soils. In addition, the removal of vegetation for testing and replacement of this vegetation is not likely to result in compaction and flooding.

The proposed clearing is unlikely exacerbate the incidence or intensity of flooding and therefore not at variance to this principle.

Methodology

- Biological Survey (Biota 2024 & 2025)
- BoM Website (Accessed March 2024)
- PGL Preliminary Design Principles (HP 2024)

6. Environmental Avoidance and Management

PGL has consistently applied the disturbance hierarchy of avoid, minimise, reduce, and rehabilitate to mitigate potential environmental impacts associated with the Investigation Activities. This approach was integrated into the design and planning phase (Section 6.1) and will continue through field activities (Section 6.2), guided by Environmental Management Plans (EMPs).

6.1 Impact avoidance and minimisation through design and planning

6.1.1 Flora, Vegetation and Fauna

Flora and vegetation loss was avoided through design of the investigation locations and access tracks within the PGL Surveyed Area by avoiding vegetation clearing where possible. Avoidance and minimisation strategies included:

- When considering placement of the Investigation Pads, a site-by-site assessment with engineers, geotechnicians, hydrogeologists, heritage, land-tenure, and environmental disciplines occurred, to identify constraints and the most practicable location of the Investigation Pads and access tracks.
- Potential impacts to flora and vegetation have been minimised by selecting existing access tracks where possible.
- Where testing requirements for each investigation type is feasible, co-location of proposed Investigation Activities has occurred to minimise clearing required.
- The location of the Investigation Pads and access tracks have avoided known heritage locations

Impacts to native fauna and fauna habitat were avoided during the design of the DE within the PGL Surveyed Area by avoiding areas where threatened species were recorded during the Biota (2024 & 2025) surveys. This included the adoption of the PGL Preliminary Design Principles which outlines buffer areas required to avoid and minimise disturbances to each recorded threatened species (Figure 7).

6.2 Impact avoidance and minimisation measures applied on site

The following avoidance and minimisation management measures will be adopted onsite during the Investigation Activities. These management measures aim to reduce the loss of vegetation, flora and fauna species habitat affected by the direct and indirect potential impacts outlined in Section 4.

Aspect	Management measure
Extent of clearing	<ul style="list-style-type: none"> – Clearing of native vegetation will be restricted to the Indicative Impact Areas selected within the DE – The investigation teams will utilise existing tracks and roads where possible. – New tracks will be established using the shortest appropriate distance from an existing track to where the pad is required within the DE. – A maximum width of 10 m of clearing has been allocated to access tracks. This is an upper limit value and is unlikely to be needed for most of the proposed access tracks required, with a typical access track estimated to be 3m to 4m wide if active vegetation clearing is required. – For new access tracks to pads containing geotechnical test pits and/or boreholes, where practicable, low vegetation will be driven over rather than cleared by blade to minimise vegetation disturbance / clearing – The DE is to be demarcated (e.g. flagging tape or similar) prior to clearing activities. No clearing of native vegetation or ground disturbance is to occur outside of this area.

PROTECTED

Aspect	Management measure
	<ul style="list-style-type: none"> – Upon the conclusion of each Investigation Activity, the same access track will be used to return to the main road. – Preference will be given to previously disturbed or already cleared vegetation when selecting access tracks where terrain allows.
Flora and Vegetation	<ul style="list-style-type: none"> – Areas that are sparsely vegetated and/or previously cleared will be used preferentially for the location of pads and access tracks required for the Investigation Activities. – Minimise disturbance by avoiding large trees and shrubs and where feasible, leaving rootstock in the ground to assist with stabilisation and natural regeneration. – The investigation team will be provided material that identifies known Environmentally sensitive areas within the DE to be avoided. – Boreholes are to be capped, and the cleared drill pad area will have stripped topsoil and vegetation respread to facilitate natural regrowth. – Topsoil (i.e. typically the top 10 mm of soil) and cleared vegetation will be separately stockpiled and re-spread over temporarily cleared areas – Test pits will be backfilled with the excavated material on completion in accordance with the Western Australian Dept. of Energy, Mines, Industry Regulations and Safety's Exploration and Prospecting Rehabilitation Guidance, including mounding over the backfilled hole to facilitate water shedding away from the backfilled areas to the surface and then cover with recovered topsoil and vegetation. – Drill pads and test pits will be rehabilitated back to original contour, so that the surface of the rehabilitated area is consistent with the shape of the surrounding uncleared land
Fauna and Fauna Habitat	<ul style="list-style-type: none"> – Clearing of native vegetation will be undertaken in a slow, progressive manner in one direction to allow fauna to move away from the clearing area. – Construction personnel will be provided with fauna identification materials for relevant conservation significant species for on-site identification and active avoidance. – Construction personnel are not permitted to touch, feed or otherwise directly interact with fauna. – Injury or mortality of fauna will be recorded as an environmental incident. – Any excavations left open overnight will include fauna egress and be inspected at the start and end of each day for fauna – Vehicle and machinery speeds within the Indicative Impact Area will be restricted to reduce the likelihood of fauna strike. – All waste containers will have lids to prevent fauna from eating food scraps or becoming trapped in waste containers. – Any tree branches and rocks originally removed as part of pad establishment will be used in rehabilitation to promote ground stability and to provide potential fauna habitat.
Heritage	<ul style="list-style-type: none"> – No construction activities are permitted to occur within areas designated as Registered Aboriginal Sites without the appropriate permit and/or stakeholder consultation – Aboriginal heritage surveys of the DE will be undertaken prior to the Investigation Activities – Heritage sites within the DE are to be appropriately demarcated. – A New Finds process will be included in the EMP, including the discovery of potential human bones.
Weeds	<ul style="list-style-type: none"> – Standard biosecurity measures will be developed in the EMP and be implemented to mitigate the risk of weeds entering the site or spreading. – Prior to vehicles, plant and equipment coming to the Site, and moving between pads, they shall be free of plant material and soil clumps.
Erosion	<ul style="list-style-type: none"> – Standard management measures regarding erosion and sediment control (including topsoil management) will be implemented during the Investigation Activities.
Dust	<ul style="list-style-type: none"> – Standard dust control and mitigation measures will be implemented during clearing. This may include the use of a water truck(s). – Ground disturbance and/or clearing of vegetation will be restricted during high winds if dust cannot be adequately controlled. – Review of weather forecasts will be undertaken prior to native vegetation clearing to identify periods of extreme weather conditions likely to result in increased dust emissions so that additional mitigation measures can be implemented.
Fire	<ul style="list-style-type: none"> – Basic fire-fighting equipment such as fire extinguishers will be provided in all motor vehicles and mobile plant and during any hot works.

Aspect	Management measure
	<ul style="list-style-type: none"> – Fires will not be permitted on sites.
Noise	<ul style="list-style-type: none"> – Standard construction noise management measures will be implemented. – Complaints regarding noise will be recorded and investigated.
Waste	<ul style="list-style-type: none"> – Management strategies will be implemented to ensure the generation of waste during the Investigation Activities is minimised. – Rubbish will be disposed of in appropriate containers with lids and all waste will be removed from the site and disposed of lawfully.
Acid Sulphate Soils	<ul style="list-style-type: none"> – ASS management measures will be incorporated into an EMP to appropriately manage the associated Investigation Activities and will ensure that any potential disturbance of ASS will be appropriately managed.

7. Other approvals

Following review of approval requirements under the EP Act, the following approvals are not considered necessary for the purpose of these Investigation Activities:

- *Planning and Development Act 2005*
- Works Approval or Licence under Part V of the EP Act
- Referral to DCCEEW for assessment under the EPBC Act
- State or municipal heritage approvals
- Aboriginal Sites of significance under the *Aboriginal Heritage Act 1972*

7.1 Referral to EPA

The Investigation Activities described in this application are not considered to be a significant impact under s38 of the *EP Act* and therefore will not be referred to the EPA.

7.2 Approvals under the RiWI Act

PGL has liaised with DWER on the potential approvals required to drill groundwater bores and abstract groundwater, including for monitoring purposes. Appropriate approvals will be sought in accordance with the RiWI Act.

7.3 Aboriginal Heritage

The *Aboriginal Heritage Act 1972* (AH Act) states that compliance means that the proponent is obligated to identify and protect all Aboriginal heritage places as defined by Section 5 of the AH Act before development occurring. If protection is not an option, then consent to affect such places can be sought from the Department of Planning, Land and Heritage (DPLH) and the Minister of Aboriginal Affairs under Sections 16 and 18 of the AH Act, or Regulation 10 of the *Aboriginal Heritage Regulations 1974*.

PGL is responsible for the necessary cultural heritage surveys and any subsequent approvals for all Investigation Activities. PGL has engaged with the relevant Traditional Owner groups and knowledge holders associated with the Alignment Area. This engagement has also included the attendance of Cultural Safety Monitors during biological surveys.

The Investigation Activities shall avoid all identified exclusion areas of heritage significance included in a proposed Heritage Management Plan. All areas of heritage significance identified within the vicinity of each Investigation Activity shall be appropriately protected in consultation with the relevant Tradition Owner representatives and in accordance with the proposed Heritage Management Plan requirements so that the Contractor's personnel are aware of the area to be avoided. During the Investigation Activities the Contractor shall inform its personnel of the heritage areas to be avoided which have been identified, and the process to protect those additional areas that have been identified during the undertaking of the Investigation Activities.

8. Offsets

Environmental offsets are conservation actions that provide environmental benefits intended to counterbalance significant residual environmental impacts associated with a proposal (GoWA 2014).

PGL have considered requirements to counterbalance the residual impacts through environmental offsets for Investigation Activities. Consideration has been given to requirements of the Western Australian Government's Environmental Offset Policy (GoWA 2011) and the Western Australian Offsets Guidelines (GoWA 2014).

PGL operates on a hierarchy of avoid, minimise, reduce, rehabilitate and offset for identified environmental impacts. This hierarchy is achieved primarily through changes in scope and design, development and implementation of the environmental management plans or strategies and finally, if required, development of an offset proposal.

Application of the management hierarchy has been applied throughout the selection of investigation sites and described in this document. The assessment against the ten clearing principles concluded the proposed clearing is not or unlikely to be, at variance with any of the ten clearing principles. Therefore, offsets are not proposed to compensate for the residual impacts associated with the proposed clearing.

9. References

- Atlas of Living Australia. 2025a. *Tribulopsis angustifolia* R.Br.. <https://bie.ala.org.au/species/https://id.biodiversity.org.au/node/apni/2913483>
- Atlas of Living Australia. 2025b. *Croton aridus* P.I.Forst. <https://bie.ala.org.au/species/https://id.biodiversity.org.au/node/apni/2916090#overview>
- Biota Environmental Sciences. 2024. *Pilbara Green Link Project Link 1 Biological Survey*. Prepared for GHD on behalf of Horizon Power
- Biota Environmental Sciences. 2025. *Pilbara Green Link Project Gap Areas Biological Assessment*. Prepared for GHD on behalf of Horizon Power.
- Bureau of Meteorology. 2024a. *Monthly climate data*. Australian Government. <http://www.bom.gov.au/climate/data/>.
- Bureau of Meteorology. 2024b. *Groundwater Dependent Ecosystem Atlas*. Australian Government. <http://www.bom.gov.au/water/groundwater/gde/map.shtml>.
- Commonwealth of Australia. 2001. *National Objectives and Targets for Biodiversity Conservation 2001–2005*.
- Department of Biodiversity, Conservation and Attraction. <https://library.dbca.wa.gov.au/static/FullTextFiles/020395.pdf>
- Department of Agriculture and Water Resources. 2017. *Australian Weeds Strategy 2017-2027*. Commonwealth of Australia. <https://www.agriculture.gov.au/sites/default/files/sitecollectiondocuments/pests-diseases-weeds/consultation/aws-final.pdf>
- Department of Biodiversity, Conservation and Attractions. 2017. *Ramsar sites (DBCA-010)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/ramsar-sites>
- Department of Biodiversity, Conservation and Attractions. 2018. *Directory of Important Wetlands in Australia – Western Australia (DBCA-045)*. Australian Government. <https://catalogue.data.wa.gov.au/dataset/directory-of-important-wetlands-in-western-australia>
- Department of Biodiversity, Conservation and Attractions. 2022. *Threatened Ecological Communities (DBCA-038)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/threatened-ecological-communities>
- Department of Climate Change, Energy, Environment and Water. 2021. *Interim Biogeographic Regionalisation for Australia (IBRA 7) Codes*. Australian Government. <https://www.dcceew.gov.au/environment/land/nrs/science/ibra/ibra7-codes>
- Department of Climate Change, Energy, the Environment and Water. 2024. *EPBC Act List of Threatened Fauna*. Australian Government. <https://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl?wanted=fauna>
- Department of Conservation and Land Management. 2002. *A biodiversity audit of Western Australia's 53 biogeographical subregions in 2002*. Government of Western Australia. <https://library.dbca.wa.gov.au/static/FullTextFiles/021927.pdf>
- Department of Energy, Mines, Industry, Regulation and Safety. 2018. *1:500 000 State interpreted bedrock geology (DMIRS-016)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/1-500-000-state-interpreted-bedrock-geology-dmirs-016>
- Department of Primary Industries and Regional Development. 2019. *Pre-European Vegetation (DPIRD-006)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/pre-european-dpird-006>
- Department of Water and Environmental Regulation. 2014. *Acid Sulfate Soil Risk Map, Pilbara Coastline (DWER-053)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/acid-sulfate-soil-risk-map-pilbara-coastline-dwer-053>

- Department of Water and Environmental Regulation. 2018a. *RIWI Act, Groundwater Areas (DWER-034)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/riwi-act-groundwater-areas>
- Department of Water and Environmental Regulation. 2018b. *De Grey River Water Reserve*. Government of Western Australia. [De-Grey-River-Water-Reserve-drinking-water-source-protection-review-WRP-140.pdf](https://catalogue.data.wa.gov.au/dataset/de-grey-river-water-reserve-drinking-water-source-protection-review-WRP-140.pdf)
- Department of Water and Environmental Regulation. 2018c. *Hydrography, Linear (Hierarchy) (DWER-031)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/hydrography-linear-hierarchy>
- Department of Water and Environmental Regulation. 2018d. *Hydrographic Catchments – Basins (DWER_027)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/hydrographic-catchments-basins>
- Department of Water and Environmental Regulation. 2021. *Clearing Regulations - Environmentally Sensitive Areas (DWER-046)*, Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/clearing-regulations-environmentally-sensitive-areas-dwer-046>
- Department of Water and Environmental Regulation. 2024a. *Contaminated Sites Database (DWER-059)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/contaminated-reported-sites-dwer-059>
- Department of Water and Environmental Regulation. 2024b. *Public Drinking Water Source Areas (DWER-033)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/public-drinking-water-source-areas>
- Environmental Protection Act 1986 (WA)* pt 5 div 2 s 51(E). https://www.legislation.wa.gov.au/legislation/statutes.nsf/main_mrtitle_304_homepage.html
- Electricity Corporations Act 2005 (WA)*. https://www.legislation.wa.gov.au/legislation/statutes.nsf/main_mrtitle_286_homepage.html
- Environmental Protection Authority. 2016. *Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment*. Government of Western Australia. https://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA%20Technical%20Guidance%20-%20Flora%20and%20Vegetation%20survey_Dec13.pdf
- Environmental Protection Authority. 2020. *Technical Guidance: Terrestrial vertebrate fauna surveys for environmental impact assessment*. Government of Western Australia. https://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/2020.09.17%20-%20EPA%20Technical%20Guidance%20-%20Vertebrate%20Fauna%20Surveys%20-%20Final.pdf
- Environmental Protection Authority. 2021. *Environmental Impact Assessment (Part IV Divisions 1 and 2) Procedures Manual*. Government of Western Australia. https://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EIA%20%28Part%20IV%20Divisions%201%20and%202%29%20Procedures%20Manual_1.pdf
- Landgate. 2024. *Reserves (LGATE-277)*. Government of Western Australia. <https://catalogue.data.wa.gov.au/dataset/reserves>
- Reeves, J. M., De Deckker, P. & Halse, S. A. (2007), *Groundwater Ostracods from the arid Pilbara region of northwestern Australia: distribution and water chemistry*. *Hydrobiologia*. 585:99- 118
- van Vreeswyk, A.M.E., Payne, A.L., Hennig, P., and Leighton, K.A. 2004, *An Inventory and Condition Survey of the Pilbara Region*. Western Australia. Department of Agriculture, Western Australia. https://library.dpird.wa.gov.au/cgi/viewcontent.cgi?article=1006&context=tech_bull

Appendix A

Investigation sites and activities

Table 21 Definition of acronyms

Acronym	Term
GT	Geotechnical
GW	Groundwater
TP	Test Pit
BH	Geotechnical Borehole
DS	Groundwater Borehole

Table 22 Investigation Pads and activities

Investigation Pad	Name	Investigation Activity	Discipline	Total Clearing proposed (ha)
1	DS_01a	Borehole	Groundwater	0.49
2	MRD_01	Borehole	Groundwater	0.49
3	DS_01b	Borehole	Groundwater	0.49
4	TP13	Test pit	Geotechnical	1.58
	BH07	Borehole	Geotechnical	
	Access Track for TP13	Access track		
5	DS_03a	Borehole	Groundwater	0.49
6	TP18	Test pit	Geotechnical	0.02
7	BH12	Borehole	Geotechnical	3.71
	TP23	Test pit	Geotechnical	
	Access Track for TP23	Access track		
8	DS_05	Borehole	Groundwater	0.49
9	Camp_01	Borehole	Groundwater	0.49
10	Camp_02	Borehole	Groundwater	0.91
	Access Track between Camp_01 and Camp_02	Access track		
11	Camp TP1	Test pit	Geotechnical	0.02
12	Camp TP2	Test pit	Geotechnical	0.02
13	Camp TP3	Test pit	Geotechnical	0.02
14	Camp TP4	Test pit	Geotechnical	0.02
15	Camp TP5	Test pit	Geotechnical	0.85
	Access track for camp	Access track		
16	CB_01	Borehole	Groundwater	0.49
17	TP24	Test pit	Geotechnical	0.02
18	TP29a	Test pit	Geotechnical	0.10
	BH15a	Borehole	Geotechnical	
19	TP30a	Test pit	Geotechnical	0.02
20	TP31	Test pit	Geotechnical	7.86
	BH16	Borehole	Geotechnical	
	DS_07 (gt)	Borehole	Groundwater	
	Access track for TP31	Access track		
21	DS_07a	Borehole	Groundwater	0.49

PROTECTED

Investigation Pad	Name	Investigation Activity	Discipline	Total Clearing proposed (ha)
22	TP32a	Test pit	Geotechnical	0.02
23	TP33a	Test pit	Geotechnical	0.10
	BH17a	Borehole	Geotechnical	
24	BH18	Borehole	Geotechnical	0.10
	TP35	Test pit	Geotechnical	
25	TP36	Test pit	Geotechnical	1.53
	Access track for TP36	Access track		
26	TP39	Test pit	Geotechnical	2.28
	BH20	Borehole	Geotechnical	
	Access track TP39	Access track		
27	TP40	Test pit	Geotechnical	2.19
	Access track to TP40	Access track		
28	DS_08 (gt)	Borehole	Groundwater	2.78
	TP41	Test pit	Geotechnical	
	BH21	Borehole	Geotechnical	
	Access track to TP41	Access track		
29	TP43	Test pit	Geotechnical	1.71
	BH22	Borehole	Geotechnical	
	Access track to TP43	Access track		
30	TP44	Test pit	Geotechnical	3.45
	DS_10	Borehole	Groundwater	
	Access track to TP44	Access track		
31	Access track between TP44 and TP45	Access track		10.70
	TP45	Test pit	Geotechnical	
	BH23	Borehole	Geotechnical	
	Access track to TP45	Access track		
32	TP46	Test pit	Geotechnical	2.74
	DS_11	Borehole	Groundwater	
	Access track to TP46	Access track		
33	TP47	Test pit	Geotechnical	3.63
	BH24	Borehole	Geotechnical	
	Access track between TP46 and TP47	Access track		
34	TP48	Test pit	Geotechnical	11.36
	Access track for TP48 (pt1)	Access track		
	Access track for TP48 (pt 2)	Access track		
	Access track between TP48 and TP49	Access track		
35	TP49	Test pit	Geotechnical	0.41
	BH25	Borehole	Geotechnical	
	Access track for TP49	Access track		
36	DS_12	Borehole	Groundwater	0.49

PROTECTED

Investigation Pad	Name	Investigation Activity	Discipline	Total Clearing proposed (ha)
37	Access track between TP49 and TP50	Access track		25.60
	TP50	Test pit	Geotechnical	
	DS_12a (gt)	Borehole	Groundwater	
	Access track for TP50	Access track		
38	TP51	Test pit	Geotechnical	5.07
	BH26	Borehole	Geotechnical	
	DS_13	Borehole	Groundwater	
	Access track for TP51	Access track		

Appendix B

Land parcel clearing

Table 23 *Impacts of proposed clearing on Land Parcels*

Cadastre			Proposed Indicative Impact Area ha)	Investigation Pad
Pi Parcel	Land type	Usage Description		
P238018 101	Crown	Land Act (Type 2)	7.25	26, 27, 28
P238025 100	Crown	Land Act (Type 2)	2.74	20, 21, 22, 23, 24, 25
P238657 252	Crown	Land Act (Type 2)	2.07	2, 4
P241586 42	Crown	Land Act (Type 2)	7.37	20
LPL N050027	Lease	Lease (Type 3 L)	0.25	6, 7, 18, 19
LPL N050324	Lease	Lease (Type 3 L)	6.95	7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17
P037092 414	Road	Road Isolation (Type 3 P)	0.49	3
P037093 417	Road	Road Isolation (Type 3 P)	0.49	5
P042630 301	Freehold	Transfer of Land Act (Type 1)	0.49	1
V CROWN LAND	Other	Unallocated Crown Land (Type 3 V)	65.15	29, 30, 31, 32, 33, 34, 35, 36, 37, 38

Appendix C

Dandjoo Database Search Results

Row Labels	Count of TAXON
CR	153
Animalia	153
BIRD	153
<i>Calidris ferruginea</i>	46
<i>Calidris tenuirostris</i>	56
<i>Limosa lapponica</i> subsp. <i>menzbier</i>	12
<i>Numenius madagascariensis</i>	39
EN	997
Animalia	997
BIRD	67
<i>Calidris canutus</i>	30
<i>Charadrius mongolus</i>	37
MAMMAL	930
<i>Dasyurus hallucatus</i>	930
MI	1257
Animalia	1257
BIRD	1257
<i>Actitis hypoleucos</i>	69
<i>Apus pacificus</i>	4
<i>Arenaria interpres</i>	82
<i>Calidris acuminata</i>	40
<i>Calidris alba</i>	56
<i>Calidris melanotos</i>	2
<i>Calidris ruficollis</i>	96
<i>Calidris subminuta</i>	6
<i>Charadrius veredus</i>	30
<i>Chlidonias leucopterus</i>	27
<i>Fregata ariel</i>	10
<i>Gallinago megala</i>	2
<i>Gallinago stenura</i>	4
<i>Gelochelidon nilotica</i>	41
<i>Glareola maldivarum</i>	32
<i>Hirundo rustica</i>	18
<i>Hydroprogne caspia</i>	86
<i>Limicola falcinellus</i>	18
<i>Limnodromus semipalmatus</i>	14
<i>Limosa lapponica</i>	75
<i>Limosa limosa</i>	17
<i>Numenius minutus</i>	28
<i>Numenius phaeopus</i>	71
<i>Oceanites oceanicus</i>	2
<i>Onychoprion anaethetus</i>	2
<i>Pandion cristatus</i>	61
<i>Phalaropus lobatus</i>	11
<i>Philomachus pugnax</i>	1
<i>Plegadis falcinellus</i>	12
<i>Pluvialis fulva</i>	40
<i>Pluvialis squatarola</i>	51
<i>Sterna dougallii</i>	2
<i>Sterna hirundo</i>	23
<i>Sternula albifrons</i>	40
<i>Thalasseus bergii</i>	44
<i>Tringa glareola</i>	23
<i>Tringa nebularia</i>	58
<i>Tringa stagnatilis</i>	31
<i>Xenus cinereus</i>	28
MI & P4	76
Animalia	76

BIRD	76
Tringa brevipes	76
OS	7
Animalia	7
BIRD	6
Falco peregrinus	6
MAMMAL	1
Dugong dugon	1
P1	40
Plantae	40
DICOT	38
Corchorus sp. Yarrie (J. Bull & D. F	2
Heliotropium parviantrum	1
Tephrosia rosea var. Port Hedland	35
MONOCOT	2
Fimbristylis sp. H Kimberley Flora (1
Fimbristylis sp. Shay Gap (K.R. Ne	1
P2	9
Animalia	2
REPTILE	2
Lerista separanda	2
Plantae	7
DICOT	7
Gomphrena pusilla	7
P3	131
Animalia	17
INVERT	1
Antipodogomphus hodgkini	1
REPTILE	16
Ctenotus angusticeps	16
Plantae	114
DICOT	98
Abutilon sp. Pritzelianum (S. van L	28
Bonamia oblongifolia	3
Croton aridus	2
Euphorbia clementii	7
Gomphrena leptophylla	1
Gymnanthera cunninghamii	5
Heliotropium muticum	31
Indigofera ammobia	2
Nicotiana umbratica	3
Polymeria distigma	2
Rothia indica subsp. australis	9
Terminalia kumpaja	5
MONOCOT	16
Eragrostis crateriformis	15
Fuirena incrassata	1
P4	272
Animalia	261
MAMMAL	261
Dasycercus blythi	210
Dasycercus cristicauda	3
Lagorchestes conspicillatus subsp.	2
Leggadina lakedownensis	5
Notoryctes caurinus	7
Pseudomys chapmani	32
Rhinonictes aurantia	2
Plantae	11
DICOT	8

Goodenia nuda	7
Ptilotus mollis	1
MONOCOT	3
Bulbostylis burbridgeae	3
VU	2521
Animalia	2521
BIRD	70
Charadrius leschenaultii	66
Falco hypoleucos	4
MAMMAL	175
Lagostrophus fasciatus subsp. fasc	1
Macroderma gigas	65
Macrotis lagotis	106
Rhinonictes aurantia (Pilbara)	3
REPTILE	2276
Chelonia mydas	2
Liasis olivaceus subsp. barroni	4
Natator depressus	2268
Pogona minor subsp. minima	2
(blank)	19814
Animalia	16189
ALGA	6
Lobophora sp.	1
Turbinaria spp	5
AMPHI	235
Cyclorana australis	10
Cyclorana longipes	1
Cyclorana maini	19
Limnodynastes spenceri	2
Litoria caerulea	1
Litoria nasuta	2
Litoria rothii	1
Litoria rubella	24
Neobatrachus aquilonius	21
Neobatrachus sutor	1
Notaden nicholli	76
Platyplectrum spenceri	9
Uperoleia glandulosa	36
Uperoleia micromeles	14
Uperoleia russelli	2
Uperoleia saxatilis	4
Uperoleia talpa	12
BIRD	8970
Acanthagenys rufogularis	4
Accipiter cirrocephalus	17
Accipiter fasciatus	20
Accipiter fasciatus subsp. fasciatus	3
Acrocephalus australis	2
Aegotheles cristatus	19
Amytornis striatus	3
Anas gracilis	104
Anas rhynchotis	1
Anas superciliosa	108
Anhinga melanogaster	2
Anhinga melanogaster subsp. nova	2
Anhinga novaehollandiae	45
Anthus australis	17
Anthus australis subsp. australis	2
Aquila audax	37

<i>Aquila morphnoides</i>	2
<i>Ardea alba</i>	1
<i>Ardea alba</i> subsp. <i>modesta</i>	3
<i>Ardea garzetta</i>	3
<i>Ardea garzetta</i> subsp. <i>nigripes</i>	2
<i>Ardea ibis</i>	1
<i>Ardea intermedia</i>	11
<i>Ardea modesta</i>	69
<i>Ardea novaehollandiae</i>	8
<i>Ardea pacifica</i>	47
<i>Ardeotis australis</i>	70
<i>Arenaria interpres</i> subsp. <i>interpres</i>	8
<i>Artamus cinereus</i>	166
<i>Artamus cinereus</i> subsp. <i>melanops</i>	1
<i>Artamus leucorhynchus</i>	131
<i>Artamus leucorhynchus</i> subsp. <i>leucorhynchus</i>	6
<i>Artamus minor</i>	18
<i>Artamus personatus</i>	24
<i>Artamus superciliosus</i>	2
<i>Aythya australis</i>	42
<i>Barnardius zonarius</i>	40
<i>Burhinus grallarius</i>	39
<i>Butorides striata</i>	30
<i>Butorides striatus</i>	2
<i>Cacatua roseicapilla</i>	24
<i>Cacatua roseicapilla</i> subsp. <i>assimilis</i>	4
<i>Cacatua sanguinea</i>	108
<i>Cacatua sanguinea</i> subsp. <i>westralis</i>	3
<i>Cacomantis pallidus</i>	39
<i>Calidris fuscicollis</i>	1
<i>Calyptorhynchus banksii</i>	3
<i>Centropus phasianinus</i>	21
<i>Certhionyx niger</i>	2
<i>Certhionyx variegatus</i>	6
<i>Charadrius leschenaultii</i> subsp. <i>leschenaultii</i>	3
<i>Charadrius melanops</i>	15
<i>Charadrius ruficapillus</i>	110
<i>Chenonetta jubata</i>	7
<i>Cheramoeca leucosterna</i>	3
<i>Cheramoeca leucosternus</i>	1
<i>Chroicocephalus novaehollandiae</i>	82
<i>Chrysococcyx basalis</i>	4
<i>Chrysococcyx osculans</i>	1
<i>Cincloramphus cruralis</i>	26
<i>Cincloramphus mathewsi</i>	52
<i>Circus approximans</i>	15
<i>Circus assimilis</i>	38
<i>Cladorhynchus leucocephalus</i>	17
<i>Climacteris melanura</i>	21
<i>Colluricincla harmonica</i>	34
<i>Colluricincla harmonica</i> subsp. <i>brui</i>	1
<i>Colluricincla harmonica</i> subsp. <i>ruficeps</i>	1
<i>Columba livia</i>	20
<i>Coracina novaehollandiae</i>	223
<i>Coracina novaehollandiae</i> subsp. <i>saxatilis</i>	1
<i>Corvus bennetti</i>	14
<i>Corvus coronoides</i>	32
<i>Corvus orru</i>	139
<i>Corvus orru</i> subsp. <i>ceciliae</i>	3

<i>Coturnix pectoralis</i>	3
<i>Coturnix ypsilophora</i>	35
<i>Coturnix ypsilophora</i> subsp. <i>austral</i>	2
<i>Cracticus nigrogularis</i>	131
<i>Cracticus tibicen</i>	11
<i>Cracticus torquatus</i>	4
<i>Cuculus pallidus</i>	10
<i>Cygnus atratus</i>	55
<i>Dacelo leachii</i>	77
<i>Daphoenositta chrysoptera</i>	1
<i>Dendrocygna arcuata</i>	2
<i>Dendrocygna eytoni</i>	39
<i>Dicaeum hirundinaceum</i>	1
<i>Dromaius novaehollandiae</i>	9
<i>Egretta garzetta</i>	46
<i>Egretta novaehollandiae</i>	59
<i>Egretta sacra</i>	43
<i>Elanus axillaris</i>	14
<i>Elanus caeruleus</i>	9
<i>Elanus caeruleus</i> subsp. <i>axillaris</i>	7
<i>Elseyornis melanops</i>	101
<i>Emblema pictum</i>	62
<i>Eolophus roseicapillus</i>	71
<i>Eopsaltria pulverulenta</i>	4
<i>Ephippiorhynchus asiaticus</i>	40
<i>Epthianura aurifrons</i>	4
<i>Epthianura tricolor</i>	18
<i>Eremiornis carteri</i>	18
<i>Erythronys cinctus</i>	22
<i>Esacus magnirostris</i>	2
<i>Esacus neglectus</i>	2
<i>Eurostopodus argus</i>	29
<i>Falco berigora</i>	109
<i>Falco berigora</i> subsp. <i>berigora</i>	5
<i>Falco cenchroides</i>	148
<i>Falco cenchroides</i> subsp. <i>cenchroi</i>	1
<i>Falco longipennis</i>	24
<i>Fulica atra</i>	36
<i>Fulica atra</i> subsp. <i>australis</i>	2
<i>Gallirallus philippensis</i>	3
<i>Gallirallus philippensis</i> subsp. <i>melk</i>	1
<i>Gavialis virescens</i>	68
<i>Gelochelidon nilotica</i> subsp. <i>affinis</i>	1
<i>Geopelia cuneata</i>	134
<i>Geopelia humeralis</i>	28
<i>Geopelia placida</i>	1
<i>Geopelia striata</i>	183
<i>Geopelia striata</i> subsp. <i>placida</i>	12
<i>Geophaps plumifera</i>	54
<i>Gerygone fusca</i>	2
<i>Gerygone tenebrosa</i>	23
<i>Grallina cyanoleuca</i>	300
<i>Grus rubicunda</i>	16
<i>Haematopus fuliginosus</i>	18
<i>Haematopus longirostris</i>	107
<i>Haliaeetus leucogaster</i>	47
<i>Haliastur indus</i>	71
<i>Haliastur sphenurus</i>	170
<i>Hamirostra melanosternon</i>	4

<i>Heteromunia pectoralis</i>	3
<i>Heteroscelus brevipes</i>	8
<i>Hieraaetus morphnoides</i>	27
<i>Himantopus himantopus</i>	79
<i>Hirundo ariel</i>	8
<i>Hirundo neoxena</i>	15
<i>Hirundo nigricans</i>	24
<i>Lalage tricolor</i>	24
<i>Larus novaehollandiae</i>	17
<i>Larus novaehollandiae</i> subsp. novae	6
<i>Lichenostomus keartlandi</i>	37
<i>Lichenostomus penicillatus</i>	236
<i>Lichenostomus plumulus</i>	1
<i>Lichenostomus virescens</i>	200
<i>Lichmera indistincta</i>	106
<i>Lichmera indistincta</i> subsp. indistin	1
<i>Lophoictinia isura</i>	1
<i>Malacorhynchus membranaceus</i>	19
<i>Malurus lamberti</i>	61
<i>Malurus lamberti</i> subsp. assimilis	2
<i>Malurus leucopterus</i>	97
<i>Malurus splendens</i>	1
<i>Manorina flavigula</i>	214
<i>Megalurus cruralis</i>	2
<i>Melanodryas cucullata</i>	2
<i>Melithreptus gularis</i>	13
<i>Melithreptus gularis</i> subsp. laetior	1
<i>Melopsittacus undulatus</i>	148
<i>Merops ornatus</i>	282
<i>Microcarbo melanoleucos</i>	30
<i>Milvus migrans</i>	98
<i>Milvus migrans</i> subsp. affinis	2
<i>Mirafra javanica</i>	118
<i>Motacilla flava</i> subsp. simillima	2
<i>Neochmia ruficauda</i>	16
<i>Neochmia ruficauda</i> subsp. subclai	1
<i>Ninox connivens</i>	20
<i>Ninox novaeseelandiae</i>	30
<i>Numenius phaeopus</i> subsp. varieg.	1
<i>Nycticorax caledonicus</i>	17
<i>Nycticorax caledonicus</i> subsp. hilli	1
<i>Nymphicus hollandicus</i>	67
<i>Ocyphaps lophotes</i>	188
<i>Onychoprion fuscata</i>	1
<i>Oreoica gutturalis</i>	12
<i>Pachycephala lanioides</i>	35
<i>Pachycephala melanura</i>	8
<i>Pachycephala rufiventris</i>	22
<i>Pandion haliaetus</i>	3
<i>Pandion haliaetus</i> subsp. cristatus	4
<i>Pardalotus rubricatus</i>	75
<i>Pardalotus striatus</i>	8
<i>Pardalotus striatus</i> subsp. murchisi	2
<i>Passer montanus</i>	30
<i>Pelecanus conspicillatus</i>	116
<i>Peneoenanthe pulverulenta</i>	2
<i>Petrochelidon ariel</i>	97
<i>Petrochelidon nigricans</i>	83
<i>Petroica goodenovii</i>	12

Phalacrocorax carbo	9
Phalacrocorax melanoleucos	5
Phalacrocorax sulcirostris	44
Phalacrocorax varius	53
Phaps chalcoptera	1
Phaps elegans	1
Phaps histrionica	2
Philemon citreogularis	2
Platalea flavipes	6
Platalea regia	34
Platycercus spurius	1
Platycercus zonarius	2
Podargus strigoides	9
Poliocephalus poliocephalus	11
Pomatostomus temporalis	67
Porphyrio porphyrio	1
Porzana fluminea	3
Ptilonorhynchus guttatus	20
Ptilonorhynchus maculatus	2
Ptilonorhynchus maculatus subsp.	5
Ptilotula keartlandi	6
Ptilotula penicillata	1
Ptilotula penicillatus	23
Purnella albifrons	1
Recurvirostra novaehollandiae	29
Rhipidura albiscapa	6
Rhipidura fuliginosa	1
Rhipidura leucophrys	166
Rhipidura leucophrys subsp. leuco	1
Rhipidura phasiana	20
Smicromis brevirostris	8
Sterna albifrons	8
Sterna albifrons subsp. sinensis	3
Sterna bengalensis	22
Sterna bergii	14
Sterna caspia	11
Sterna hirundo subsp. longipennis	4
Sterna hybrida	9
Sterna hybrida subsp. javanica	1
Sterna leucoptera	11
Sterna nereis	2
Sterna nilotica	3
Stiltia isabella	17
Sugomel niger	11
Tachybaptus novaehollandiae	46
Tachybaptus novaehollandiae subs	4
Taeniopygia guttata	315
Thalasseus bengalensis	15
Threskiornis molucca	56
Threskiornis spinicollis	79
Todiramphus chloris	6
Todiramphus chloris subsp. pilbara	1
Todiramphus pyrrhopygia	14
Todiramphus pyrrhopygius	85
Todiramphus sanctus	87
Todiramphus sanctus subsp. sanct	1
Tribonyx ventralis	1
Tringa cinerea	3
Tringa hypoleucos	2

Turnix pyrrhothorax	3
Turnix velox	80
Tyto alba	1
Tyto alba subsp. delicatula	1
Tyto delicatula	1
Vanellus miles	39
Vanellus tricolor	2
Zosterops luteus	51
FISH	119
? ?	2
Abudefduf bengalensis	1
Acanthopagrus australis	1
Acanthopagrus latus	5
Amniataba caudavittata	1
Amphiprion clarkii	1
Apogon rueppellii	2
Arrhamphus sclerolepis	1
Atelomycterus sp.	2
Bathygobius fuscus	2
Batrachomoeus dahli	2
Butis butis	1
Caranx ignobilis	1
Caranx sp.	1
Centrogenys vaigiensis	1
Cephalopholis boenak	1
Chiloscyllium punctatum	2
Choerodon cyanodus	1
Craterocephalus cuneiceps	2
Dactyloptena papilio	1
Drepane punctata	1
Eleutheronema tetradactylum	1
Epinephelus areolatus	1
Epinephelus malabaricus	1
Epinephelus quoyanus	2
Filicampus tigris	1
Gerres subfasciatus	1
Glossamia aprion	1
Halophryne diemensis	5
Herklotsichthys koningsbergeri	3
Herklotsichthys lippa	1
Hyporhamphus quoyi	1
Hyporhamphus sp.	1
Hypseleotris compressa	1
Hypseleotris compressa?	1
Ichthyoscopus spinosus	2
Labracinus sp.	1
Leiopotherapon unicolor	4
Leptobrama muelleri	1
Liza vaigiensis	2
Lophiocharon trisignatus	2
Lutjanus carponotatus	2
Megalops cyprinoides	2
Melanotaenia australis	5
Monodactylus argenteus	3
Mugil cephalus	2
Nematalosa erebi	1
Nematalosa vlaminghi	1
Neoarius graeffei	1
Neosilurus hyrtlai	1

Omobranchus sp.	1
Ophichthus rutidoderma	1
Opistognathus inornatus	1
Ostracion sp.	1
Paraplotosus albilabris	2
Parascorpaena picta	1
Pentapodus porosus	1
Periophthalmodon freycineti	1
Protonibea diacanthus	1
Rastrelliger serventyi (invalid)	1
Rendahlia jaubertensis	1
Rhynchostracion nasus	1
Salarias sexfilum	1
Salarias sp.	1
Scaevius milii	1
Scomberoides commersonnianus	1
Scomberomorus semifasciatus	1
Selenotoca multifasciata	4
Sillago analis	3
Sillago schomburgkii	1
Strongylura strongylura	2
Synanceia horrida	2
Synodus sageneus	1
Terapon jarbua	1
Yongleichthys nebulosus	2
Zabidius novemaculeatus	2
INVERT	2651
?Simarus sp2	2
Acariformes sp.	8
Achnanthes exigua Grun.	4
Achnanthes exilis K???tz.	2
Achnanthidium minutissima (K???t	10
Adversaeschna brevistyla	1
Aelosoma sp. 1 (PSS)	2
Aeshnidae sp.	1
Aganippe myg207	1
Aganippe sp.2	1
Agraptocorixa parvipunctata	1
Albia sp.	1
Alleculinae Genus 1 sp3	1
Allodessus bistrigatus	7
Allonais pectinata	3
Alluaudomyia sp.	1
Alona cf. verrucosa	1
Alona 'davidi vermiculata'	2
Alona rectangula novaezealandiae	2
Alona rigidicaudis	1
Amblystomus sp1	1
Ameriana sp. P1 (PSW)	2
amphipod Genus 2 sp. B14	2
'Amphitritecandona' 'prima'(pss)	3
Amphora coffeaeformis (Ag.) K???t	1
Amphora veneta K???tz.	6
Aname ellenae	1
Aname sp.1	1
Aname sp.14	1
Aname sp.15	2
Aname sp.3	1
Anax papuensis	4

Ancylidae sp.	2
Anisops canaliculatus	11
Anisops elstoni	1
Anisops gratus	3
Anisops nasutus	5
Anisops sp.	2
Anisops stali	4
Anisops thienemanni	2
Anomoeoneis brachysira (Br???b.)	2
Anomoeoneis sphaerophora (Ehr.)	1
Anomoeoneis styriaca (Grun.) Husi	2
Anomoeoneis vitrea	1
Anomotarus crudelis	1
Anopheles annulipes s.l.	10
ant sp.	24
Arcella sp.	1
Arcella sp. P1	2
Areacandona cf. 'iuno' (PSS)	1
Areacandona 'fortescueiensis' (PS)	1
Areacandona 'incogitata' (PSS)	1
Areacandona 'iuno' (PSS)	2
Areacandona 'jessicae' (PSS)	1
Argiocnemis rubescens	3
Armatalona macrocopa	1
Arrenurus (Brevicadaturus) sp. 18 (1
Arrenurus sp. nov. 2 (PSS)	1
Arthrorhabdus paucispinus	1
Asadipus banjiiwarn	1
Asadipus barlee	1
Asplanchna sieboldi	1
Asplanchna sp.	1
Aulacoseira ambigua	2
Aulodrilus pigueti	5
Austracantha minax	1
Australiobates queenslandensis	2
Australiobates sp. P3 (nr crassiset)	1
Australobolbus impressicollis	1
Australobolbus pseudobscurius	1
Australutica sp.1	3
Australutica sp.3	2
Austraturus sp. P3 (PRP)	1
Austroagrion pindrina/Ischnura het	1
Austroepigomphus (Xerogomphus)	3
Austrogomphus mjobergi	7
Austrolestes analis	1
Austrolestes aridus	1
Austropeplea lessoni	2
Backobourkia collina	1
Baetidae sp.	3
Batrachomatus wingi	2
Bdelloidea sp. 2:2	5
Bennelongia australis lineage	1
Bennelongia australis OrdX (PSW)	1
Bennelongia barangaroo lineage	3
Bennelongia nimala	1
Bennelongia strellyensis	1
Berosus josephenae	1
Berosus nr josephenae (was Pilbar	4
Berosus pulchellus	4

Berosus sp.	6
Bezzia sp. P3 (PSW)	1
Bidessodes denticulatus	1
Blackburnium neocavicolle	1
Blackburnium sp. nov. nr. pauperc	1
Blackburnium sp. nr. neocavicolle	1
Boeckella triarticulata	4
Bogidiellidae sp.	2
Bolbobaineus planiceps	1
Bolboleus truncatus	4
Boreosaragus sp1	16
Boreosaragus sp3	1
Brachionus angularis	1
Brachionus calyciflorus	1
Brachionus cf. forficula	1
Brachionus dichotomus	1
Brachionus quadridentatus	4
Branchinella mcraei	1
Buddelundia '14'	1
Buddelundia sp. 10	1
Buddelundia sp. 13	8
Buddelundia sp. 14RE	3
Buddelundia sp. 17	10
Buddelundia sp. 19	9
Buddelundia sp. 44	1
Buddelundia sp. nov. 10	11
Buddelundia sp. nov. 14	31
Buddelundia sp. nov. 17	9
Buddelundia sp. nov. 19	15
Buddelundia sp. nov. 31	3
Caedius sp4	1
Caedius sp5	1
Caenidae sp.	4
Calamoecia baylyi (Cue form) (ex r	4
Caloneis molaris (Grun.) Krammer	1
Caloneis pulchra Messikommer	1
Caloneis silicula (Ehr.) Cl.	4
Camponotus cf. evae Forel (sp. JD	6
Camponotus cowlei Froggatt	1
Camponotus crozieri McArthur & L	1
Camponotus discors Forel	9
Camponotus fideae Forel	7
Camponotus novaehollandiae Mayr	4
Camponotus wiederkehri Forel	1
Campylodiscus clypeus Ehr.	2
Candonopsis kimberleyi	1
Cardiocondyla nuda (Mayr)	1
Carenum nr. emarginatum sp2	3
Carenum pulchrum	4
Carenum sp10	1
Carenum sp1-black globular	1
Carenum sp8	6
Carenum venustum	1
Catadromus lacordairei	2
Cavisternum clavatum	1
Cephalodella gibba	1
Cephalodella sp. P1 (very long toes	1
Cerapachys fervidus (Wheeler)	1
Ceratopogonidae sp.	4

<i>Ceriodaphnia cornuta</i>	6
<i>Chalcopteroides</i> sp5	1
<i>Chaoborus punctilliger</i>	2
<i>Cheumatopsyche dostinei</i>	1
<i>Cheumatopsyche wellsae</i>	2
<i>Chilibathynella</i> sp.	1
<i>Chimarra</i> sp AV17 (PSW)	1
<i>Chironominae</i> sp.	3
<i>Chironomus</i> aff. <i>alternans</i> (V24) (C	4
<i>Chlaenius</i> ? <i>darlingensis</i>	1
<i>Chlaenius australis</i>	5
<i>Chydorus eurynotus</i>	1
<i>Clinotanypus crux</i>	5
<i>Cloeon</i> sp.	17
<i>Cnephia</i> nr <i>aurantiacum</i>	1
<i>Cnephia tonnoiri</i>	1
<i>Cocconeis placentula</i> var. <i>euglypta</i>	2
<i>Coelopynia pruinosa</i>	17
<i>Coenagrionidae</i> sp.	5
<i>Conchostraca</i> (unident.)	3
<i>Conochilus dossuarius</i>	1
<i>Conopterum pyripenne</i>	1
<i>Copelatus nigrolineatus</i>	3
<i>Corbicula</i> sp.	7
<i>Corixidae</i> sp.	5
<i>Cormocephalus strigosus</i>	2
<i>Corynoneura</i> sp.	1
<i>Craticula cuspidata</i> (Grun. ex. Van	2
<i>Craticula halophila</i> (Grun. ex. Van t	1
<i>Crematogaster frivola</i> Forel	1
<i>Crematogaster queenslandica</i> gp. :	8
<i>Crocothemis nigrifrons</i>	1
<i>Cryptochironomus griseidorsum</i>	10
<i>Cryptoerithrus occultus</i>	1
<i>Cryptoerithrus</i> sp.8	2
<i>Culex</i> (<i>Culex</i>) <i>annulirostris</i>	3
<i>Culex crinicauda</i>	3
<i>Culex</i> nr. <i>crinicauda</i> (PSW)	1
<i>Culex</i> sp.	1
<i>Culicidae</i> sp.	1
<i>Culicoides</i> sp.	1
<i>Culicoides</i> sp. P2 (PSW)	1
<i>Cybister tripunctatus</i>	8
<i>Cylotella stelligera</i> Cl. & Grun.	1
<i>Cymbella affinis</i> K???tz.	4
<i>Cymbella cymbiformis</i> Ag.	1
<i>Cymbella delicatula</i> K???tz.	3
<i>Cypretta baylyi</i>	5
<i>Cypretta seurati</i>	5
<i>Cypretta</i> sp PSW074	4
<i>Cypretta</i> sp. PSW018 (PSW)	1
<i>Cypricercus salinus</i>	4
<i>Cypricercus</i> sp. 422 (CB)	1
<i>Cypricercus</i> sp. 444/885 (CB)	1
<i>Cyprinotis</i> 'maximus' n. sp.	2
<i>Cyprinotus cingalensis</i> (ex kimberle	3
<i>Daphnia carinata</i>	1
<i>Daphnia</i> cf. <i>cephalata</i>	1
<i>Darwinula stevensoni</i>	1

Dasyhelea sp.	1
Dasyheleinae sp. P1 (PSW)	4
Dasyheleinae sp. P2 (PSW)	2
Dero furcata	1
Dero nivea	3
Diacyclops cockingi	5
Diacyclops einslei	1
Diacyclops humphreysi humphreys	5
Diacyclops humphreysi s. str X uni	1
Diacyclops scanloni	7
Diacyclops sobeprolatus	2
Diacyclops sp.	1
Diaphanosoma unguiculatum	2
Dicranophorus halbachi	2
Dicrotendipes jobetus	6
Dicrotendipes sp P4 (PSW)	1
Diffugia sp. P1	4
Diffugia sp. P2	1
Dineutus australis	4
Diplacodes bipunctata	4
Diplacodes haematodes	11
Diploneis pseudovalis Hust.	1
Diplonychus eques	1
Dolichopodidae sp.	3
Dolichopodidae sp. A (SAP)	1
Dunhevedia crassa	2
Dytiscidae sp.	3
Ecnomidae sp.	3
Ecnomus pilbarensis	16
Ecnomus sp. AV16 (PSW)	2
Ectocyclops phaleratus	2
Egadroma sp1 (? piceum)	2
Elaphoidella humphreysi	9
Encentridophorus sarasini	4
Enchytraeidae sp.	4
Encyonema gracile Rabh.	1
Enochrus deserticola	5
Enochrus elongatulus	3
Enochrus maculiceps/deserticola (f	1
Enochrus sp.	2
Enteroplea cf. lacustris (PSW)	1
Eodiaptomus lumholtzi	5
Eolimna minima (Grun.) Lange-Ber	1
Eolimna miniscula (Grun.) Lange-B	1
Eolimna subminiscula (Grun.) Lang	4
Ephemeroporus barroisi s.l.	1
Epistylis sp	2
Epithemia adnata (K???tz.) Br???b	1
Epithemia smithii Carruthers	3
Eretes australis	7
Ergasilidae sp.	1
Ethmostigmus curtipes	6
Ethmostigmus parkeri	1
Ethmostigmus rubripes	1
Euasteron sp.1	1
Euchlanis dilatata	8
Euchlanis incisa	1
Euchlanis oropha	3
Eucyclops australiensis	2

Euglypha sp.	4
Eunotia bilunaris (Ehr.) Mills.	3
Eunotia pectinatus (Dillw.) Rabh.	9
Euonicellus intermedius	1
Euryscaphus waterhousei	2
Eurysticta coolawanyah	8
Eylais sp.	3
Flosculariidae sp.	1
Forcypomyia sp.	1
Fragilaria capucina Desm.	1
Fragilaria capucina var. vaucheriae	1
Fragilaria nitzschoides	1
Fragilaria ulna (Nitz.) Lange Bertalc	5
Gamasomorpha sp.1	2
Gamasomorpha sp.2	2
Geoscaptus laevisimus	2
Gerridae sp.	1
Gigadema sulcatum	3
Glyptophysa sp	1
Gnathaphanus aridus	5
Gnathaphanus melbournensis	1
Gnathaphanus multipunctatus	5
Gomphema gracile Ehr.	1
Gomphidae sp.	1
Gomphonema parvulum (K??tz.) I	6
Gonocephalum sp1	2
Gonocephalum sp2	1
Grayenulla australensis	1
Grayenulla sp. 1	3
Grayenulla sp. 11	1
Grayenulla sp. 19	1
Grayenulla sp. 5	1
Grayenulla waldockae	1
Gretacarus nsp. P1 (PSW)	2
Grymeus sp.10	1
Grymeus sp.12	1
Gyraulus hesperus	14
Gyrosigma attenuatum (K??tz.) R	1
Gyrosigma fonticolum Hust. (in Foç	1
Habronestes sp.8	1
Halicyclops (Rochacyclops) calm	9
Halicyclops (Rochacyclops) roachi	1
Halplidae sp.	1
Halplus halsei	1
Halplus sp.	1
Hantzschia amphioxys (Ehr.) Grun.	6
Hantzschia virgata	2
Harpacticoida sp	1
Helea sp5	5
Hellyethira sp.	1
Helochares/E mastersi larvae	2
Hemicordulia tau	7
Heteronyx beltanae/frenchi	1
Heteronyx parvulus	1
Heteronyx pellucida	1
Heteronyx tepperi	1
Hexarthra cf brandorffi (PSW)	3
Hexarthra mira	1
Hogna sp.2	1

Hogna sp.5	2
Holconia neglecta	1
Holoplatys meda	1
Holoplatys sp. 7	2
Humphreyscandona 'capillus' (PSS)	7
Humphreyscandona sp.	1
Humphreyscandona waldockae	1
Hydaticus quadrivittatus	1
Hydra sp.	3
Hydrachna sp.	2
Hydrachna sp. 4/5 (PSW)	3
Hydraena barbipes	6
Hydraena cf. rudallensis (PSW)	3
Hydraena sp.	1
Hydrobiidae sp P1 (not assimineid)	1
Hydrochus burdekinensis	1
Hydrochus group 3 "black" (PSW)	3
Hydrochus interioris	1
Hydrochus macroaquilonius	1
Hydrochus sp. P1 (PSW)	2
Hydrochus sp. P5	1
Hydroglyphus basalis	3
Hydroglyphus basalis var fuscoline:	1
Hydroglyphus grammopterus (=trilii	9
Hydroglyphus leai	4
Hydroglyphus orthogrammus	4
Hydrophilidae sp.	2
Hydrophilus brevispina	2
Hydrovatus rufoniger	3
Hydrovatus weiri	1
Hypharpax sp2	1
Hypharpax sp3	1
Hypharpax sp4	1
Hypharpax sp5	2
Hyphydrus lyratus	8
Hyphydrus sp.	1
Ilyocryptus raridentatus	3
Ilyocypris australiensis	2
Ilyocypris perigundi	1
Ilyocypris 'spiculata' (ms name) (S/	2
Ilyodromus sp. PB	4
Indolpium sp.	13
Iridomyrmex anceps (Roger)	5
Iridomyrmex bicknelli azureus Vieh	1
Iridomyrmex chasei complex sp. JC	1
Iridomyrmex chasei concolor Forel	5
Iridomyrmex chasei Forel	1
Iridomyrmex hartmeyer Forel	1
Iridomyrmex hartmeyer gp sp. JDM	10
Iridomyrmex sanguineus Forel	1
Iridomyrmex sp. JDM 133	4
Iridomyrmex sp. JDM 137 (incl. '31:	4
Iridomyrmex sp. JDM 319	6
Iridomyrmex sp. JDM 843	6
Ischnura aurora aurora	13
Ischnura heterosticta heterosticta	2
Isidorella egraria	6
Isocypris williamsi (ex Ilyodromus s	4
Karayevia clevei cf.	2

<i>Keratella australis</i>	1
<i>Keratella procurva</i>	6
<i>Keratella slacki</i>	1
<i>Keratella</i> sp. nov. (aff. <i>australis</i> grp)	1
<i>Keratella tropica</i>	4
<i>Knoelle clara</i>	5
<i>Kurzia longirostris</i>	1
<i>Kwonkan myg007</i>	4
<i>Kwonkan</i> sp.1	1
<i>Kymberlia</i> sp.1	1
<i>Laccobius matthewsi</i>	2
<i>Laccophilus clarki</i>	1
<i>Laccophilus sharpi</i>	9
<i>Laccotrephes tristis</i>	1
<i>Lagynochthonius leemouldi</i>	7
<i>Lampona ampeinna</i>	4
<i>Lamponina scutata</i>	19
<i>Larsia albiceps</i>	19
<i>Latonopsis australis</i>	2
<i>Latonopsis brehmi</i>	1
<i>Latrodectus hasseltii</i>	1
<i>Leberis</i> cf. <i>diaphanus</i> (striate) (PSV)	1
<i>Lecane bulla</i>	5
<i>Lecane</i> cf. <i>spenceri</i> (PSW)	1
<i>Lecane crepida</i>	1
<i>Lecane decipiens</i>	1
<i>Lecane hornemanni</i>	1
<i>Lecane ludwigi</i> form P1	1
<i>Lecane luna</i>	2
<i>Lecane obtusa</i>	1
<i>Lecane papuana</i>	2
<i>Lecane ungulata</i>	1
<i>Lecanomerus</i> sp1	1
' <i>Leicacandona</i> ' ' <i>gyralea</i> ' (PSS)	1
' <i>Leicacandona</i> ' ' <i>mookae</i> ' (PSS)	1
' <i>Leicacandona</i> ' ' <i>quasimookae</i> ' (PSS)	3
<i>Lepadella amphitropis</i>	1
<i>Lepadella ovalis</i>	2
<i>Lepadella patella</i>	1
<i>Lepidoptera</i> (non-pyralid) Pilbara sp.	1
<i>Lepidoptera</i> sp.	1
<i>Leptoceridae</i> sp.	3
<i>Lesquereusia spiralis</i>	1
<i>Lethocerus distinctifemur</i>	3
<i>Leydigia australis</i>	4
<i>Libellulidae</i> sp.	3
<i>Limbodessus compactus</i>	2
<i>Limnebius</i> sp.	1
<i>Limnesia</i> sp.	1
<i>Limnesia</i> sp. 4 (PSW)	3
<i>Limnichidae</i> sp.	1
<i>Limnochares australica</i>	3
<i>Limnocythere dorsosicula</i>	3
<i>Limnocythere</i> sp BOS068	1
<i>Limnocythere stationis</i>	1
<i>Limnocytheridae</i> n.gen. sp 419 (CE)	2
<i>Limnogonus luctuosus</i>	2
<i>Liparetrus</i> sp. nr. <i>atrox</i>	1
<i>Loxandrus laevigatus</i>	3

<i>Loxandrus micantior</i>	3
<i>Loxandrus</i> WA n sp1	2
<i>Luticola goeppertiana</i> (Bleisch) Ma	1
<i>Luticola mutica</i> (K???tz.) Mann	1
<i>Lychas</i> ???bituberculatus???	3
<i>Lychas</i> 'adonis'	2
<i>Lychas annulatus</i>	4
<i>Lychas annulatus</i> ' Glauert, 1925	1
<i>Lychas bituberculatus</i>	8
<i>Lychas</i> 'harveyi'	5
<i>Lychas</i> sp. 2	4
<i>Lychas</i> sp. 3	1
<i>Lychas</i> sp. 4	4
<i>Lychas</i> sp. 6	1
<i>Lychas</i> sp. 8	2
<i>Lycidas</i> sp. 1	1
<i>Lycidas</i> sp. 11	4
<i>Lynceus argillaphilus</i>	1
<i>Macrobrachium</i> sp. P1	1
<i>Macrochaetus altamirai</i>	3
<i>Macrodiplax cora</i>	4
<i>Macrothrix capensis</i>	1
<i>Macrothrix</i> cf. <i>pectinata</i>	1
<i>Macrothrix indistincta</i>	1
<i>Macrothrix</i> sp.	1
<i>Masasteron tealei</i>	5
<i>Mastogloia elliptica</i> (Ag.) Cl.	4
<i>Mastogloia elliptica</i> var. <i>danseii</i> (th	3
<i>Mastogloia smithii</i> Thwaites	3
<i>Mayamaea atomus</i>	1
<i>Megaporus ruficeps</i>	1
<i>Megaporus</i> sp.	1
<i>Melaps</i> sp2	1
<i>Melitidae</i> sp.	1
<i>Melitidae</i> sp. 1 (PSS)	4
<i>Melophorus bagoti</i> Lubbock	1
<i>Melophorus froggatti</i> Forel	1
<i>Melophorus ludius sulla</i> Forel	4
<i>Melophorus</i> nr. <i>aeneovirens</i> (Lowne	2
<i>Melophorus turneri</i> Forel	2
<i>Melophorus wheeleri</i> complex sp. J	1
<i>Melophorus wheeleri</i> Forel	1
<i>Mesocyclops brooksi</i>	10
<i>Mesocyclops darwini</i>	1
<i>Mesomorphus</i> sp1	2
<i>Mesovelia horvathi</i>	1
<i>Mesovelia hungerfordi</i>	2
<i>Mesovelia vittigera</i>	1
<i>Metacyclops/Pescecyclops</i> sp.	2
<i>Metistete</i> sp1	1
<i>Microcerberidae</i> sp.	4
<i>Microchironomus</i> 'K1' (PSW)	1
<i>Microcyclops varicans</i>	10
<i>Micronecta adelaidae</i> (ex P4)	3
<i>Micronecta gracilis</i>	2
<i>Micronecta micra</i>	11
<i>Micronecta</i> n. sp. P3 (PSW)	3
<i>Micronecta robusta</i>	4
<i>Micronecta</i> sp.	1

Microvelia (Austromicrovelia) austr:	1
Microvelia (Austromicrovelia) peran	1
Microvelia (Pacifcovelia) oceanica	2
Microvelia sp.	1
Minasteron minusculum	10
Miralona victoriensis	1
Missulena rutaspina	1
Missulena sp.3	1
Moina aff weismanni	2
Moina cf. australiensis (CB)	1
Moina micrura s.l.	3
Monomorium carinatum Heterick	1
Monomorium disetigerum Heterick	11
Monomorium fieldi Forel	4
Monomorium insolescens Wheeler	1
Monomorium laeve Mayr	11
Monomorium rothsteini Forel	5
Monomorium sydneyense Forel (se	1
Monopylephorus n. sp. WA29 (ex F	10
Morebilus diversus	4
Muscidae sp. A (SAP)	3
Muscidae sp. N	1
Myrmopopaea sp.13	2
Myrmopopaea sp.15	1
Myrmopopaea sp.8	1
Mytilina ventralis macracantha	3
Naididae (ex Tubificidae)	11
Nais communis	1
Navicula absoluta	1
Navicula bryophila Petersen	2
Navicula cryptocephala K???tz.	5
Navicula cryptonella Lange-Bertalo	1
Navicula elginensis (Greg.) Ralfs.	2
Navicula kriegerii	1
Navicula molestiformis Hust.	7
Navicula muraliformis	1
Navicula rhynchocephala K???tz.	2
Navicula schroeterii Meister	1
Navicula similis Krasske	1
Navicula subrhynchocephala Hust.	4
Navicula tenelloides Hust.	1
Navicula veneta K???tz.	1
Nedsia nr hurlberti	2
Nedsia sp.	12
Nematoda sp.	3
Nematoda sp. P2/P4 (PSW)	2
Nematoda sp. P6 (PSW)	1
Nematoda sp. P8 (PSW)	1
Neocarenum blackburni	2
Neocarenum sp2	3
Neostorena sp.5	1
Neothrix superarmata	3
Nephila edulis	1
Nilobezzia sp.	1
Nilobezzia sp. P2 (PSW)	5
Nitzschia amphibia Grun.	4
Nitzschia calida Grun.	1
Nitzschia capitellata	1
Nitzschia compressa (Bailey) Boye	2

Nitzschia compressa var. elongata	1
Nitzschia constricta (Greg.) Grun.	3
Nitzschia desertorum Hust.	1
Nitzschia dissipata (K???tz.) Grun.	2
Nitzschia filiformis (W. Sm.) Van H.	5
Nitzschia frustulum (K???tz.) Grun.	4
Nitzschia intermedia Hantz.	1
Nitzschia levidensis var. victoriae (G.	1
Nitzschia linearis (Ag.) W. Sm.	1
Nitzschia microcephala Grun.	2
Nitzschia palea (K???tz.) W. Sm.	9
Nitzschia perminuta (Grun.) M. Per	1
Nitzschia sigma (K???tz.) W. Sm.	7
Nitzschia sublinearis Hust.	1
Nitzschia umbonata (Ehr.) Lange-B	2
No invertebrates	21
Nomindra leeuwenii	3
Notobathynella sp.	2
Notobia sp1	3
Notommata cf. pachyura (PSW)	1
Notommata nr cerberus (PSW)	1
Notonectidae sp.	4
Nototarus sp. nov. 14 (MB)	1
Nototarus sp. nov. 57 (MB)	1
Nototarus sp. nov. 69 (MB)	1
Ochetellus flavipes (Kirby)	5
Ochthebius sp. P1 (PSW)	1
Ochthebius sp. P2 (PSW)	2
Odontomachus ruficeps Smith	6
Oecetis sp.	1
Oecetis sp. Pilbara 2 (PSW)	1
Oecetis sp. Pilbara 4 (PSW)	3
Oecetis sp. Pilbara 5 (PSW)	10
Oecetis sp. Pilbara 6 (PSW)	1
Oecetis sp. Pilbara 8 (PSW)	1
Oligochaeta sp.	1
Onthophagus consentaneus	1
Onthophagus gazella	12
Onthophagus margaretensis	9
Onthophagus minusculus	3
Onthophagus mjobergi	1
Onthophagus pugnacior	1
Oodes sp1	1
Oodes sp4	1
Oodes sp5	2
Opisthopsis haddoni rufoniger Fore	4
Opopaea sp.17	2
Opopaea sp.2	3
Oribatida group 1 (PSS)	2
Oribatida sp. 4 (PSW)	1
Orthetrum caledonicum	13
Orthetrum pruinatum migratum	2
Orthocladinae sp.	1
Ostracoda (unident.)	11
Ovatalona cf. cambouei	2
Oxus orientalis	1
Ozestheria packardii	5
Pachycondyla (Brachyponera) lutea	3
Palorinae sp2	1

<i>Pantala flavescens</i>	2
<i>Paracladopelma</i> sp. P1 (nr M1) (P.	1
<i>Paracladopelma</i> sp. P2 (nr M2) (P.	3
<i>Paracyclops chiltoni</i>	3
<i>Paracymus pygmaeus</i>	2
<i>Paracymus spenceri</i>	5
<i>Paramelitidae</i> cf. sp. 2 (PSS)	2
<i>Paramelitidae</i> sp.	2
<i>Paramelitidae</i> sp. 2 (PSS)	4
<i>Paramelitidae</i> sp. 6 (PSS)	2
<i>Paramelitidae</i> sp. 8 (PSS)	1
<i>Parametriocnemus ornaticornis</i>	1
<i>Parametriocnemus</i> sp P1 (PSW)	1
<i>Paranacaena</i> sp. P1	1
<i>Paranyctiophylax</i> sp AV5 (KIM-UW	1
<i>Parastenocarididae</i> sp.	1
<i>Parastenocaris jane</i>	1
<i>Paratanytarsus</i> sp. P1 (PSW)	1
<i>Paratendipes</i> sp. 'K1' (PSW)	1
<i>Paratrechina braueri</i> glabrior (Forel	1
<i>Paratrechina minutula</i> (Forel)	2
<i>Pellenes bitaeniata</i>	1
<i>Pentaneurini</i> sp. P3 (PSW)	1
<i>Pentaneurini</i> sp. P6 (PSW)	1
<i>Pescecyclops</i> sp. P1	1
<i>Pheidole</i> sp. JDM 1068	2
<i>Pheidole</i> sp. JDM 1176	1
<i>Pheidole</i> sp. JDM 280	2
<i>Pheidole</i> sp. JDM 337	1
<i>Pheidole</i> sp. JDM 536	3
<i>Phorticosomus</i> ?grandis	1
<i>Phorticosomus gularis</i>	13
<i>Phorticosomus</i> sp1	2
<i>Phorticosomus</i> sp2	1
<i>Phorticosomus</i> sp3	3
<i>Phreodrilid</i> with dissimilar ventral cl	6
<i>Phreodrilid</i> with similar ventral cha	2
<i>Pilbarascutigera incola</i>	2
<i>Pilbarus millsii</i>	1
<i>Pinnularia brevicostata</i> Cl.	1
<i>Pinnularia divergens</i> W. Sm.	5
<i>Pinnularia gibba</i> Ehr.	1
<i>Pinnularia obscura</i>	1
<i>Pinnularia subcapitata</i> Greg.	1
<i>Pinnularia subrostrata</i> (A. Cl.) Cl.-E	2
<i>Planorbidae</i> sp.	4
<i>Planothidium lanceolata</i> var. rostra	1
<i>Platonus patulus</i>	2
<i>Platycoelus melliei</i>	2
<i>Platycoelus</i> sp1	1
<i>Platyias quadricornis</i>	2
<i>Platynectes decempunctatus</i> var d	2
<i>Pleidae</i> sp.	3
<i>Pleurosigma delicatulum</i> W. Sm.	1
<i>Plotiopsis australis</i>	13
<i>Polyarthra dolichoptera</i>	5
<i>Polypedilum griseoguttatum</i>	1
<i>Polypedilum leei</i>	10
<i>Polypedilum</i> nr vespertinus (M2) (S	1

<i>Polypedilum nubifer</i>	4
<i>Polypedilum watsoni</i>	6
<i>Polyrhachis</i> (<i>Campomyrma</i>) <i>incons</i>	2
<i>Polyrhachis</i> (<i>Campomyrma</i>) sp. JD	1
<i>Polyrhachis</i> (<i>Chariomyrma</i>) 'aurea'	2
<i>Polyrhachis</i> (<i>Chariomyrma</i>) nr. <i>heir</i>	1
<i>Polyrhachis</i> (<i>Chariomyrma</i>) <i>senilis</i>	2
<i>Polyrhachis</i> (<i>Hagiomyrma</i>) <i>crawley</i>	1
<i>Polyrhachis</i> (<i>Hagiomyrma</i>) sp. JDM	1
<i>Polyrhachis</i> (<i>Hagiomyrma</i>) sp. JDM	1
<i>Prethopalpus eberhardi</i>	2
<i>Pristina longiseta</i>	3
<i>Pristina sima</i>	1
<i>Procladius paludicola</i>	6
<i>Procordulia affinis</i>	1
<i>Pseudagrion aureofrons</i>	10
<i>Pseudagrion microcephalum</i>	8
<i>Pseudectinosoma galassiae</i>	2
<i>Pseudholophylla</i> sp. nov.	1
<i>Pseudocloeon hypodelum</i> (ex Baet	5
<i>Pseudomonospilus diporus</i>	1
<i>pseudoscorpion austrohorus</i>	5
<i>pseudoscorpion</i> Genus 7/4	6
<i>pseudoscorpion indolpium</i>	267
<i>pseudoscorpion</i> sp.	4
<i>Ptygura</i> sp.	2
<i>Pyalidae</i> nr. sp. 39/40 of JHH (SAI	2
<i>Pyalidae</i> Pilbara sp 2 (PSW)	5
<i>Pyalidae</i> Pilbara sp 5 (PSW)	2
<i>Pyalidae</i> Pilbara sp 6 (PSW)	1
<i>Pyalidae</i> sp.	3
<i>Recifella</i> sp.	1
<i>Regimbartia attenuata</i>	7
<i>Reimeria sinutata</i>	1
<i>Renneria kamouni</i>	1
<i>Rhagada</i> cf. <i>richardsonii</i>	7
<i>Rhagada</i> sp. (juv)	2
<i>Rhagadotarsus anomalus</i>	1
<i>Rheocricotopus</i> sp. P1 (PSW)	2
<i>Rheotanytarsus christinae</i>	1
<i>Rheotanytarsus</i> sp.	1
<i>Rheotanytarsus trivittatus</i>	2
<i>Rhopalodia gibba</i> (Ehr.) O. Mull.)	5
<i>Rhytidoponera crassinoda</i> (Forel)	7
<i>Rhytidoponera taurus</i> Forel	2
<i>Rhytidoponera tyloxys</i> Brown & Doi	4
<i>Rhytidoponera violacea</i> (Forel)	1
<i>Rhytisternus large</i> sp2	1
<i>Rhytisternus medium</i> sp1	1
<i>Rhytisternus small</i> sp1	1
<i>Rotifera</i> sp.	1
<i>Saldidae</i> sp.	3
<i>Sarothrocrepis benefica</i>	1
<i>Scaridium bostjani</i>	1
<i>Schizopera</i> sp.	1
<i>Scirtidae</i> sp.	3
<i>Scolopendra laeta</i>	8
<i>Scolopendra morsitans</i>	7
<i>Scopodes</i> sp (<i>denticollis</i> -gr)	1

<i>Sellephora pupula</i> (K???tz) Meresc	1
<i>Sellophora seminulum</i> (Grun.) Man	1
<i>Simulium ornatipes</i>	1
<i>Skusella</i> nr "V12 ex-WA" (Cranstor	4
<i>Sobas</i> ?minor	3
<i>Spercheus platycephalus</i>	1
<i>Spinasteron arenarium</i>	1
<i>Spinasteron</i> sp.2	2
<i>Spongillidae</i> sp.	2
<i>Staphylinidae</i> sp.	4
<i>Stauroneis anceps</i> Ehr.	4
<i>Stauroneis phoenicenteron</i> (Nitz.) E	3
<i>Stauroneis producta</i>	1
<i>Staurosira construens</i> Ehr.	2
<i>Stenaspidius</i> sp. nov. nr. <i>albosetos</i>	2
<i>Stereomyrmex anderseni</i> (Taylor)	2
<i>Sternolophus australis</i>	1
<i>Sternopriscus multimaculatus</i>	3
<i>Sternopriscus pilbarensis</i>	2
<i>Sternopriscus</i> sp.	4
<i>Stigmacros</i> sp. JDM 829	1
<i>Stilobezzia</i> sp P1 (PSW)	1
<i>Strandesia</i> sp.	2
<i>Stratiomyidae</i> sp.	4
<i>Stygonitocrella bispinosa</i>	5
<i>Stygonitocrella trispinosa</i>	9
<i>Stygonitocrella unispinosa</i>	1
<i>Supunna</i> sp.1	7
<i>Supunna</i> sp.11	1
<i>Supunna</i> sp.13	2
<i>Supunna</i> sp.16	1
<i>Surirella ovalis</i> Br???b.	1
<i>Surirella striatula</i> Turp.	1
<i>Synothele</i> sp.2	1
<i>Tabanidae</i> sp.	7
<i>Tamopsis facialis</i>	1
<i>Tanypodinae</i> sp.	4
<i>Tanytarsus fuscithorax/semibarbita</i>	4
<i>Tanytarsus</i> sp. D (SAP)	1
<i>Tanytarsus</i> sp. G (SAP)	2
<i>Tanytarsus</i> sp. P10 (PSW)	1
<i>Tanytarsus</i> sp. P2 (PSW)	1
<i>Tanytarsus</i> sp. P4 (PSW)	3
<i>Tapinoma</i> sp. JDM 78	5
<i>Tasmanocoenis arcuata</i>	15
<i>Tasmanocoenis</i> sp. E (PSW)	1
<i>Tesserodon granulatum</i>	1
<i>Tesserodon novaehollandiae</i>	4
<i>Testudinella</i> cf <i>trilobata</i> (=sp P3 PS	1
<i>Testudinella</i> cf. <i>elliptica</i> (PSW)	1
<i>Testudinella patina</i>	6
<i>Testudinella</i> sp P2 (PSW)	1
<i>Tetramorpha alaus</i>	6
<i>Tetramorium sjostedti</i> Forel	3
<i>Tetramorium spininode</i> Bolton	7
<i>Tetramorium striolatum</i> gp. sp. JDM	1
<i>Tetramorium striolatum</i> Viehmeyer	6
<i>Thermocyclops decipiens</i>	2
<i>Thiaridae</i> sp.	3

Thienemanniella sp. P1 (PSW)	1
Tiporus tambreyi	4
Tipulidae sp.	1
Tipulidae type P1 (PSW)	2
Tramea stenoloba	4
Triaenodes sp. P1=P2 (PSW)	8
Trichocarenum cylindricum	2
Trichocerca braziliensis	1
Trichocerca similis	7
Trichocycclus gnalooma	3
Triplectides australis	8
Tropocyclops confinis (ex Paracycl	1
Tubificidae stygo type 4	1
Turbellaria sp.	2
Tyrannochthonius aridus	1
Unidentati genus 5 sp. 1	3
Unionicola crassipalpis	2
Unionicola nr minutissima (PSW)	1
Unionicola nr vidrinei (PSW)	1
Unionicola sp P1 (PSW)	1
Unionicola vidrinei	1
Urodacus armatus	1
Urodacus hoplurus	2
Urodacus sp. 2	1
Urodacus sp. 5	1
Urodacus sp. 6	2
Urodacus varians	1
Urodacus yaschenkoi	1
Veliidae sp.	1
Venator sp.5	1
Vestalenula matildae	2
Xanthagrion erythroneurum	3
Xenochironomus sp P3 (PSW)	1
Yilgarnia sp.2	1
Zebraplatys keyserlingi	5
Zenodorus orbiculatus	1
Zonocyprretta kalimna	2
Zyxomma elgneri	2
MAMMAL	1721
Antechinomys laniger	1
Bos taurus	8
Camelus dromedarius	2
Canis dingo	2
Canis familiaris	1
Canis lupus	7
Canis lupus subsp. dingo	2
Canis lupus subsp. familiaris	5
Chaerephon jobensis	4
Chalinolobus gouldii	9
Dasyercus sp.	17
Dasykaluta rosamondae	138
Dasykaluta rosemondade	1
Equus caballus	2
Felis catus	67
Macropus agilis	2
Macropus robustus	29
Macropus robustus subsp. erubesc	11
Macropus rufus	18
Mormopterus (Ozimops) cobourgia	5

Mormopterus loriae	1
Mormopterus loriae subsp. cobourgi	1
Mus musculus	127
Ningauia timealeyi	17
Notomys alexis	482
Nyctophilus arnhemensis	2
Nyctophilus geoffroyi	1
Nyctophilus geoffroyi subsp. pallasi	1
Oryzomys cuniculus	3
Onychomys robustus	3
Ozomys cobourgi	1
Petrogale rothschildi	14
Planigale ingrami	8
Planigale maculata	1
Planigale Sp.1 (WAM)	13
Pseudantechinus roryi	2
Pseudantechinus woolleyae	9
Pseudomys delicatulus	10
Pseudomys desertor	72
Pseudomys hermannsburgensis	348
Pseudomys nanus	4
Pteropus scapulatus	4
Rattus rattus	2
Rhinonictes aurantius	14
Saccolaimus flaviventris	6
Scotorepens greyii	4
Sminthopsis macroura	25
Sminthopsis youngsoni	56
Sousa chinensis	3
Tachyglossus aculeatus	7
Tadarida australis	1
Taphozous georgianus	40
Taphozous hilli	1
Tursiops aduncus	1
Tursiops sp.	1
Vespadelus finlaysoni	37
Vulpes vulpes	8
Zyomys argurus	60
REPTILE	2487
Acanthophis GT NOTHERN species	1
Acanthophis pyrrhus	16
Acanthophis wellsi	1
Aipysurus laevis	2
Amphibolurus gilberti	5
Amphibolurus longirostris	23
Anilius ammodytes	11
Antaresia perthensis	8
Antaresia stimsoni	6
Antaresia stimsoni subsp. stimsoni	3
Aspidites melanocephalus	6
Aspidites ramsayi	9
Brachyurophis approximans	3
Brachyurophis fasciolatus subsp. fasciolatus	1
Carlia munda	19
Carlia triacantha	19
Chelonia sp.	3
Crenadactylus ocellatus subsp. horneri	3
Cryptoblepharus buecananii	1
Cryptoblepharus plagiocephalus	1

<i>Cryptoblepharus ustulatus</i>	4
<i>Ctenophorus caudicinctus</i>	31
<i>Ctenophorus caudicinctus</i> subsp. c	27
<i>Ctenophorus isolepis</i>	95
<i>Ctenophorus isolepis</i> subsp. gularis	1
<i>Ctenophorus isolepis</i> subsp. isolep	69
<i>Ctenophorus nuchalis</i>	73
<i>Ctenophorus reticulatus</i>	2
<i>Ctenotus brooksi</i>	1
<i>Ctenotus colletti</i>	1
<i>Ctenotus duricola</i>	26
<i>Ctenotus duricola/piankai</i>	8
<i>Ctenotus dux</i>	1
<i>Ctenotus grandis</i>	21
<i>Ctenotus grandis</i> subsp. grandis	4
<i>Ctenotus grandis</i> subsp. titan	5
<i>Ctenotus hanloni</i>	8
<i>Ctenotus helenae</i>	42
<i>Ctenotus pantherinus</i>	80
<i>Ctenotus pantherinus</i> subsp. ocellif	40
<i>Ctenotus piankai</i>	16
<i>Ctenotus quattuordecimlineatus</i>	7
<i>Ctenotus robustus</i>	1
<i>Ctenotus rubicundus</i>	3
<i>Ctenotus rufescens</i>	8
<i>Ctenotus rutilans</i>	2
<i>Ctenotus saxatilis</i>	103
<i>Ctenotus schomburgkii</i>	9
<i>Ctenotus schomburgkii</i> Western Pi	2
<i>Ctenotus serventyi</i>	31
<i>Cyclodomorphus melanops</i>	2
<i>Cyclodomorphus melanops</i> subsp.	3
<i>Delma borea</i>	1
<i>Delma butleri</i>	6
<i>Delma desmosa</i>	21
<i>Delma elegans</i>	2
<i>Delma haroldi</i>	11
<i>Delma nasuta</i>	8
<i>Delma pax</i>	31
<i>Delma</i> sp.	2
<i>Delma tincta</i>	16
<i>Demansia angusticeps</i>	1
<i>Demansia psammophis</i>	2
<i>Demansia psammophis</i> subsp. cup	1
<i>Demansia psammophis</i> subsp. psa	1
<i>Demansia rufescens</i>	7
<i>Demansia</i> sp.	1
<i>Demansia torquata</i>	1
<i>Diplodactylus conspicillatus</i>	95
<i>Diplodactylus laevis</i>	12
<i>Diplodactylus savagei</i>	7
<i>Diplodactylus stenodactylus</i>	1
<i>Diporiphora paraconvergens</i>	2
<i>Diporiphora pindan</i>	26
<i>Diporiphora valens</i>	1
<i>Diporiphora vescus</i>	10
<i>Diporiphora winneckeii</i>	5
<i>Egernia depressa</i>	3
<i>Egernia epsisolus</i>	4

<i>Egernia striata</i>	1
<i>Ephalophis greyae</i>	1
<i>Eremiascincus fasciolatus</i>	6
<i>Eremiascincus isolepis</i>	3
<i>Eremiascincus musivus</i>	30
<i>Eremiascincus pallidus</i>	5
<i>Eremiascincus richardsonii</i>	4
<i>Eretmochelys imbricata</i> subsp. bis	1
<i>Fordonia leucobalia</i>	4
<i>Furina ornata</i>	7
<i>Gehyra 'fenestra'</i>	1
<i>Gehyra pilbara</i>	28
<i>Gehyra punctata</i>	40
<i>Gehyra purpurascens</i>	5
<i>Gehyra</i> sp.	5
<i>Gehyra variegata</i>	50
<i>Gehyra variegata/purpurascens</i>	5
<i>Hemidactylus frenatus</i>	1
<i>Heteronotia binoei</i>	26
<i>Heteronotia spelea</i>	5
<i>Hydrelaps darwiniensis</i>	1
<i>Hydrophis elegans</i>	2
<i>Hydrophis kingii</i>	1
<i>Hydrophis stokesii</i>	2
<i>Lerista aff. bipes</i>	15
<i>Lerista bipes</i>	309
<i>Lerista clara</i>	23
<i>Lerista ips</i>	6
<i>Lerista jacksoni</i>	19
<i>Lerista muelleri</i>	2
<i>Lerista</i> sp.	3
<i>Lialis burtonis</i>	19
<i>Liopholis striata</i>	11
<i>Lophognathus gilberti</i>	1
<i>Lophognathus longirostris</i>	4
<i>Lucasium stenodactylum</i>	20
<i>Lucasium wombeyi</i>	6
<i>Lucasium 'woodwardi'</i>	5
<i>Menetia greyii</i>	57
<i>Menetia surda</i> subsp. <i>surda</i>	2
<i>Moloch horridus</i>	2
<i>Morethia ruficauda</i>	24
<i>Morethia ruficauda</i> subsp. <i>exquisita</i>	7
<i>Morethia ruficauda</i> subsp. <i>ruficauda</i>	9
<i>Nephurus laevis</i>	2
<i>Nephurus levis</i>	4
<i>Nephurus levis</i> subsp. <i>pilbarensis</i>	10
<i>Notoscincus ornatus</i>	10
<i>Notoscincus ornatus</i> subsp. <i>ornatus</i>	7
<i>Oedura fimbria</i>	4
<i>Oedura marmorata</i>	14
<i>Pogona minor</i>	16
<i>Pogona minor</i> subsp. <i>minor</i>	6
<i>Pogona minor</i> subsp. <i>mittelli</i>	15
<i>Proablepharus reginae</i>	2
<i>Pseudechis australis</i>	13
<i>Pseudonaja mengdeni</i>	11
<i>Pseudonaja modesta</i>	24
<i>Pseudonaja nuchalis</i>	4

Pygopus nigriceps	12
Ramphotyphlops ammodytes	40
Ramphotyphlops braminus	1
Ramphotyphlops grypus	21
Ramphotyphlops GT NOTHERN s	1
Ramphotyphlops pilbarensis	8
Rhynchoedura ornata	22
Simoselaps anomalus	20
Strophurus ciliaris	12
Strophurus ciliaris subsp. aberrans	22
Strophurus ciliaris subsp. ciliaris	1
Strophurus elderi	4
Strophurus jeanae	29
Suta fasciata	1
Suta punctata	10
Tiliqua multifasciata	26
Varanus acanthurus	62
Varanus brevipinna	30
Varanus bushi	1
Varanus eremius	34
Varanus giganteus	26
Varanus gilleni	1
Varanus gouldii	18
Varanus panoptes	1
Varanus panoptes subsp. rubidus	1
Varanus pilbarensis	18
Fungi	14
FUNGUS	12
Anthracoecystis paraneurachnis	1
Cercospora ipomoeae	1
Corioloopsis brunneo-leuca	1
Curvularia sp.	1
Ganoderma steyaertanum	1
Macalpinomyces eriachnes	2
Phyllachora sp.	1
Pisolithus tinctorius	1
Polyporus hartmannii	1
Triodiomyces altilis	1
Triodiomyces lituanus	1
LICHEN	2
Xanthoparmelia taractica	2
Plantae	3592
ALGA	41
Acanthophora spicifera	3
Anadyomene plicata	1
Asparagopsis taxiformis	2
Caulerpa brachypus	2
Caulerpa chemnitzia	4
Caulerpa cylindracea	1
Caulerpa lamourouxii	3
Caulerpa lentillifera	2
Caulerpa racemosa forma laxa	1
Caulerpa sertularioides	1
Chaetomorpha melagonium	1
Dichotomaria obtusata	1
Dictyosphaeria cavernosa	1
Galaxaura rugosa	1
Gelidiella acerosa	1
Heterosiphonia crassipes	3

Neomeris bilimbata	1
Neomeris van-bosseae	5
Sebdenia flabellata	1
Udotea argentea	5
Udotea glaucescens	1
DICOT	2737
Abrus precatorius	1
Abutilon amplum	1
Abutilon fraseri	1
Abutilon indicum var. australiense	1
Abutilon lepidum	2
Abutilon macrum	2
Abutilon otocarpum	3
Abutilon oxycarpum subsp. Prostra	1
Abutilon sp. Dioicum (A.A. Mitchell	1
Abutilon sp. Pilbara (W.R. Barker 2	1
Acacia acradenia	13
Acacia adoxa var. adoxa	17
Acacia adoxa var. subglabra	3
Acacia ampliceps	4
Acacia ampliceps x bivenosa	3
Acacia anaticeps	15
Acacia ancistrocarpa	26
Acacia bivenosa	11
Acacia bivenosa x sclerosperma sl	1
Acacia colei	2
Acacia colei var. colei	25
Acacia colei var. ileocarpa	1
Acacia coriacea	2
Acacia dictyophleba	2
Acacia drepanocarpa subsp. drepa	1
Acacia eriopoda	7
Acacia eriopoda x tumida var. pilba	1
Acacia glaucocaesia	4
Acacia hilliana	14
Acacia hilliana x stellaticeps	1
Acacia inaequilatera	11
Acacia maitlandii	1
Acacia monticola	11
Acacia monticola x tumida var. pilb	2
Acacia orthocarpa	10
Acacia pruinocarpa	1
Acacia ptychophylla	3
Acacia pyrifolia var. pyrifolia	4
Acacia retivenea subsp. clandestin	1
Acacia robeorum	7
Acacia sabulosa	8
Acacia sclerosperma subsp. sclero	5
Acacia sericophylla	9
Acacia sp.	1
Acacia sp. Nalgi (N.T. Burbidge 13	5
Acacia sp. Ripon Hills (B.R. Maslin	1
Acacia sphaerostachya	11
Acacia spondylophylla	1
Acacia stellaticeps	47
Acacia synchronicia	4
Acacia trachycarpa	22
Acacia translucens	1
Acacia trudgeniana	3

<i>Acacia tumida</i>	1
<i>Acacia tumida</i> var. <i>kulpari</i>	6
<i>Acacia tumida</i> var. <i>pilbarensis</i>	16
<i>Acacia tumida</i> var. <i>tumida</i>	1
<i>Achyranthes aspera</i>	5
<i>Adriana tomentosa</i>	1
<i>Adriana tomentosa</i> var. <i>tomentosa</i>	7
<i>Aerva javanica</i>	20
<i>Aeschynomene indica</i>	7
<i>Albizia lebbek</i>	1
<i>Alstonia linearis</i>	1
<i>Alstonia spectabilis</i>	1
<i>Alternanthera angustifolia</i>	5
<i>Alternanthera denticulata</i>	2
<i>Alternanthera nana</i>	5
<i>Alternanthera nodiflora</i>	1
<i>Alysicarpus muelleri</i>	8
<i>Amaranthus pallidiflorus</i>	6
<i>Amaranthus undulatus</i>	4
<i>Ammannia baccifera</i>	5
<i>Ammannia muelleri</i>	2
<i>Ammannia multiflora</i>	4
<i>Amyema preissii</i>	2
<i>Amyema sanguinea</i> var. <i>pulchra</i>	2
<i>Argemone ochroleuca</i>	1
<i>Argemone ochroleuca</i> subsp. <i>ochroleuca</i>	3
<i>Atalaya hemiglauc</i>	3
<i>Atriplex semilunaris</i>	1
<i>Avicennia marina</i>	6
<i>Avicennia marina</i> subsp. <i>marina</i>	1
<i>Basilicum polystachyon</i>	4
<i>Bauhinia cunninghamii</i>	9
<i>Bergia ammannioides</i>	2
<i>Bergia henshallii</i>	1
<i>Bergia pedicellaris</i>	3
<i>Bergia perennis</i>	2
<i>Bergia perennis</i> subsp. <i>obtusifolia</i>	1
<i>Bergia perennis</i> subsp. <i>perennis</i>	1
<i>Bergia trimera</i>	5
<i>Blumea tenella</i>	6
<i>Boerhavia burbridgeana</i>	1
<i>Boerhavia coccinea</i>	9
<i>Boerhavia gardneri</i>	1
<i>Boerhavia paludosa</i>	1
<i>Boerhavia repleta</i>	1
<i>Bonamia alatisemina</i>	13
<i>Bonamia erecta</i>	12
<i>Bonamia linearis</i>	5
<i>Bonamia media</i>	7
<i>Bonamia pannosa</i>	4
<i>Bonamia pilbarensis</i>	1
<i>Bonamia</i> sp.	1
<i>Bougainvillea glabra</i>	1
<i>Bruguiera exaristata</i>	4
<i>Buchnera linearis</i>	2
<i>Byblis filifolia</i>	3
<i>Byblis liniflora</i>	2
<i>Byblis</i> sp.	1
<i>Cajanus cinereus</i>	7

<i>Cajanus marmoratus</i>	8
<i>Cajanus pubescens</i>	3
<i>Calandrinia pentavalvis</i>	14
<i>Calandrinia ptychosperma</i>	1
<i>Calandrinia pumila</i>	4
<i>Calandrinia</i> sp. Pinga (T.R. Lally Tf	2
<i>Calandrinia stagnensis</i>	6
<i>Calandrinia strophiolata</i>	2
<i>Calandrinia tepperiana</i>	6
<i>Calotis hispidula</i>	1
<i>Calotis plumulifera</i>	2
<i>Calytrix carinata</i>	4
<i>Canavalia rosea</i>	12
<i>Capparis spinosa</i> subsp. nummulari	2
<i>Carissa lanceolata</i>	8
<i>Cassytha capillaris</i>	12
<i>Cassytha filiformis</i>	3
<i>Centaurium clementii</i>	1
<i>Centipeda minima</i>	4
<i>Centipeda minima</i> subsp. macroce	4
<i>Centipeda minima</i> subsp. minima	1
<i>Ceriops australis</i>	3
<i>Ceriops tagal</i>	1
<i>Chenopodium auricomum</i>	1
<i>Chrysocephalum apiculatum</i> subsp	2
<i>Citrullus amarus</i>	3
<i>Cleome oxalidea</i>	1
<i>Cleome uncifera</i>	9
<i>Cleome uncifera</i> subsp. uncifera	8
<i>Cleome viscosa</i>	17
<i>Clerodendrum tomentosum</i>	1
<i>Clerodendrum tomentosum</i> var. lar	1
<i>Clerodendrum tomentosum</i> var. mc	1
<i>Clerodendrum tomentosum</i> var. tor	1
<i>Clitoria ternatea</i>	1
<i>Coccinia grandis</i>	1
<i>Codonocarpus cotinifolius</i>	8
<i>Conyza bonariensis</i>	1
<i>Conyza parva</i>	1
<i>Corchorus carnarvonensis</i>	1
<i>Corchorus elachocarpus</i>	11
<i>Corchorus incanus</i>	9
<i>Corchorus incanus</i> subsp. incanus	9
<i>Corchorus laniflorus</i>	2
<i>Corchorus lasiocarpus</i>	1
<i>Corchorus parviflorus</i>	3
<i>Corchorus sidoides</i>	2
<i>Corchorus sidoides</i> subsp. sidoides	1
<i>Corchorus sidoides</i> subsp. vermicu	3
<i>Corchorus tectus</i>	2
<i>Corchorus tridens</i>	11
<i>Corchorus trilocularis</i>	1
<i>Corchorus walcottii</i>	4
<i>Corymbia aspera</i>	4
<i>Corymbia candida</i>	1
<i>Corymbia candida</i> / <i>flavescens</i>	1
<i>Corymbia candida</i> subsp. <i>laetifolia</i>	6
<i>Corymbia deserticola</i> subsp. <i>deser</i>	1
<i>Corymbia flavescens</i>	24

<i>Corymbia hamersleyana</i>	14
<i>Corymbia opaca</i>	3
<i>Corymbia zygophylla</i>	15
<i>Crotalaria crispata</i>	1
<i>Crotalaria cunninghamii</i>	11
<i>Crotalaria dissitiflora</i> subsp. <i>benthamii</i>	1
<i>Crotalaria medicaginea</i> var. <i>neglecta</i>	10
<i>Crotalaria ramosissima</i>	7
<i>Crotalaria spectabilis</i> subsp. <i>spectabilis</i>	1
<i>Cucumis argenteus</i>	2
<i>Cucumis maderaspatanus</i>	7
<i>Cucumis melo</i>	4
<i>Cucumis melo</i> subsp. <i>agrestis</i>	2
<i>Cucumis</i> sp.	1
<i>Cucumis variabilis</i>	7
<i>Cucurbita pepo</i>	1
<i>Cullen cinereum</i>	2
<i>Cullen lachnostachys</i>	6
<i>Cullen leucanthum</i>	6
<i>Cullen leucochaetes</i>	1
<i>Cullen martinii</i>	6
<i>Cullen pustulatum</i>	1
<i>Cullen stipulaceum</i>	6
<i>Cyanostegia cyanocalyx</i>	4
<i>Cyanthillium cinereum</i> var. <i>cinereum</i>	1
<i>Cynanchum floribundum</i>	1
<i>Dampiera candidans</i>	7
<i>Datura metel</i>	2
<i>Dentella asperata</i>	4
<i>Dentella minutissima</i>	1
<i>Desmodium filiforme</i>	14
<i>Desmodium muelleri</i>	1
<i>Desmodium scorpiurus</i>	1
<i>Desmodium</i> sp.	1
<i>Dichrostachys spicata</i>	1
<i>Dicrastylis cordifolia</i>	2
<i>Diplopeltis eriocarpa</i>	2
<i>Dissocarpus paradoxus</i>	1
<i>Distimake davenportii</i>	3
<i>Distimake dissectus</i> var. <i>dissectus</i>	2
<i>Dodonaea coriacea</i>	9
<i>Dodonaea hispidula</i>	1
<i>Dolichandrone heterophylla</i>	1
<i>Dolichandrone occidentalis</i>	7
<i>Drosera burmanni</i>	1
<i>Drosera finlaysonian</i>	2
<i>Drosera indica</i>	5
<i>Duboisia hopwoodii</i>	3
<i>Dysphania plantaginella</i>	6
<i>Dysphania rhadinostachya</i>	1
<i>Dysphania rhadinostachya</i> subsp. <i>indica</i>	8
<i>Dysphania sphaerosperma</i>	1
<i>Ehretia saligna</i>	3
<i>Elacholoma hornii</i>	1
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	2
<i>Erythrina vespertilio</i>	1
<i>Erythrophleum chlorostachys</i>	3
<i>Eucalyptus camaldulensis</i> subsp. <i>camaldulensis</i>	3
<i>Eucalyptus camaldulensis</i> subsp. <i>camaldulensis</i>	10

<i>Eucalyptus leucophloia</i> subsp. <i>leuc</i>	3
<i>Eucalyptus odontocarpa</i>	7
<i>Eucalyptus victrix</i>	9
<i>Euphorbia alsiniflora</i>	1
<i>Euphorbia australis</i>	2
<i>Euphorbia australis</i> var. <i>australis</i>	4
<i>Euphorbia australis</i> var. <i>subtoment</i>	5
<i>Euphorbia biconvexa</i>	1
<i>Euphorbia careyi</i>	1
<i>Euphorbia coghlanii</i>	6
<i>Euphorbia drummondii</i>	1
<i>Euphorbia fitzroyensis</i>	1
<i>Euphorbia myrtoides</i>	6
<i>Euphorbia psilosperma</i>	2
<i>Euphorbia</i> sp.	1
<i>Euphorbia tannensis</i> subsp. <i>eremo</i>	6
<i>Euphorbia tirucalli</i>	1
<i>Euphorbia trigonosperma</i>	6
<i>Euphorbia vaccaria</i> var. <i>vaccaria</i>	5
<i>Euphorbia wheeleri</i>	1
<i>Evolvulus alsinoides</i>	1
<i>Evolvulus alsinoides</i> var. <i>decumben</i>	12
<i>Evolvulus alsinoides</i> var. <i>villosicaly</i>	10
<i>Ficus aculeata</i>	2
<i>Ficus aculeata</i> var. <i>indecora</i>	3
<i>Ficus brachypoda</i>	5
<i>Ficus cerasicarpa</i>	1
<i>Ficus opposita</i>	1
<i>Ficus virens</i>	1
<i>Flaveria trinervia</i>	5
<i>Flueggea virosa</i>	1
<i>Flueggea virosa</i> subsp. <i>melanthesc</i>	2
<i>Frankenia ambita</i>	7
<i>Frankenia cordata</i>	1
<i>Gardenia pyriformis</i> subsp. <i>keartlar</i>	6
<i>Gardenia pyriformis</i> subsp. <i>pyriforn</i>	1
Genus D sp.2	1
<i>Glinus lotoides</i>	4
<i>Glinus oppositifolius</i>	2
<i>Glossostigma diandrum</i>	3
<i>Glycine canescens</i>	1
<i>Glycine</i> sp.	1
<i>Glycine tomentella</i>	3
<i>Gompholobium simplicifolium</i>	7
<i>Gomphrena affinis</i> subsp. <i>pilbarens</i>	13
<i>Gomphrena canescens</i> subsp. <i>can</i>	6
<i>Gomphrena celosioides</i>	1
<i>Gomphrena cunninghamii</i>	3
<i>Gomphrena leptoclada</i>	3
<i>Gomphrena leptoclada</i> subsp. <i>leptc</i>	6
<i>Gomphrena sordida</i>	2
<i>Gonocarpus ephemerus</i>	1
<i>Goodenia armitiana</i>	2
<i>Goodenia azurea</i> subsp. <i>hesperia</i>	4
<i>Goodenia forrestii</i>	6
<i>Goodenia lamprosperma</i>	13
<i>Goodenia microptera</i>	15
<i>Goodenia muelleriana</i>	4
<i>Goodenia scaevolina</i>	2

<i>Goodenia</i> sp.	1
<i>Goodenia stobbsiana</i>	5
<i>Gossypium australe</i>	4
<i>Gossypium hirsutum</i>	2
<i>Gossypium robinsonii</i>	1
<i>Grevillea eriostachya</i>	1
<i>Grevillea pyramidalis</i>	6
<i>Grevillea pyramidalis</i> subsp. <i>leucac</i>	3
<i>Grevillea refracta</i> subsp. <i>refracta</i>	10
<i>Grevillea wickhamii</i>	2
<i>Grevillea wickhamii</i> subsp. <i>aprica</i>	2
<i>Grevillea wickhamii</i> subsp. <i>hispidul</i>	7
<i>Grevillea wickhamii</i> subsp. <i>macrod</i>	2
<i>Gyrocarpus americanus</i>	3
<i>Gyrostemon tepperi</i>	3
<i>Hakea chordophylla</i>	1
<i>Hakea lorea</i>	3
<i>Hakea lorea</i> subsp. <i>lorea</i>	4
<i>Hakea macrocarpa</i>	7
<i>Hakea stenophylla</i>	1
<i>Halgania gustafsenii</i>	1
<i>Halgania solanacea</i>	1
<i>Halgania solanacea</i> var. <i>Mt Doreen</i>	1
<i>Halgania solanacea</i> var. <i>solanacea</i>	6
<i>Haloragis gossei</i>	2
<i>Helichrysum luteoalbum</i>	2
<i>Heliotropium ammophilum</i>	2
<i>Heliotropium chrysocarpum</i>	1
<i>Heliotropium conocarpum</i>	1
<i>Heliotropium crispatum</i>	5
<i>Heliotropium cunninghamii</i>	4
<i>Heliotropium curassavicum</i>	1
<i>Heliotropium europaeum</i>	2
<i>Heliotropium foliatum</i>	3
<i>Heliotropium leptaleum</i>	1
<i>Heliotropium ovalifolium</i>	1
<i>Heliotropium pachyphyllum</i>	5
<i>Heliotropium</i> sp.	2
<i>Heliotropium transforme</i>	3
<i>Heliotropium vestitum</i>	3
<i>Hemichroa diandra</i>	2
<i>Hibiscus apodus</i>	4
<i>Hibiscus austrinus</i> var. <i>austrinus</i>	6
<i>Hibiscus brachychlaenus</i>	2
<i>Hibiscus burtonii</i>	2
<i>Hibiscus goldsworthii</i>	1
<i>Hibiscus leptocladus</i>	5
<i>Hibiscus sturtii</i> var. <i>campylochlamy</i>	7
<i>Hibiscus sturtii</i> var. <i>platychlamys</i>	2
<i>Hibiscus verdcourtii</i>	1
<i>Hybanthus aurantiacus</i>	14
<i>Hybanthus enneaspermus</i> subsp. <i>ε</i>	1
<i>Hypertelis cerviana</i>	1
<i>Indigastrium parviflorum</i>	2
<i>Indigofera boviparda</i>	1
<i>Indigofera boviparda</i> subsp. <i>bovi</i> pe	1
<i>Indigofera colutea</i>	11
<i>Indigofera hirsuta</i>	5
<i>Indigofera hochstetteri</i>	3

<i>Indigofera linifolia</i>	13
<i>Indigofera linnaei</i>	6
<i>Indigofera monophylla</i>	7
<i>Indigofera oblongifolia</i>	11
<i>Indigofera rugosa</i>	2
<i>Indigofera sessiliflora</i>	3
<i>Indigofera trita</i>	8
<i>Ipomoea coptica</i>	5
<i>Ipomoea costata</i>	1
<i>Ipomoea diamantinensis</i>	1
<i>Ipomoea muelleri</i>	12
<i>Ipomoea pes-caprae</i>	1
<i>Ipomoea pes-caprae</i> subsp. <i>brasiliensis</i>	3
<i>Ipomoea polymorpha</i>	3
<i>Isotropis atropurpurea</i>	6
<i>Jacksonia aculeata</i>	12
<i>Jacquemontia</i> sp.	1
<i>Jasminum calcareum</i>	2
<i>Jasminum didymum</i> subsp. <i>lineare</i>	1
<i>Jatropha gossypifolia</i>	1
<i>Josephinia eugeniae</i>	1
<i>Josephinia</i> sp.	1
<i>Lepidium muelleri-ferdinandii</i>	1
<i>Leptopus decaisnei</i>	1
<i>Leptosema anomalum</i>	9
<i>Leucaena leucocephala</i>	1
<i>Lobelia arnhemiaca</i>	1
<i>Lotus cruentus</i>	5
<i>Ludwigia perennis</i>	3
<i>Lysiana spathulata</i> subsp. <i>parvifolia</i>	3
<i>Maireana tomentosa</i> subsp. <i>tomentosa</i>	3
<i>Maireana villosa</i>	2
<i>Melaleuca alsophila</i>	2
<i>Melaleuca argentea</i>	6
<i>Melaleuca glomerata</i>	5
<i>Melaleuca lasiandra</i>	7
<i>Melaleuca linophylla</i>	1
<i>Melhania oblongifolia</i>	4
<i>Melochia pyramidata</i>	2
<i>Microstachys chamaelea</i>	1
<i>Mimulus gracilis</i>	4
<i>Minuria</i> sp.	2
<i>Mirbelia viminalis</i>	2
<i>Mitrasacme connata</i>	3
<i>Mitrasacme exserta</i>	3
<i>Mollugo molluginea</i>	1
<i>Moringa oleifera</i>	1
<i>Muellerolimon salicorniaceum</i>	6
<i>Myoporum montanum</i>	4
<i>Neobassia astrocarpa</i>	13
<i>Neptunia dimorphantha</i>	7
<i>Neptunia monosperma</i>	1
<i>Newcastelia cladotricha</i>	6
<i>Nicotiana benthamiana</i>	4
<i>Nicotiana heterantha</i>	1
<i>Nicotiana occidentalis</i> subsp. <i>obliqua</i>	1
<i>Nicotiana occidentalis</i> subsp. <i>occidentalis</i>	1
<i>Nicotiana rosulata</i> subsp. <i>rosulata</i>	1
<i>Notoleptopus decaisnei</i>	4

<i>Nymphoides indica</i>	4
<i>Oldenlandia crouchiana</i>	3
<i>Oldenlandia galioides</i>	3
<i>Oldenlandia pterospora</i>	2
<i>Operculina aequisejala</i>	4
<i>Osbornia octodonta</i>	1
<i>Owenia reticulata</i>	5
<i>Parkinsonia aculeata</i>	1
<i>Passiflora foetida</i> var. <i>hispida</i>	2
<i>Peplidium aithocheilum</i>	1
<i>Peplidium muelleri</i>	4
<i>Peripleura virgata</i>	2
<i>Petalostylis labicheoides</i>	4
<i>Phyla nodiflora</i> var. <i>nodiflora</i>	1
<i>Phyllanthus aridus</i>	1
<i>Phyllanthus eremicus</i>	2
<i>Phyllanthus erwinii</i>	1
<i>Phyllanthus exilis</i>	1
<i>Phyllanthus maderaspatensis</i>	8
<i>Phyllanthus reticulatus</i>	1
<i>Phyllanthus</i> sp.	1
<i>Phyllanthus virgatus</i>	2
<i>Physalis angulata</i>	2
<i>Pimelea ammocharis</i>	2
<i>Pittosporum angustifolium</i>	1
<i>Pluchea dentex</i>	2
<i>Pluchea ferdinandi-muelleri</i>	4
<i>Pluchea rubelliflora</i>	11
<i>Pluchea tetranthera</i>	15
<i>Polycarpaea corymbosa</i>	3
<i>Polycarpaea corymbosa</i> var. <i>corymbosa</i>	12
<i>Polycarpaea holtzei</i>	2
<i>Polycarpaea involucrata</i>	1
<i>Polycarpaea longiflora</i>	4
<i>Polygala galeocephala</i>	8
<i>Polygala glaucifolia</i>	3
<i>Polygala saccopectala</i>	1
<i>Polygala</i> sp.	1
<i>Polymeria ambigua</i>	2
<i>Polymeria lanata</i>	4
<i>Polymeria</i> sp.	2
<i>Portulaca australis</i>	1
<i>Portulaca conspicua</i>	1
<i>Portulaca cyclophylla</i>	2
<i>Portulaca decipiens</i>	2
<i>Portulaca digyna</i>	1
<i>Portulaca oleracea</i>	21
<i>Portulaca pilosa</i>	8
<i>Portulaca</i> sp.	2
<i>Prosopis pallida</i>	1
<i>Pseudognaphalium luteoalbum</i>	2
<i>Pterocaulon intermedium</i>	6
<i>Pterocaulon serrulatum</i>	1
<i>Pterocaulon sphacelatum</i>	8
<i>Ptilotus appendiculatus</i>	1
<i>Ptilotus arthrolasius</i>	8
<i>Ptilotus astrolasius</i>	14
<i>Ptilotus auriculifolius</i>	1
<i>Ptilotus axillaris</i>	14

<i>Ptilotus calostachyus</i>	9
<i>Ptilotus divaricatus</i>	2
<i>Ptilotus exaltatus</i>	5
<i>Ptilotus fusiformis</i>	13
<i>Ptilotus gomphrenoides</i>	3
<i>Ptilotus helipteroides</i>	2
<i>Ptilotus incanus</i>	7
<i>Ptilotus lanatus</i>	1
<i>Ptilotus macrocephalus</i>	1
<i>Ptilotus murrayi</i>	5
<i>Ptilotus nobilis</i>	1
<i>Ptilotus obovatus</i>	4
<i>Ptilotus polystachyus</i>	4
<i>Ptilotus polystachyus</i> var. <i>arthrotrich</i>	1
<i>Ptilotus villosiflorus</i>	6
<i>Pupalia lappacea</i>	1
<i>Rhagodia eremaea</i>	6
<i>Rhizophora stylosa</i>	8
<i>Rhynchosia minima</i>	14
<i>Ricinus communis</i>	3
<i>Roepera compressa</i>	2
<i>Rostellularia adscendens</i> var. <i>clem</i>	3
<i>Rotala diandra</i>	9
<i>Salsola australis</i>	2
<i>Samolus repens</i>	2
<i>Sauropus</i> sp.	1
<i>Scaevola amblyanthera</i>	1
<i>Scaevola amblyanthera</i> var. <i>centra</i>	8
<i>Scaevola browniana</i>	2
<i>Scaevola browniana</i> subsp. <i>brownii</i>	1
<i>Scaevola crassifolia</i>	1
<i>Scaevola spinescens</i>	1
<i>Schenkia clementii</i>	3
<i>Sclerolaena bicornis</i> var. <i>bicornis</i>	5
<i>Sclerolaena costata</i>	1
<i>Sclerolaena densiflora</i>	1
<i>Sclerolaena glabra</i>	1
<i>Sclerolaena hostilis</i>	3
<i>Sclerolaena uniflora</i>	1
<i>Senna artemisioides</i> subsp. <i>helmsi</i>	1
<i>Senna artemisioides</i> subsp. <i>helmsi</i>	1
<i>Senna artemisioides</i> subsp. <i>oligoph</i>	15
<i>Senna bicapsularis</i>	1
<i>Senna costata</i>	1
<i>Senna curvistyla</i>	6
<i>Senna glutinosa</i>	2
<i>Senna glutinosa</i> subsp. <i>glutinosa</i>	3
<i>Senna notabilis</i>	13
<i>Senna occidentalis</i>	2
<i>Senna stricta</i>	2
<i>Senna symonii</i>	1
<i>Senna venusta</i>	3
<i>Seringia elliptica</i>	6
<i>Seringia nephrosperma</i>	5
<i>Sesbania cannabina</i>	13
<i>Sesbania formosa</i>	6
<i>Sesuvium portulacastrum</i>	1
<i>Sida arenicola</i>	4
<i>Sida cardiophylla</i>	2

<i>Sida clementii</i>	3
<i>Sida echinocarpa</i>	1
<i>Sida fibulifera</i>	3
<i>Sida macropoda</i>	10
<i>Sida rohlena</i>	8
<i>Sida rohlena</i> subsp. <i>rohlena</i>	5
<i>Sida</i> sp.	3
<i>Sida</i> sp. Articulation below (A.A. Mi	5
<i>Sida</i> sp. Pilbara (A.A. Mitchell PRP	12
<i>Sida</i> sp. Pindan (B.G. Thomson 33	5
<i>Sida</i> sp. Rabbit Flat (B.J. Carter 62	3
<i>Sida spinosa</i>	1
<i>Solanum chippendalei</i>	1
<i>Solanum cleistogamum</i>	2
<i>Solanum dioicum</i>	5
<i>Solanum dioicum</i> sens. lat.	1
<i>Solanum diversiflorum</i>	11
<i>Solanum esuriale</i>	6
<i>Solanum horridum</i>	3
<i>Solanum lasiophyllum</i>	5
<i>Solanum lucani</i>	3
<i>Solanum nigrum</i>	2
<i>Solanum phlomoides</i>	10
<i>Spermacoce hillii</i>	2
<i>Spermacoce</i> sp.	1
<i>Stackhousia intermedia</i>	1
<i>Stemodia grossa</i>	9
<i>Stemodia kingii</i>	2
<i>Stemodia lathraia</i>	7
<i>Stemodia</i> sp.	5
<i>Stemodia</i> sp. Shay Gap (B. Cook 7	6
<i>Stemodia viscosa</i>	5
<i>Stenopetalum decipiens</i>	1
<i>Streptoglossa bubakii</i>	1
<i>Streptoglossa cylindriceps</i>	1
<i>Streptoglossa decurrens</i>	8
<i>Streptoglossa macrocephala</i>	2
<i>Streptoglossa odora</i>	5
<i>Streptoglossa</i> sp.	1
<i>Streptoglossa tenuiflora</i>	5
<i>Striga curviflora</i>	1
<i>Striga</i> sp.	1
<i>Striga squamigera</i>	2
<i>Stylidium desertorum</i>	3
<i>Stylobasium spathulatum</i>	3
<i>Stylosanthes guianensis</i> var. <i>guian</i>	3
<i>Stylosanthes hamata</i>	6
<i>Suaeda arbusculoides</i>	2
<i>Surreya diandra</i>	2
<i>Swainsona formosa</i>	2
<i>Swainsona laciniata</i>	1
<i>Swainsona pterostylis</i>	9
<i>Swainsona tanamiensis</i>	1
<i>Symphyotrichum squamatum</i>	1
<i>Synaptantha tillaeacea</i> var. <i>tillaeac</i>	4
<i>Tecticornia auriculata</i>	21
<i>Tecticornia halocnemoides</i>	10
<i>Tecticornia halocnemoides</i> subsp. '	1
<i>Tecticornia indica</i> subsp. <i>bidens</i>	1

<i>Tecticornia indica</i> subsp. <i>indica</i>	1
<i>Tecticornia indica</i> subsp. <i>leiostachy</i>	9
<i>Tecticornia pterygosperma</i> subsp. <i>i</i>	3
<i>Templetonia hookeri</i>	2
<i>Tephrosia clementii</i>	2
<i>Tephrosia flammea</i>	1
<i>Tephrosia leptoclada</i>	13
<i>Tephrosia rosea</i>	6
<i>Tephrosia rosea</i> var. <i>clementii</i>	5
<i>Tephrosia rosea</i> var. <i>clementii</i> / <i>ros</i>	1
<i>Tephrosia rosea</i> var. <i>Fortescue cre</i>	1
<i>Tephrosia rosea</i> var. <i>rosea</i>	6
<i>Tephrosia rosea</i> var. <i>venulosa</i>	4
<i>Tephrosia simplicifolia</i>	2
<i>Tephrosia</i> sp. B Kimberley Flora (C	11
<i>Tephrosia</i> sp. Bungaroo Creek (M.I	16
<i>Tephrosia</i> sp. Carnarvon (J.H. Ros	1
<i>Tephrosia</i> sp. clay soils (S. van Lee	1
<i>Tephrosia</i> sp. D Kimberley Flora (R	21
<i>Tephrosia</i> sp. Fortescue (A.A. Mitc	3
<i>Tephrosia</i> sp. NW Eremaean (S. v:	7
<i>Tephrosia supina</i>	4
<i>Tephrosia virens</i>	6
<i>Terminalia canescens</i>	6
<i>Terminalia circumalata</i>	1
<i>Threlkeldia diffusa</i>	4
<i>Tinospora smilacina</i>	5
<i>Trachymene oleracea</i> subsp. <i>olera</i>	6
<i>Trachymene pilbarensis</i>	1
<i>Trianthema cusackianum</i>	4
<i>Trianthema oxycalyptum</i> var. <i>oxyc</i>	1
<i>Trianthema pilosum</i>	14
<i>Trianthema portulacastrum</i>	7
<i>Trianthema triquetra</i>	1
<i>Trianthema triquetrum</i>	14
<i>Trianthema turgidifolia</i>	1
<i>Trianthema turgidifolium</i>	9
<i>Tribulopsis angustifolia</i>	8
<i>Tribulus cistoides</i>	1
<i>Tribulus hirsutus</i>	2
<i>Tribulus occidentalis</i>	6
<i>Tribulus platypterus</i>	1
<i>Tribulus</i> sp.	1
<i>Tribulus</i> sp. long-styled eichlerianu:	1
<i>Trichodesma zeylanicum</i>	3
<i>Trichosanthes cucumerina</i> var. <i>cuc</i>	2
<i>Tridax procumbens</i>	1
<i>Trigastrotheca molluginea</i>	31
<i>Trigonella suavissima</i>	3
<i>Triumfetta appendiculata</i>	4
<i>Triumfetta chaetocarpa</i>	3
<i>Triumfetta clementii</i>	5
<i>Triumfetta deserticola</i>	2
<i>Triumfetta incana</i>	1
<i>Triumfetta johnstonii</i>	3
<i>Triumfetta maconochieana</i>	3
<i>Triumfetta propinqua</i>	2
<i>Triumfetta ramosa</i>	11
<i>Uvedalia linearis</i> var. <i>linearis</i>	2

Velleia panduriformis	2
Vigna lanceolata	3
Vigna lanceolata var. lanceolata	3
Vigna sp.	1
Vigna sp. Hamersley Clay (A.A. Mii	1
Wahlenbergia queenslandica	1
Wahlenbergia tumidifructa	5
Waltheria indica	7
Xanthium occidentale	1
Zaleya galericulata subsp. galericul	2
Zornia albiflora	1
Zornia chaetophora	3
Zornia muelleriana	1
Zornia muelleriana subsp. congesta	3
FERN	18
Cheilanthes brownii	1
Cheilanthes sieberi subsp. sieberi	1
Marsilea drummondii	1
Marsilea exarata	3
Marsilea hirsuta	4
Marsilea sp.	3
Marsilea spp.	5
LIVERWORT	2
Riccia crystallina	1
Riccia singularis	1
MONOCOT	794
Agave americana	1
Amphipogon caricinus var. caricinu	1
Amphipogon sericeus	4
Andropogon gayanus	1
Aristida contorta	11
Aristida holathera	12
Aristida holathera var. holathera	3
Aristida hygrometrica	6
Aristida inaequiglumis	6
Aristida latifolia	3
Astrebla pectinata	2
Bothriochloa ewartiana	1
Brachyachne convergens	1
Bulbostylis barbata	24
Bulbostylis turbinata	2
Cenchrus ciliaris	19
Cenchrus setaceus	1
Cenchrus setiger	6
Centrolepis banksii	1
Chloris barbata	5
Chloris pectinata	1
Chloris pumilio	8
Chloris virgata	1
Chrysopogon fallax	19
Commelina ensifolia	2
Corynotheca micrantha	2
Corynotheca micrantha var. micran	2
Corynotheca pungens	7
Corynotheca sp.	2
Cymbopogon ambiguus	6
Cymbopogon bombycinus	1
Cymbopogon oblectus	5
Cynodon convergens	2

<i>Cynodon dactylon</i>	6
<i>Cynodon radiatus</i>	1
<i>Cyperus bifax</i>	2
<i>Cyperus blakeanus</i>	4
<i>Cyperus bulbosus</i>	5
<i>Cyperus castaneus</i> var. <i>brevimucro</i>	2
<i>Cyperus concinnus</i>	1
<i>Cyperus conicus</i>	5
<i>Cyperus cunninghamii</i>	1
<i>Cyperus cunninghamii</i> subsp. <i>cunn</i>	1
<i>Cyperus difformis</i>	3
<i>Cyperus gymnocaulos</i> / <i>vaginatus</i>	1
<i>Cyperus hesperius</i>	1
<i>Cyperus iria</i>	8
<i>Cyperus ixiocarpus</i>	4
<i>Cyperus macrostachyos</i>	7
<i>Cyperus microcephalus</i> subsp. <i>mic</i>	1
<i>Cyperus microcephalus</i> subsp. <i>sax</i>	1
<i>Cyperus polystachyos</i>	1
<i>Cyperus pulchellus</i>	4
<i>Cyperus pygmaeus</i>	1
<i>Cyperus rigidellus</i>	1
<i>Cyperus squarrosus</i>	8
<i>Cyperus vaginatus</i>	3
<i>Dactyloctenium aegyptium</i>	1
<i>Dactyloctenium radulans</i>	8
<i>Dichanthium fecundum</i>	3
<i>Dichanthium sericeum</i> subsp. <i>humi</i>	2
<i>Dichanthium sericeum</i> subsp. <i>polys</i>	1
<i>Digitaria brownii</i>	3
<i>Digitaria ciliaris</i>	2
<i>Digitaria ctenantha</i>	2
<i>Diplachne fusca</i>	1
<i>Diplachne fusca</i> subsp. <i>fusca</i>	4
<i>Echinochloa colona</i>	7
<i>Eleocharis atropurpurea</i>	4
<i>Elytrophorus spicatus</i>	8
<i>Enneapogon caeruleus</i>	5
<i>Enneapogon lindleyanus</i>	2
<i>Enneapogon pallidus</i>	2
<i>Enneapogon polyphyllus</i>	4
<i>Enneapogon purpurascens</i>	5
<i>Enneapogon robustissimus</i>	3
<i>Enteropogon ramosus</i>	5
<i>Eragrostis cumingii</i>	25
<i>Eragrostis dielsii</i>	5
<i>Eragrostis elongata</i>	1
<i>Eragrostis eriopoda</i>	12
<i>Eragrostis falcata</i>	5
<i>Eragrostis olida</i>	1
<i>Eragrostis pergracilis</i>	1
<i>Eragrostis pilosa</i>	1
<i>Eragrostis speciosa</i>	1
<i>Eragrostis tenellula</i>	11
<i>Eragrostis xerophila</i>	9
<i>Eriachne aristidea</i>	16
<i>Eriachne benthamii</i>	9
<i>Eriachne ciliata</i>	4
<i>Eriachne flaccida</i>	7

<i>Eriachne glauca</i>	2
<i>Eriachne glauca</i> var. <i>glauca</i>	11
<i>Eriachne helmsii</i>	1
<i>Eriachne lanata</i>	8
<i>Eriachne melicacea</i>	2
<i>Eriachne mucronata</i>	3
<i>Eriachne obtusa</i>	11
<i>Eriachne pulchella</i>	6
<i>Eriachne</i> sp.	4
<i>Eriachne sulcata</i>	1
<i>Eriocaulon cinereum</i>	2
<i>Eriochloa procera</i>	3
<i>Eriochloa pseudoacrotricha</i>	2
<i>Eulalia aurea</i>	11
<i>Fimbristylis ammobia</i>	1
<i>Fimbristylis caespitosa</i>	1
<i>Fimbristylis dichotoma</i>	9
<i>Fimbristylis elegans</i>	1
<i>Fimbristylis littoralis</i>	9
<i>Fimbristylis microcarya</i>	5
<i>Fimbristylis neilsonii</i>	5
<i>Fimbristylis nuda</i>	1
<i>Fimbristylis oxystachya</i>	5
<i>Fimbristylis rara</i>	3
<i>Fimbristylis simulans</i>	4
<i>Fuirena ciliaris</i>	1
<i>Halodule uninervis</i>	2
<i>Iseilema dolichotrichum</i>	2
<i>Iseilema eremaeum</i>	1
<i>Iseilema membranaceum</i>	4
<i>Iseilema vaginiflorum</i>	1
<i>Lamarckia aurea</i>	2
<i>Leptochloa digitata</i>	3
<i>Leptochloa fusca</i> subsp. <i>fusca</i>	1
<i>Leptochloa</i> sp.	1
<i>Lipocarpa microcephala</i>	7
<i>Murdannia graminea</i>	4
<i>Panicum decompositum</i>	5
<i>Paraneurachne muelleri</i>	2
<i>Paspalidium clementii</i>	3
<i>Paspalidium rarum</i>	5
<i>Paspalidium tabulatum</i>	6
<i>Paspalum fasciculatum</i>	1
<i>Perotis rara</i>	6
<i>Potamogeton tricarinatus</i>	1
<i>Schizachyrium fragile</i>	3
<i>Schoenoplectiella dissachantha</i>	1
<i>Schoenoplectiella laevis</i>	2
<i>Schoenoplectiella lateriflora</i>	1
<i>Schoenoplectus dissachanthus</i>	4
<i>Schoenoplectus laevis</i>	3
<i>Schoenoplectus lateriflorus</i>	1
<i>Setaria dielsii</i>	3
<i>Setaria italica</i>	1
<i>Setaria sphacelata</i>	1
<i>Setaria surgens</i>	3
<i>Setaria verticillata</i>	1
<i>Sorghum interjectum</i>	2
<i>Sorghum plumosum</i>	8

Sorghum stipoideum	1
Spinifex longifolius	8
Sporobolus actinocladus	1
Sporobolus australasicus	9
Sporobolus mitchellii	4
Sporobolus virginicus	3
Stuckenia pectinata	4
Thalassia hemprichii	1
Themeda avenacea	1
Themeda triandra	4
Triodia biflora	2
Triodia bitextura	1
Triodia epactia	51
Triodia lanigera	4
Triodia longiceps	2
Triodia schinzii	8
Triodia secunda	1
Triodia sp.	1
Triodia wiseana	5
Tripogon loliiformis	1
Triraphis mollis	4
Typha domingensis	1
Urochloa holosericea	1
Urochloa holosericea subsp. velutir	3
Urochloa occidentalis var. occident	3
Urochloa panicoides	1
Urochloa piligera	5
Vallisneria nana	4
Whiteochloa cymbiformis	14
Xerochloa barbata	5
Xerochloa imberbis	1
Yakirra australiensis	16
Yakirra majuscula	1
Protozoa	19
ALGA	19
Colpomenia sinuosa	2
Dictyopteris australis	5
Dictyota ciliolata	1
Hormophysa cuneiformis	2
Hydroclathrus clathratus	1
Lobophora variegata	2
Spatoglossum macrodontum	5
Stoechospermum polypodioides	1
(blank)	
(blank)	
(blank)	
Grand Total	25277



ghd.com

→ **The Power of Commitment**