

**Forrestania Nickel Operations
Lounge Lizard
Vegetation, Flora and Fauna Biological Assessment**

October 2014

Prepared for
Western Areas Limited



Astron Environmental Services

129 Royal Street

East Perth WA 6004

Phone: (08) 9421 9600

Fax: (08) 9421 9699

Email: perth@astron.com.au

Report Reference: 17603-14-BISR-1Rev0_141203

Forrestania Nickel Operations Lounge Lizard – Vegetation, Flora and Fauna Biological Assessment

Prepared for
Western Areas Limited

Job Number: 17603-14

Reference: 17603-14-BISR-1Rev0_141203

Revision Status

Rev	Date	Description	Author(s)	Reviewer
A	13/11/2014	Draft Issued for Client Review	N. Cadd S. Stapleton	V. Clarke
B	25/11/2014	Revised Draft Issued for Client Review	N. Cadd S. Stapleton	V. Clarke
0	3/12/2014	Issued for Information	N. Cadd S. Stapleton	V. Clarke

Approval

Rev	Date	Issued to	Authorised by	
			Name	Signature
A	13/11/2014	Phillip Knapton	S. Pearse	
B	25/11/2014	Phillip Knapton	S. Pearse	
0	3/12/2014	Phillip Knapton	S. Pearse	



© Copyright 2014 Astron Environmental Services Pty Ltd. All rights reserved.

This document and information contained in it has been prepared by Astron Environmental Services under the terms and conditions of its contract with its client. The report is for the clients use only and may not be used, exploited, copied, duplicated or reproduced in any form or medium whatsoever without the prior written permission of Astron Environmental Services or its client.

Abbreviations and Definitions

Abbreviation	Definition
Astron	Astron Environmental Services
Biota	Biota Environmental Sciences
BOM	Bureau of Meteorology
Botanica	Botanica Consulting
CALM	Conservation and Land Management
DAFWA	Department of Agriculture and Food
DEC	Department of Environment and Conservation
DoE	Department of the Environment
DRF	Declared Rare Flora
EPA	Environmental Protection Authority
EPBC Act	<i>Environment Protection and Biodiversity Conservation Act 1999</i>
ESA	Environmentally Sensitive Areas
GIS	Geographical Information System
Ha	Hectares
IBRA	Interim Biogeographic Regionalisation for Australia
Km	Kilometres
m	Metres
mm	Millimetres
NVCP	Native Vegetation Clearing Permit
P	Priority
Parks and Wildlife	Department of Parks and Wildlife
PEC	Priority Ecological Community
RNE	Register of National Estate
‘Survey area’	The proposed Lounge Lizard sand pit, which is 145 hectares and lies immediately west of the existing Flying Fox operations.
T	Threatened
TEC	Threatened Ecological Community
TPFL	Threatened and Priority Flora Database
TP list	Threatened and Priority Flora Species List
WAHerb	Western Australian Herbarium
WAM	Western Australian Museum
WC Act	<i>Wildlife Conservation Act 1950</i>
WSA	Western Areas Limited

Executive Summary

Western Areas Limited commissioned Astron Environmental Services to conduct a Level 2 flora and vegetation survey and Level 1 fauna assessment within the Lounge Lizard mining area. The survey area comprises approximately 145 hectares.

Three vegetation associations were defined within the survey area: sandplain heath, and two mallee woodlands; none of which represent a threatened or priority ecological community. Remnant vegetation was rated as ‘excellent’ condition, with very little disturbance noted.

One hundred and twenty-eight taxa from 32 families and 72 genera were recorded within the survey area. The most represented families were Myrtaceae, Proteaceae and Fabaceae. No threatened flora was recorded during the survey. One priority flora species was recorded; *Eutaxia hirsuta* P2.

No introduced flora species were recorded within the survey area.

Two broad fauna habitats based on landform and vegetation type, namely a *Banksia* low heath and *Eucalyptus* mallee woodland with *Melaleuca* low scrub were recorded in the survey area. Both of these fauna habitat types have been previously recorded within the local vicinity and can be broadly found within the Western Mallee subregion. The *Banksia* low heath is 87.8 ha of the survey area and is considered foraging habitat for the State and Federally-listed Carnaby’s cockatoo (*Calyptorhynchus latirostris*).

Twenty-seven fauna species, comprised of 26 bird species and one mammal species, were recorded during the field survey. This included one fauna species of conservation significance; the crested bellbird (southern subspecies) (*Oreoica gutturalis gutturalis*). This species was recorded within the *Eucalyptus* mallee woodland with *Melaleuca* low scrub fauna habitat type.

Nine fauna species identified by desktop review were considered to have a high likelihood of occurring in the survey area. The majority of these species are birds and therefore highly mobile and can move to similar habitats within the local vicinity and therefore not solely reliant upon fauna habitats of the survey area. However, the clearing of more than 1 ha of quality foraging habitat (*Banksia* dominated heath) is considered under Federal guidelines to comprise a significant impact to the Carnaby’s cockatoo and therefore requires referral under guidelines stipulated by the Federal Department of Environment.

This page has been left blank intentionally.

Table of Contents

1	Introduction.....	1
1.1	Project Background.....	1
1.2	Scope and Objectives.....	1
1.3	Environmental Context.....	5
1.3.1	Climate.....	5
1.3.2	Geology and Soils.....	5
1.3.3	Surface Water and Hydrology.....	6
1.3.4	Vegetation and Flora.....	6
1.3.4.1	Pre-European Vegetation.....	6
1.3.5	Interim Biogeographic Regionalisation of Australia.....	6
1.3.6	Land Use and Tenure.....	6
2	Methodology.....	7
2.1	Desktop Assessment.....	7
2.1.1	Database Searches.....	7
2.1.1.1	Vegetation and Flora.....	8
2.1.1.2	Fauna and Fauna Habitat.....	8
2.1.2	Literature Review.....	8
2.2	Field Survey.....	9
2.2.1	Vegetation and Flora.....	10
2.2.2	Fauna Assessment.....	11
2.3	Taxonomy and Nomenclature.....	11
2.3.1	Flora.....	11
2.3.2	Fauna.....	12
2.4	Limitations.....	12
3	Results.....	13
3.1	Desktop Study.....	13
3.1.1	Environmentally Sensitive Areas.....	13
3.1.2	Vegetation and Flora.....	13
3.1.3	Fauna.....	14

3.2	Vegetation and Flora.....	16
3.2.1	Vegetation.....	16
3.2.2	Conservation Significance of Vegetation	19
3.2.3	Vegetation Condition	19
3.2.4	Flora	19
3.2.5	Conservation Significance of Flora.....	19
3.2.6	Post-survey Likelihood of Occurrence of Conservation Significant Flora	20
3.2.7	Introduced Flora (Weeds)	20
3.3	Fauna.....	20
3.3.1	Fauna Habitats and Condition.....	20
3.3.2	Fauna Observations	21
3.3.3	Likelihood of Occurrence for Conservation Significant Fauna.....	22
4	Discussion	25
5	References.....	27

List of Figures

Figure 1: Survey area location.	3
Figure 2: Climate data for Hyden (Station 10568). Mean annual rainfall data has been calculated from 1928-2014 and mean maximum temperature has been calculated from 1970-2014 (BOM 2014).	5
Figure 3: Mean (1928-2014) rainfall (mm) and total recorded rainfall (mm) received at Hyden station in the 12 months preceding the survey (BOM 2014).....	10

List of Plates

Plate 1: Vegetation representing sandplain heath – facing south-east from GPS coordinate 751662mE and 6409206mS (MGA Zone 50 H).	17
Plate 2: Vegetation representing Eucalyptus mallee woodlands – facing east from GPS coordinate 751727mE and 6408117mS (MGA Zone 50 H).	18
Plate 3: Vegetation representing Eucalyptus mallee woodlands – facing south-east from GPS coordinate 751849mE and 6408430mS (MGA Zone 50 H).....	18
Plate 4: <i>Eutaxia hirsuta</i> P2.	20
Plate 5: Sandplain habitat in which <i>Eutaxia hirsuta</i> P2 was recorded.....	20

Plate 6: <i>Banksia</i> low heath.....	21
--	----

Plate 7: <i>Eucalyptus</i> mallee woodland with <i>Melaleuca</i> low scrub.....	21
---	----

List of Tables

Table 1: Extent of pre-European vegetation in the survey area (Government of Western Australia 2013a).	6
Table 2: Database searches requested.	7
Table 3: Criteria used pre-survey to assess the likely presence of conservation significant flora in the survey area.....	8
Table 4: Criteria used pre and post-survey to assess the likely presence of conservation significant fauna in the survey area.	8
Table 5: Literature review results from surveys conducted within the vicinity of the survey area.	14
Table 6: Literature review results from surveys conducted within the vicinity of the survey area.	15
Table 7: Vegetation associations described for the survey area.	17
Table 8: Taxa most frequently recorded in the survey area.....	19
Table 9: Conservation significant flora recorded during the survey.....	19
Table 10: Fauna species recorded during the survey.	22

List of Appendices

Appendix A: Conservation Categories for Flora, Fauna and Ecological Communities, and Categories for Introduced Flora and Fauna

Appendix B: Vegetation Classification and Condition Scales, and Fauna Habitat Condition Scale

Appendix C: Threatened and Priority Flora and Fauna Species Likelihood of Occurrence within the Survey Areas

Appendix D: Fauna Literature Review and Database Search Results

Appendix E: Vegetation Association Mapping

Appendix F: Quadrat and Relevé Site Data

Appendix G: Vegetation Condition Mapping

Appendix H: Vascular Flora Species List

This page has been left blank intentionally.

1 Introduction

1.1 Project Background

Astron Environmental Services (Astron) was engaged by Western Areas Limited (WSA) to complete a vegetation, flora and fauna assessment to support a Native Vegetation Clearing Permit (NVCP) application, for the proposed Lounge Lizard sand pit (herein referred to as the 'survey area'). The survey area is approximately 145 hectares (ha) and lies immediately west of the existing Flying Fox operations, approximately 75 kilometres (km) east of Hyden, Western Australia (Figure 1).


1.2 Scope and Objectives

The objective of the survey was to provide a Level 2 flora and vegetation survey and Level 1 fauna assessment through a desktop assessment and field survey, to contribute data to a report of sufficient standard to support an NVCP application.

Astron conducted the survey in accordance with the Environmental Protection Authority (EPA) *Position Statement No. 3* (EPA 2002), *Guidance Statement No. 51* (EPA 2004a), *Guidance Statement No. 56* (EPA 2004b), and the *Technical Guide – Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment* (EPA and Department of Environment and Conservation (DEC) 2010). The scope of works was to undertake:

- A desktop assessment, including database searches and literature review of available resources.
- A vegetation and flora assessment, to include:
 - the presence of threatened (T) (formerly known as declared rare flora) and priority (P) flora, and weeds
 - mapping of dominant vegetation associations (with supporting photos of each type and location data) including dominant species, structure and condition
 - discussion of the significance of any vegetation recorded
 - a list of the vascular plant species recorded with the survey area
 - discussion of the levels of disturbance and erosion within the survey area, including vegetation condition mapping.
- A fauna and fauna habitat assessment, to include:
 - habitat assessments for each fauna habitat observed including landform, vegetation structure and condition, and photographs
 - the likely presence of fauna and their conservation significance
 - opportunistic fauna sightings, and their conservation significance
 - mapping of significant fauna habitats (with supporting photos of each type and location data), and discussion of their representation in a regional context
 - an assessment of the significance of any fauna habitat recorded.
- Preparation of a final report which addresses the tasks outlined above, relevant contextual information, methodology, timing and limitations.

This page has been left blank intentionally.



Western Areas - Forresteria Nickel Operations
Lounge Lizard Vegetation, Flora and Fauna Biological Assessment

Figure 1: Survey Area Location

Author: N. Cadd

Date: 07-11-2014

Drawn: C. Dyde

Figure Ref: 17603-14-BIDR-1RevA_141107_Fig01

Datum: GDA 1994 - Projection: MGA Zone 50

0 5 10 15 20 Km



astron
environmental services

This page has been left blank intentionally.

1.3 Environmental Context

1.3.1 Climate

The Mallee region in which the survey area is found, is described as having a dry warm Mediterranean climate, with winter precipitation and an annual rainfall between 300 to 500 millimetres mm (Beard 1990).

Based on long-term climatic data from the nearest weather station at Hyden (Station 10568), (approximately 75 km west of the survey area), the mean annual rainfall since 1928 is 342.3 mm (Bureau of Meteorology (BOM) 2014). The mean maximum daily temperatures range between 16.4 degrees Celsius (°C) and 33.7°C (BOM 2014) (Figure 2).




Figure 2: Climate data for Hyden (Station 10568). Mean annual rainfall data has been calculated from 1928-2014 and mean maximum temperature has been calculated from 1970-2014 (BOM 2014).

1.3.2 Geology and Soils

The Mallee region is described as Archaean and Proterozoic granites overlain in the east by early Tertiary sediments (Beard 1990).

The survey area is mapped as soil landscape zone 250 – *South Eastern Zone of Ancient Drainage of the Avon Province* (Tille et al 2004). The geology is described as gently undulating terrain, dominated by salt lake chains and areas of prominent granite outcrops, on deeply weathered mantle and alluvium over granite rocks of the Yilgarn Craton. Soil is comprised of sandy duplexes that are often alkaline with ironstone gravelly soils and loamy earths that are often calcareous and some loamy duplexes, sandy earths, deep sands, and saline wet soils (Tille et al 2004).

1.3.3 Surface Water and Hydrology

The survey area is located within the Swan Avon-Lockhart catchment of the Avon River Basin. Several tributaries from the surrounding Lake Cronin and Lake Carmody systems are located within the vicinity of the survey area. A minor drainage flat intersects the southern end of the survey area running from east to west.

1.3.4 Vegetation and Flora

1.3.4.1 Pre-European Vegetation

The survey area is located within the Roe Botanical District of the Southwest Province as described by Beard (1990). The Roe Botanical District is dominated by mallee, with *Eucalyptus eremophila* the most consistent species. Patches of eucalypt woodland occur on lower ground, and scrub heath and *Casuarina* thickets on residual plateau soils (Beard 1990).

One vegetation association, Forrestania 2048 (Shepherd et al. 2002), is associated with the survey area. This vegetation unit is described in Table 1, including the current and pre-European extent of the vegetation unit within the survey area and the Mallee region.

Table 1: Extent of pre-European vegetation in the survey area (Government of Western Australia 2013a).

Vegetation association	Extent in survey area (ha)	Current extent in Mallee bioregion (ha)	Pre-European extent (ha)	Proportion of Pre-European extent remaining (%)	Vegetation Description
Forrestania 2048	145	154,944.50	313,728.20	43.39	Shrublands, scrub-heath

1.3.5 Interim Biogeographic Regionalisation of Australia

The Interim Biogeographic Regionalisation for Australia (IBRA version 7) divides the Australian continent into 89 bioregions and 419 subregions (Department of the Environment (DoE) 2014a). The IBRA regions represent a landscape-based approach to classifying the land surface, including attributes of climate, geomorphology, landform, lithology, and characteristic flora and fauna. The survey area occurs in the Mallee bioregion, of which approximately 15-30% is represented in the national reserve system (DoE 2014b).

The biodiversity of the 53 subregions recognised in Western Australia in 2002 was documented as part of a national audit to provide priorities for conservation action (Department of Conservation and Land Management 2002). The survey area occurs within Western Mallee subregion of the Mallee region. The subregion was described in the audit as follows:

Western Mallee MAL2 – has more relief than its eastern counterpart, with main surface-types comprising of clays and silts underlain by Kankar, exposed granite, sandplains and laterite pavements. Vegetation is comprised of mallee communities that occur on a variety of surfaces, *Eucalyptus* woodlands that occur mainly on fine-textured soils with scrub-heath on sand and laterite (Beecham and Danks 2001).

1.3.6 Land Use and Tenure

The survey area is located within the Shire of Kondinin on unallocated Crown land subject to mineral exploration. The local area is primarily used for mineral exploration and mining activity.

2 Methodology

2.1 Desktop Assessment

2.1.1 Database Searches

A search for Environmentally Sensitive Areas (ESA) in the vicinity of the survey area was conducted using the WA Atlas (Landgate 2014) and Register of the National Estate (RNE) spatial database (Australian Government 2008).

Database searches were conducted to identify listed conservation significant ecological communities and flora and fauna species within or in close proximity to the survey area that are listed under the *Wildlife Conservation Act 1950* (WC Act), the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) or are endorsed by the Western Australian Minister for Environment; Heritage. Conservation categories for flora, fauna and ecological communities are presented in Appendix A. Search details are summarised in Table 2. Introduced flora species were compared to the DAFWA list to determine if any have been listed as declared pests (DAFWA 2014) and the Weeds of National Significance (WONS) list (Australian Weeds Committee 2012). Introduced flora and fauna categories are presented in Appendix A.

Table 2: Database searches requested.

Database name	Date search results received	Search focus	Search area
Protected Matters Search Tool (DoE 2014c)	3/10/2014	Flora and fauna of national environmental significance	20 km radius surrounding point defined by the coordinates: -32°29'36", 119°42'00" (GDA94).
Threatened and Priority Ecological Communities Database (Parks and Wildlife 2014a)	9/10/2014	Listed threatened and priority ecological communities.	30 km radius surrounding point defined by the coordinates: -32°29'37.3", 119°41'58.1"
Threatened and Priority Flora Database (TPFL) (Parks and Wildlife 2014b)	13/10/2014	Conservation listed flora species.	
Western Australian Herbarium flora (WAHerb) (Parks and Wildlife 2014c)			
Threatened and Priority Flora Species List (TP list) (Parks and Wildlife 2014d)			
NatureMap (Parks and Wildlife 2014e)	1/10/2014	Terrestrial flora and fauna of conservation significance	20 km radius surrounding point defined by the coordinates: -32°29'36", 119°42'00" (GDA94).
Threatened Fauna Database (Parks and Wildlife 2014f)	9/10/2014	Threatened and priority fauna species	30 km radius surrounding point defined by the coordinates: -32°29'37.3", 119°41'58.1"
BirdLife Australia (Birdlife Australia 2014)	6/10/2014	Bird Species	One degree, coordinates: 32.03636, 120.16847

2.1.1.1 Vegetation and Flora

The 21 Priority Ecological Communities (PECs) listed for the Wheatbelt region (Department of Parks and Wildlife 2014f) were reviewed to determine if any were analogous with ecological communities recorded during the survey area.

Prior to conducting the field survey, aerial imagery was interpreted to identify potential habitat to support conservation significant flora. The conservation significant flora species returned from the database searches were then categorised according to the criteria listed in Table 3 for potential occurrence within the survey area.

Table 3: Criteria used pre-survey to assess the likely presence of conservation significant flora in the survey area.

Likelihood of occurrence	Pre-survey
Likely	Species previously recorded within 10 km of the survey area and suitable habitat appears to be present in the survey area.
Potential	Species previously recorded within 10 - 20 km of the survey area and/or suitable habitat appears to be present in the survey area.
Unlikely	No preferable habitat appears to be present in the survey area.

Following the field survey, the conservation significant flora species which were identified during the desktop study as having the highest potential to occur within the survey area, but were not recorded during the current survey, were again assessed to determine their likelihood of occurrence within the survey area. Post-field survey likelihood of occurrence was primarily based on validating the presence, and thorough inspection of, suitable habitats within the survey area, combined with life form, habit and flowering information for each flora species.

2.1.1.2 Fauna and Fauna Habitat

Conservation significant fauna species returned from the above database searches (Table 2) were categorised pre-survey according to the criteria in Table 4 for occurrence within the survey area.

Table 4: Criteria used pre and post-survey to assess the likely presence of conservation significant fauna in the survey area.

Likelihood of occurrence	Pre-survey
High	Species has been recorded within the survey area or within 20 km of the survey area and preferred habitat appears to be present.
Moderate	Species has not been recorded from within the survey area, however species has been recorded within 20 km of the survey area and suitable habitat appears to be present.
Low	Species recorded within 20 km of the survey area but suitable habitat does not appear to be present.

2.1.2 Literature Review

A number of biological surveys have previously been commissioned by WSA within the vicinity of the survey area and supplied to Astron for the purpose of the desktop assessment. This includes two flora and vegetation assessments, and five fauna assessments. The reports reviewed as part of this assessment are:

- Biota Environmental Sciences Pty Ltd (Biota) 2006a, 'Forrestania Fauna Survey', unpublished report to Western Areas NL.
- Biota 2006b, 'Forrestania Water Disposal Pipeline Fauna Survey', unpublished report to Western Areas NL.
- Biota 2007, 'Forrestania Fauna Monitoring Survey – Flying Fox Phases III and IV', unpublished report to Western Areas NL.
- Biota 2009, 'Forrestania Targeted Malleefowl Survey', unpublished report to Western Areas NL.
- Biota 2010, 'Spotted Quoll Haul Road – Single Phase Fauna Survey', unpublished report to Western Areas NL.
- Botanica Consulting (Botanica) 2013a, 'Flora and Vegetation Survey of the Greater Flying Fox Area', unpublished report to Western Areas NL.
- Botanica 2013b, 'Flora and Vegetation Survey of the Greater New Morning/Spotted Quoll Area', unpublished report to Western Areas NL.
- Craig and Bamford 2004, 'Site Survey of the Proposed Flying Fox, New Morning, Cosmic Boy and Digger Rocks Nickel Mines near Middle Ironcap', unpublished report to Western Areas NL.

2.2 Field Survey

The field survey was conducted from 14 to 16 October 2014 by Astron Senior Environmental Scientist, Natalie Cadd and Scientist, Samantha Stapleton. The daily weather observations recorded at Hyden (station 10568), approximately 75 km west from the survey area, were used to identify local rainfall preceding the survey (Figure 3). A total of 278 mm was recorded at Hyden in the 12 months preceding the survey. This is 62.1 mm below the long term average of 340.1 mm at this site (BOM 2014). In the three months preceding the field survey, approximately 110 mm of rainfall was recorded for Hyden, which is largely consistent with the average rainfall received for the same period (Figure 3).




Figure 3: Mean (1928-2014) rainfall (mm) and total recorded rainfall (mm) received at Hyden station in the 12 months preceding the survey (BOM 2014).

2.2.1 Vegetation and Flora

The vegetation and flora field survey was undertaken in accordance with the requirements outlined in the EPA guidance documents (EPA 2002, 2004a).

Information acquired during the desktop study assisted in the execution and site familiarisation for the field survey.

Seven 20 metre (m) x 20 m quadrats and two relevés (unbounded sample areas of approximately 20 x 20 m) were surveyed in representative vegetation associations within the survey area.

The following information was collected at each quadrat:

- **Location** – coordinates measured using a handheld GPS (MGA50, GDA94). One set of coordinates taken from a central location of each quadrat and relevé.
- **Recorder and date**—personnel involved in sampling that location and the survey date.
- **Species and foliar cover** – all vascular plant species present including conservation significant and introduced species. Species that could not be identified in the field were collected for later identification at the Astron herbarium or the WA Herbarium. Percentage cover for each plant species was estimated to the nearest per cent.
- **Weeds** – the coordinates, density, photographs, and associated habitat/landscape element of any introduced (weed) flora.
- **Vegetation description** – vegetation was described according to Muir's (1977) vegetation classification system and the National Vegetation Information System, level 5 (DoE 2014d)(Appendix B). At this level, vegetation is described to 'association' where up to three

dominant genera for each of the upper, mid and ground strata are categorised based on dominant growth form, cover and height.

- **Vegetation condition** – assessed according to the M.E. Trudgen (Keighery 1994) vegetation condition classification (Appendix B).
- **Habitat** – a broad description of the surrounding landscape based on landform, topography and soil.
- **Disturbances** – records of any obvious disturbances such as fire, tracks or grazing.
- **Photographs** – a photograph was taken of each quadrat and relevé.

Location, population size and photographs for conservation significant flora species identified in the field were also recorded.

The survey area was accessed by 4WD vehicle then traversed on foot. An ArcPad geographic information system (GIS) with the survey area uploaded, plus hard copies of colour aerial photography on A3 maps at a scale of 1: 7,000 were used to assist in navigation, as well as delineating vegetation boundaries.

2.2.2 Fauna Assessment

The fauna survey was undertaken in accordance with relevant EPA guidance (EPA 2004b, EPA and DEC 2010). This guidance advises field observers to describe fauna habitats within the survey area in order to provide a comprehensive indication of fauna that can reasonably be expected to occur. A comprehensive fauna list can only be provided after a detailed Level 2 fauna survey, i.e. a survey that includes the installation and monitoring of traps over several days during at least two seasons.

Therefore, the aim of this fauna survey was to record opportunistic fauna observations, assess the potential for the survey area to support conservation significant fauna species or habitat and, assess and describe the fauna habitat types observed within the survey area.

The survey area was traversed on foot by Natalie Cadd and Samantha Stapleton, concurrent with the botanical assessment. Fauna habitat assessments for each fauna habitat type present within the survey area were completed, with habitat descriptions based on landform and vegetation type. Habitat condition was assessed based on the presence of anthropogenic (human-induced) disturbances, and using the descriptors suggested by Thompson and Thompson (2010) (Appendix B). All vertebrate fauna species observed opportunistically by each field team member were recorded and compiled into one comprehensive list.

2.3 Taxonomy and Nomenclature

2.3.1 Flora

Plant specimens that were not identified in the field were identified in Perth by Astron Principal Scientist, Vanessa Clarke and Senior Botanist, Natalie Cadd. Specimens which were cryptic or difficult to determine were taken to the WA Herbarium for full determination. The assigned nomenclature is consistent with the current listing of scientific names recognised by the WA Herbarium and was used for the species list and associated species information collected. Data from each mapping note were entered into a customised Access database.

2.3.2 Fauna

Nomenclature and sequence for amphibians, reptiles and mammals within this report is as per Western Australian Museum (WAM) *Checklist of the Vertebrates of Western Australia* (WAM 2014). Birds are delineated according to Christidis and Boles (2008).

For species identified in the desktop assessment where there is doubt to their true taxonomy (through subsequent name changes or taxonomic reviews) every effort was made to determine the current scientific name for each taxon. In addition, some taxon names may be followed by 'sp.', meaning that the species name was not given in the data source or the identification is in doubt.

2.4 Limitations

No major limitations were encountered while conducting the vegetation, flora and fauna survey. The region is well documented and seven surveys previously conducted within close proximity to the survey area provided contextual information. There were no access limitations, with the entire survey area able to be traversed on foot. The survey was conducted following average monthly rainfall for the three months prior to the survey. Despite this, many flora species were no longer in flower; the majority however still retained sufficient diagnostic characteristics for positive identification. All fauna data collected from the field were reviewed and validated by Astron Senior Ecologist Matthew Love, who is familiar with the fauna and habitats of the Western Mallee region. The habitats recorded are considered widespread and the species observed are regionally common so competency and experience are not considered to be limiting factors.

3 Results

3.1 Desktop Study

3.1.1 Environmentally Sensitive Areas

Approximately 104 ha of the survey area lies within the buffer of the Lake Cronin ESA. The Lake Cronin area is listed on the RNE (DoE 2014b).

3.1.2 Vegetation and Flora

No State or Commonwealth listed Threatened Ecological Communities (TECs) or PECs were identified within the survey area (Parks and Wildlife 2014a; DoE 2014c). The buffer of the State listed terrestrial PEC “Ironcap Hills vegetation complexes (Mt Holland, Middle, North and South Ironcap Hill, Digger Rock and Hatter Hill) (banded ironstone formation)” is however located approximately 250 m south-east of the survey area.

The Parks and Wildlife TPFL (Parks and Wildlife 2014b), WAHerb (Parks and Wildlife 2014c) and TP list (Parks and Wildlife 2014d) database searches indicated that six species listed as threatened (T) have been recorded within 10 km of the search area:

- *Banksia sphaerocarpa* var. *dolichostyla*
- *Boronia revoluta*
- *Eucalyptus steedmanii*
- *Grevillea involucrata*
- *Grevillea scapigera*
- *Paragoodia crenulata* (Appendix C).

Eighty-eight priority flora species have been previously recorded within 30 km of the search area (Appendix D) (Parks and Wildlife 2014b, 2014c, 2014d, 2014e). Of these, 22 have a priority (P) one (P1) status, 13 have a P2 status, 40 have a P3 status and the remaining 13 have a P4 status. These species were assessed, as part of the desktop study, to determine their likelihood of occurrence within the survey area. Assessment was based on habitat preferences, combined with previous location data and associated vegetation information. The pre-survey desktop assessment indicated that 19 of the listed threatened and priority flora species were considered ‘likely to occur’ in the survey area. A further 34 species were considered to have the ‘potential to occur’ in the survey area (Appendix C).

Results of the literature review identified one threatened, two P2, five P3 and two P4 flora species as having been previously recorded within close proximity of the survey area (Table 5) (Botanica Consulting 2013a,b). The nearest record of a currently listed priority flora species is *Microcorys* sp. Forrestania (V. English 2004) P4. No TECs were recorded, however the North and Middle Ironcap Hills of the “Ironcap Hills Vegetation complexes” PEC have been recorded in previous surveys in the vicinity.

Table 5: Literature review results from surveys conducted within the vicinity of the survey area.

Author (year)	Survey area	Survey focus	Significant conservation values recorded within survey area boundary
Botanica (2013a)	Greater Flying Fox area (267 ha)	Vegetation and flora	<i>Baeckea</i> sp. North Ironcap (R.J. Cranfield 10580) (P2) <i>Boronia westringioides</i> (P2) <i>Baeckea</i> sp. Parker Range (M. Hislop & F. Hort MH 2968) (P3) <i>Daviesia elongata</i> subsp. <i>implexa</i> (P3) <i>Dillwynia acerosa</i> (P3)^ <i>Eutaxia acanthoclada</i> (P3) <i>Verticordia mitodes</i> (P3) <i>Microcorys</i> sp. Forrestania (V. English 2004) (P4) North Ironcap region of the <i>Ironcap Hills Vegetation Complexes</i> (PEC)
Botanica (2013b)	Greater New Morning/Spotted Quoll area (178.5 ha)	Vegetation and flora	<i>Eucalyptus steedmanii</i> (T) <i>Stylidium sejunctum</i> (P2)* (now P3) <i>Eremophila racemosa</i> (P4) <i>Microcorys</i> sp. Forrestania (V. English 2004) (P4) North Ironcap and Middle Ironcap region of the <i>Ironcap Hills Vegetation Complexes</i> (PEC)

*Priority listing revised since completion of report

^Species no longer listed on the priority flora list.

3.1.3 Fauna

The fauna database searches and literature review identified 234 terrestrial vertebrate fauna species recorded within the vicinity of the survey area. This included eight amphibian species, 59 reptile species, 140 bird species and 27 mammal species (including seven introduced), of which 21 species are deemed to have conservation significance. Results are presented in Appendix D. Table 6 presents the conservation significant fauna species identified from prior surveys in the vicinity of the survey areas.

Appendix D lists those species of conservation significance identified by the EPBC Protected Matters (DoE 2014c), *Naturemap* (Parks and Wildlife 2014e), Parks and Wildlife Threatened fauna database (2014f) and *Birddata* (Birdlife Australia 2014) database searches. Aquatic and marine species have been excluded from the report as no marine or aquatic habitats occur within the survey area.

Table 6: Literature review results from surveys conducted within the vicinity of the survey area.

Author (year)	Survey area	Survey focus	Significant conservation values recorded within survey area boundary
Biota (2006a)	Flying Fox area	Fauna	<i>Calyptorhynchus latirostris</i> (Carnaby's black cockatoo) (En, S1) <i>Dasyurus geoffroii</i> (Western quoll/chuditch) (Vu, S1) <i>Platycercus icterotis xanthogenys</i> (Western rosella) (S1) <i>Macropus irma</i> (Western brush wallaby) (P4) <i>Pomatostomus superciliosus ashbyi</i> (White-browed babbler) (P4) <i>Morelia spilota imbricata</i> (Carpet python) (S4) <i>Oreoica gutturalis gutturalis</i> (Crested bellbird) (P4) <i>Hylacola cauta whitlocki</i> (Shy groundwren (western)) (P4) <i>Merops ornatus</i> (Rainbow bee-eater) (Mi, S3)
Biota (2006b)	Flying Fox mine to Diggers Rock	Fauna	<i>Leipoa ocellata</i> (Malleefowl) (S1, Vu) <i>Platycercus icterotis xanthogenys</i> (Western rosella) (S1) <i>Falco peregrinus</i> (Peregrine falcon) (S4) <i>Hylacola cauta whitlocki</i> (Shy groundwren (western)) (P4) <i>Calamanthus campestris montanellus</i> (Rufous fieldwren) (P4) <i>Oreoica gutturalis gutturalis</i> (Crested bellbird) (P4)
Biota (2007)	Flying Fox area	Fauna	<i>Calyptorhynchus latirostris</i> (Carnaby's black cockatoo) (En, S1) <i>Dasyurus geoffroii</i> (Western quoll/chuditch) (Vu, S1) <i>Platycercus icterotis xanthogenys</i> (Western rosella) (S1) <i>Falco peregrinus</i> (Peregrine falcon) (S4) <i>Hylacola cauta whitlocki</i> (Shy groundwren (western)) (P4) <i>Calamanthus campestris montanellus</i> (Rufous fieldwren) (P4) <i>Oreoica gutturalis gutturalis</i> (Crested bellbird) (P4) <i>Pomatostomus superciliosus ashbyi</i> (White-browed babbler) (P4) <i>Macropus irma</i> (Western brush wallaby) (P4)
Biota (2009)	Forrestania area	Targeted Fauna Search	<i>Leipoa ocellata</i> (Malleefowl) (S1, Vu) (inactive mounds were recorded, active mound and sightings in the vicinity of the survey area)
Biota (2010)	Spotted Quoll haul road	Fauna	<i>Calyptorhynchus latirostris</i> (Carnaby's black cockatoo) (En, S1) <i>Leipoa ocellata</i> (Malleefowl) (S1, Vu) (inactive mounds were recorded in the vicinity of a site) <i>Hylacola cauta whitlocki</i> (Shy groundwren (western)) (P4) <i>Pomatostomus superciliosus ashbyi</i> (White-browed babbler) (P4) <i>Oreoica gutturalis gutturalis</i> (Crested bellbird) (P4) <i>Dasyurus geoffroii</i> (Western quoll/chuditch) (Vu, S1)
Craig and Bamford (2004)	Flying Fox, New Morning, Cosmic Boy and Digger Rocks mine areas	Fauna	<i>Morelia spilota imbricata</i> (Carpet python) (S4) <i>Leipoa ocellata</i> (Malleefowl) (S1, Vu) <i>Falco peregrinus</i> (Peregrine falcon) (S4) <i>Calyptorhynchus latirostris</i> (Carnaby's black cockatoo) (En, S1) <i>Platycercus icterotis xanthogenys</i> (Western rosella) (S1) <i>Merops ornatus</i> (Rainbow bee-eater) (Mi, S3) <i>Hylacola cauta whitlocki</i> (Shy groundwren (western)) (P4)

3.2 Vegetation and Flora

3.2.1 Vegetation

Three vegetation associations were described and mapped for the survey area. Vegetation associations recorded, and the extent of each within the survey area, are summarised in Table 7. Vegetation association mapping is presented in Appendix E and survey site data is presented in Appendix F.

Table 7: Vegetation associations described for the survey area.

Vegetation association code and description	Habitat	Area within survey area (proportion of survey area)	Site(s)	Vegetation condition	Representative photograph
SpH1 <i>Banksia cirsioides</i> open low scrub over mixed dwarf scrub. Associated species: <i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>, <i>Beaufortia interstans</i>, <i>Hakea corymbosa</i>, <i>Grevillea cagiana</i>, <i>Allocasuarina microstachya</i>, <i>Verticordia chrysanthella</i>, <i>Verticordia inclusa</i>, <i>Melaleuca sericata</i>, <i>Mesomelaena preissii</i>.	Sandplain heath	87.8 ha (61%)	FOR01, FOR02, FOR01	Excellent	 Plate 1: Vegetation representing sandplain heath – facing south-east from GPS coordinate 751662mE and 6409206mS (MGA Zone 50 H).

Vegetation association code and description	Habitat	Area within survey area (proportion of survey area)	Site(s)	Vegetation condition	Representative photograph
EMW1 <i>Eucalyptus eremophila</i> open tree mallee over mixed <i>Melaleuca</i> species open dwarf scrub to heath. Associated species: <i>Eucalyptus platycorys</i> , <i>Melaleuca hamata</i> , <i>Melaleuca eleuterostachya</i> , <i>Melaleuca adnata</i> , <i>Melaleuca glaberrima</i> , <i>Dodonaea bursarifolia</i> , <i>Phebalium tuberculosum</i> , <i>Lepidosperma viscidum</i> .	Mallee Woodland	10.7 ha (7%)	FOR03, FOR06	Excellent	 Plate 2: Vegetation representing Eucalyptus mallee woodlands – facing east from GPS coordinate 751727mE and 6408117mS (MGA Zone 50 H).
EMW2 <i>Eucalyptus platycorys</i> open tree mallee to tree mallee over <i>Melaleuca hamata</i> open low scrub to low scrub over mixed open dwarf scrub. Associated species: <i>Eucalyptus rigidula</i> , <i>Melaleuca seriata</i> , <i>Melaleuca glaberrima</i> , <i>Leptospermum erubescens</i> , <i>Gastrolobium spinosum</i> , <i>Hibbertia gracilipes</i> , <i>Lepidosperma viscidum</i> .	Mallee Woodland	44.0 ha (30%)	FOR04, FOR05, FOR07, FORr02	Excellent	 Plate 3: Vegetation representing Eucalyptus mallee woodlands – facing south-east from GPS coordinate 751849mE and 6408430mS (MGA Zone 50 H).

3.2.2 Conservation Significance of Vegetation

The vegetation recorded in the survey area is considered representative of the vegetation expected on comparable landforms in the broader region (Beecham and Danks 2001). The vegetation associations recorded in the survey area have been previously described for the broader Lounge Lizard and Flying Fox project areas (Botanica 2006, 2007, 2009, 2013a,b; EPA 2009a).

None of the vegetation associations were considered analogous with a listed TEC or PEC.

One-hundred and four hectares of the survey area lies within the buffer of the Lake Cronin ESA.

3.2.3 Vegetation Condition

Remnant vegetation within the survey area was considered to be in 'excellent' (Keighery 1994) condition (Appendix G). No weed species; recent indication of fire; or evidence of erosion was recorded.

Approximately 2 ha (1%) of the survey area, located immediately south of the adjoining Flying Fox operations, was recorded as being in 'good' condition (Keighery 1994) due to previous partial clearing.

3.2.4 Flora

One hundred and twenty-eight taxa from 32 families and 72 genera were recorded within the survey area. The dominant family was Myrtaceae with 29 taxa recorded and Melaleuca was the most frequently recorded genus (Table 8). A species list for the survey area is presented in Appendix H.

Table 8: Taxa most frequently recorded in the survey area.

Family	Number of taxa
Myrtaceae	29
Proteaceae	23
Fabaceae	15
Family	Number of taxa
Melaleuca	9
Grevillea	6
Acacia	6

3.2.5 Conservation Significance of Flora

Targeted searches in potential habitat identified one state-listed priority flora species within the survey area; *Eutaxia hirsuta* P2 (Table 9) (**Plate 4**). This species was recorded only once, and was associated with the sandplain heath vegetation of vegetation association SpH1 (**Plate 5**).

Table 9: Conservation significant flora recorded during the survey.

Species	Conservation Status	Abundance	Habitat	Location (GPS coordinates)
<i>Eutaxia hirsuta</i>	P2	1	Gently undulating sandplain	751662mE/6409206mS



Plate 4: *Eutaxia hirsuta* P2.



Plate 5: Sandplain habitat in which *Eutaxia hirsuta* P2 was recorded.

3.2.6 Post-survey Likelihood of Occurrence of Conservation Significant Flora

Prior to the field reconnaissance and surveys, threatened and priority flora species recorded from within the proximity of the survey area were reviewed and rated according to their perceived potential to occur with the survey area. Following the field survey, and with greater understanding of the soils, landforms and habitats within the survey area, this list was reviewed (Appendix C). More informed and qualified ratings were then assigned, which reduced the number of species with potential to occur within the survey area from 53 down to 23.

3.2.7 Introduced Flora (Weeds)

No introduced flora species were recorded within the survey area.

3.3 Fauna

3.3.1 Fauna Habitats and Condition

Based on the site inspection and the identified vegetation associations, two broad fauna habitat types were identified within the survey area (Plate 6 and Plate 7). The fauna habitats types recorded were:

- *Banksia* low heath
- *Eucalyptus* mallee woodland with *Melaleuca* low scrub

Fauna habitats were considered to be 'high quality' habitat (Thompson and Thompson 2010), as they appeared to be generally unaffected by grazing herbivores or fire. These fauna habitats types are generally recorded across the Mallee region and are not unique for the survey area.



Plate 6: *Banksia* low heath.



Plate 7: *Eucalyptus* mallee woodland with *Melaleuca* low

3.3.2 Fauna Observations

Twenty-seven fauna species, comprised of 26 bird species and one mammal species, were recorded during the field survey either through direct observation or indirect evidence (Table 10). One fauna of conservation significance was recorded during the field survey, crested bellbird (southern subsp.) (*Oreoica gutturalis gutturalis*). This species was recorded at MGA Zone 50 H 751648mE 6408356mN within the *Eucalyptus* mallee woodland with *Melaleuca* low scrub fauna habitat type. All the vertebrate fauna species recorded in the survey area are known and typical for the Mallee bioregion. No species were outside their expected distribution and all are common for the local vicinity.

Table 10: Fauna species recorded during the survey.

Species name	Common name	Observation type
Birds		
<i>Aquila audax</i>	Wedge-tailed eagle	Individual(s)
<i>Falco berigora</i>	Brown falcon	Individual(s)
<i>Eolophus roseicapillus</i>	Galah	Individual(s), calls
<i>Barnardius zonarius</i>	Australian ringneck	Individual(s), calls
<i>Phaps chalcoptera</i>	Common bronzewing	Individual(s)
<i>Lichenostomus virescens</i>	Singing honeyeater	Individual(s), calls
<i>Lichmera indistincta indistincta</i>	Brown honeyeater	Individual(s), calls
<i>Glyciphila melanops</i>	Tawny-crowned honeyeater	Individual(s), calls
<i>Manorina flavigula</i>	Yellow-throated miner	Individual(s), calls
<i>Acanthagenys rufogularis</i>	Spiny-cheeked honeyeater	Individual(s), calls
<i>Gerygone fusca</i>	Western gerygone	Individual(s)
<i>Anthochaera carunculata</i>	Red wattlebird	Individual(s)
<i>Coracina novaehollandiae</i>	Black-faced cuckoo shrike	Individual(s)
<i>Myiagra inquieta</i>	Restless flycatcher	Calls
<i>Oreoica gutturalis</i>	Crested bellbird	Individual(s), calls
<i>Malurus pulcherrimus</i>	Blue-breasted fairy-wren	Calls
<i>Smicronis brevirostris</i>	Weebill	Individual(s), calls
<i>Pardalotus striatus</i>	Striated pardalote	Individual(s), calls
<i>Acanthiza uropygialis</i>	Chestnut-rumped thornbill	Individual(s), calls
<i>Acanthiza apicalis</i>	Inland thornbill	Individual(s), calls
<i>Cracticus tibicen</i>	Australian magpie	Individual(s)
<i>Rhipidura leucophrys</i>	Willie wagtail	Individual(s)
<i>Corvus coronoides</i>	Australian raven	Individual(s)
<i>Grallina cyanoleuca</i>	Magpie-lark	Individual(s)
<i>Petrochelidon nigricans</i>	Tree martin	Individual(s), calls
<i>Anthus cervinus</i>	Red-throated pipit	Individual(s), calls
Mammals		
<i>Macropus fuliginosus</i>	Western grey kangaroo	Scats, tracks

3.3.3 Likelihood of Occurrence for Conservation Significant Fauna

A description of habitat preferences for each of the State and Federally-listed terrestrial vertebrate fauna of conservation significance identified during the desktop study, and an assessment of the likelihood of the species occurring within the survey area prior to and following the field survey, is provided in Appendix D.

Of the conservation significant fauna identified during the desktop study, six species were deemed to have a moderate likelihood of occurring in the survey area and seven species were considered to have a low likelihood of being present in the survey area due to limited presence of preferred

habitat. Nine species were considered to have a high likelihood of occurring in the survey area, including the crested bellbird (southern spp.) which was recorded during the current assessment. These include the carpet python, Lake Cronin snake, rainbow bee-eater, Carnaby's cockatoo, western rosella, white-browed babbler, shy heathwren and chuditch, which have a historical presence in the survey area along with suitable habitat being present. These species are discussed in further detail below.

Carpet python, *Morelia spilota imbricata* (WC Act Schedule 4)

The Carpet Python is found in south-west Western Australia in most habitats. The species shelters in hollow trunks and limbs, disused burrows, caves, crevices and beneath boulders (Wilson and Swan 2010). The carpet python was previously recorded within the vicinity of the survey area at Flying Fox (Biota 2006a) and Cosmic Bay and Digger Rocks Mine (Craig and Bamford 2004). This species would be expected to occur within the survey area despite not being recorded given the preferred habitats present and preferred habitat present. The lack of record during the current assessment probably reflects the low densities at which it occurs, combined with its cryptic behaviour. The species would be expected to occur in most or all habitats within the study area at a low density.

Lake Cronin snake, *Paroplocephalus atriceps* (WC Act Priority 3)

The Lake Cronin snake is a poorly understood elapid that is endemic to the Lake Cronin area with less than five specimens having been found. Very little is known of their ecological requirements with some anecdotal evidence suggesting this species of snake is generally active at night and is possibly arboreal. The Lake Cronin snake has not been recorded in any of the previous fauna assessments detailed in Table 6, however the little known records are within 9 km of Lake Cronin near to the current survey area.

Rainbow bee-eater, *Merops ornatus* (WC Act Schedule 3, EPBC Act Migratory)

The rainbow bee-eater is seasonally common and widespread in Western Australia. It is a resident, breeding visitor, passage migrant and winter visitor, wintering from the Gascoyne north to Indonesia. It migrates south mainly late September and early October and north from February to April. It is scarce to very common across its range. This species prefers lightly wooded, preferably sandy, country near water (Johnstone and Storr 2004). This species usually nest in excavated burrows in the banks of rivers, creeks or dams, in roadside cuttings, in the walls of gravel pits or quarries, in mounds of gravel, or in cliff-faces (Boland 2004, Forshaw and Cooper 1987, Fry 1984, Higgins 1999, Lill 1993). Nests can be reused (Lill 1993), however mating pairs usually construct new nests at the beginning of each breeding season (Boland 2004, Lill 1993, Morris 1977). The rainbow bee-eater was previously recorded within the vicinity of the survey area at Flying Fox (Biota 2006a) and Cosmic Bay and Digger Rocks Mine (Craig and Bamford 2004). No nests were recorded within the survey area, however, this species is likely to be present during its annual seasonal migration.

Carnaby's black-cockatoo, *Calyptorhynchus latirostris* (WC Act Schedule 1, EPBC Act Endangered)

Carnaby's black-cockatoo is found only in the south-west of Western Australia, and since 1970 its range has reduced by at least 50% (Cale 2003). Carnaby's black-cockatoo feeds in eucalypt woodlands and kwongan heaths, mainly eating the seeds of *Banksia*, *Hakea*, *Grevillea* and *Eucalyptus* species (Cale 2003). Nesting mainly occurs in hollows of salmon gum (*Eucalyptus salmonophloia*) and wandoo (*E. wandoo*) in the Wheatbelt region (Cale 2003). There have been numerous records of the Carnaby's cockatoo within the vicinity of the survey area, specifically at Flying Fox (Biota 2006a, 2007), Spotted Quoll (Biota 2010) and Cosmic Bay and Digger Rocks Mine

(Craig and Bamford 2004). The *Banksia* heath fauna habitat of the survey area represents a potential foraging habitat resource for this species.

Western Rosella (inland subspecies), *Platycercus icterotis xanthogenys* (WC Act Priority 4)

The inland subspecies of the western rosella prefers open eucalypt forest and timbered areas, with a heath understorey, but can also be found in cultivated land and orchards. The survey area represents a cross over in distribution with the other south west subspecies *Platycercus icterotis icterotis*. There have been numerous records of the western rosella within the vicinity of the survey area, specifically at Flying Fox (Biota 2006a, b, 2007) and Cosmic Bay and Digger Rocks Mine (Craig and Bamford 2004). The subspecies is probably widespread in the area, particularly in drier woodland areas and the survey area probably represents a small part of a much larger home range for this species.

White-browed babbler (western Wheatbelt), *Pomatostomus superciliosus ashbyi* (WC Act Priority 4)

The white-browed babbler is mainly found in arid and semiarid zones in habitat dominated by dry sclerophyll woodlands with a shrubby understorey. The subspecies has been recorded in a range of habitat types but typically woodlands with a dense understorey (Johnstone and Storr 2004). This species has a historical presence within the vicinity of the survey area with records of the white-browed babbler at Flying Fox (Biota 2006a, 2007) and at Spotted Quoll (Biota 2010). The white-browed babbler could potentially occur in all fauna habitat types of the survey area.

Crested bellbird (southern subspecies), *Oreoica gutturalis gutturalis* (WC Act Priority 4)

The southern subspecies of the crested bellbird inhabits woodland, mallee and Acacia shrublands (Higgins and Peter 2002) and has previously been recorded within the local vicinity of the survey area at Flying Fox (Biota 2006a, b, 2007) and at Spotted Quoll (Biota 2010). This species was recorded within the current survey area at 50 H 751648mE 6408356mN within the *Eucalyptus* mallee woodland with *Melaleuca* low scrub fauna habitat type. It is likely to be widespread throughout the area in low densities.

Shy heathwren, *Hylacola cauta whitlocki* (WC Act Priority 4)

The shy heathwren inhabits eucalypt mallee woodlands that have relatively dense shrub and heath understorey (Johnstone and Storr 2004). There have been numerous records of the shy heathwren within the vicinity of the survey area, specifically at Flying Fox (Biota 2006a, b, 2007), Spotted Quoll (Biota 2010) and Cosmic Bay and Digger Rocks Mine (Craig and Bamford 2004). It is likely to be widespread throughout the area in low densities.

Chuditch, *Dasyurus geoffroii* (WC Act Schedule 1, EPBC Act Vulnerable)

The chuditch can occupy a wide variety of habitats including woodland, dry sclerophyll forests in both moist, densely vegetated, steeply sloping forest and drier, open, gently sloping forests (Orell and Morris 1994). The former range of the chuditch has contracted markedly since European colonisation due a range of pressures but primarily due to land clearing (Orell and Morris 1994). The chuditch has been recorded close to survey area at Flying Fox (Biota 2006a, 2007), and at Spotted Quoll (Biota 2010). This species is likely to use fauna habitats primarily in a foraging capacity as part of much large home range. No large logs and significant trees were recorded in the survey area as potential den sites.

4 Discussion

The survey area occurs on gently undulating sandplains within the Mallee region. This landform is dominated by mixed low heath and open mallee woodlands.

The vegetation recorded within the survey area is considered typical of what might be expected on these landforms in the Western Mallee subregion and all vegetation associations described within the survey area have been documented for the broader Lounge Lizard and Flying fox mining areas (Botanica 2006, 2007, 2009, 2013; EPA 2009a).

No Commonwealth or State listed TECs or PECs were recorded within the survey area (DoE 2014c; Parks and Wildlife 2014a). The buffer of the State listed PEC “*Ironcap Hills vegetation complexes (Mt Holland, Middle, North and South Ironcap Hill, Digger Rock and Hatter Hill) (Banded ironstone formation)*” is located approximately 250 m south-east of the survey area. The survey area does not contain banded ironstone formation or vegetation complexes associated with this PEC. Aside from a small area of previously disturbed vegetation adjacent to the existing Flying Fox operations, vegetation within the survey area was in ‘excellent’ condition.

A large proportion of the survey area lies within the buffer of the Lake Cronin ESA. This ESA is centered on the Lake Cronin wetland. The buffer zone associated with the Lake Cronin ESA has been described as supporting extensive contiguous shrubland, sandplain and woodland environments, including excellent representation of a range of vegetation types that are now extensively cleared in the Wheatbelt (EPA 2009b). As such, vegetation within the buffer area may be considered locally and regionally significant.

The suite of flora recorded was considered representative of what may be expected in the local area (Botanica 2009, 2013a, b). The survey followed three months of rainfall consistent with the average, however a number of flora species were no longer in flower during the field survey. Despite this, a reasonable inventory of flora was captured and the majority of species still retained sufficient diagnostic characteristics for positive identification.

No threatened flora was identified in the survey area and, based on comprehensive searches of the survey area, none are considered likely to occur. One priority flora species, *Eutaxia hirsuta* P2, was recorded. One individual of this species was identified in association with sandplain heath vegetation. There are currently five vouchered records of this species at the Western Australian Herbarium (2014), with two records listed as occurring within conservation reserves; Lake Barker Reserve and Wogarl Reserve. The closest record occurs some 80 km north-east of the survey area, near Lake Barker (Western Australian Herbarium 2014). Given the current known spatial distribution of this species, the occurrence of specimens within conservation reserves and the extent of suitable habitat outside of the survey area, it is not considered likely that the proposed clearing will have a significant impact on the regional conservation status of this taxa. A permit to take is not required prior to clearing, as this taxon is not specially protected under any State or Commonwealth legislation.

Despite undertaking a comprehensive targeted search for conservation significant flora species, there remains potential for 23 conservation significant species to occur within the survey area. Although the survey area was traversed comprehensively, vegetation at times was dense and some species are cryptic. Should any of these species be present within the survey area, the proposed clearing is not considered likely to significantly impact the conservation status of any of the species, based on the number of locations outside of the survey area that they are currently known from.

No weed species were recorded within the survey area.

The survey area contains two broad fauna habitats based on landform and vegetation type, namely a *Banksia* low heath and *Eucalyptus* mallee woodland with *Melaleuca* low scrub. Both of these fauna habitat types have been previously recorded within the local vicinity (Biota 2006a, b, 2007, 2010), are typical of the Western Mallee subregion, and are therefore not unique to the survey area. However, these fauna habitats potentially provide foraging and shelter for nine conservation significant species considered to have a high likelihood of occurrence; carpet python, Lake Cronin snake, rainbow bee-eater, Carnaby's cockatoo, western rosella, white-browed babbler, crested bellbird, shy heathwren and chuditch. The majority of these conservation significant species would generally exist in low densities within the fauna habitat types of the survey area and are not likely to be solely reliant on the survey area. No breeding activities were recorded within the survey area. In addition the majority of these species are birds and therefore highly mobile, being able to easily move to similar habitats within the vicinity of the survey area. The carpet python and the Lake Cronin snake may suffer localised mortalities due to clearing activities however it is not expected that this would reduce the population of either species in a regional sense. Overall, the direct impact to fauna within the survey area, from the proposed clearing, is expected to be minimal.

Twenty-seven fauna species, comprised of 26 bird species and one mammal species, were recorded during the field survey either through direct observation or indirect evidence including one fauna species of conservation significance the crested bellbird (southern subspecies) (*Oreoica gutturalis gutturalis*).

The *Banksia* low heath habitat (87.8 ha) and potentially the *Eucalyptus* mallee woodland habitats (54.7 ha) within the survey area are considered potential foraging habitat for Carnaby's black cockatoo. This species could be expected to occur on an infrequent basis, foraging within the survey area during their annual migration activities. However similar foraging habitat exists outside of the survey area within the local area and no breeding hollows or potential breeding trees were recorded within the survey area. The potential impacts to Carnaby's black cockatoo from clearing of foraging habitat within the survey area should be assessed against the EPBC Act referral guidelines (DSEWPaC 2012). These guidelines state that clearing of greater than 1 ha of quality foraging habitat constitutes a high risk of significant impact and therefore referral to the Federal environment minister is recommended.

5 References

- Australian Government 2008, Register of the National Estate - Spatial Database (RNESDB), viewed 23 September 2014, <http://www.environment.gov.au/metadataexplorer/explorer.jsp>
- Australian Weeds Committee 2012, *Weeds of National Significance 2012*, Department of Agriculture, Fisheries and Forestry, Canberra.
- Birdlife International Birddata 2014, Birddata database search, viewed 6 October 2014, <http://www.birddata.com.au/homecontent.do>
- Boland, CRJ 2004, 'Breeding biology of Rainbow Bee-eaters (*Merops ornatus*): a migratory, colonial, cooperative bird'. *Auk*, vol. 121, pp. 811-823.
- Bureau of Meteorology (BOM) 2014a, Climate Data Online, viewed 24 October 2014, <http://www.bom.gov.au/climate/data/>
- Beard, JS 1990, *Plant life of Western Australia*, Kangaroo Press, Kenthurst, NSW.
- Beecham, B and Danks, A 2001, *Mallee 2 (MAL2 – Western Mallee subregion), A Biodiversity Audit of Western Australia's 53 Biogeographical Subregions in 2002*, CALM, Perth.
- Biota Environmental Sciences Pty Ltd (Biota) 2006a, 'Forrestania Fauna Survey', unpublished report to Western Areas NL.
- Biota 2006b, 'Forrestania Water Disposal Pipeline Fauna Survey', unpublished report to Western Areas NL.
- Biota 2007, 'Forrestania Fauna Monitoring Survey – Flying Fox Phases III and IV', unpublished report to Western Areas NL.
- Biota 2009, 'Forrestania Targeted Malleefowl Survey', unpublished report to Western Areas NL.
- Biota 2010, 'Spotted Quoll Haul Road – Single Phase Fauna Survey', unpublished report to Western Areas NL.
- Birdlife Australia 2014, *Birddata*, viewed 22 September 2014, <http://www.birddata.com.au/homecontent.do>
- Botanica Consulting (Botanica) 2006, 'Flora and Vegetation Survey of the Flying Fox North East Exploration Area', unpublished report to Western Areas NL.
- Botanica 2007, 'Flora and Vegetation Survey within the Greater Flying Fox mine site, prepared for Western Areas', unpublished report to Western Areas NL.
- Botanica 2009, 'Flora and Vegetation Survey within the lounge Lizard/Flying Fox area, Proposed Gravel Pit, Prepared for Western Areas', unpublished report to Western Areas NL.
- Botanica Consulting 2013a, 'Flora and Vegetation Survey of the Greater Flying Fox Area', unpublished report to Western Areas NL.

Botanica Consulting 2013b, 'Flora and Vegetation Survey of the Greater New Morning/Spotted Quoll Area', unpublished report to Western Areas NL.

Cale, B 2003, 'Carnaby's Black-Cockatoo (*Calyptrorhynchus latirostris*) Recovery Plan', Department of Conservation and Land Management, Perth.

Craig and Bamford 2004, 'Site Survey of the Proposed Flying Fox, New Morning, Cosmic Boy and Digger Rocks Nickel Mines near Middle Ironcap', unpublished report to Western Areas NL

Department of Conservation and Land Management (CALM) 2002, *Bioregional Summary of the 2002 Audit for Western Australia*, printed under CALM, Perth.

Department of the Environment (DoE) 2014a, Australia's Bioregions (IBRA), viewed 23 October 2014, <http://www.environment.gov.au/topics/land/nrs/science-maps-and-data/ibra/ibra7-codes>

DoE 2014b, National Reserve System, viewed 24 September 2014, <http://www.environment.gov.au/topics/land/national-reserve-system>

DoE 2014c, Protected Matters Search Tool, viewed 23 October 2014, <http://www.environment.gov.au/erin/ert/epbc/index.html>

DoE 2014d, *Australian Vegetation Attribute Manual*, National Vegetation Information System Version 6, viewed June 2014, <http://www.environment.gov.au/erin/nvis/publications/avam/section-2-1.html#table1>

Department of Parks and Wildlife (Parks and Wildlife) 2014a, *Threatened and Priority Ecological Community database*, Parks and Wildlife, Kensington.

Department of Sustainability, Environment, Water, Population and Communities (DSEWPaC) 2012, EPBC Act referral guidelines for three threatened black cockatoo species, DSEWPaC, Canberra.

Environmental Protection Authority (EPA) 2002, *Terrestrial Biological Surveys as an Element of Biodiversity Protection*, Position Statement 3, EPA, Perth.

EPA 2004a, *Terrestrial Flora and Vegetation Surveys for Environmental Impact Assessment in Western Australia*, Guidance Statement 51, EPA, Perth.

EPA 2004b, *Terrestrial Fauna Surveys for Environmental Impact Assessment in Western Australia*. Guidance Statement 56, EPA Perth.

EPA 2009a, *Spotted Quoll Open Pit Nickel Mine; Report and recommendations of the Environmental Protection Authority*, EPA Perth.

EPA 2009b, *Advice on Conservation Values and Review of Nature Reserve Proposals in the Lake Cronin Region*, EPA Perth.

EPA and DEC 2010, *Technical Guide – Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment* (eds B.M. Hyder, J. Dell, and M.A. Cowan), Perth, Western Australia.

Forshaw, JM and Cooper, WT 1987, *Kingfishers and Related Birds: Todidae, Momotidae, Meropidae*, Lansdowne Editions, Melbourne.

Fry, CH 1984, *The Bee-eaters*, Poyser, Calton, England.

Higgins, PJ (ed.) 1999, *Handbook of Australian, New Zealand and Antarctic Birds*. Volume Four - Parrots to Dollarbird, Melbourne, Oxford University Press.

Higgins, PJ and Peter, JM (eds) 2002, *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 6: Pardalotes to Shrike-thrushes, Melbourne, Oxford University Press.

Johnstone, RE and Storr, GM, 2004, *Handbook of Western Australian Birds Volume II: Passerines (Blue-winged Pitta to Goldfinch)*, Western Australian Museum, Perth.

Keighery, BJ 1994, *Bushland Plant Survey: A Guide to Plant Community Surveys for the Community*, Wildflower Society of Western Australia Inc., Nedlands.

Landgate 2014, WA Atlas, viewed 24 October 2014
<https://www2.landgate.wa.gov.au/bmvf/app/waatlas/>

Lill, A 1993, 'Breeding of Rainbow Bee-eaters in Southern Victoria', *Corella*, vol. 17, pp. 100-106.

Morris, IC 1977, 'More observations of Rainbow Bee-eaters *Merops ornatus* in the Warby Ranges', *Victorian Naturalist*, vol. 94, pp. 158-160.

Muir, BG 1977, *Biological Survey of the Western Australian Wheatbelt Part 2 Vegetation and Habitat of Bendering Reserve*, Records of the Western Australian Museum Supplement 3.

Orell, P and Morris, K 1994, *Chuditch Recovery Plan 1992-2001*, Western Australian Department of Conservation and Land Management Western Australian Wildlife Research Centre, PO Box 51, Wanneroo, WA 6065

Parks and Wildlife 2014b, *Threatened and Priority Flora database*, Parks and Wildlife, Kensington.

Parks and Wildlife 2014c, *Western Australian Herbarium database*, Parks and Wildlife, Kensington.

Parks and Wildlife 2014d, *Threatened and Priority Flora List*, Parks and Wildlife, Kensington.

Parks and Wildlife 2014e, *NatureMap database search*, viewed 24 October 2014,
<http://naturemap.dec.wa.gov.au>

Parks and Wildlife 2014f, *Priority Ecological Communities for Western Australia Version 21*, viewed 24 October 2014, <http://www.Parks and Wildlife.wa.gov.au/images/documents/plants-animals/threatened-species/tecs/Priority ecological community list 20 May2014.pdf>

Shepherd, DP, Beeston, GR and Hopkins, AJM 2002, *Native Vegetation in Western Australia: Resource Management Technical Report 249*, Department of Agriculture, Western Australia.

Thompson, SA, and Thompson, GG 2010, *Terrestrial Vertebrate Fauna Assessments for Ecological Impact Assessment*, Terrestrial Ecosystems, Mt Claremont.

Tille, P, Schoknecht, N and Purdie, B 2004, *Soil-landscapes Mapping in South-Western Australia*, Resource Management Technical Report 280, Department of Agriculture and Food, Perth.

Western Australian Herbarium 2014, *FloraBase* – the Western Australian Flora, viewed 28 October 2014, <http://florabase.dpaw.wa.gov.au/search/advanced>

Western Australian Museum 2014, *Western Australian Museum Checklist of the Vertebrates of Western Australia*, viewed 24 October 2014, <http://museum.wa.gov.au/research/departments/terrestrial-zoology/checklist-terrestrialvertebrate-fauna-western-australia>

Wilson, S and Swan, G 2010, *A Complete Guide to Reptiles of Australia*, 3rd Edition, New Holland Publishers, Sydney.

Appendix A: Conservation Categories for Flora, Fauna and Ecological Communities, and Introduced Categories for Flora and Fauna

This page has been left blank intentionally.

Table A.1: Categories of threatened ecological communities (DEC 2010).

PD: Presumed Destroyed
An ecological community that has been adequately searched for but for which no representative occurrences have been located. The community has been found to be totally destroyed or so extensively modified throughout its range that no occurrence of it is likely to recover its species composition and/or structure in the foreseeable future. An ecological community will be listed as presumed totally destroyed if there are no recent records of the community being extant and either of the following applies (A or B): A) Records within the last 50 years have not been confirmed despite thorough searches of known or likely habitats or B) All occurrences recorded within the last 50 years have since been destroyed.
CR : Critically Endangered
An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or that was originally of limited distribution and is facing severe modification or destruction throughout its range in the immediate future, or is already severely degraded throughout its range but capable of being substantially restored or rehabilitated. An ecological community will be listed as Critically Endangered when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future. This will be determined on the basis of the best available information, by it meeting any one or more of the following criteria (A, B or C): A) The estimated geographic range, and/or total area occupied, and/or number of discrete occurrences since European settlement have been reduced by at least 90% and either or both of the following apply (i or ii): i) geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is imminent (within approximately 10 years); ii) modification throughout its range is continuing such that in the immediate future (within approximately 10 years) the community is unlikely to be capable of being substantially rehabilitated. B) Current distribution is limited, and one or more of the following apply (i, ii or iii): i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the immediate future (within approximately 10 years); ii) there are very few occurrences, each of which is small and/or isolated and extremely vulnerable to known threatening processes; iii) there may be many occurrences but total area is very small and each occurrence is small and/or isolated and extremely vulnerable to known threatening processes. C) The ecological community exists only as highly modified occurrences that may be capable of being rehabilitated if such work begins in the immediate future (within approximately 10 years).

En: Endangered
An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or was originally of limited distribution and is in danger of significant modification throughout its range or severe modification or destruction over most of its range in the near future. An ecological community will be listed as Endangered when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future. This will be determined on the basis of the best available information by it meeting any one or more of the following criteria (A, B, or C): A) The geographic range, and/or total area occupied, and/or number of discrete occurrences have been reduced by at least 70% since European settlement and either or both of the following apply (i or ii): i) the estimated geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is likely in the short term future (within approximately 20 years); ii) modification throughout its range is continuing such that in the short term future (within approximately 20 years) the community is unlikely to be capable of being substantially restored or rehabilitated. B) Current distribution is limited, and one or more of the following apply (i, ii or iii): i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the short term future (within approximately 20 years); ii) there are few occurrences, each of which is small and/or isolated and all or most occurrences are very vulnerable to known threatening processes; iii) there may be many occurrences but total area is small and all or most occurrences are small and/or isolated and very vulnerable to known threatening processes. C) The ecological community exists only as very modified occurrences that may be capable of being substantially restored or rehabilitated if such work begins in the short-term future (within approximately 20 years).
VU: Vulnerable
An ecological community that has been adequately surveyed and is found to be declining and/or has declined in distribution and/or condition and whose ultimate security has not yet been assured and/or a community that is still widespread but is believed likely to move into a category of higher threat in the near future if threatening processes continue or begin operating throughout its range. An ecological community will be listed as Vulnerable when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction or significant modification in the medium to long-term future. This will be determined on the basis of the best available information by it meeting any one or more of the following criteria (A, B or C): A) The ecological community exists largely as modified occurrences that are likely to be capable of being substantially restored or rehabilitated. B) The ecological community may already be modified and would be vulnerable to threatening processes, is restricted in area and/or range and/or is only found at a few locations. C) The ecological community may be still widespread but is believed likely to move into a category of higher threat in the medium to long term future because of existing or impending threatening processes.

Table A.2: Definitions and criteria for priority ecological communities (DEC 2010).

Possible threatened ecological communities that do not meet survey criteria or that are not adequately defined are added to the Priority Ecological Community Lists under Priorities 1, 2 and 3. Ecological Communities that are adequately known, and are rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

P1: Priority One – Poorly-known ecological communities
Ecological communities with apparently few, small occurrences, all or most not actively managed for conservation (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) and for which current threats exist. Communities may be included if they are comparatively well-known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.
P2: Priority Two – Poorly-Known ecological communities
Communities that are known from few small occurrences, all or most of which are actively managed for conservation (e.g. within national parks, conservation parks, nature reserves, State forest, unallocated Crown land, water reserves, etc.) and not under imminent threat of destruction or degradation. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.
P3: Priority Three – Poorly-Known ecological communities
(i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or: (ii) communities known from a few widespread occurrences, which are either large or within significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or; (iii) communities made up of large, and/or widespread occurrences, that may or not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, and inappropriate fire regimes. Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.
P4: Priority Four
Ecological communities that are adequately known, rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring. Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands. Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable. (c) Ecological communities that have been removed from the list of threatened communities during the past five years.
P5: Priority Five – Conservation dependent ecological communities
Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

Table A.3: Definitions and criteria for threatened ecological communities (DoE 2014).

Three categories exist for listing threatened ecological communities under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). An ecological community may be categorised:

Categories of ecological communities	
Critically endangered	If, at that time, it is facing an extremely high risk of extinction in the wild in the immediate future.
Endangered	If, at that time, it is not critically endangered and is facing a very high risk of extinction in the wild in the near future.
Vulnerable	If, at that time, it is not critically endangered or endangered, and is facing a high risk of extinction in the wild in the medium-term future.

Table A.4: Conservation codes for Western Australian flora and fauna (DPaW 2013).

Code	Conservation category	Definition
X	Schedule 2 of the Wildlife Conservation (Specially Protected Fauna) Notice and Wildlife Conservation (Rare Flora) Notice under the Wildlife Conservation Act 1950. Presumed Extinct Fauna Presumed Extinct Flora (Declared Rare Flora – Extinct)	Taxa which have been adequately searched for and there is no reasonable doubt that the last individual has died, and have been gazetted as such.
T	Schedule 1 of the Wildlife Conservation (Specially Protected Fauna) Notice and Wildlife Conservation (Rare Flora) Notice under the Wildlife Conservation Act 1950. Threatened Fauna (Fauna that is rare or is likely to become extinct) Threatened Flora (Declared Rare Flora - Extant)	Taxa that have been adequately searched for and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection, and have been gazetted as such.
IA	Schedule 3 of the Wildlife Conservation (Specially Protected Fauna) Notice under the Wildlife Conservation Act 1950. Birds protected under an international agreement	Birds that are subject to an agreement between governments of Australia and Japan relating to the protection of migratory birds and birds in danger of extinction.
S	Schedule 4 of the Wildlife Conservation (Specially Protected Fauna) Notice under the Wildlife Conservation Act 1950. Other specially protected fauna	Fauna that is in need of special protection, otherwise than for the reasons mentioned in the above schedules.
	Threatened fauna and flora (Schedule 1) are further ranked by the Department according to their level of threat using IUCN Red List criteria: CR: Critically Endangered - considered to be facing an extremely high risk of extinction in the wild. EN: Endangered - considered to be facing a very high risk of extinction in the wild. VU: Vulnerable - considered to be facing a high risk of extinction in the wild.	

Table A.5: Priority species under Western Australian Wildlife Conservation Act 1950

Taxa that have not yet been adequately surveyed to be listed under Schedule 1 or 2 are added to the Priority Flora and Priority Fauna Lists under Priorities 1, 2 or 3. These three categories are ranked in order of priority for survey and evaluation of conservation status so that consideration can be given to their declaration as threatened flora or fauna. Taxa that are adequately known, are rare but not threatened, or meet criteria for Near Threatened, or that have been recently removed from the threatened list for other than taxonomic reasons, are placed in Priority 4. These taxa require regular monitoring. Conservation Dependent species are placed in Priority 5.

P1: Priority One – Poorly known taxa
Taxa that are known from one or a few collections or sight records (generally less than five), all on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, Shire, Westrail and Main Roads WA road, gravel and soil reserves, and active mineral leases and under threat of habitat destruction or degradation. Taxa may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes.
P2: Priority Two – Poorly known taxa
Taxa that are known from one or a few collections or sight records, some of which are on lands not under imminent threat of habitat destruction or degradation, e.g. national parks, conservation parks, nature reserves, State forest, vacant Crown land, water reserves, etc. Taxa may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes.
P3: Priority Three – Poorly known taxa
Taxa that are known from collections or sight records from several localities not under imminent threat, or from few but widespread localities with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Taxa may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and known threatening processes exist that could affect them.
P4: Priority Four: Rare, near threatened and other taxa in need of monitoring
(a) Rare. Taxa that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These taxa are usually represented on conservation lands. (b) Near Threatened. Taxa that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable. (c) Taxa that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.
P5: Priority Five: Conservation dependent taxa
Taxa that are not threatened but are subject to a specific conservation program, the cessation of which would result in the taxa becoming threatened within five years.

Table A.6: Categories and definitions for EPBC Act listed flora and fauna species (DoE 2014).

Conservation category	Definition
Extinct	Taxa not definitely located in the wild during the past 50 years.
Extinct in the wild	Taxa known to survive only in captivity.
Critically endangered (CR)	Taxa facing an extremely high risk of extinction in the wild in the immediate future.
Endangered (E)	Taxa facing a very high risk of extinction in the wild in the near future.
Vulnerable (V)	Taxa facing a high risk of extinction in the wild in the medium term.
Near threatened (NT)	Taxa that risk becoming Vulnerable in the wild.
Conservation dependent (CD)	Taxa whose survival depends upon ongoing conservation measures. Without these measures, a conservation dependent taxon would be classified as Vulnerable or more severely threatened.
Data deficient (insufficiently known) (DD)	Taxa suspected of being Rare, Vulnerable or Endangered, but whose true status cannot be determined without more information.
Least concern (LC)	Taxa that are not considered threatened.

Table A.7: Declared pests categories as gazetted under the Biosecurity and Agriculture Management Act 2007 (Department of Agriculture and Food 2013).

The management of introduced flora species in Western Australia is now regulated through the Biosecurity and Agriculture Management Act 2007 (BAM Act). A list of declared pests, including ‘pest’ plants is provided under the BAM Act, which has been updated to incorporate a number of other Acts that are administered by DAFWA (2013). Declared pests can fall into two categories: one that relates to the prevention of introducing the species or eradicating it; and the other relates to managing the species and whether it can be kept (i.e. for scientific purposes, education or other purpose).

The threat and risk posed to site-specific biodiversity values, influences to rehabilitation success, primary production, infrastructure assets or human health will differ depending on the unique characteristics of each site and the associated land management practice or operation. Therefore site or project specific weed assessments and priorities should be reviewed for each project.

As per introduced flora species, the BAM Act seeks to establish a modern biosecurity regulatory scheme to prevent serious animal pests from entering the State and becoming established, and to minimise the spread and impact of any that are already present within the State. Declared animal pests fall into three categories as Gazetted under the BAM Act (DAFWA 2013). These categories are outlined in Table A.7.

Category	Description
C1 (Exclusion)	Pests will be assigned to this category if they are not established in Western Australia and control measures are to be taken, including border checks, in order to prevent them entering and establishing in the State.
C2 (Eradication)	Pests will be assigned to this category if they are present in Western Australia in low enough numbers or in sufficiently limited areas that their eradication is still a possibility.
C3 (Management)	Pests will be assigned to this category if they are established in Western Australia but it is feasible, or desirable, to manage them in order to limit their damage. Control measures can prevent a C3 pest from increasing in population size or density or moving from an area in which it is established into an area which currently is free of that pest.

This page has been left blank intentionally.

Appendix B: Vegetation Classification and Condition Scales, and Fauna Habitat Condition Scale

This page has been left blank intentionally.

Table B.1: Muir life form/height class Muir (1977).

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70%-100%	MID DENSE 30%-70%	SPARSE 10%-30%	VERY SPARSE 2%-10%
Trees > 30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
Trees 15 – 30m	Dense Forest	Forest Low	Woodland	Open Woodland
Trees 5 – 15m	Dense Low Forest A	Forest A	Low Woodland A	Open Low Woodland A
Trees < 5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
Mallee Tree Form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
Mallee Shrub Form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
Shrubs > 2m	Dense Thicket	Thicket	Scrub	Open Scrub
Shrubs 1.5 – 2m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
Shrubs 1 – 1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
Shrubs 0.5 – 1m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
Shrubs 0 – 0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
Mat Plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
Hummock Grass	Dense Hummock Grass	Mid-dense Hummock	Hummock Grass	Open Hummock Grass
Bunch Grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
Bunch Grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
Sedges > 0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
Mosses, Liverworts	Dense Mosses	Mosses	Open Mosses	Very Open Mosses

Table B.2: Vegetation condition scale Keighery (1994).

Condition	Description
Pristine	Pristine or nearly so, no obvious signs of disturbance.
Excellent	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species.
Very Good	Vegetation structure altered obvious signs of disturbance. Disturbance to vegetation structure covers repeated fire, aggressive weeds, dieback, logging, grazing.
Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. Disturbance to vegetation structure covers frequent fires, aggressive weeds at high density, partial clearing, dieback and grazing.
Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. Disturbance to vegetation structure includes frequent fires, presence of very aggressive weeds, partial clearing, dieback and grazing.
Completely Degraded	The structure of the vegetation is no longer intact and the area is completely or almost completely without native species. These areas often described as “parkland cleared” with the flora comprising weed or crop species with isolated native trees or shrubs.

Modified from Trudgen, 1991, by Keighery for the Swan Coastal Plain Survey. Found in *Bushland Plant Survey: a Guide to Plant Community Survey for the Community*.

Table B.3: Fauna habitat condition scale (Thompson and Thompson 2010).

Habitat condition	Condition description
High Quality Fauna Habitat	These areas closely approximate the vegetation mix and quality that would have been in the area prior to any human induced disturbance. The habitat has connectivity with other habitats and is likely to support the most natural vertebrate fauna assemblage.
Very Good Fauna Habitat	These areas show minimal signs of human induced disturbance (e.g. grazing, clearing, fragmentation, weeds) and retain almost all of the characteristics of the habitat had it not been disturbed. The habitat has connectivity with other habitats, and fauna assemblages in these areas are likely to be minimally effected by disturbance.
Good Fauna Habitat	These areas show signs of human induced disturbance (e.g. grazing, clearing, fragmentation, weeds) but generally retain many of the characteristics of the habitat had it not been disturbed. The habitat still retains some connectivity with other habitats but fauna assemblages in these areas are likely to be affected by disturbance. Fauna assemblages in these areas are likely to be similar to what might be expected in this habitat.
Disturbed Fauna Habitat	These areas show signs of human induced significant disturbance (e.g. mining, clearing, tracks and roads). Many of the trees, shrubs and undergrowth have died or have been cleared. These areas may be in the early succession and regeneration stages. Areas may show signs of significant grazing, contain an abundance of weeds or have been damaged by vehicles or machinery. Habitats are fragmented or have limited connectivity with other fauna habitats. Fauna assemblages in these areas are likely to differ significantly from what might be expected in the area had the disturbance not occurred.
Highly Degraded Fauna Habitat	These areas often have a significant human induced loss of vegetation, and / or a large number of vehicle tracks and / or have been completely cleared, and / or areas have been heavily grazed or farmed. There is limited or no fauna habitat connectivity. Fauna assemblages in these areas are likely to differ significantly from what existed prior to the disturbance, and are often depleted compared to what existed prior to the disturbance.

This page has been left blank intentionally

Appendix C: Threatened and Priority Flora, and Fauna Species Likelihood of Occurrence within the Survey Area.

This page has been left blank intentionally.

Table C.1: Likelihood of occurrence of threatened and priority flora recorded within 20 km of the survey area (TPFL (Parks and Wildlife 2014b); WAHerb (Parks and Wildlife 2014c); TP List (Parks and Wildlife 2014d); *NatureMap* (Parks and Wildlife 2014e)). Note: The TP List database is searched using place names. As a result, many of the records obtained from this database may occur beyond 50 km of the survey area (Parks and Wildlife 2014d).

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
Threatened						
<i>Banksia sphaerocarpa</i> var. <i>dolichostyla</i>	Lignotuberous shrub, 1-3 m high.	Perennial	Yellow-orange flowers from March - May.	Lateritic gravel, grey sand.	Potential	Unlikely
<i>Boronia revoluta</i>	Shrub, 0.4-0.8 m high.	Perennial	Pink flowers from July - August.	Stony sandy loam or sand. Plains, hillsides & summits.	Potential	Potential
<i>Eucalyptus steedmanii</i>	Tree, 2-8(-12) m high, bark smooth.	Perennial	White flowers from January - March.	Gravelly loam over ironstone, sand. Low hills, undulating plains.	Likely	Unlikely
<i>Grevillea involucrata</i>	Prostrate to low-domed open shrub, 0.15-0.3 m high, up to 2 m wide.	Perennial	Pink/pink-red flowers in June - October.	Amongst medium trees, or low trees, or low (sclerophyll) shrubland; in rocky or stony soil, or sand, or loam; occupying lateritic ridge.	Unlikely	Unlikely
<i>Grevillea scapigera</i>	Suckering, prostrate to weakly ascending shrub, 0.15-0.4 m high, up to 1.8 m wide.	Perennial	White/yellow-green flowers in February, or October - November.	Amongst medium trees, or low trees; in gravelly soil, or sand; occupying heathlands, sand plains; growing in gravel pits.	Unlikely	Unlikely
<i>Paragoodia crenulata</i>	Prostrate shrub	Annual	Yellow and brown flowers.	Hill slopes, gullies. Stony red loams, clays.	Potential	Unlikely
Priority 1						
<i>Aotus lanea</i>	Prostrate shrub, to 0.2 m high.	Perennial	Red/orange/brown, flowers in April or October.	Grey clayey sand, yellow clay, deep siliceous sand. Edges of salt lakes, valleys.	Unlikely	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Baeckea</i> sp. Blue Haze Mine (P. Armstrong 06/910)	Shrub, approximately 0.7m high.	Perennial	Pink flowers. September - October	Undulating plains, granite hills. Loams and sandy clay-loams.	Potential	Unlikely
<i>Baeckea</i> sp. Crossroads (B.L. Rye & M.E. Trudgen 241186)	Open straggly shrub to 1.0 m high.	Perennial	Pink flowers in November	Gentle slopes and small hills. Yellow-brown silty sands.	Potential	Unlikely
<i>Baeckea</i> sp. Forrestania (K.R. Newbey 1105)	Shrub, to 0.6 m high.	Perennial	No information available.	Sand. Plains.	Potential	Unlikely
<i>Baeckea</i> sp. Lake Cronin (K.R. Newbey 9191)	Upright, spreading, moderately open shrub.	Perennial	White/pink flowers in October.	Well-drained gravelly sands. Moderately exposed, gently undulating plain.	Likely	Unlikely
<i>Brachyloma nguba</i>	Erect, compact to spreading, mid-dense shrub, to 0.8 m high.	Perennial	Red flowers from April - May.	White to brown sandy clay, shallow sandy loam. Open mallee woodland, mallee scrub, flat plains.	Potential	Potential
<i>Brachyloma</i> sp. Forrestania White (M. Hislop & F. Hort MH 2591)	Erect shrub to approximately 1.5 m high.	Perennial	White flowers. May.	Plains, yellow sandy loams.	Potential	Unlikely
<i>Dampiera scaevolina</i>	Erect to ascending herb or shrub, 0.2-0.5 m high.	Perennial	Blue/white flowers from September - November.	Sandy and gravelly soils.	Unlikely	Unlikely
<i>Dicrastylis capitellata</i>	Low spreading shrub, 0.2-0.25 m high.	Perennial	Blue-purple flowers in May.	Loamy sand, sandy loam.	Unlikely	Unlikely
<i>Dillwynia</i> sp. Capel (P.A. Jurjevich 1771)	Erect, open, spreading shrub, to 2 m high.	Perennial	Yellow-orange-red - pink flowers from September - October.	Littered grey loamy sand, rocky soils. Valleys, rangelands.	Likely	Unlikely
<i>Drosera grieveri</i>	Fibrous-rooted herb, to 0.03 m high.	Perennial	White flowers in September.	Clayey sand.	Unlikely	Unlikely
<i>Eremophila adenotricha</i>	Erect, bushy, viscid shrub, 0.6-1.2 m high.	Perennial	Blue flowers in September.	Red/brown earth, clay.	Unlikely	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Eremophila lucida</i>	Shrub, to 1.8 m high.	Perennial	Cream-yellow flowers in October.	Clay loam, sandy loam. Adjacent to samphire flats and granite outcrops.	Unlikely	Unlikely
<i>Eucalyptus myriadena</i> subsp. <i>parviflora</i>	Mallee or tree, 3-10 m high, bark rough, coarse and flaky on trunk, smooth above.	Perennial	White flowers, April	Grey-whitish loams, brown sandy clays. Edges of swamps, slopes, plains.	Likely	Potential
<i>Eucalyptus phenax</i> subsp. <i>compressa</i>	Robust, erect (mallee), to 9 m high, bark rough, ribbony, tan to grey bark to 2m, smooth light grey over tan above.	Perennial	White flowers from September - December or January.	Gravelly clays to loams. Hilly sites.	Unlikely	Unlikely
<i>Hemigenia</i> sp. Newdegate (E. Bishop 75)	Spindly, erect to spreading shrub, 0.2-0.45 m high, to 0.5 m wide.	Perennial	Blue/purple flowers from September - October.	Upper slopes, Clay loam. Disturbed sites.	Potential	Unlikely
<i>Lepidosperma amantiferrum</i>	Tufted rhizomatous, herb (sedge), culms and leaves distichous.	Perennial	No information available.	Yellow sandy loam with banded ironstone gravel and rocks. Gentle lower slopes, broad flats	Potential	Unlikely
<i>Lepidosperma ferriculmen</i>	Tufted rhizomatous, herb (sedge), culms and leaves distichous.	Perennial	July	Well-drained orange-red sandy loam with banded ironstone gravel and rocks. Stony slopes.	Potential	Unlikely
<i>Leucopogon</i> sp. Yellowdine (M. Hislop & F. Hort MH 3194)	Shrub.	Perennial	White flowers.	Mixed heath, undulating sand plains, yellow sands.	Unlikely	Unlikely
<i>Stenanthemum liberum</i>	Dwarf shrub, ca 0.5 m high.	Perennial	No information available.	Slopes, Banded ironstone formation, yellow sandy loam over laterite.	Potential	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Stylidium validum</i>	Caespitose herb, 0.06-0.3 m high,	Perennial	White/pink flowers from September - October.	Clayey sand or loam, ironstone, greenstone gravel. Hill slopes and hilltops. Eucalypt woodland, mallee shrubland.	Potential	Unlikely
<i>Xanthoparmelia sargentii</i>	Prostrate Lichen	Lichen	No information available.	Granite outcrops	Unlikely	Unlikely
Priority 2						
<i>Acacia asepala</i>	Diffuse, much-branched shrub, 0.5-1.5 m high.	Perennial	Yellow flowers in August.	Red-brown sandy loam. Slopes, undulating plains, along drainage lines.	Likely	Unlikely
<i>Acacia gemina</i>	Erect, open shrub, 0.4-1.2 m high.	Perennial	Yellow flowers.	Lateritic gravelly soils, sandstone. Creek lines and gullies, scree slopes.	Potential	Unlikely
<i>Acacia kerryana</i>	Low, spreading, domed shrub, 0.3-1 m high.	Perennial	Yellow flowers from October - December or January - February.	Granitic loamy sand, stony clayey loam or clayey sand. Low stony ridges, undulating plains.	Potential	Unlikely
<i>Acacia tuberculata</i>	Diffuse, slender shrub, 0.3-1.8 m high.	Perennial	Yellow flowers in September.	Granite Outcrops.	Unlikely	Unlikely
<i>Baeckea</i> sp. North Ironcap (R.J. Cranfield 10580)	Erect, open shrub, to 0.4 m high.	Perennial	White/pink flowers in October.	Red clay. Slopes, Gently undulating sand plains.	Likely	Potential
<i>Boronia westringioides</i>	Erect shrub, 0.2-0.75 m high.	Perennial	Pink flowers from July - September.	Loamy sand. Plains.	Likely	Potential
<i>Conospermum sigmoideum</i>	Erect shrub, 0.2-0.5 m high.	Perennial	Blue flowers from August - September.	Yellow, white sands. Hillisides, plains, sand dunes.	Potential	Unlikely
<i>Eutaxia lasiocalyx</i>	Low, spreading, multi-stemmed shrub, to 0.15 m high.		Yellow flowers.	Red sandy loam, laterite and quartz gravel. Gentle lower slopes.	Potential	Unlikely
<i>Kulinia eludens</i>	Rhizomatous, tufted herb (sedge-like).	Perennial	Brown flowers in November.	Deep yellow sand. Openings in open shrubland.	Unlikely	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Olearia lacinifolia</i>	Erect, few-stemmed shrub, 0.6-1.2 m high.	Perennial	Blue/purple and white/yellow flowers from May - Sep.	White sand. Around playa lakes.	Unlikely	Unlikely
<i>Stypandra jamesii</i>	Effuse, spreading rhizomatous herb, to 0.3 m high.	Perennial	Cream-white flowers from October to November.	Shallow soils. Crevices and fissures in granite rocks, around edges of outcrops.	Unlikely	Unlikely
<i>Synaphea cervifolia</i>	Shrub, to 0.3 m high.	Perennial	Yellow flowers from June - October.	Sandy clay and gravel.	Unlikely	Unlikely
Priority 3						
<i>Acacia lanei</i>	Spreading shrub, 1.5-2.3 m high.	Perennial	Yellow flowers from July to September.	Clay, clay loam, gravelly loam. Along drainage lines and creeks.	Unlikely	Unlikely
<i>Acacia obesa</i>	Low spreading shrub, 0.3-0.6(-1) m high.	Perennial	Yellow flowers from July to September.	Yellow sand, gravelly loam.	Unlikely	Unlikely
<i>Acacia undosa</i>	Dense, spreading shrub, 0.3-1.5 m high.	Perennial	Yellow flowers from July to September.	Sandy clay loam, clayey sand. Undulating plains, low-lying areas.	Potential	Unlikely
<i>Baeckea</i> sp. Hyden (J.M. Brown 141)	Shrub, 0.25-0.75 m high.	Perennial	Pink-white flowers in May or August - October.	Sandy gravel, sandy clay, gravelly loam.	Unlikely	Unlikely
<i>Baeckea</i> sp. Parker Range (M. Hislop & F. Hort MH 2968)	Straggly, erect shrub, to 1.5 m high.	Perennial	Pink flowers in June.	Yellow to orange sandy loam with laterite gravel. Hill slopes, gently undulating sand plains, low ridges, road verges.	Likely	Potential
<i>Banksia rufa</i> subsp. <i>flavescens</i>	Prostrate, lignotuberous shrub, to 0.45 m high.	Perennial	Cream-yellow flowers from July to August.	Sandy loam or sand with gravel.	Potential	Potential
<i>Banksia viscida</i>	Densely branched, non-lignotuberous shrub, 0.4-1 m high.	Perennial	Yellow-orange flowers from July - October.	Gravelly soils. Lateritic rises.	Potential	Potential

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Bossiaea flexuosa</i>	Compact shrub, to 0.6 m high.	Perennial	Yellow-orange-red-brown flowers from September - November.	Deep sandy soil.	Potential	Unlikely
<i>Comesperma calcicola</i>	Soft herb, to 0.3 m high.	Perennial	Pink flowers from October - December or January.	Calcareous or semi-saline clay loams, limestone. Areas around saline water.	Unlikely	Unlikely
<i>Cryptandra dielsii</i>	Intricately branched, spreading shrub, 0.2-0.6 m high.	Perennial	White flowers from July - September.	Sand, often over laterite. Sand plains.	Unlikely	Unlikely
<i>Cryptandra polyclada</i> subsp. <i>polyclada</i>	Mat-forming or upright shrub, 0.1-0.7 m high.	Perennial	White/cream flowers from January - May or August - October.	Gravel, Sand. Sand plains.	Likely	Potential
<i>Daviesia elongata</i> subsp. <i>implexa</i>	Spreading or sprawling shrub, 0.4-1 m high.	Perennial	Yellow/orange & red flowers in September.	Sand and laterite.	Likely	Potential
<i>Elatine macrocalyx</i>	Prostrate, glabrous, mat-forming herb.	Annual	White flowers from May - October (probably opportunistic).	Shallow sands over clay. Margins of playa lakes and clay pans.	Unlikely	Unlikely
<i>Eucalyptus exigua</i>	Mallee, 2-5 m high, bark smooth.	Perennial	White-cream flowers in March.	Sandy loam, white sand. Sand plains.	Potential	Unlikely
<i>Eucalyptus ornata</i>	Tree, 6-10 m high, bark smooth, grey.	Perennial	White flowers.	Laterite. Ridges.	Unlikely	Unlikely
<i>Eutaxia acanthoclada</i>	Compact, mat-forming, prostrate shrub, to 0.3 m high.	Perennial	Yellow/orange/red, flowers from October - November.	Light brown sandy clay, shallow sandy loam, red clay over banded ironstone, gravel. Gently undulating plains.	Likely	Potential

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Eutaxia nanophylla</i>	Straggly, rounded shrub, to 0.35 m high.	Perennial	Yellow - orange - red flowers from October - November.	Clayey sand, red clay, stoney clayey loam. Low-lying areas, damp flats, slopes, undulating plains, low stony ridges.	Likely	Potential
<i>Eutaxia rubricarina</i>	Straggling shrub, to 0.5 m high.	Perennial	Orange and yellow and brown flowers in August or October.	Gravelly sand, grey to pinkish-white sandy clay, red loam. Flats, slopes, valley floors, road verges.	Likely	Potential
<i>Frankenia drummondii</i>	Prostrate shrub.	Perennial	White flowers.	Sand. Lake edges.	Unlikely	Unlikely
<i>Grevillea insignis</i> subsp. <i>elliottii</i>	Erect, bushy, non-lignotuberous shrub, 1-2 m high.	Perennial	Red/pink - cream - white flowers in October.	Gravelly sand or loam over ironstone. Hilltops or rises.	Potential	Unlikely
<i>Grevillea pilosa</i> subsp. <i>redacta</i>	Spreading to prostrate, non-lignotuberous shrub, 0.4-1.2 m high.	Perennial	Red flowers in February or October or December.	Amongst medium trees, or low trees, or low (sclerophyll) shrubland; in gravelly soil, or sand, or loam; occupying gravel rises.	Potential	Unlikely
<i>Hakea pendens</i>	Shrub, 2-3 m high, 2.5-3.1 m wide.	Perennial	Pink-white in September.	Stony loam. Ironstone ridges.	Unlikely	Unlikely
<i>Hibbertia pachyphylla</i>	Shrub, to 0.5 m high.	Perennial	Yellow flowers from September - November.	White to yellow sand, brown sandy gravel, gravelly loam, laterite, granite, quartz. Undulating plains, low rises, valley floors.	Potential	Potential
<i>Isolepis australiensis</i>	Grass-like or herb (sedge), 0.03-0.055 m high.	Annual	Flowers in June or September.	Silty sand, sandy clay. Lake margins, pools.	Unlikely	Unlikely
<i>Keraudrenia adenogyna</i>	Erect shrub, ca 0.4 m high.	Perennial	Purple-blue flowers in September.	Heavy loamy clay, loamy gravelly soils. Low-lying area.	Potential	Unlikely
<i>Melaleuca ochroma</i>	Shrub.	Perennial	Pale pink/mauve	Sandy loams and clays	Unlikely	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
			flowers in October/November.			
<i>Microseris scapigera</i>	Erect tuberous herb, 0.15-0.8 m high, summer-dormant.	Perennial	Yellow flowers from September - October.	Sandy soils. Margins of salt lakes, granite rock areas, samphire flats.	Unlikely	Unlikely
<i>Mirbelia densiflora</i>	Erect or straggling shrub, 0.2-1 m high.	Perennial	Yellow-orange flowers in October or January.	Stony loam, loamy sand. Small ridges, breakaways, undulating plains.	Unlikely	Unlikely
<i>Oxymyrrhine plicata</i>	Shrub.	Perennial			Unlikely	Unlikely
<i>Persoonia cymbifolia</i>	Erect, spreading shrub, 0.2-0.6(-1) m high.		Yellow flowers in December or January.	Sandy soils. On flats or in rock crevices.	Potential	Unlikely
<i>Phebalium brachycalyx</i>	Shrub, 0.4-1.5 m high.		Yellow-cream-white flowers from August - September.	Sand, gravelly soils. Lateritic uplands, hills.	Unlikely	Unlikely
<i>Phebalium drummondii</i>	Upright shrub, 0.6-1.5 m high.		Yellow flowers from July to September.	Gravelly sandy or clayey soils. Flats, roadsides.	Unlikely	Unlikely
<i>Pityrodia scabra</i> subsp. <i>dendrotricha</i>	Shrub.	Perennial	White flowers.	Low plains near water.	Potential	Unlikely
<i>Prostanthera nanophylla</i>	Shrub, 0.1-1 m high.		Blue-purple-white flowers from August to November.	Yellow sand over laterite, rocky loam. Sand plains.	Unlikely	Unlikely
<i>Pultenaea daena</i>	Dense, prostrate, domed shrub, to 0.07 m high.		Yellow flowers in March.	White to yellow sand or sandy loam, sandy or loamy clay, gravel, limestone, dolomite, laterite. Gently undulating plains, adjacent to salt lakes, in disturbed areas.	Likely	Potential

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Sphaerolobium validum</i>	Erect shrub, to 0.9 m high.	Perennial	Yellow and red flowers in September.	White-grey sand, red-brown clayey sand, laterite gravel and quartz pebbles. Gently undulating areas, flats, roadsides.	Potential	Potential
<i>Stylidium rhipidium</i>	Slender herb, ca 0.05 m high.	Annual	White flowers from October - November.	Sandy soils. Wet creek flats, swamps, granite outcrops.	Potential	Unlikely
<i>Stylidium sejunctum</i>	Caespitose herb, 0.25-0.45 m high.	Perennial	White/pink-purple flowers from September - November.	Clayey sand or loam, laterite. Outcrops, upper slopes, breakaways. Mallee and Allocasuarina shrubland.	Likely	Potential
<i>Thelymitra variegata</i>	Tuberous herb, 0.1-0.35 m high.	Perennial	Orange - red - purple - pink flowers from June - September	Sandy clay, sand, laterite.	Unlikely	Unlikely
<i>Thysanotus cymosus</i>	Caespitose herb (with fibrous roots with ellipsoidal tubers), to 0.3 m high.	Perennial	Purple flowers from September - October	Clay, granitic or lateritic sand.	Unlikely	Unlikely
<i>Verticordia gracilis</i>	Low, slender shrub, 0.15-0.6 m high.		Pink flowers from October - November.	Yellow sand, gravelly sand, sandy loam.	Likely	Potential
<i>Verticordia mitodes</i>	Spreading shrub, 0.15-0.7 m high.		Pink-purple flowers from October - December or January.	Yellow sand. Undulating plains.	Likely	Potential

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
Priority 4						
<i>Banksia shanklandiorum</i>	Upright, non-lignotuberous shrub, 0.4-2.5 m high, to 3 m wide.	Perennial	June - August.	White/yellow sand with lateritic gravel.	Unlikely	Unlikely
<i>Calamphoreus inflatus</i>	Erect, spreading shrub, 0.4-1.6 m high, to 2 m wide.		Blue-purple/green flowers from October - December or February - March.	Clay loam with ironstone gravel. Flats, slopes, disturbed sites.	Potential	Unlikely
<i>Daviesia oxylobium</i>	Glaucous shrub, 0.5-1 m high.	Perennial	Yellow - red - pink flowers from July - August.	Sandy lateritic soils. Undulating plains.	Unlikely	Unlikely
<i>Drosera graniticola</i>	Erect tuberous herb, 0.1-0.2 m high.	Perennial	White flowers from August - September.	Granite outcrops.	Unlikely	Unlikely
<i>Eremophila biserrata</i>	Prostrate shrub, to 3 m wide.	Perennial	Green/yellow-green flowers from September - November or March.	Sandy or sandy clay soils. Alluvial flats, salt flats and lakes.	Unlikely	Unlikely
<i>Eremophila racemosa</i>	Erect shrub, 0.5-1.7 m high.	Perennial	Purple-pink-red/white flowers in March or August - December.	Sandy or stony loam, clay loam. Hills, undulating plains, roadsides.	Potential	Unlikely
<i>Eremophila serpens</i>	Prostrate, creeping, forming large patches shrub, 0.03-0.4 m high, forming large patches to 2 m wide.	Perennial	Green/yellow-green flowers from September - December or March to May.	White/grey sand, alluvium, loam. Winter-wet depressions, sub-saline flats, drainage lines, salt lakes.	Unlikely	Unlikely

Species	Habit	Life Form	Flowering time	Habitat	Likelihood of Occurrence	
					Pre-survey	Post-survey
<i>Eucalyptus georgei</i> subsp. <i>fulgida</i>	Tree, 4-20 m high, bark smooth, often hanging in ribbons.	Perennial	Cream-white flowers.	Sandy loam, clayey sand. Slight depressions.	Potential	Unlikely
<i>Grevillea prostrata</i>	Loose, prostrate shrub, 0.04-0.1 m high, 0.8-1.2 m wide.	Perennial	Cream-white/pink-red flowers from August - December or January.	White, grey or yellow sand, gravel. Sand plains.	Likely	Potential
<i>Gyrostemon ditrigynus</i>	Shrub, 0.4-1.5 m high.		Creamy green flowers, May	Sand, sandy clay, loam. Plains, low ironstone ridges.	Potential	Potential
<i>Haegiela tatei</i>	Ascending to erect herb, 0.02-0.08(-0.2) m high.	Annual	White-yellow flowers from August to November.	Clay, sandy loam, gypsum. Saline habitats.	Unlikely	Unlikely
<i>Microcorys</i> sp. <i>Forrestania</i> (V. English 2004)	Prostrate or erect shrub, 0.35-0.4 m high.	Perennial	White/purple flowers in January or April.	Yellow sandy clay or red-brown clay. Open woodland or cleared areas.	Likely	Potential
<i>Sowerbaea multicaulis</i>	Tufted herb, 0.075-0.25 m high.	Perennial	Purple-violet flowers from October - December or January.	Yellow-brown sand.	Potential	Potential
Likely	Species previously recorded within 10 km of the survey area and suitable habitat appears to be present in the survey area.					
Potential	Species previously recorded within 10 - 20 km of the survey area and/or suitable habitat appears to be present in the survey area.					
Unlikely	No preferable habitat appears to be present in the survey area.					

Table C.2: Likelihood of occurrence of conservation significant fauna species listed as potentially occurring in the vicinity of the survey area (*NatureMap* (DPaW 2014e), DPaW Threatened Fauna Database (DPaW 2014f), PMST (DoE, 2014c).

Scientific name (common name)	Conservation codes			Preferred habitat	Likelihood of Occurrence
	EPBC	WC	DPaW		
Reptiles					
<i>Morelia spilota</i> subsp. <i>imbricata</i> (carpet python)		S4		Semi-arid coastal and inland habitats, banksia woodland, eucalypt woodlands, and grasslands and commonly utilises hollow logs for shelter.	High
<i>Paroplocephalus atriceps</i> (Lake Cronin snake)			P3	Semi-arid southern interior of Western Australia (Lake Cronin, Peak Eleanor and Maggi Hayes Hills).	High
Birds					
<i>Apus pacificus</i> (fork-tailed swift)	Mi	S3		Largely aerial species independent of the terrestrial environment.	Low
<i>Ardea modesta</i> (eastern great egret)	Mi			Shallow freshwaters and salt waters; rarely dry pastures.	Low
<i>Ardea ibis</i> (cattle egret)	Mi	S3		Largely wetland species however can exploit drier open habitats more than other heron species.	Low
<i>Charadrius rubricollis</i> (hooded plover)			P4	Inhabits sandy, ocean beaches, with highest densities on beaches with large amounts of washed up seaweed and open dunes. Also inhabits coastal and inland saltlakes.	Low
<i>Calidris acuminata</i> (sharp-tailed sandpiper)	Mi	S3		Muddy edges of shallow fresh/brackish wetlands with emergent sedges, saltmarsh, grass and low vegetation	Low
<i>Falco peregrinus</i> (peregrine falcon)		S4		Found throughout most of Australia across most habitat types, but especially areas with cliffs and rocky outcrops.	Moderate

Scientific name (common name)	Conservation codes			Preferred habitat	Likelihood of Occurrence
	EPBC	WC	DPaW		
<i>Ardeotis australis</i> (Australian bustard)			P4	Grassland, open woodland and open agricultural country across northern Australia.	Moderate
<i>Leipoa ocellata</i> (malleefowl)	Vu	S1		Largely confined to arid and semi-arid woodland that is dominated by mallee eucalypts on sandy soils with less than 430 mm of rainfall annually.	Moderate
<i>Merops ornatus</i> (rainbow bee-eater)	Mi	S3		Lightly wooded, preferably sandy country near water.	High
<i>Calyptorhynchus baudinii</i> (Baudin's cockatoo)	Vu	S1		Eucalypt forest, where it feeds on mainly marri seeds, flowers, nectar and buds. Also feed on seeds of <i>Eucalyptus</i> , <i>Hakea</i> , <i>Banksia</i> and pine species.	Moderate
<i>Calyptorhynchus latirostris</i> (Carnaby's cockatoo)	En	S1		Woodland or forest that contains live or dead trees of Salmon Gum, Wandoo, Tuart, Jarrah, Flooded Gum, Karri or Marri. Feeds on seeds, flowers and nectar of native proteaceous plant species (e.g. <i>Banksia</i> spp., <i>Dryandra</i> spp., <i>Grevillea</i> spp.), eucalypts and <i>Pinus</i> spp.	High
<i>Platycercus icterotis</i> subsp. <i>xanthenys</i> (western rosella (inland))			P4	Open eucalypt forest and timbered areas, including cultivated land and orchards. Also found in drier woodland, with a heath understorey.	High
<i>Pomatostomus superciliosus</i> subsp. <i>ashbyi</i> (white-browed babbler (western Wheatbelt))			P4	Dry sclerophyll woodlands with shrubby understorey, mulga, acacias, mallee, Cyprus pine scrubs, timber, scrub along watercourses and saltbush.	High
<i>Oreoica gutturalis</i> subsp. <i>gutturalis</i> (crested bellbird (southern))			P4	Most types of scrubs and thickets including open <i>Banksia</i> scrubs and heathland with emergent <i>Eucalyptus tottiana</i> .	High-recorded

Scientific name (common name)	Conservation codes			Preferred habitat	Likelihood of Occurrence
	EPBC	WC	DPaW		
<i>Falcunculus frontatus</i> subsp. <i>leucogaster</i> (crested shrike-tit (south-western))			P4	Open eucalypt forest and woodland.	Moderate
<i>Hylacola cauta</i> subsp. <i>whitlocki</i> (shy heathwren (western))			P4	Found in shrublands, including understorey of eucalypt woodlands and scrubs, post fire regeneration and uncleared road verges and remnants in farmlands. It favours scrubs with stony soils.	High
Mammals					
<i>Dasyurus geoffroii</i> (chuditch, western quoll)	Vu	S1		Wandoo and Salmon Gum woodland, mallee, Jarrah forest and mixed Marri/Jarrah forest.	High
<i>Macropus irma</i> (western brush wallaby)			P4	Open forest or woodland, particularly favouring open, seasonally wet flats with low grasses and open scrubby thickets.	Moderate
<i>Phascogale calura</i> (red-tailed phascogale)	En	S1		Wandoo and Sheoak woodland associations, with populations most dense in the latter vegetation type.	Low
Recorded	N/A				
High	Species has been recorded within the survey area or within 20 km of the survey area and preferred habitat appears to be present.				
Moderate	Species has not been recorded from within the survey area, however species has been recorded within 20 km of the survey area and suitable habitat appears to be present.				
Low	Species recorded within 20 km of the survey area but suitable habitat does not appear to be present.				

This page has been left blank intentionally.

Appendix D: Fauna Database Search and Literature Review Results

This page has been left blank intentionally.

Table D.1: List of amphibian fauna identified by the literature review and database searches as recorded within the vicinity of the survey area.

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D
			EPBC Act	WC Act	DPaW				
Limnodynastidae									
<i>Heleioporus albopunctatus</i>	Western spotted frog					x			
<i>Limnodynastes dorsalis</i>	Western banjo frog					x			
<i>Neobatrachus albipes</i>	White-footed trilling frog					x			
<i>Neobatrachus kunapalari</i>	Kunapalari frog					x			
<i>Neobatrachus pelobatoides</i>	Humming frog					x			
<i>Neobatrachus sutor</i>	Shoemaker frog					x			
Myobatrachidae									
<i>Crinia pseudinsignifera</i>	Bleating froglet					x		x	
<i>Pseudophryne occidentalis</i>	Western toadlet					x		x	
Key: A = Listed in NatureMap or DPaW Threatened Fauna Search; B = Listed in the EPBC Protected Search Matters Tool; C = Recorded in previous surveys within the survey area; D = Recorded in current assessment.									

Table D.2: List of reptilian fauna identified by the literature review and database searches as recorded within the vicinity of the survey area.

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D
			EPBC Act	WC Act	DPaW				
Agamidae									
<i>Ctenophorus adelaidensis</i>	Western heath dragon							X	
<i>Ctenophorus chapmani</i>	Eastern heath dragon					X			
<i>Ctenophorus cristatus</i>	Bicycle dragon					X		X	
<i>Ctenophorus maculatus</i>	Spotted military dragon					X		X	
<i>Ctenophorus maculatus badius</i>									
<i>Ctenophorus maculatus dualis</i>									
<i>Ctenophorus maculatus griseus</i>						X		X	
<i>Ctenophorus ornatus</i>	Ornate crevice dragon					X			
<i>Ctenophorus salinarum</i>	Salt pan dragon					X			
<i>Moloch horridus</i>	Thorny devil					X		X	
<i>Pogona minor</i>								X	
<i>Pogona minor minor</i>						X		X	
Diplodactylidae									
<i>Crenadactylus ocellatus</i>	Clawless gecko					X		X	
<i>Crenadactylus ocellatus ocellatus</i>						X		X	
<i>Diplodactylus calcicolus</i>	South coast gecko					X			
<i>Diplodactylus granariensis</i>						X		X	
<i>Diplodactylus granariensis granariensis</i>						X		X	
<i>Diplodactylus pulcher</i>						X		X	
<i>Hesperoedura reticulata</i>						X		X	
<i>Lucasium maini</i>						X		X	
<i>Strophurus spinigerus</i>								X	
<i>Strophurus spinigerus inornatus</i>						X			
Carphodactylidae									
<i>Underwoodisaurus milii</i>	Southern barking gecko					X		X	
Gekkonidae									

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D
			EPBC Act	WC Act	DPaW				
<i>Gehyra variegata</i>						x		x	
Pygopodidae									
<i>Delma australis</i>						x		x	
<i>Delma fraseri</i>						x		x	
<i>Lialis burtonis</i>						x		x	
<i>Pygopus lepidopodus</i>	Common scaly foot							x	
Scincidae									
<i>Cryptoblepharus buchananii</i>						x		x	
<i>Cryptoblepharus plagiocephalus</i>								x	
<i>Ctenotus atlas</i>						x			
<i>Ctenotus impar</i>						x		x	
<i>Ctenotus schomburgkii</i>						x		x	
<i>Cyclodomorphus melanops elongatus</i>						x			
<i>Egernia richardi</i>						x		x	
<i>Hemiergis initialis</i>						x		x	
<i>Hemiergis initialis brookeri</i>								x	
<i>Hemiergis initialis initialis</i>						x			
<i>Lerista distinguenda</i>						x		x	
<i>Lerista dorsalis</i>								x	
<i>Lerista picturata</i>						x		x	
<i>Liopholis multiscutata</i>						x		x	
<i>Menetia greyii</i>						x			
<i>Morethia butleri</i>						x			
<i>Morethia obscura</i>						x		x	
<i>Tiliqua occipitalis</i>	Western bluetongue					x		x	
<i>Tiliqua rugosa</i>						x		x	
Varanidae									
<i>Varanus gouldii</i>	Bungarra or sand monitor					x		x	
<i>Varanus rosenbergi</i>	Heath monitor							x	

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D
			EPBC Act	WC Act	DPaW				
Typhlopidae									
<i>Ramphotyphlops australis</i>						x		x	
<i>Ramphotyphlops bituberculatus</i>						x			
Boidae									
<i>Morelia spilota imbricata</i>	Carpet python			S4		x		x	
Elapidae									
<i>Neelaps bimaculatus</i>	Black-naped snake					x			
<i>Parasuta gouldii</i>						x		x	
<i>Parasuta nigriceps</i>						x		x	
<i>Paroplocephalus atriceps</i>	Lake Cronin snake				P3	x			
<i>Pseudonaja affinis</i>	Dugite					x		x	
<i>Pseudonaja affinis affinis</i>						x			
<i>Simoselaps bertholdi</i>	Jan's banded snake					x		x	
Key: A = Listed in NatureMap or DPaW Threatened Fauna Search; B = Listed in the EPBC Protected Search Matters Tool; C = Recorded in previous surveys within the survey area; D = Recorded in current assessment.									

Table D.3: List of avian fauna identified by the literature review and database searches as recorded within the vicinity of the survey area.

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
Casuariidae										
<i>Dromaius novaehollandiae</i>	Emu					X		X		X
Megapodiidae										
<i>Leipoa ocellata</i>	Malleefowl		VU	S1		X	X	X	X	
Anatidae										
<i>Biziura lobata</i>	Musk duck					X				
<i>Stictonetta naevosa</i>	Freckled duck					X				
<i>Cygnus atratus</i>	Black swan					X				
<i>Tadorna tadornoides</i>	Australian shelduck					X		X		
<i>Chenonetta jubata</i>	Australian wood duck					X				
<i>Malacorhynchus membranaceus</i>	Pink-eared duck					X				
<i>Anas rhynchotis</i>	Australasian shoveler					X				
<i>Anas gracilis</i>	Grey teal					X		X		
<i>Anas castanea</i>	Chestnut teal					X				
<i>Anas superciliosa</i>	Pacific black duck					X				
<i>Aythya australis</i>	Hardhead					X				
Podicipedidae										
<i>Tachybaptus novaehollandiae</i>	Australasian grebe					X				
<i>Poliiocephalus poliocephalus</i>	Hoary-headed grebe					X				
<i>Podiceps cristatus</i>	Great crested grebe					X				
Columbidae										
<i>Phaps chalconptera</i>	Common bronzewing					X		X	X	X
<i>Phaps elegans</i>	Brush bronzewing					X		X	X	
<i>Ocyphaps lophotes</i>	Crested pigeon					X		X		
Podargidae										
<i>Podargus strigoides</i>	Tawny frogmouth					X		X	X	
Eurostopodidae										
<i>Eurostopodus argus</i>	Spotted nightjar							X	X	X

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
Aegothelidae										
<i>Aegotheles cristatus</i>	Australian owl-nightjar							x	x	
Apodidae										
<i>Apus pacificus</i>	Fork-tailed swift			S3			x	x		
Ardeidae										
<i>Ardea modesta</i>	Eastern great egret			S3			x			
<i>Ardea ibis</i>	Cattle egret			S3			x			
<i>Egretta novaehollandiae</i>	White-faced Heron							x		
Threskiornithidae										
<i>Threskiornis molucca</i>	Australian white ibis					x				
<i>Threskiornis spinicollis</i>	Straw-necked Ibis					x				
Accipitridae										
<i>Lophoictinia isura</i>	Square-tailed kite							x	x	
<i>Haliastur spheunurus</i>	Whistling kite							x		
<i>Accipiter fasciatus</i>	Brown goshawk							x		
<i>Accipiter cirrocephalus</i>	Collared sparrowhawk					x		x	x	
<i>Aquila audax</i>	Wedge-tailed eagle					x		x	x	x
<i>Hieraetus morphnoides</i>	Little eagle							x		
Falconidae										
<i>Falco cenchroides</i>	Nankeen kestrel					x		x	x	
<i>Falco berigora</i>	Brown falcon					x		x	x	x
<i>Falco longipennis</i>	Australian hobby					x		x		
<i>Falco peregrinus</i>	Peregrine falcon			S4		x		x	x	
Rallidae										
<i>Fulica atra</i>	Eurasian coot					x				
Recurvirostridae										
<i>Cladorhynchus leucocephalus</i>	Banded stilt							x		
Charadriidae										
<i>Charadrius ruficapillus</i>	Red-capped plover							x		

Scientific Name	Common Name	Introduced	Conservation Codes				A	B	C	D	E
			EPBC Act	WC Act	DPaW						
<i>Thinornis (Charadrius) rubricollis</i>	Hooded plover				P4			x			
Scolopacidae											
<i>Calidris acuminata</i>	Sharp-tailed sandpiper			S3			x				
Turnicidae											
<i>Turnix velox</i>	Little button-quail						x			x	
Cacatuidae											
<i>Calyptorhynchus baudinii</i>	Baudin's black-cockatoo		VU	S1			x				
<i>Calyptorhynchus latirostris</i>	Carnaby's black-cockatoo		EN	S1			x	x	x	x	
<i>Eolophus roseicapillus</i>	Galah								x	x	x
<i>Nymphicus hollandicus</i>	Cockatiel						x				
Psittacidae											
<i>Glossopsitta porphyrocephala</i>	Purple-crowned lorikeet						x		x	x	
<i>Polytelis anthopeplus</i>	Regent parrot						x		x	x	
<i>Platycercus icterotis</i>	Western rosella						x		x	x	
<i>Platycercus icterotis xanthogenys</i>	Western rosella (Inland ssp)				P4		x				
<i>Barnardius zonarius</i>	Australian ringneck						x		x	x	x
<i>Psephotus varius</i>	Mulga parrot						x				
<i>Melopsittacus undulatus</i>	Budgerigar						x		x		
<i>Neophema elegans</i>	Elegant parrot						x		x	x	
Cuculidae											
<i>Chalcites basalus</i>	Horsfield's bronze-cuckoo								x		
<i>Chalcites osculans</i>	Black-eared cuckoo								x		
<i>Chalcites lucidus</i>	Shining bronze-cuckoo								x	x	
<i>Cacomantis pallidus</i>	Pallid cuckoo						x		x		
<i>Cacomantis flabelliformis</i>	Fan-tailed cuckoo						x		x	x	
Strigidae											
<i>Ninox novaeseelandiae</i>	Southern boobook						x		x	x	
Halcyonidae											
<i>Todiramphus pyrrhopygius</i>	Red-backed kingfisher						x		x		

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
<i>Todiramphus sanctus</i>	Sacred kingfisher					x		x		
Meropidae										
<i>Merops ornatus</i>	Rainbow bee-eater					x	x	x	x	
Climacteridae										
<i>Climacteris rufa</i>	Rufous treecreeper					x		x	x	
Maluridae										
<i>Malurus splendens</i>	Splendid fairy-wren					x				
<i>Malurus leucopterus</i>	White-winged fairy-wren					x		x		
<i>Malurus pulcherrimus</i>	Blue-breasted fairy-wren					x		x	x	x
<i>Stipiturus malachurus</i>	Southern emu-wren								x	
Acanthizidae										
<i>Sericornis frontalis</i>	White-browed scrubwren					x		x		
<i>Hylacola cauta</i>	Shy heathwren					x		x	x	
<i>Hylacola cauta whitlocki</i>	Shy heathwren (Western ssp)				P4	x			x	
<i>Calamanthus campestris</i>	Rufous fieldwren					x		x	x	
<i>Pyrrholaemus brunneus</i>	Redthroat					x		x	x	
<i>Smicrornis brevirostris</i>	Weebill					x		x	x	x
<i>Gerygone fusca</i>	Western gerygone					x		x	x	x
<i>Acanthiza chrysorrhoa</i>	Yellow-rumped thornbill					x		x	x	
<i>Acanthiza uropygialis</i>	Chestnut-rumped thornbill					x		x	x	x
<i>Acanthiza apicalis</i>	Inland thornbill					x		x	x	x
Pardalotidae										
<i>Pardalotus punctatus</i>	Spotted pardalote					x			x	
<i>Pardalotus striatus</i>	Striated pardalote					x		x	x	x
Meliphagidae										
<i>Certhionyx variegatus</i>	Pied honeyeater							x	x	
<i>Lichenostomus virescens</i>	Singing honeyeater							x	x	x
<i>Lichenostomus leucotis</i>	White-eared honeyeater					x		x	x	
<i>Lichenostomus cratitius</i>	Purple-gaped honeyeater					x		x	x	

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
<i>Lichenostomus ornatus</i>	Yellow-plumed honeyeater							x	x	
<i>Lichenostomus plumulus</i>	Grey-fronted honeyeater								x	
<i>Purnella albifrons</i>	White-fronted honeyeater							x	x	
<i>Manorina flavigula</i>	Yellow-throated miner					x		x	x	x
<i>Acanthagenys rufogularis</i>	Spiny-cheeked honeyeater					x		x	x	x
<i>Anthochaera lunulata</i>	Western wattlebird					x				
<i>Anthochaera carunculata</i>	Red wattlebird					x		x	x	x
<i>Epthianura albifrons</i>	White-fronted chat					x				
<i>Sugomel niger</i>	Black honeyeater							x		
<i>Glyciphila melanops</i>	Tawny-crowned honeyeater							x	x	x
<i>Lichmera indistincta</i>	Brown honeyeater					x		x	x	x
<i>Phylidonyris novaehollandiae</i>	New Holland honeyeater					x				
<i>Phylidonyris niger</i>	White-cheeked honeyeater							x	x	
<i>Melithreptus brevirostris</i>	Brown-headed honeyeater					x		x	x	
Pomatostomidae										
<i>Pomatostomus superciliosus</i>	White-browed babbler					x		x	x	
<i>Pomatostomus superciliosus</i> subsp. <i>ashbyi</i>	White-browed babbler (Western wheatbelt)				P4	x				
Eupetidae										
<i>Cinclosoma castanotum</i>	Chestnut quail-thrush							x	x	
Neosittidae										
<i>Daphoenositta chrysoptera</i>	Varied sittella					x		x	x	
Campephagidae										
<i>Coracina novaehollandiae</i>	Black-faced cuckoo-shrike					x		x	x	x
<i>Lalage sueurii</i>	White-winged triller							x	x	
Pachycephalidae										
<i>Falcunculus frontatus leucogaster</i>	Crested shrike-tit (southwestern ssp.)				P4			x		
<i>Pachycephala inornata</i>	Gilbert's whistler					x		x	x	

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
<i>Pachycephala pectoralis</i>	Golden whistler					x		x	x	
<i>Pachycephala rufiventris</i>	Rufous whistler					x		x		
<i>Colluricincla harmonica</i>	Grey shrike-thrush					x		x	x	
<i>Oreoica gutturalis</i>	Crested bellbird					x		x	x	
<i>Oreoica gutturalis</i> subsp. <i>gutturalis</i>	Crested bellbird (southern ssp.)				P4	x				x
Artamidae										
<i>Artamus personatus</i>	Masked woodswallow					x		x		
<i>Artamus cinereus</i>	Black-faced woodswallow					x		x	x	
<i>Artamus cyanopterus</i>	Dusky woodswallow					x		x	x	
<i>Cracticus torquatus</i>	Grey butcherbird					x		x	x	
<i>Cracticus nigrogularis</i>	Pied butcherbird					x		x	x	
<i>Cracticus tibicen</i>	Australian magpie					x		x	x	x
<i>Strepera versicolor</i>	Grey currawong					x		x	x	
Rhipiduridae										
<i>Rhipidura albiscapa</i>	Grey fantail					x		x	x	
<i>Rhipidura leucophrys</i>	Willie wagtail					x		x	x	x
Corvidae										
<i>Corvus coronoides</i>	Australian raven					x		x	x	x
<i>Corvus bennetti</i>	Little crow							x	x	
<i>Corvus orru</i>	Torresian crow							x		
Monarchidae										
<i>Myiagra inquieta</i>	Restless flycatcher					x				x
<i>Grallina cyanoleuca</i>	Maggie-lark					x				x
Petroicidae										
<i>Microeca fascians</i>	Jacky winter					x		x	x	
<i>Petroica goodenovii</i>	Red-capped robin					x		x		
<i>Melanodryas cucullata</i>	Hooded robin							x	x	
<i>Eopsaltria griseogularis</i>	Western yellow robin					x		x	x	

Scientific Name	Common Name	Introduced	Conservation Codes			A	B	C	D	E
			EPBC Act	WC Act	DPaW					
<i>Drymodes brunneopygia</i>	Southern scrub-robin					x		x	x	
Megaluridae										
<i>Cincloramphus mathewsi</i>	Rufous songlark					x		x		
Timaliidae										
<i>Zosterops lateralis</i>	Silvereye					x		x	x	
Hirundinidae										
<i>Hirundo neoxena</i>	Welcome swallow					x		x		
<i>Petrochelidon nigricans</i>	Tree martin							x	x	x
Nectariniidae										
<i>Dicaeum hirundinaceum</i>	Mistletoebird							x		
Estrildidae										
<i>Taeniopygia guttata</i>	Zebra finch					x				
Motacillidae										
<i>Anthus novaeseelandiae</i>	Australasian pipit							x	x	x
<i>Anthus cervinus</i>	Red-throated pipit					x				
Key: A = Listed in NatureMap or DPaW Threatened Fauna Search; B = Listed in the EPBC Protected Search Matters Tool; C = Listed by Birdlife Birddata; D = Recorded in previous surveys within the survey area; E = Recorded in current assessment.										

Table D.4: List of mammalian fauna identified by the literature review and database searches as recorded within the vicinity of the survey area.

Scientific Name	Common name	Introduced	Conservation Codes				A	B	C	D
			EPBC Act	WC Act	DPaW					
Tachyglossidae										
<i>Tachyglossus aculeatus</i>	Short-beaked echidna								X	
Dasyuridae										
<i>Dasyurus geoffroii</i>	Western quoll, chuditch		VU	S1		X	X	X		
<i>Ningaui yvonneae</i>	Southern ningau					X				
<i>Phascogale calura</i>	Red-tailed phascogale		EN	S1			X			
<i>Sminthopsis crassicaudata</i>	Fat-tailed dunnart					X		X		
<i>Sminthopsis gilberti</i>	Gilbert's dunnart					X		X		
<i>Sminthopsis granulipes</i>	White-tailed dunnart					X		X		
<i>Sminthopsis griseoventer</i>	Grey-bellied dunnart							X		
Macropodidae										
<i>Macropus fuliginosus</i>	Western grey kangaroo					X		X		X
<i>Macropus irma</i>	Western brush wallaby				P4	X		X		
Burramyidae										
<i>Cercartetus concinnus</i>	Western pygmy-possum, mundarda					X		X		
Tarsipedidae										
<i>Tarsipes rostratus</i>	Honey possum, noolbenger							X		
Vespertilionidae										
<i>Chalinolobus gouldii</i>	Gould's wattled bat					X		X		
<i>Chalinolobus morio</i>	Chocolate wattled bat					X		X		
<i>Nyctophilus geoffroyi</i>	Lesser long-eared bat					X		X		
<i>Vespadelus regulus</i>	Southern forest bat					X		X		
Molossidae										
<i>Mormopterus planiceps</i>	Southern freetail-bat					X		X		
<i>Tadarida australis</i>	White-striped freetail-bat							X		
Muridae										
<i>Mus musculus</i>	House mouse	*				X	X	X		
<i>Notomys mitchelli</i>	Mitchell's hopping-mouse					X		X		

Scientific Name	Common name	Introduced	Conservation Codes			A	B	C	D
			EPBC Act	WC Act	DPaW				
<i>Pseudomys albocinereus</i>	Ash-grey mouse					x		x	
Leporidae									
<i>Oryctolagus cuniculus</i>	European rabbit	*				x	x	x	
Canidae									
<i>Canis lupus familiaris</i>	Dog	*						x	
<i>Vulpes vulpes</i>	Red fox	*				x	x	x	
Felidae									
<i>Felis catus</i>	Cat	*				x	x	x	
Suidae									
<i>Sus scrofa</i>	Pig	*					x		
Camelidae									
<i>Camelus dromedarius</i>	Dromedary, camel	*					x		

Key: A = Listed in NatureMap or DPaW Threatened Fauna Search; B = Listed in the EPBC Protected Search Matters Tool; C = Recorded in previous surveys within the survey area; D = Recorded in current assessment.

This page has been left blank intentionally.

Appendix E: Vegetation Association Mapping and Priority Flora Location

This page has been left blank intentionally.

751000

751500

752000

752500

Vegetation Mapping

Sandplain Heath

SpH1

Banksia cirsioides open low scrub over mixed dwarf scrub.

Associated species: *Petrohile ericifolia* subsp. *ericifolia*, *Beaufortia interstans*, *Hakea corymbosa*, *Grevillea cagiana*, *Allocasuarina microstachya*, *Verticordia chrysanthella*, *Verticordia inclusa*, *Melaleuca seriata*, *Mesomelaena preissii*.

Eucalyptus Mallee Woodlands

EMW1

Eucalyptus eremophila open tree mallee over mixed *Melaleuca* species open dwarf scrub to heath.


Associated species: *Eucalyptus platycorys*, *Melaleuca hamata*, *Melaleuca eleuterostachya*, *Melaleuca adnata*, *Melaleuca glaberrima*, *Dodonaea bursarifolia*, *Phebalium tuberculatum*, *Lepidosperma viscidum*.

EMW2

Eucalyptus platycorys open tree mallee to tree mallee over *Melaleuca hamata* open low scrub to low scrub over mixed open dwarf scrub.

Associated species: *Eucalyptus rigidula*, *Melaleuca seriata*, *Melaleuca glaberrima*, *Leptospermum erubescens*, *Gastrolobium spinosum*, *Hibbertia gracilipes*, *Lepidosperma viscidum*.

Cleared



6409500

6409000

6408500

6408000

Legend

— Haul Road

Survey Area

Lake Cronin ESA

Priority Flora

+ *Eutaxia hirsuta* (P2)

TEC/PEC Boundaries

Ironcap Hills vegetation complexes (Mt Holland, Middle, North and South Ironcap Hills, Digger Rock and Hatter Hill) (banded ironstone formation)

Eutaxia hirsuta (P2)

SpH1

SpH1

EMW2

EMW2

EMW2

EMW1

EMW1

Western Areas - Forrestania Nickel Operations
Lounge Lizard Vegetation, Flora and Fauna Biological Assessment

Appendix E: Vegetation Association Mapping and Priority Flora Location

Author: N. Cadd

Date: 10-11-2014

Drawn: C. Dyde

Figure Ref: 17603-14-BIDR-1RevA_141110_AppE

Datum: GDA 1994 - Projection: MGA Zone 50 - Scale: 1:7,000 (A3)

0 100 200 300 400 500 Metres



astron
environmental services

This page has been left blank intentionally

This page has been left blank intentionally.

WSA Forrestania**Site FOR01****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 14/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751662 mE**Northing:** 6409206 mN**Habitat:** Gently undulating sandplain.**Soil:** Yellow beige sands.**Rock Type:** N/A.**Vegetation:** *Banksia cirsioides* open low scrub over mixed dwarf scrub.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Species List**

Name	Cover (%)	Height (m)
? <i>Stylidium clavatum</i>	+	0.20
<i>Acacia acoma</i>	+	0.20
<i>Adenanthos argyreus</i>	+	0.30
<i>Allocasuarina humilis</i>	+	1.00
<i>Allocasuarina microstachya</i>	2	0.30
<i>Allocasuarina spinosissima</i>	+	0.60
<i>Aotus</i> sp. Tortile (G.J. Keighery 3767)	+	0.40
<i>Banksia cirsioides</i>	1	1.00
<i>Banksia laevigata</i> subsp. <i>fuscolutea</i>	+	0.60
<i>Banksia violacea</i>	+	0.80
<i>Beaufortia interstans</i>	4	0.40
<i>Boronia coerulescens</i> subsp. <i>spicata</i>	+	0.20
<i>Cassytha glabella</i>	+	climber
<i>Caustis dioica</i>	+	0.15
<i>Comesperma scoparium</i>	+	0.40
<i>Conospermum brownii</i>	+	0.50

<i>Eutaxia hirsuta</i>	+	0.15
<i>Grevillea cagiana</i>	1	0.80
<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>	+	0.90
<i>Hakea corymbosa</i>	1	0.50
<i>Hemigenia westringioides</i>	+	0.30
<i>Isopogon scabriusculus</i>	+	0.45
<i>Lepidobolus preissianus</i>	+	0.15
<i>Lepidosperma</i> sp.	+	0.30
<i>Leptospermum fastigiatum</i>	+	1.20
<i>Leptospermum spinescens</i>	+	0.60
<i>Lysinema ciliatum</i>	+	0.50
<i>Melaleuca cordata</i>	+	0.60
<i>Melaleuca seriata</i>	1	0.50
<i>Mesomelaena preissii</i>	1	0.30
<i>Neurachne alopecuroidea</i>	+	0.05
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	4	0.40
<i>Petrophile stricta</i>	+	1.00
<i>Schoenus</i> sp.	+	0.08
<i>Stylidium</i> sp.	+	0.04
<i>Synaphea interioris</i>	+	0.15
<i>Tetratheca efoliata</i>	+	0.15
<i>Verticordia chrysanthella</i>	2	0.50
<i>Verticordia inclusa</i>	+	0.40
<i>Verticordia picta</i>	+	0.30

WSA Forrestania**Site FOR02****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 14/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751281 mE**Northing:** 6409186 mN**Habitat:** Open sandplain.**Soil:** Beige sands with scattered pea gravel.**Rock Type:** Scattered surface pea gravel.**Vegetation:** *Banksia cirsioides* open low scrub over mixed dwarf scrub.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 10-20%.

Occasional patches of *Eucalyptus ?subangusta* subsp. *angusta*, *Melaleuca hamata* and *Melaleuca glaberrima* occurred in this vegetation association.

**Species List**

Name	Cover (%)	Height (m)
<i>?Stylidium clavatum</i>	+	0.10
<i>Acacia acoma</i>	+	0.20
<i>Allocasuarina microstachya</i>	2	0.25
<i>Allocasuarina spinosissima</i>	+	0.40
<i>Aotus</i> sp. Tortile (G.J. Keighery 3767)	+	0.30
<i>Banksia cirsioides</i>	1	1.00
<i>Beaufortia interstans</i>	5	0.40
<i>Callitris columellaris</i>	+	0.40
<i>Conospermum brownii</i>	+	0.10
<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>	+	0.90
<i>Hakea corymbosa</i>	4	0.60
<i>Hakea platysperma</i>	1	0.60

<i>Halgania lavandulacea</i>	+	0.10
<i>Hemigenia westringioides</i>	+	0.10
<i>Isopogon scabriusculus</i>	+	0.60
<i>Leptospermum fastigiatum</i>	+	1.00
<i>Leucopogon dielsianus</i>	+	0.30
<i>Melaleuca cordata</i>	+	0.30
<i>Melaleuca seriata</i>	1	0.40
<i>Mesomelaena preissii</i>	1	0.20
<i>Neurachne alopecuroidea</i>	+	0.05
<i>Petrophile ?divaricata</i>	+	0.30
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	1	0.45
<i>Petrophile stricta</i>	1	0.50
<i>Scaevola restiacea</i>	+	0.30
<i>Schoenus</i> sp.	+	0.05
<i>Stylidium</i> sp.	+	0.05
<i>Synaphea interioris</i>	+	0.10
<i>Tetratheca efoliata</i>	+	0.15
<i>Thysanotus</i> sp.	+	climber
<i>Verticordia chrysanthella</i>	+	0.40
<i>Verticordia inclusa</i>	1	0.40
<i>Verticordia picta</i>	+	0.60

WSA Forrestania**Site FOR03****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 15/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751142 mE**Northing:** 6408372 mN**Habitat:** Very gently undulating/sloping plain.**Soil:** Grey-beige sand.**Rock Type:** N/A.**Vegetation:** *Eucalyptus eremophila* and *Eucalyptus platycorys* open tree mallee over *Melaleuca hamata* and *Melaleuca eleuterostachya* low scrub over *Melaleuca glaberrima* open dwarf scrub with *Lepidosperma viscidum* very open tall sedges.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 20-50%.**Species List**

Name	Cover (%)	Height (m)
<i>Allocasuarina spinosissima</i>	+	0.40
<i>Baeckea</i> sp.	+	0.40
<i>Beyeria sulcata</i> ? var.	+	0.45
<i>Callitris columellaris</i>	+	1.00
<i>Cassytha glabella</i>	+	climber
<i>Darwinia</i> sp. Lake Cobham (K. Newbey 3262)	+	0.20
<i>Daviesia lancifolia</i>	+	0.40
<i>Dodonaea viscosa</i> subsp. <i>spatulata</i>	+	0.60
<i>Eucalyptus eremophila</i>	10	2.20
<i>Eucalyptus platycorys</i>	5	1.70
<i>Eucalyptus rigidula</i>	+	1.80
<i>Gahnia ancistrophylla</i>	+	0.15
<i>Gastrolobium spinosum</i>	+	0.40

<i>Grevillea huegelii</i>	+	0.70
<i>Guichenotia micrantha</i>	+	0.35
<i>Hakea commutata</i>	+	0.60
<i>Lepidosperma viscidum</i>	1	0.65
<i>Melaleuca eleuterostachya</i>	2	1.40
<i>Melaleuca glaberrima</i>	5	0.40
<i>Melaleuca hamata</i>	20	1.10
<i>Melaleuca laxiflora</i>	+	0.60
<i>Melaleuca</i> sp.	+	0.40
<i>Phebalium tuberosum</i>	1	0.30
<i>Santalum acuminatum</i>	+	1.50
<i>Verticordia plumosa</i> var. <i>incrassata</i>	+	0.40

WSA Forrestania**Site FOR04****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 15/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751421 mE**Northing:** 6408407 mN**Habitat:** Gently sloping plain.**Soil:** Greyish orange (pale) sands.**Rock Type:** Pea gravel pebbles - very scattered.**Vegetation:** *Eucalyptus platycorys* tree mallee over *Melaleuca hamata* low scrub over *Melaleuca seriata* and *Melaleuca glaberrima* open dwarf scrub with *Lepidosperma viscidum* very open tall sedges.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 20-50%.**Species List**

Name	Cover (%)	Height (m)
<i>Banksia cirsioides</i>	+	0.50
<i>Callitris columellaris</i>	+	0.70
<i>Calytrix</i> sp.	+	0.40
<i>Cassytha glabella</i>	+	climber
<i>Darwinia</i> sp. Lake Cobham (K. Newbey 3262)	+	0.20
<i>Daviesia lancifolia</i>	+	0.30
<i>Desmocladius flexuosus</i>	+	0.10
<i>Eucalyptus platycorys</i>	26	2.20
<i>Eucalyptus rigidula</i>	+	2.00
<i>Gahnia ancistrophylla</i>	+	0.40
<i>Gastrolobium spinosum</i>	+	0.50
<i>Grevillea oncogyne</i>	+	1.10
<i>Guichenotia micrantha</i>	+	0.30

<i>Hibbertia gracilipes</i>	+	0.30
<i>Lepidosperma viscidum</i>	1	0.50
<i>Leptomeria preissiana</i>	+	1.20
<i>Lomandra mucronata</i>	+	0.10
<i>Lysinema ciliatum</i>	+	0.60
<i>Melaleuca glaberrima</i>	1	0.40
<i>Melaleuca hamata</i>	10	1.0
<i>Melaleuca seriata</i>	6	0.40
<i>Neurachne alopecuroidea</i>	+	0.03
<i>Olax benthamiana</i>	+	0.40
<i>Petrophile ?divaricata</i>	+	0.40
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	+	0.30
<i>Phebalium tuberosum</i>	+	0.40
<i>Verticordia chrysanthella</i>	+	0.40
<i>Verticordia inclusa</i>	+	0.40
<i>Verticordia plumosa</i> var. <i>incrassata</i>	+	0.45

WSA Forrestania**Site FOR05****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 15/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751849 mE**Northing:** 6408430 mN**Habitat:** Gently sloping sandplain.**Soil:** Beige - fawn fine loamy sand.**Rock Type:** Fine scattered pea gravel pebbles.**Vegetation:** *Eucalyptus platycorys* open tree mallee over *Melaleuca hamata* and *Leptospermum erubescens* low scrub over *Melaleuca glaberrima*, *Gastrolobium spinosum* and *Hibbertia gracilipes* dwarf scrub with *Lepidosperma viscidum* very open low sedges.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 50-90%.**Species List**

Name	Cover (%)	Height (m)
<i>Allocasuarina spinosissima</i>	+	1.10
<i>Banksia cirsioides</i>	+	0.80
<i>Calytrix</i> sp.	+	0.40
<i>Darwinia</i> sp. Lake Cobham (K. Newbey 3262)	+	0.20
<i>Daviesia benthamii</i>	+	1.00
<i>Daviesia lancifolia</i>	+	0.30
<i>Eucalyptus eremophila</i>	+	2.20
<i>Eucalyptus platycorys</i>	15	1.50
<i>Gastrolobium spinosum</i>	1	0.40
<i>Grevillea cagiana</i>	+	1.00
<i>Guichenotia micrantha</i>	+	0.30
<i>Hakea multilineata</i>	+	1.40
<i>Hibbertia gracilipes</i>	1	0.30

<i>Hibbertia pungens</i>	+	0.15
<i>Isopogon scabriusculus</i>	+	0.50
<i>Lepidosperma viscidum</i>	1	0.40
<i>Leptomeria preissiana</i>	+	1.00
<i>Leptospermum erubescens</i>	1	1.00
<i>Leucopogon</i> sp.	+	0.20
<i>Lysinema ciliatum</i>	+	0.60
<i>Melaleuca glaberrima</i>	4	0.35
<i>Melaleuca hamata</i>	15	1.0
<i>Micromyrtus erichsenii</i>	+	0.40
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	+	0.45
<i>Phebalium tuberosum</i>	+	0.20
<i>Stylidium</i> sp.	+	0.03
<i>Verticordia chrysanthella</i>	+	0.40

WSA Forrestania**Site FOR06****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 15/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751727 mE**Northing:** 6408117 mN**Habitat:** Undulating sandplain.**Soil:** Orange-brown loamy sand.**Rock Type:** N/A.**Vegetation:** *Eucalyptus eremophila* open tree mallee over *Melaleuca hamata* and *Melaleuca adnata* heath.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 20-50%.**Species List**

Name	Cover (%)	Height (m)
<i>Astartea aspera</i>	+	0.60
<i>Daviesia benthamii</i>	+	1.00
<i>Dodonaea bursariifolia</i>	1	0.50
<i>Eucalyptus eremophila</i>	12	3.00
<i>Hakea commutata</i>	+	1.10
<i>Lepidosperma viscidum</i>	+	0.40
<i>Melaleuca adnata</i>	5	1.10
<i>Melaleuca eleuterostachya</i>	+	1.00
<i>Melaleuca hamata</i>	40	1.50
<i>Melaleuca lateriflora</i>	+	1.00
<i>Melaleuca laxiflora</i>	+	0.90
<i>Phebalium tuberosum</i>	+	0.50
<i>Santalum acuminatum</i>	+	1.10

WSA Forrestania**Site FOR07****Location:** Lounge lizard sand pit area.**Type:** Quadrat 20 x 20m**Date:** 21/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 752265 mE**Northing:** 6408463 mN**Habitat:** Gently sloping sandplain.**Soil:** Pale orange/yellow/grey sands.**Rock Type:** N/A.**Vegetation:** *Eucalyptus platycorys* open tree mallee over *Melaleuca hamata* low scrub.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 50-90%.**Species List**

Name	Cover (%)	Height (m)
<i>Banksia cirsioides</i>	+	0.80
<i>Beaufortia interstans</i>	+	0.40
<i>Darwinia</i> sp.	+	0.20
<i>Daviesia lancifolia</i>	+	0.40
<i>Eucalyptus platycorys</i>	10	2.10
<i>Eucalyptus rigidula</i>	1	1.10
<i>Gahnia ancistrophylla</i>	+	0.40
<i>Gastrolobium spinosum</i>	+	0.40
<i>Gompholobium viscidulum</i>	+	0.10
<i>Grevillea cagiana</i>	+	
<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>	+	0.70
<i>Guichenotia micrantha</i>	+	0.40
<i>Hakea corymbosa</i>	+	0.50
<i>Lepidosperma viscidum</i>	+	0.60

<i>Leptomeria preissiana</i>	+	1.00
<i>Leptospermum fastigiatum</i>	+	0.60
<i>Lysinema ciliatum</i>	+	0.70
<i>Melaleuca hamata</i>	30	1.10
<i>Melaleuca seriata</i>	+	0.50
<i>Phebalium tuberosum</i>	+	0.20
<i>Verticordia chrysanthella</i>	+	0.50

WSA Forrestania**Site****FORR01****Location:** Lounge lizard sand pit area.**Type:** Relevé**Date:** 16/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751245 mE**Northing:** 6408833 mN**Habitat:** Crest of very low sandplain rise.**Soil:** Yellow-grey sands.**Rock Type:** N/A.**Vegetation:** *Banksia cirsioides* open low scrub over mixed open dwarf scrub.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Notes:** Bare ground: 50-90%.**Species List**

Name	Cover (%)	Height (m)
<i>Adenanthos argyreus</i>	+	0.20
<i>Allocasuarina microstachya</i>	2	0.20
<i>Allocasuarina thuyoides</i>	+	0.80
<i>Banksia cirsioides</i>	2	1.00
<i>Banksia erythrocephala</i>	+	0.80
<i>Banksia laevigata</i> subsp. <i>fuscolutea</i>	+	0.80
<i>Beaufortia interstans</i>	1	0.40
<i>Grevillea cagiana</i>	+	0.50
<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>	+	0.80
<i>Hakea corymbosa</i>	1	0.50
<i>Isopogon scabriusculus</i>	+	0.50
<i>Lepidobolus preissianus</i>	+	0.15
<i>Leptospermum spinescens</i>	+	0.50
<i>Lysinema ciliatum</i>	+	0.90

<i>Melaleuca cordata</i>	+	0.50
<i>Melaleuca seriata</i>	1	0.40
<i>Mesomelaena preissii</i>	1	0.20
<i>Neurachne alopecuroidea</i>	+	0.05
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	2	0.30
<i>Schoenus</i> sp.	+	0.05
<i>Synaphea interioris</i>	+	0.20
<i>Verticordia chrysanthella</i>	+	0.50
<i>Verticordia inclusa</i>	+	0.40

WSA Forrestania**Site****FORR02****Location:** Lounge lizard sand pit area.**Type:** Relevé**Date:** 16/10/2014**Described by:** NC/SS**Seasonal Conditions:** Excellent**MGA Zone:** 50**Easting:** 751498 mE**Northing:** 6408689 mN**Habitat:** Sandplain.**Soil:** Pale grey-yellow/orange sands.**Rock Type:** N/A.**Vegetation:** *Eucalyptus platycorys* tree mallee over *Melaleuca hamata* open low scrub.**Veg Condition:** Excellent.**Fire Age:** >10 years.**Species List**

Name	Cover (%)	Height (m)
<i>Acacia sphacelata</i> subsp. <i>sphacelata</i>	+	0.50
<i>Banksia cirsioides</i>	+	0.80
<i>Calytrix</i> sp.	+	0.50
<i>Desmocladius flexuosus</i>	+	0.15
<i>Eucalyptus platycorys</i>	25	2.20
<i>Gahnia ancistrophylla</i>	+	0.20
<i>Gompholobium viscidulum</i>	+	0.15
<i>Grevillea cagiana</i>	+	0.90
<i>Hibbertia gracilipes</i>	1	0.35
<i>Lepidosperma viscidum</i>	+	0.70
<i>Leptospermum fastigiatum</i>	+	1.10
<i>Lomandra mucronata</i>	+	0.10
<i>Lysinema ciliatum</i>	+	0.70
<i>Melaleuca hamata</i>	8	1.10
<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	+	0.30

Verticordia inclusa

+

0.50

WSA Forrestania

Site

OPP

Location: Lounge lizard sand pit area.

Type: Opportunistic Collections

Date: 20/10/2014

Species List


Name

<i>Acacia assimilis</i> subsp. <i>assimilis</i>	<i>Grevillea oncogyne</i>
<i>Acacia burkittii</i>	<i>Hakea corymbosa</i>
<i>Acacia erinacea</i>	<i>Hakea platysperma</i>
<i>Acacia poliochroa</i>	<i>Halgania lavandulacea</i>
<i>Acacia sphacelata</i> subsp. <i>sphacelata</i>	<i>Lachnostachys bracteosa</i>
<i>Allocasuarina acutivalvis</i> subsp. <i>acutivalvis</i>	<i>Laxmannia paleacea</i>
<i>Allocasuarina thuyoides</i>	<i>Leptomeria preissiana</i>
<i>Amphipogon turbinatus</i>	<i>Leucopogon</i> sp. Coujinup (M.A. Burgman 1085)
<i>Aotus</i> sp. Tortile (G.J. Keighery 3767)	<i>Logania tortuosa</i>
<i>Astroloma serratifolium</i>	<i>Lomandra effusa</i>
<i>Austrostipa hemipogon</i>	<i>Lomandra mucronata</i>
<i>Banksia elderiana</i>	<i>Melaleuca glaberrima</i>
<i>Banksia erythrocephala</i>	<i>Melaleuca hamata</i>
<i>Beaufortia micrantha</i> var. <i>micrantha</i>	<i>Melaleuca lateriflora</i>
<i>Callitris columellaris</i>	<i>Microcorys exserta</i>
<i>Conospermum stoechadis</i> subsp. <i>sclerophyllum</i>	<i>Neurachne alopecuroidea</i>
<i>Conostylis petrophiloides</i>	<i>Olearia ramosissima</i>
<i>Dampiera juncea</i>	<i>Patersonia rudis</i> subsp. <i>velutina</i>
<i>Daviesia benthamii</i>	<i>Petrophile ?divaricata</i>
<i>Daviesia benthamii</i> subsp. <i>acanthoclona</i>	<i>Petrophile stricta</i>
<i>Desmocladius flexuosus</i>	<i>Pimelea suaveolens</i>
<i>Dianella revoluta</i> var. <i>divaricata</i>	<i>Pimelea subvillifera</i>
<i>Eremophila scoparia</i>	<i>Santalum murrayanum</i>
<i>Eucalyptus ?subangusta</i> subsp. <i>subangusta</i>	<i>Templetonia rossii</i>
<i>Eucalyptus platycorys</i>	<i>Verticordia picta</i>
<i>Eutaxia neurocalyx</i> subsp. <i>papillosa</i>	<i>Verticordia plumosa</i> var. <i>incrassata</i>
<i>Exocarpos aphyllus</i>	<i>Verticordia roei</i> subsp. <i>roei</i>
<i>Exocarpos sparteus</i>	<i>Westringia cephalantha</i>
<i>Glischrocaryon roei</i>	
<i>Grevillea excelsior</i>	
<i>Grevillea incurva</i>	

This page has been left blank intentionally

Appendix G: Vegetation Condition Mapping

This page has been left blank intentionally.



Western Areas - Forrestania Nickel Operations
Lounge Lizard Vegetation, Flora and Fauna Biological Assessment

Appendix G: Vegetation Association Mapping

Author: N. Cadd

Date: 07-11-2014

Drawn: C. Dyde

Figure Ref: 17603-14-BIDR-1RevA_141110_AppG

Datum: GDA 1994 - Projection: MGA Zone 50 - Scale: 1:7,000 (A3)

0 100 200 300 400 500 Metres



This page has been left blank intentionally

This page has been left blank intentionally

Appendix H: Vascular Flora Species List

This page has been left blank intentionally.

Table F.1: Vascular flora species list for the survey area.


Family	Species Name	Conservation Status
Asparagaceae	<i>Laxmannia paleacea</i>	
	<i>Lomandra effusa</i>	
	<i>Lomandra mucronata</i>	
	<i>Thysanotus</i> sp. indet.	
Asteraceae	<i>Olearia ramosissima</i>	
Boraginaceae	<i>Halgania lavandulacea</i>	
Casuarinaceae	<i>Allocasuarina acutivalvis</i> subsp. <i>acutivalvis</i>	
	<i>Allocasuarina humilis</i>	
	<i>Allocasuarina microstachya</i>	
	<i>Allocasuarina spinosissima</i>	
	<i>Allocasuarina thuyoides</i>	
Cupressaceae	<i>Callitris columellaris</i>	
Cyperaceae	<i>Caustis dioica</i>	
	<i>Gahnia ancistrophylla</i>	
	<i>Lepidosperma</i> sp. indet.	
	<i>Lepidosperma viscidum</i>	
	<i>Mesomelaena preissii</i>	
	<i>Schoenus</i> sp. indet.	
Dilleniaceae	<i>Hibbertia gracilipes</i>	
	<i>Hibbertia pungens</i>	
Elaeocarpaceae	<i>Tetratheca efoliata</i>	
Ericaceae	<i>Astroloma serratifolium</i>	
	<i>Leucopogon dielsianus</i>	
	<i>Leucopogon</i> sp. indet.	
	<i>Leucopogon</i> sp. Coujinup (M.A. Burgman 1085)	
	<i>Lysinema ciliatum</i>	
Euphorbiaceae	<i>Beyeria sulcata</i>	
Fabaceae	<i>Acacia acoma</i>	
	<i>Acacia assimilis</i> subsp. <i>assimilis</i>	
	<i>Acacia burkittii</i>	
	<i>Acacia erinacea</i>	
	<i>Acacia poliochroa</i>	
	<i>Acacia sphacelata</i> subsp. <i>sphacelata</i>	
	<i>Aotus</i> sp. <i>Tortile</i> (G.J. Keighery 3767)	
	<i>Daviesia benthamii</i>	
	<i>Daviesia benthamii</i> subsp. <i>acanthoclona</i>	
	<i>Daviesia lancifolia</i>	
	<i>Eutaxia hirsuta</i>	P2
	<i>Eutaxia neurocalyx</i> subsp. <i>papillosa</i>	
	<i>Gastrolobium spinosum</i>	
	<i>Gompholobium viscidulum</i>	
	<i>Templetonia rossii</i>	
Goodeniaceae	<i>Dampiera juncea</i>	
	<i>Scaevola restiacea</i>	
Haemodoraceae	<i>Conostylis petrophiloides</i>	
Haloragaceae	<i>Glischrocaryon roei</i>	
Hemerocallidaceae	<i>Dianella revoluta</i> var. <i>divaricata</i>	

Family	Species Name	Conservation Status
Iridaceae	<i>Patersonia rudis</i> subsp. <i>velutina</i>	
Lamiaceae	<i>Hemigenia westringioides</i>	
	<i>Lachnostachys bracteosa</i>	
	<i>Microcorys exserta</i>	
	<i>Westringia cephalantha</i>	
Lauraceae	<i>Cassytha glabella</i>	
Loganiaceae	<i>Logania tortuosa</i>	
Malvaceae	<i>Guichenotia micrantha</i>	
Myrtaceae	<i>Astartea aspera</i>	
	<i>Baeckea</i> sp. indet.	
	<i>Beaufortia interstans</i>	
	<i>Beaufortia micrantha</i> var. <i>micrantha</i>	
	<i>Calytrix</i> sp. indet.	
	<i>Darwinia</i> sp. indet.	
	<i>Darwinia</i> sp. Lake Cobham (K. Newbey 3262)	
	<i>Eucalyptus</i> ? <i>subangusta</i> subsp. <i>subangusta</i>	
	<i>Eucalyptus eremophila</i>	
	<i>Eucalyptus platycorys</i>	
	<i>Eucalyptus rigidula</i>	
	<i>Leptospermum erubescens</i>	
	<i>Leptospermum fastigiatum</i>	
	<i>Leptospermum spinescens</i>	
	<i>Melaleuca adnata</i>	
	<i>Melaleuca cordata</i>	
	<i>Melaleuca eleuterostachya</i>	
	<i>Melaleuca glaberrima</i>	
	<i>Melaleuca hamata</i>	
	<i>Melaleuca lateriflora</i>	
	<i>Melaleuca laxiflora</i>	
	<i>Melaleuca seriata</i>	
	<i>Melaleuca</i> sp. indet.	
	<i>Micromyrtus erichsenii</i>	
	<i>Verticordia chrysanthella</i>	
	<i>Verticordia inclusa</i>	
	<i>Verticordia picta</i>	
	<i>Verticordia plumosa</i> var. <i>incrassata</i>	
	<i>Verticordia roei</i> subsp. <i>roei</i>	
Olacaceae	<i>Olax benthamiana</i>	
Poaceae	<i>Amphipogon turbinatus</i>	
	<i>Austrostipa hemipogon</i>	
	<i>Neurachne alopecuroidea</i>	
Polygalaceae	<i>Comesperma scoparium</i>	
Proteaceae	<i>Adenanthos argyreus</i>	
	<i>Banksia cirsioides</i>	
	<i>Banksia elderiana</i>	
	<i>Banksia erythrocephala</i>	
	<i>Banksia laevigata</i> subsp. <i>fuscolutea</i>	
	<i>Banksia violacea</i>	
	<i>Conospermum brownii</i>	

Family	Species Name	Conservation Status
Proteaceae	<i>Conospermum stoechadis</i> subsp. <i>sclerophyllum</i>	
	<i>Grevillea cagiana</i>	
	<i>Grevillea excelsior</i>	
	<i>Grevillea huegelii</i>	
	<i>Grevillea incurva</i>	
	<i>Grevillea oncogyne</i>	
	<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>	
	<i>Hakea commutata</i>	
	<i>Hakea corymbosa</i>	
	<i>Hakea multilineata</i>	
	<i>Hakea platysperma</i>	
	<i>Isopogon scabriusculus</i>	
	<i>Petrophile divaricata</i>	
	<i>Petrophile ericifolia</i> subsp. <i>ericifolia</i>	
	<i>Petrophile stricta</i>	
	<i>Synaphea interioris</i>	
Restionaceae	<i>Desmocladus flexuosus</i>	
	<i>Lepidobolus preissianus</i>	
Rutaceae	<i>Boronia coerulescens</i> subsp. <i>spicata</i>	
	<i>Phebalium tuberculosum</i>	
Santalaceae	<i>Exocarpos aphyllus</i>	
	<i>Exocarpos sparteus</i>	
	<i>Leptomeria preissiana</i>	
	<i>Santalum acuminatum</i>	
	<i>Santalum murrayanum</i>	
Sapindaceae	<i>Dodonaea bursariifolia</i>	
	<i>Dodonaea viscosa</i> subsp. <i>spatulata</i>	
Scrophulariaceae	<i>Eremophila scoparia</i>	
Stylidiaceae	? <i>Stylidium clavatum</i>	
	<i>Stylidium</i> sp. indet.	
Thymelaeaceae	<i>Pimelea suaveolens</i>	
	<i>Pimelea subvillifera</i>	



APPENDIX C – FLORA, VEGETATION AND FAUNA ASSESSMENT (BOTANICA 2021)



**Flora/ Vegetation and Fauna
Assessment
Lounge Lizard Sand Pits
Prepared For
Western Areas Limited**



**October 2021
Version 1**

Prepared by:
Botanica Consulting Pty Ltd
33 Brewer Street
Perth WA 6000



Disclaimer

This document and its contents are to be treated as confidential and are published in accordance with and subject to an agreement between Botanica Consulting (BC) and the client for whom it has been prepared and is restricted to those issues that have been raised by the client in its engagement of BC. Neither this document nor its contents may be referred to or quoted in any manner (report or other document) nor reproduced in part or whole by electronic, mechanical or chemical means, including photocopying, recording or any information storage system, without the express written approval of the client and/or BC.

This document and its contents have been prepared utilising the standard of care and skill ordinarily exercised by Environmental Scientists in the preparation of such documents. All material presented in this document is published in good faith and is believed to be accurate at the time of writing. Any person or organisation who relies on or uses the document and its contents for purposes or reasons other than those agreed by BC and the client without primarily obtaining the prior written consent of BC, does so entirely at their own risk. BC denies all liability in tort, contract or otherwise for any loss, damage or injury of any kind whatsoever (whether in negligence or otherwise) that may be endured as a consequence of relying on this document and its contents for any purpose other than that agreed with the client.

Quality Assurance

An internal quality review process has been implemented to each project task undertaken by BC. Each document and its contents are carefully reviewed by core members of the Consultancy team and signed off at Director Level prior to issue to the client. Draft documents are submitted to the client for comment and acceptance prior to final production.

Document Job Number: 2021/105

Prepared by: Lauren Pick
Senior Environmental Consultant
Botanica Consulting Pty Ltd

Reviewed by: Andrea Williams
Director
Botanica Consulting Pty Ltd

Approved by: Jim Williams
Director
Botanica Consulting Pty Ltd

Contents	Page No.
1 Introduction	1
1.1 Project Description	1
1.2 Objectives	1
2 Regional Biophysical Environment	4
2.1 Regional Environment	4
2.2 Soils and Landscape Systems	6
2.3 Remnant Vegetation	8
2.4 Climate	10
2.5 Hydrology	10
2.6 Land Use	12
3 Survey Methodology	14
3.1 Desktop Assessment	14
3.2 Field Assessment	15
3.2.1 Reconnaissance Flora Survey	16
3.2.2 Targeted Flora Survey	16
3.2.3 Basic Fauna Survey	16
3.2.4 Targeted Fauna Survey	17
3.2.5 Personnel involved	17
3.2.6 Scientific licences	17
3.3 Survey limitations and constraints	19
4 Results	21
4.1 Desktop Assessment	21
4.1.1 Flora/Vegetation	21
4.1.2 Fauna	26
4.2 Field Assessment	31
4.2.1 Vegetation Types	31
4.2.2 Vegetation Condition	35
4.2.3 Fauna Habitat	37
4.2.4 Introduced Species	40
4.2.5 Significant Flora	41
4.2.6 Significant Vegetation	45
4.2.7 Significant Fauna	46
4.3 Matters of National Environmental Significance	48
4.3.1 <i>Environment Protection and Biodiversity Conservation Act 1999</i>	48
4.4 Relevant State Legislation	48
4.4.1 <i>Environmental Protection Act WA 1986</i>	48
4.4.2 <i>Biodiversity Conservation Act 2016</i>	49
4.4.3 Conservation Reserves	49
4.5 Native Vegetation Clearing Principles	49
5 Recommendations	52
6 Bibliography	53

Appendices

Appendix 1: Conservation Ratings BC Act and EPBC Act	57
Appendix 2: Regional map of the survey area in relation to conservation areas	61
Appendix 3: List of species identified within each vegetation type	62
Appendix 4: Vegetation Condition Rating	66
Appendix 5: Conservation Significant Flora within 40km of the survey area	67
Appendix 6: Potential Fauna Species List	74

Tables

Table 2-1: Soil Landscape System within the survey area	6
Table 2-2: Pre-European Vegetation Association within the survey area	8
Table 2-3: Potential Terrestrial Groundwater Dependent Ecosystems (BoM, 2020).....	10
Table 3-1: Scientific Licences of Botanica Staff coordinating the survey	17
Table 3-2: Limitations and constraints associated with the survey	19
Table 4-1: Introduced flora within 40km of the survey area	21
Table 4-2: Flora of Conservation Significance identified as possible or known to occur within the survey area.....	23
Table 4-3: Likelihood of Occurrence – Fauna Species of Conservation Significance	28
Table 4-4: Summary of vegetation types within the survey area	32
Table 4-5: Main Terrestrial Fauna Habitats within the survey area	38
Table 4-6: Summary of Potential Vertebrate Fauna Species	40
Table 4-7: Significant Flora recorded within the survey area.....	42
Table 4-8: Assessment of development within the survey area against native vegetation clearing principles	50

Figures

Figure 1-1: Flying Fox sand pit survey area.....	2
Figure 1-2: Regional map of the survey area.....	3
Figure 2-1: Map of IBRA Bioregions in relation to the survey area	5
Figure 2-2: Map of Soil Landscape Systems within the survey area	7
Figure 2-3: Pre-European Vegetation Associations within the survey area	9
Figure 2-4: Monthly rainfall and mean monthly rainfall (January 2020 to September 2021) for the Hyden weather station #10568 (BoM, 2021)	10
Figure 2-5: Surface Hydrology of the survey area	11
Figure 2-6: Location of survey area within the Great Western Woodlands	13
Figure 3-1: GPS Tracks recorded within the survey area	18
Figure 4-1: Vegetation types within the survey area.....	34
Figure 4-2: Vegetation condition within the survey area	36
Figure 4-3: Main Terrestrial Fauna Habitats within the survey area	39
Figure 4-4: Significant flora recorded within the survey area.....	44

Glossary

Acronym	Description
ANCA	Australian Nature Conservation Agency.
BA	Birdlife Australia.
BAM Act	<i>Biosecurity and Agriculture Management Act 2007</i> , WA Government.
BC Act	<i>Biodiversity Conservation Act 2016</i> , WA Government.
Botanica	Botanica Consulting Pty Ltd.
BoM	Bureau of Meteorology.
CAMBA	China Australia Migratory Bird Agreement 1998.
DAFWA	Department of Agriculture and Food (now DPIRD), WA Government.
DAWE	Department of Agriculture, Water and the Environment (formerly DotEE), Australian Government.
DBCA	Department of Biodiversity, Conservation and Attractions (formerly DPaW), WA Government.
DMIRS	Department of Mines, Industry Regulation and Safety (formerly DMP), WA Government
DotEE	Department of the Environment and Energy (now DAWE), Australian Government.
DPIRD	Department of Primary Industries and Regional Development, WA Government
DWER	Department of Water and Environmental Regulation (formerly EPA, DER and DoW), WA Government
EP Act	Environmental Protection Act 1986, WA Government.
EP Regulations	Environmental Protection (Clearing of Native Vegetation) Regulations 2004, WA Government.
EPA	Environmental Protection Authority, WA Government.
EPBC Act	<i>Environment Protection and Biodiversity Conservation Act 1999</i> , Australian Government.
ESA	Environmentally Sensitive Area.
Ha	Hectare (10,000 square meters).
IBRA	Interim Biogeographic Regionalisation for Australia.
IUCN	International Union for the Conservation of Nature and Natural Resources – commonly known as the World Conservation Union.
JAMBA	<i>Japan Australia Migratory Bird Agreement 1981</i> .
Km	Kilometer (1,000 meters).
MVG	Major Vegetation Groups.
NVIS	National Vegetation Information System.
PEC	Priority Ecological Community.
ROKAMBA	Republic of Korea-Australia Migratory Bird Agreement 2007.
SRE	Short Range Endemic.
SSC	Species Survival Commission, International.
TEC	Threatened Ecological Community.
WA	Western Australia.
WAHERB	Western Australian Herbarium.
WAM	Western Australian Museum, WA Government.

Executive Summary

Botanica Consulting Pty Ltd (Botanica) was commissioned by Western Areas Limited to undertake a reconnaissance flora/ vegetation survey, basic fauna survey and targeted flora/ fauna survey of proposed new borrow pit developments within tenements M77/545 and M77/911 (referred to as the 'survey area'). The survey area is located at Lounge Lizard, an area adjacent and to the south of the Flying Fox mine site, owned and operated by Western Areas Limited, located approximately 78km east of Hyden, Western Australia. The survey was conducted on the 21st and 22nd September 2021. The total survey area encompassed an approximate area of 383 ha. The target survey area encompassed an approximate area of 25.7 ha

Four vegetation types were identified within the survey area. These vegetation types were located within three landform types and comprised of three major vegetation groups, which were represented by a total of 30 Families, 75 Genera and 179 Taxa. No introduced flora were identified within the survey area. Based on the vegetation condition rating scale adapted from Keighery, 1994 and Trudgen, 1988 vegetation was rated as 'very good'.

The broad scale terrestrial fauna habitats within the survey area have been identified as comprising a mosaic of clay-loam plain and sand-loam plain. Results of the literature review identified 32 mammals (including 9 bat species), 121 birds, 65 reptiles and 22 frog species that have previously been recorded in the general area, some of which have the potential to occur, subject to the identified habitats being suitable.

No Threatened flora or Threatened Ecological Communities (TEC) as listed under the Western Australian *Biodiversity Conservation (BC) Act 2016* or Commonwealth *Environment Protection and Biodiversity Conservation (EPBC) Act 1999* were identified within the survey area. The survey area is located adjacent to the spatially mapped 500m buffer of the *Ironcap Hills Vegetation Complexes* (North Ironcap) obtained from the Department of Biodiversity, Conservation and Attractions (DBCA) communities database search. This Priority Ecological Community is listed by the DBCA as a Priority 3 Ecological Community. No banded ironstone was identified within the survey area and no vegetation representative of this Priority Ecological Community was identified within the survey area. Three Priority flora taxa were identified within the survey area; *Eutaxia hirsuta* (P2), *Microcorys* sp. Forrestania (V. English 2004) (P4) and *Rinzia triplex* (P3).

A review of the EPBC Act Threatened fauna list, DBCA's Threatened Fauna Database and Priority List, unpublished reports and scientific publications identified a number of specially protected, migratory or priority fauna species as having been previously recorded or as being potentially present in the general vicinity of the survey area. However, no fauna of conservation significance is likely to be significantly impacted on by the proposed development. This conclusion is primarily based on the lack of suitable habitats, the known local extinction of some species, the relatively small size of the impact footprint and the extensive habitat connectivity with adjoining areas. Impacts on fauna and fauna habitat are therefore anticipated to be localised, small/negligible and as a consequence manageable.

The survey area does not contain any world or national heritage places and does not occur within a Bush Forever site. There are no wetlands of international importance (Ramsar Wetlands), national importance (Australian Nature Conservation Agency (ANCA) Wetlands) or conservation category wetlands within the survey area.

The survey area is located within an Environmentally Sensitive Area (ESA) as listed under the *Environmental Protection (EP) Act 1986*. This ESA is associated with Lake Cronin which is located approximately 4.5km north-east of the survey area.

The survey area is not located within a vested Conservation Reserve, however according to the EPA (2009) Advice on Conservation Values and Review of Nature Reserve Proposals in the Lake Cronin Region, the survey area occurs within a 56,750ha area within the mineralised greenstone belt in the Lake Cronin Region which is proposed to be managed under Section 33(2) of the *Conservation and Land Management Act 1984* (CALM Act) but not formally reserved.

Botanica have undertaken an assessment against the ten native vegetation Clearing Principles contained in Schedule 5 of the EP Act. Based on the information provided in this report, this assessment considered that the proposed sand pit works in the survey area is unlikely to be at variance with these ten Clearing Principles.

1 Introduction

1.1 Project Description

Botanica Consulting Pty Ltd (Botanica) was commissioned by Western Areas Limited to undertake a reconnaissance flora/ vegetation survey, basic fauna survey and targeted flora/ fauna survey of proposed new borrow pit developments within tenements M77/545 and M77/911 (referred to as the 'survey area') . The survey area is located within the Flying Fox mine site, owned and operated by Western Areas Limited, located approximately 78km east of Hyden, Western Australia (Figure 1-2). The survey was conducted on the 21st and 22nd September 2021. The total survey area encompassed an approximate area of 383 ha. The target survey area encompassed an approximate area of 25.7 ha (Figure 1-1).

1.2 Objectives

The flora assessment was conducted in accordance with the requirements of a reconnaissance flora survey as defined in *Technical Guidance - Flora and Vegetation Surveys for Environmental Impact Assessment – December 2016* (EPA, 2016a). The objectives of the assessment were to:

- gather background information on flora and vegetation in the target area (literature review, database and map-based searches);
- identify significant flora, vegetation/ecological communities and assess the potential sensitivity to impact;
- conduct a field survey to verify / ground truth the desktop assessment findings;
- undertake floristic community mapping to a scale appropriate for the bioregion and described according to the National Vegetation Information System (NVIS) structure and floristics;
- undertake vegetation condition mapping;
- assess the project area's plant species diversity, density, composition, structure and weed cover, using NVIS classification system for vegetation description;
- assess Matters of National Environmental Significance (MNES) and indicate whether potential impacts on MNES as protected under the EPBC Act are likely to require referral of the project to the Commonwealth DAWE; and
- determine the State legislative context of environmental aspects required for the assessment.

The fauna assessment was conducted in accordance with the requirements of a basic terrestrial fauna survey as defined in *Technical Guidance - Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment – June 2020* (EPA, 2020). The objectives of the assessment were to:

- Gather background information on fauna in the survey area (literature review, database and map-based searches);
- Delineate and characterise the faunal assemblages and fauna habitats present in the survey area;
- Document and map locations of any Threatened or Priority listed fauna species located; and
- Assess the regional and local conservation status of fauna species and fauna habitats within the survey area.




Figure 1-1: Flying Fox sand pit survey area




Figure 1-2: Regional map of the survey area

2 Regional Biophysical Environment

2.1 Regional Environment

The survey area lies within the South-West and Interzone Province of Western Australia. Based on the Interim Biogeographic Regionalisation of Australia (IBRA), Version 7 (DotEE, 2012), the survey area is located on the border of the Coolgardie and Mallee Bioregions. The Coolgardie and Mallee Bioregions are further divided into subregions with the survey area located within the Western Mallee subregion (MAL2) of the Mallee Bioregion and the Southern Cross subregion (COO2) of the Coolgardie Bioregion (Figure 2-1).

The Coolgardie Bioregion forms part South-West and Interzone of Western Australia in a region known as the Coolgardie Botanical District (Beard, 1990). The Coolgardie Bioregion is located within the Yilgarn Craton and is characterised by a granite basement which includes Archaean Greenstone intrusions in parallel belts. Drainage is occluded. The Southern Cross subregion comprises gently undulating uplands on granite strata and broad valleys with bands of low greenstone hills (McKenzie, J.E. May and S. McKenna, 2002).

The Mallee Bioregion also forms part of the South-West and Interzone Province of Western Australia in a region known as the Roe Botanical District. The Mallee Bioregion is located in the south-eastern part of Yilgarn Craton which is gently undulating, with partially occluded drainage. The Western Mallee subregion has more relief than its eastern counterpart. Its main surface-types comprise clays and silts underlain by kankar, exposed granite, sandplains, isolated uplands of laterite pavements and Salt Lake systems on a granite basement (McKenzie, J.E. May and S. McKenna, 2002).




Figure 2-1: Map of IBRA Bioregions in relation to the survey area

2.2 Soils and Landscape Systems

Based on geographic information provided by the Department of Agriculture and Food Western Australia (DAFWA, 2014), the survey area is located within the South-eastern Zone of Ancient Drainage (250) of the Avon Province (25).

The Avon Province is characterised as a laterised plateau (dissected at fringes and with saline drainage lines inland) on deeply weathered mantle and alluvium over granitic rocks of the Yilgarn Craton (and Albany-Fraser Orogen). Soil are comprised of sandy duplexes soils and ironstone gravelly soils with loamy earths, loamy duplexes, sandy earths, deep sands and wet soils. Vegetation is dominated by York gum-wandoo-salmon gum-morrel gimlet woodland and jarrah-marri-karri-wandoo woodlands/forests (with some mallee scrub, tammar-wodjil thickets and scrub-heath). This Province is located in the south-west, between Nannup, Denmark, Jerramungup, Southern Cross, Lake Moore, Carnamah and the Perth Hills (Tille, 2006).

The South-western Zone of Ancient Drainage (250) is characterised by gently undulating terrain (with some salt lake chains and areas of prominent granitic outcrops) on deeply weathered mantle and alluvium over granitic rocks of the Yilgarn Craton. Soils include sandy duplexes (often alkaline) with ironstone gravelly soils and loamy earths (often calcareous) and some loamy duplexes, sandy earths, deep sands and saline wet soils. Mallee scrub and salmon gum-gimlet-morrel woodlands (and some scrub-heath). This zone is located in the southern Wheatbelt between Kondinin, Lake Grace, Gnowangerup, Frank Hann National Park and Mt Holland (Tille, 2006).

These South-western Zone of Ancient Drainage is further divided into soil landscape systems, with the survey area located within the soil landscape system described in Table 2-1 and Figure 2-2 below.

Table 2-1: Soil Landscape System within the survey area

Soil Landscape System	Description
Ms8	Gently undulating plains with broad shallow drainage depressions




Figure 2-2: Map of Soil Landscape Systems within the survey area

2.3 Remnant Vegetation

The Department of Primary Industries and Regional Development (DPIRD) Vegetation Association GIS file (2018) indicates that the survey area is located within Pre-European Beard vegetation association Forrestania 511 of the Southern Cross subregion and Forrestania 2048 of the Western Mallee subregion. The extent of these vegetation associations, as specified in the 2018 Statewide Vegetation Statistics (DBCA, 2019) are provided in Table 2-2 and Figure 2-3.

Areas retaining less than 30% of their pre-European vegetation extent generally experience exponentially accelerated species loss, while areas with less than 10% are considered “endangered” (EPA, 2000). Development within the survey area will not significantly reduce the extent of pre-European vegetation.

Table 2-2: Pre-European Vegetation Association within the survey area

Region	Pre-European Extent (ha)	Pre-European extent remaining (%)	% of Current extent within DBCA managed lands	Extent within the survey area (ha)
Forrestania 511: Medium woodland; salmon gum & morrel				
Southern Cross Subregion	153,641.65	99.58	9.68	50
Western Australia	163,919.47	99.59	9.07	
Forrestania 2048: Shrublands; scrub-heath in the Mallee Region				
Western Mallee Subregion	5,735.92	97.56	0	4
Western Australia	7,829.67	97.92	0.64	



Figure 2-3: Pre-European Vegetation Associations within the survey area

2.4 Climate

The climate of the Southern Cross subregion is characterised as arid to semi-arid Mediterranean with an annual rainfall of 200-300mm (Beard, 1990; Cowan, 2001). The climate of the Western Mallee subregion is characterised as dry warm Mediterranean with an annual rainfall of 300-500mm (Beard, 1990; Beecham & Danks, 2001). Rainfall data for the Hyden weather station (#10568) located approximately 78km west of the survey area is shown in Figure 2-4 (BoM, 2021). Mean monthly rainfall ranges from 48.2 mm in June to 14.4 mm in December, whilst the mean annual rainfall is 339.6 mm. Annual rainfall received in 2020 was below average. Survey work was undertaken in September 2021 within the EPA recommended timing for primary surveys of the South-West and Interzone Province (i.e. Spring) (EPA, 2016).




Figure 2-4: Monthly rainfall and mean monthly rainfall (January 2020 to September 2021) for the Hyden weather station #10568 (BoM, 2021)

2.5 Hydrology

According to the Geoscience Australia database (2015), there are no perennial or ephemeral inland waters or drainage lines within the survey area (Figure 2-5).

Groundwater Dependent Ecosystems (GDE) includes biological assemblages of species such as wetlands or woodlands that use groundwater either opportunistically or as their primary water source. For the purposes of this report, a GDE is defined as any vegetation community that derives part of its water budget from groundwater and must be assumed to have some degree of groundwater dependency. According to the BoM *Atlas of Groundwater Dependent Ecosystems* (BoM, 2019) database, there are no known or potential aquatic GDEs located within the survey area (Figure 2-5). The survey area has moderate potential to contain the following terrestrial GDE listed in Table 2-3 and shown spatially in Figure 2-5 (BoM, 2020). As the proposed developments within the survey relate to sand pit activities and will not impact the groundwater table, impacts to potential GDE vegetation are unlikely to occur.

Table 2-3: Potential Terrestrial Groundwater Dependent Ecosystems (BoM, 2020)

GDE Description	Potential GDE according to BoM (2019b)
Undulating plains with some sandplains, ferruginous breakaways; ridges of metamorphic rocks and granitic hills and rises; calcretes, large salt lakes and dunes along valleys.	Moderate potential GDE




Figure 2-5: Surface Hydrology of the survey area

2.6 Land Use

The dominant land uses of the Southern Cross subregion includes native pastures (17%), Conservation Reserves (11.53%), UCL & Crown Reserves (66.74%) and Cultivation – Dry Land agriculture (2.27%) (Cowan, 2001). The dominant land uses of the Western Mallee subregion includes Dry Land agriculture, UCL & Crown Reserves, roads and other easements.

The survey area also lies within the Great Western Woodlands. The Great Western Woodlands is considered by The Wilderness Society of WA to be of global biological and conservation importance as one of the largest and healthiest temperate woodlands on Earth, containing many endemic species. The region covers almost 16 million hectares, 160,000 square kilometers, from the southern edge of the Western Australian Wheatbelt to the pastoral lands of the Mulga country in the north, the inland deserts to the northeast, and the treeless Nullarbor Plain to the east (Figure 2-6).

The area provides an eastward connection between southwest forests and inland deserts (Gondwana Link) as well as linking the north-west passage to Shark Bay. The majority of the Great Western Woodlands is unallocated crown land (61.1%) with other interests including pastoral leases (20.4%), conservation reserves (15.4%) unallocated crown land ex pastoral managed by the DEC (2%) and private land (approximately 1%) (Watson *et. al.*, 2008).

No specific management strategy applies to the Great Western Woodlands, rather an approach to conservation which occurs across all land tenures and when different stakeholders work together with biodiversity in mind. The central component of this approach is to identify and conserve key large-scale, long term ecological processes that drive connectivity between ecosystems and species. The Great Western Woodlands currently includes towns, highways, roads, railways, private property, Crown Reserves, agricultural activities and mining tenements.




Figure 2-6: Location of survey area within the Great Western Woodlands

3 Survey Methodology

3.1 Desktop Assessment

Prior to the field assessment a literature review was undertaken of previous flora and fauna assessments conducted within the local region. Documents reviewed included:

- How, R. A Newbey, K.R Dell, J. Muir, B.G & Hnatiuk, R.J, (1988), *The Biological survey of the Eastern Goldfields of Western Australia: Lake Johnston-Hyden*. Western Australian Museum Supplement No. 30.
- Gibson (2004) Flora and vegetation of the Eastern Goldfields Ranges: Part 7. Middle and South Ironcap, Digger Rock and Hatter Hill. Science Division, Department of Conservation and Land Management.
- Botanica (2006), Flora and Vegetation Survey of the Flying Fox North East Exploration Area for Western Areas Limited, Botanica Consulting.
- Biota (2007a). Forrestania Monitoring Survey, Flying Fox Phases I, II, III and IV. Unpublished report for Western Areas NL.
- Botanica (2007a), Vegetation Survey of a Proposed Extension to the Current Clearing Permit Number 691/1 at the Flying Fox mine site prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2007b), Flora and Vegetation Survey within the Greater Flying Fox mine site prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2009), Flora and Vegetation Survey within the lounge Lizard/Flying Fox area, Proposed Gravel Pit, prepared for Western Areas Limited, Botanica Consulting.
- Biota (2010). Spotted Quoll Haul Road Single Phase Fauna Survey. Unpublished report for Western Areas N.L. May 2010.
- Botanica (2013), Flora and Vegetation survey of the greater Flying Fox area, prepared for Western Areas Limited, Botanica Consulting.
- Astron (2014). Forrestania Nickel Operations Lounge Lizard Vegetation, Flora and Fauna Biological Assessment, prepared for Western Areas Limited, Astron Environmental Services.
- Biota (2018), New Morning Level 1 and Targeted Terrestrial fauna Survey, prepared for Western Areas Limited, Biota Environmental Sciences.
- Botanica (2021a), Detailed Flora and Vegetation Survey and Targeted Flora survey of the New Morning Project, prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2021b), Reconnaissance Flora/ Vegetation Survey & Basic Fauna Survey Lounge Lizard East Sand Pit Extension, prepared for Western Areas Limited, Botanica Consulting.

In addition to the literature review, searches of the following databases were undertaken to aid in the compilation of a list of flora and fauna taxa within the survey area:

- DBCA Priority/ Threatened Flora Database Search (DBCA, 2018a)
- DBCA Priority/ Threatened Ecological Communities Database Search (DBCA, 2018b)
- DBCA NatureMap Database (DBCA, 2021);
- DAWE Protected Matters search tool (DAWE, 2021).

The DBCA Priority/ Threatened Flora Database Search and Priority/ Threatened Ecological Communities Database Search were conducted within a 50km radius of the survey area (DBCA, 2018a; DBCA, 2018b).

The NatureMap and Protected Matters Search were conducted for an area encompassing a 40km radius of the centre coordinates -32.4322S 119.6932E. It should be noted that these lists are based on observations from a broader area than the assessment area (40km radius) and therefore may include taxa not present. The databases also often include very old records that may be incorrect or in some cases the taxa in question have become locally or regionally extinct. Information from these sources should therefore be taken as indicative only and local knowledge and information also needs to be taken into consideration when determining what actual species may be present within the specific area being investigated. The conservation significance of flora and fauna taxa was assessed using data from the following sources:

- *Environment Protection and Biodiversity and Conservation (EPBC) Act 1999*. Administered by the Australian Government (DAWE);
- *Biodiversity Conservation (BC) Act 2016*. Administered by the WA Government (DBCA);
- Red List produced by the Species Survival Commission (SSC) of the World Conservation Union (also known as the IUCN Red List – the acronym derived from its former name of the International Union for Conservation of Nature and Natural Resources). The Red List has no legislative power in Australia but is used as a framework for State and Commonwealth categories and criteria; and
- Priority Flora/ Fauna list. A non-legislative list maintained by DBCA for management purposes (fauna list released April 2019; flora list released December 2018).

The EPBC Act also requires the compilation of a list of migratory species that are recognised under international treaties including the:

- Japan Australia Migratory Bird Agreement 1981 (JAMBA)¹;
- China Australia Migratory Bird Agreement 1998 (CAMBA);
- Republic of Korea-Australia Migratory Bird Agreement 2007 (ROKAMBA); and
- Bonn Convention 1979 (The Convention on the Conservation of Migratory Species of Wild Animals).

Most but not all migratory bird species listed in the annexes to these bilateral agreements are protected in Australia as Matters of National Environmental Significance (MNES) under the EPBC Act. Descriptions of conservation significant species and communities are provided in Appendix 1.

3.2 Field Assessment

Botanica conducted a reconnaissance flora/ vegetation survey, basic fauna survey and targeted flora/ fauna survey on the 21st and 22nd September 2021. The total survey area encompassed an approximate area of 383 ha. The target survey area encompassed an approximate area of 25.7 ha. The survey area was traversed on foot and 4WD by two personnel; Jim Williams and Michelle Luinstra. GPS tracks of the area traversed within the survey area is shown spatially in Figure 3-1.

¹ Most but not all species listed under JAMBA are also specially protected under Specially Protected Species of the BC Act.

3.2.1 Reconnaissance Flora Survey

Prior to the commencement of field work, aerial photography was inspected and obvious differences in the vegetation assemblages were identified. The different vegetation communities identified were then inspected during the field survey to assess their validity. A handheld GPS unit was used to record the coordinates of the boundaries between existing vegetation communities.

At each sample point (relevé), the following information was recorded:

- GPS location;
- Photograph of vegetation;
- Dominant taxa for each stratum;
- All vascular taxa (including annual taxa);
- Landform classification;
- Vegetation condition rating;
- Collection and documentation of unknown plant specimens; and
- GPS location, photograph and collection of flora of conservation significance if encountered.

Unknown specimens collected during the survey were identified with the aid of samples housed at the Botanica Herbarium and WAHERB. Vegetation was classified in accordance with NVIS classifications.

3.2.2 Targeted Flora Survey

A targeted search for Threatened and Priority flora was conducted within the target survey area, encompassing an approximate area of 25.7 ha. The survey area was systematically searched on foot by two Botanica staff members to identify and record the locations of Threatened and Priority flora. Any locations of Threatened and Priority flora were recorded using a hand-held GPS. For any Threatened and Priority Flora identified, a simple plant count (not differentiated between juvenile/mature plants, flowering or non-flowering plants) was conducted for each record.

3.2.3 Basic Fauna Survey

Vegetation and landform units identified during the flora assessment have been used to define broad fauna habitat types across the site. This information has been supplemented with observations made during the fauna assessment.

The main aim of the fauna habitat assessment was to determine if it was likely that any species of conservation significance would be utilising the areas that maybe impacted on as a consequence of development at the site. The habitat information obtained was also used to aid in finalising the overall potential fauna list.

As part of the desktop literature review, available information on the habitat requirements of the species of conservation significance listed as possibly occurring in the area was researched. During the field survey, the habitats within the study area were assessed and specific elements identified, if present, to determine the likelihood of listed threatened species utilising the area and its significance to them.

Opportunistic observations of fauna species were made during all field survey work which involved a series of transects across the study area during the day including observations of bird species with binoculars. Secondary evidence of a species presence such as tracks, scats, skeletal remains, foraging evidence or calls were also noted if observed/heard.

3.2.4 Targeted Fauna Survey

A targeted search for Malleefowl was conducted within the target survey area. The footprint was systematically searched on foot by two personnel to identify and record the locations of any Malleefowl activity (i.e. mounds, footprints and feathers). Any locations/ observations of Malleefowl activity were recorded using a hand-held GPS.

3.2.5 Personnel involved

Jim Williams - Environmental Consultant/ Director (Diploma of Horticulture)

Michelle Luinstra - Environmental Consultant (Bachelor of Science-Biology)

Greg Harewood - Zoologist (Bachelor of Science-Zoology)

Lauren Pick - Environmental Consultant (Bachelor of Science-Zoology/Conservation Biology)

3.2.6 Scientific licences

Table 3-1: Scientific Licences of Botanica Staff coordinating the survey

Licensed staff	Permit Number	Valid Until
Jim Williams	FB62000108 (Licence to take flora for scientific purposes)	27/05/2019-27/05/2022




Figure 3-1: GPS Tracks recorded within the survey area

3.3 Survey limitations and constraints

It is important to note that flora surveys will entail limitations notwithstanding careful planning and design. Potential limitations are listed in Table 3-2.

The conclusions presented in this report are based upon field data and environmental assessments and/or testing carried out over a limited period of time and are therefore merely indicative of the environmental condition of the site at the time of the field assessments. Also, it should be recognised that site conditions can change with time. Information not available at the time of this assessment which may subsequently become available may alter the conclusions presented.

Some species are reported as potentially occurring based on there being suitable habitat (quality and extent) within the survey area or immediately adjacent. The habitat requirements and ecology of many of the species known to occur in the wider area are however often not well understood or documented. It can therefore be difficult to exclude species from the potential list based on a lack of a specific habitats or microhabitats within the survey area. As a consequence of this limitation, the potential species list produced is most likely an overestimation of those species that actually utilise the survey area for some purpose.

In recognition of survey limitations, a precautionary approach has been adopted for this assessment. Any flora and fauna species that would possibly occur within the survey area (or immediately adjacent), as identified through ecological databases, publications, discussions with local experts/residents and the habitat knowledge of the author, has been listed as having the potential to occur.

Table 3-2: Limitations and constraints associated with the survey

Variable	Potential Impact on Survey	Details
Access problems	Not a constraint	The survey was conducted via 4WD and on foot. Numerous tracks were located within the survey area, providing ease of access.
Competency/ Experience	Not a constraint	The BC personnel that conducted the survey were regarded as suitably qualified and experienced. Coordinating Botanist/ Zoologist: Jim Williams Field Survey: Jim Williams and Michelle Luinstra Data Interpretation: Jim Williams, Greg Harewood and Lauren Pick
Timing of survey, weather & season	Minor constraint	Fieldwork was completed within the EPA's recommended primary survey time period (i.e. Spring) for the South-West and Interzone Province. The survey area has been previously surveyed by Botanica and other consultants over multiple years (2005, 2006-2008, 2012, 2014 and 2021) and multiple seasons.
Area disturbance	Not a constraint	The area has been disturbed from exploration and mining; however, vegetation was mostly intact and comprised of native vegetation.
Survey Effort/ Extent	Not a constraint	Survey intensity was appropriate for the size/ significance of the area with a reconnaissance flora survey and basic fauna completed to identify vegetation types/fauna habitats and

Variable	Potential Impact on Survey	Details
		conservation significant species/communities over the survey area. Targeted flora and fauna surveys were also conducted.
Availability of contextual information at a regional and local scale	Not a constraint	Threatened flora database searches provided by the DBCA were used to identify any potential locations of Threatened/Priority taxa. BoM, DWER, DPIRD, DBCA and DAWE databases were reviewed to obtain appropriate regional desktop information on the biophysical environment of the local region. Previous Flora/ Fauna surveys within the local area have been assessed for pertinent information and environmental context of the regional area.
Completeness	Minor constraint	In the opinion of Botanica, the survey area was covered sufficiently in order to identify vegetation assemblages. Few annual species were present during the current survey, however Botanica have previously conducted flora surveys within the survey area over multiple years (2006-2008, 2012 and 2021) and multiple seasons. All observed flora individuals were able to be identified to species level. The vegetation types for this study were based on visual descriptions of locations in the field. The distribution of these vegetation communities/ fauna habitats outside the study area is not known, however vegetation types identified were categorised via comparison to vegetation distributions throughout WA specified in the NVIS Major Vegetation Groups (DotEE, 2017b).

4 Results

4.1 Desktop Assessment

4.1.1 Flora/Vegetation

According to the results of the NatureMap search (DBCA, 2021), a total of 1,300 flora taxa have been recorded within a 40 km radius of the survey area. Dominant genera include *Acacia*, *Eucalyptus* and *Melaleuca*. Combined results of database searches (DBCA, 2021 and DAWE, 2021) identified 49 introduced taxa as potentially occurring within 40km of the survey area (Table 4-1). According to the Department of Primary Industries and Regional Development Western Australian Organism List (DPIRD, 2020), two taxa are listed as a Declared Pest under the *Biosecurity and Agriculture Management (BAM) Act 2007*. Three taxa are listed as a Weed of National Significance (WoNS).

Table 4-1: Introduced flora within 40km of the survey area

Taxon	Common Name	Declared Pest	WoNS
<i>Aira caryophyllea</i>	Silvery Hairgrass		
<i>Alyssum linifolium</i>	Flax-leaf Alyssum		
<i>Arctotheca calendula</i>	Cape Weed, African Marigold		
<i>Asparagus asparagoides</i>	Bridal Creeper	Y	Y
<i>Avellinia michelii</i>			
<i>Brassica tournefortii</i>	Mediterranean Turnip		
<i>Brassica x napus</i>			
<i>Bromus catharticus</i>	Prairie Grass		
<i>Bromus rubens</i>	Red Brome		
<i>Bupleurum semicompositum</i>			
<i>Carrichtera annua</i>	Wards Weed		
<i>Centaurea melitensis</i>	Maltese Cockspur, Malta Thistle		
<i>Centaureum erythraea</i>	Common Centaury		
<i>Centaureum tenuiflorum</i>			
<i>Cotula bipinnata</i>	Ferny Cotula		
<i>Crassula natans</i>			
<i>Crassula natans</i> var. <i>minus</i>			
<i>Ehrharta longiflora</i>	Annual Veldt Grass		
<i>Erodium cicutarium</i>	Common Storksbill		
<i>Hordeum leporinum</i>	Barley Grass		
<i>Hornungia procumbens</i>			
<i>Hypochaeris glabra</i>	Smooth Catsear		
<i>Juncus bufonius</i>	Toad Rush		
<i>Lepidium africanum</i>	Rubble Peppergrass		
<i>Lolium rigidum</i>	Wimmera Ryegrass		
<i>Lycium ferocissimum</i>	African Boxthorn		Y
<i>Lysimachia arvensis</i>	Pimpernel		
<i>Medicago sativa</i>	Alfalfa		
<i>Mesembryanthemum crystallinum</i>	Iceplant		
<i>Mesembryanthemum nodiflorum</i>	Slender Iceplant		

Taxon	Common Name	Declared Pest	WoNS
<i>Parapholis incurva</i>	Coast Barbgrass		
<i>Parentucellia latifolia</i>	Common Bartsia		
<i>Pentameris airoides</i> subsp. <i>airoides</i>			
<i>Plantago coronopus</i>	Buckshorn Plantain		
<i>Rostraria cristata</i>			
<i>Rostraria pumila</i>			
<i>Schismus barbatus</i>	Kelch Grass		
<i>Sonchus oleraceus</i>	Common Sowthistle		
<i>Spergularia diandra</i>	Lesser Sand Spurry		
<i>Spergularia rubra</i>	Sand Spurry		
<i>Stellaria pallida</i>			
<i>Tamarix aphylla</i>	Athel Pine	Y	Y
<i>Trifolium arvense</i> var. <i>arvense</i>			
<i>Trifolium campestre</i>	Hop Clover		
<i>Trifolium tomentosum</i> var. <i>tomentosum</i>			
<i>Ursinia anthemoides</i>	Ursinia		
<i>Vulpia bromoides</i>	Squirrel Tail Fescue		
<i>Vulpia myuros</i> forma <i>megalura</i>			
<i>Vulpia myuros</i> forma <i>myuros</i>			

The results of the literature review, combined search of the DBCA's Flora of Conservation Significance databases (DBCA, 2018a), NatureMap (DBCA, 2021) and DAWE protected matters search (DAWE, 2021) recorded no Threatened Flora or Priority Flora within the survey area. Thirteen Threatened Flora and 97 Priority Flora taxa were listed on the databases as occurring within a 40km radius of the survey area (Appendix 5). A map of flora locations is provided in Appendix 2. These taxa were assessed and ranked for their likelihood of occurrence² within the survey area. The rankings and criteria used were:

- Unlikely: Area is outside of the currently documented distribution for the species/no suitable habitat (type, quality and extent) was identified as being present during the field/desktop assessment.
- Possible: Area is within the known distribution of the species in question and habitat of at least marginal quality was identified as being present during the field/desktop assessment, supported in some cases by recent records being documented from within or near the area.
- Known to Occur: The species in question was positively identified as being present during current or previous field surveys.

² Based on habitat descriptions provided by the WA Herbarium (Florabase), habitat descriptions provided in previous records listed on the DBCA Threatened Flora Database (DBCA, 2018a) and DAWE Species Profile and Threats Database (DAWE, 2020b)

Of the 110 species listed as potentially occurring within 40km of the survey area (Appendix 5), 57 were considered 'possible' to occur within the survey area³ (Table 4-2). Five taxa were 'known to occur' within the survey area based on previous records by Botanica (2013), Astron (2014) and DBCA (2018a).

Table 4-2: Flora of Conservation Significance identified as possible or known to occur within the survey area

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Acacia</i> sp. Mt Holland (B. Ellery BE 1147)			P1	No description available	Possible
<i>Anticoryne melanosperma</i>			P3	No description available	Possible
<i>Austrostipa</i> sp. Carlingup Road (S. Kern & R. Jasper LCH 18459)			P1	No description available	Possible
<i>Austrostipa</i> sp. Mt Holland (W.A. Thompson & J. Allen 948)			P1	No description available	Possible
<i>Baeckea</i> sp. Blue Haze Mine (P. Armstrong 06/910)			P1	No description available	Possible
<i>Baeckea</i> sp. Crossroads (B.L. Rye & M.E. Trudgen 241186)			P1	No description available	Possible
<i>Baeckea</i> sp. Forrestania (K.R. Newbey 1105)			P1	No description available	Possible
<i>Baeckea</i> sp. Hatter Hill (K.R. Newbey 3284)			P3	Narrow, open, upright shrub, to 1.3 m high. Fl. pink, Jun to Oct. Yellow-orange coarse sandy loam with laterite gravel, red-brown sandy loam with quartz pebbles. Undulating plains.	Possible
<i>Baeckea</i> sp. Koonadgin (B.L. Rye & M.E. Trudgen BLR 241137)			P3	No description available	Possible
<i>Baeckea</i> sp. Lake Cronin (K.R. Newbey 9191)			P1	Upright, spreading, moderately open shrub. Fl. white/pink, Oct. Well-drained gravelly sands. Moderately exposed, gently undulating plain.	Possible
<i>Baeckea</i> sp. North Ironcap (R.J. Cranfield 10580)			P1	Erect, open shrub, to 0.4 m high. Fl. white/pink, Oct. Red clay. Gently undulating sandplains.	Possible
<i>Banksia rufa</i> subsp. <i>flavescens</i>			P3	Prostrate, ?lignotuberos shrub, to 0.45 m high. Fl. cream-yellow, Jul to Aug. Sandy loam or sand with gravel.	Possible
<i>Banksia viscida</i>			P3	Densely branched, non-lignotuberos shrub, 0.4-1 m high. Fl. yellow-orange, Jul to Oct. Gravelly soils. Lateritic rises.	Possible
<i>Boronia westringioides</i>			P2	Erect shrub, 0.2-0.75 m high. Fl. pink, Jul to Sep. Loamy sand. Plains.	Possible

³ Includes 27 taxa where habitat description are not available. In the absence of habitat description have been tentatively considered as 'possible' to occur

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Brachyloma nguba</i>			P1	Erect, compact to spreading, mid-dense shrub, to 0.8 m high, leaves discolorous, usually 2-3 mm long; style 0.2-0.25 mm long; disc truncate. Fl. red, Apr to May. White to brown sandy clay, shallow sandy loam. Open mallee woodland, mallee scrub, flat plains.	Possible
<i>Brachyloma stenolobum</i>			P1	No description available	Possible
<i>Calamphoreus inflatus</i>			P4	Erect, spreading shrub, 0.4-1.6 m high, to 2 m wide. Fl. blue-purple/green, Oct to Dec or Feb to Mar. Clay loam with ironstone gravel. Flats, disturbed sites.	Possible
<i>Calytrix nematoclada</i>			P3	Shrub, 0.15-0.5(-1) m high. Fl. purple-pink, Sep or Nov to Dec or Jan. Yellow or grey sand. Sandplains.	Possible
<i>Chorizema circinale</i>			P3	Prostrate, scrambling, wiry shrub, to 0.4 m high. Fl. yellow & orange & red, Sep to Dec. Yellow sand, sandy clay with gravel. Flats, margin of gravel pit.	Possible
<i>Cryptandra polyclada</i> subsp. <i>polyclada</i>			P3	Mat-forming or upright shrub, 0.1-0.7 m high. Fl. white/cream, Jan to May or Aug or Oct. Sand. Sandplains.	Possible
<i>Dampiera orchardii</i>			P2	Erect perennial, herb, 0.2-0.4 m high. Sand.	Possible
<i>Dampiera scaevolina</i>			P1	Erect to ascending perennial, herb or shrub, 0.2-0.5 m high. Fl. blue/white, Sep to Nov. Sandy & gravelly soils.	Possible
<i>Daviesia elongata</i> subsp. <i>implexa</i>			P3	Spreading or sprawling shrub, 0.4-1 m high. Fl. yellow/orange & red, Sep. Sand & laterite.	Possible
<i>Daviesia implexa</i>			P3	No description available	Possible
<i>Dicrastylis capitellata</i>			P1	Low spreading shrub, 0.2-0.25 m high. Fl. blue-purple, May. Loamy sand, sandy loam.	Possible
<i>Eremophila racemosa</i>			P4	Erect shrub, 0.5-1.7 m high. Fl. purple-pink-red/white, Mar or Aug to Dec. Sandy or stony loam, clay loam. Undulating plains, roadsides.	Possible
<i>Eucalyptus cerasiformis</i>			P4	Mallee, 2-3.5 m high, bark smooth, grey to brown. Fl. yellow, Dec or Jan to Mar. Red loamy soils.	Possible
<i>Eucalyptus deflexa</i>			P4	(Mallee), 1-3 m high, bark smooth. Fl. pink/cream-white, Mar or May to Oct. Clay loam, sandy loam, white or yellow sand, often with gravel. Flat areas & slight rises.	Possible
<i>Eucalyptus exigua</i>			P3	Mallee, 2-5 m high, bark smooth. Fl. white-cream, Mar. Sandy loam, white sand. Sandplains.	Possible
<i>Eucalyptus retusa</i>			P1	No description available	Possible
<i>Eutaxia hirsuta</i>			P2	Erect, shrub, spindly shrub (broom-like). Stems terete, glabrous; pustules or glands absent.	Known to occur ³
<i>Eutaxia</i> sp. North Ironcap (P. Armstrong PA 06/898)			P1	No description available	Possible

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Grevillea aneura</i>			P4	Dense, prickly shrub, 0.5-2.8 m high. Fl. red, Jun or Aug to Dec or Jan. Sand, sandy clay, gravel.	Possible
<i>Grevillea neodissecta</i>			P4	No description available	Possible
<i>Grevillea prostrata</i>			P4	Loose, prostrate shrub, 0.04-0.1 m high, 0.8-1.2 m wide. Fl. cream-white/pink-red, Aug to Dec or Jan. White, grey or yellow sand, gravel. Sandplains.	Possible
<i>Guichenotia anota</i>			P1	Shrubs, 1 m high; branchlets hairy, not glaucous. Leaves alternate, 10-25 mm long	Possible
<i>Guichenotia asteriskos</i>			P2	Erect, compact shrub, ca 0.35 m high. Fl. white, Sep to Oct. Sandy clay or loam with gravel.	Possible
<i>Hemigenia</i> sp. Newdegate (E. Bishop 75)			P1	Spindly, erect to spreading shrub, 0.2-0.45 m high, to 0.5 m wide. Fl. blue/purple, Sep to Oct. Clay loam. Disturbed sites.	Possible
<i>Hibbertia carinata</i>			P1	Shrub, to 0.4 m high. Fl. yellow, Aug to Sep. Well-drained gravelly sand, yellow sand with gravel.	Possible
<i>Hibbertia pachyphylla</i>			P3	Shrub, to 0.5 m high. Fl. yellow, Sep to Nov. White to yellow sand, brown sandy gravel, gravelly loam, laterite, granite, quartz. Undulating plains, low rises, valley floors.	Possible
<i>Hydrocotyle eichleri</i>			P3	No description available	Possible
<i>Hysterobaeckea pterocera</i>			P1	No description available	Possible
<i>Labichea rossii</i>			P1	No description available	Possible
<i>Melaleuca ochroma</i>			P3	No description available	Possible
<i>Microcorys</i> sp. Forrestania (V. English 2004)			P4	Prostrate or erect shrub, 0.35-0.4 m high. Fl. white/purple, Jan or Apr. Yellow sandy clay or red-brown clay. Open woodland or cleared areas.	Known to occur ²
<i>Microcorys</i> sp. Mt Holland (D. Angus DA 2397)			P1	No description available	Possible
<i>Microseris walteri</i>			P3	No description available	Possible
<i>Mirbelia taxifolia</i>			P1	Shrub, 0.6-0.9 m high. Fl. orange-yellow, Sep. Red or yellow sand.	Possible
<i>Notisia intonsa</i>			P3	No description available	Possible
<i>Orianthera exilis</i>			P2	No description available	Possible
<i>Oxymyrrhine plicata</i>			P3	No description available	Possible
<i>Pityrodia scabra</i> subsp. <i>dendrotricha</i>			P3	No description available	Possible
<i>Pityrodia</i> sp. Yilgarn (A.P. Brown 2679)			P3	No description available	Possible
<i>Pterostylis echinulata</i>			P3	No description available	Possible
<i>Rinzia torquata</i>			P3	No description available	Known to occur ¹
<i>Rinzia triplex</i>			P3	No description available	Known to occur ^{1,2}
<i>Seringia adenogyna</i>			P3	No description available	Possible

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Stylidium thylax</i>			P2	Creeping perennial, herb, 0.04-0.08 m high, Leaves adpressed to stem, ovate to lanceolate, 0.1-0.4 cm long, 0.6-1.5 mm wide, apex mucronate, margin hyaline, glabrous. Inflorescence uni-flowered, pedicels glandular. Fl. white, Oct. Sand. Gentle slopes and plains. Heath, mallee shrubland.	Possible
<i>Thysanotus lavanduliflorus</i>			P1	Caespitose perennial, herb (with tuberous roots), to 0.25 m high. Fl. purple, Nov to Dec. Sand, sandy loam.	Possible
<i>Thysanotus</i> sp. Yellowdine (A.S. George 6040)			P2	No description available	Possible
<i>Verticordia gracilis</i>			P3	Low, slender shrub, 0.15-0.6 m high. Fl. pink, Oct to Nov. Yellow sand, gravelly sand, sandy loam.	Known to occur ¹
<i>Verticordia stenopetala</i>			P3	Shrub, 0.2-0.6(-1.3) m high. Fl. pink/pink-purple-red, Oct to Dec or Jan. Yellow sand, sometimes with gravel. Undulating plains.	Possible

¹ DBCA record (2018a); ² Botanica record (2013); ³ Astron record (2014)

4.1.2 Fauna

According to the results of the NatureMap search (DBCA, 2021), a total of 243 vertebrate fauna taxa have been recorded within a 40 km radius of the survey area, including 10 amphibians, 145 bird, 25 mammals and 63 reptiles. Combined results of database searches (DBCA, 2021 and DAWE, 2021) identified eight introduced taxa as potentially occurring within the survey area, these being:

1. *Camelus dromedarius* (Camel)
2. *Canis lupus familiaris* (Dog)
3. *Capra hircus* (Goat)
4. *Felis catus* (Cat)
5. *Mus musculus* (House Mouse)
6. *Oryctolagus cuniculus* (Rabbit)
7. *Sus scrofa* (Pig)
8. *Vulpes vulpes* (Red Fox)

Vertebrate fauna of conservation significance identified during the literature review as previously being recorded in the general area were assessed and ranked for their likelihood of occurrence within the survey area itself (Table 4-3). The rankings and criteria used were:

- Would Not Occur: There is no suitable habitat for the species in the survey area and/or there is no documented record of the species in the general area since records have been kept and/or the species is generally accepted as being locally/regionally extinct (supported by a lack of recent records).
 - Locally Extinct: Populations no longer occur within a small part of the species natural range, in this case within 10 or 20km of the survey area. Populations do however persist outside of this area.

- Regionally Extinct: Populations no longer occur in a large part of the species natural range, in this case within the Southern Cross/ Western Mallee Bioregions. Populations do however persist outside of this area.
- Unlikely to Occur: The survey area is outside of the currently documented distribution for the species in question, or no suitable habitat (type, quality and extent) was identified as being present during the field assessment. Individuals of some species may occur occasionally as vagrants/transients especially if suitable habitat is located nearby but the site itself would not support a population or part population of the species.
- Possibly Occurs: Survey area is within the known distribution of the species in question and habitat of at least marginal quality was identified as likely to be present during the field survey and literature review, supported in some cases by recent records being documented in literature from within or near the survey area. In some cases, while a species may be classified as possibly being present at times, habitat may be marginal (e.g. poor quality, fragmented, limited in extent) and therefore the frequency of occurrence and/or population levels may be low.
- Known to Occur: The species in question has been positively identified as being present (for sedentary species) or as using the survey area as habitat for some other purpose (for non-sedentary/mobile species) during field surveys within or near the survey area. This information may have been obtained by direct observation of individuals or by way of secondary evidence (e.g. tracks, foraging debris, scats). In some cases, while a species may be classified as known to occur, habitat may be marginal (e.g. poor quality, fragmented, limited in extent) and therefore the frequency of occurrence and/or population levels may be low.

Table 4-3: Likelihood of Occurrence – Fauna Species of Conservation Significance

Species	Conservation Status			Habitat Description	Likelihood of Occurrence
	EPBC Act	BC Act	DBCA Priority		
Carnaby's Black Cockatoo <i>Calyptrorhynchus latirostris</i>	EN	EN	-	Carnaby's cockatoo is endemic to the south-west of WA, ranging from the Kalbarri in the north to Esperance in the south-east, and inland to Coorow, Kellerberrin and Lake Cronin. They are most common in semi-arid parts of the south-west. Carnaby's cockatoo occur in uncleared and remnant areas of woodland, shrubland and kwongan heath dominated by Proteaceous species. They breed in the semiarid and subhumid interior eucalypt woodlands, principally dominated by Salmon Gum Eucalyptus salmonophloia or Wandoo Eucalyptus wandoo. The Avon Wheatbelt bioregion is an important breeding area for the species. After breeding, flocks tend to migrate coastward in search of food, with the Swan Coastal Plain recognised as an important foraging area.	Unlikely to occur. This species is only recorded very infrequently this far north east of its main documented range. The survey area contains no hollow bearing trees suitable for this species to use for breeding purposes. Potential impacts anticipated to be non-existent/negligible
Grey Falcon <i>Falco hypoleucos</i>	VU	VU	-	Occurs in arid and semi-arid Australia. The species is mainly found where annual rainfall is less than 500 mm, except when wet years are followed by drought, when the species might become marginally more widespread, although it is essentially confined to the arid and semi-arid zones at all times. The species frequents timbered lowland plains, particularly acacia shrublands that are crossed by tree-lined water courses.	Unlikely to occur. This species is only recorded very infrequently in this general area. The survey area contains no suitable breeding habitat. No impact on this species conservation status will occur.
Malleefowl <i>Leipoa ocellata</i>	VU	VU	-	Occurs in unburned mallee and woodland with abundant litter and low scrub.	Possibly occurs however habitat appears very marginal/or unsuitable for breeding supported by lack of observations during survey. Occasional transients only. Potential impacts anticipated to be non-existent/negligible
Night Parrot <i>Pezoporus occidentalis</i>	EN	CR	-	Broad habitat requirements include areas of old-growth spinifex (<i>Triodia</i>) for roosting and nesting, together with foraging habitats that are likely to include various native grasses and herbs, and may or may not contain shrubs or low trees. (DPaW, 2017).	Would not occur. Preferred habitat absent. No records in southern goldfields/mallee region. No impact on this species conservation status will occur.
Fork-tailed Swift <i>Apus pacificus</i>	MI	MI	-	Low to very high airspace over varied habitat from rainforest to semi desert.	Possibly Occurs aerially over survey area on very rare occasions. No impact on this species conservation status will occur.
Peregrine Falcon <i>Falco peregrinus</i>	-	OS	-	Diverse from rainforest to arid shrublands, from coastal heath to alpine Mainly about cliffs along coasts, rivers and ranges and about wooded watercourses and lakes. The species utilises the ledges, cliff faces and large hollows/broken spouts of trees for nesting. It will also occasionally use the abandoned nests of other birds of prey.	Possibly Occurs aerially over survey area on very rare occasions. No suitable breeding habitat. Potential impacts anticipated to be non-existent/negligible

Species	Conservation Status			Habitat Description	Likelihood of Occurrence
	EPBC Act	BC Act	DBCA Priority		
Western Rosella (inland) <i>Platycercus icterotis</i> subsp. <i>xanthogenys</i>	-	-	P4	Western Rosellas are found in open eucalypt forest and timbered areas, including cultivated land and orchards. The inland species occurs in drier woodland, with a heath understorey.	Possibly Occurs however habitat appears marginal/or unsuitable for breeding. No impact on this species conservation status will occur.
Migratory shorebirds (various species)	MI	MI	P4	Migratory shorebirds generally prefer muddy edges of shallow fresh or brackish wetlands, with inundated or emergent sedges, grass, saltmarsh or other low vegetation. This includes lagoons, swamps, lakes and pools near the coast, and dams, waterholes, soaks, bore drains and bore swamps, salt pans and hypersaline salt lakes inland.	Would not occur. No suitable habitat. No impact on this species conservation status will occur.
Chuditch <i>Dasyurus geoffroii</i>	VU	VU	-	Previously occurred throughout arid and semi-arid Australia, but is now restricted to south-west Western Australia. It now currently inhabits a wide range of wooded habitats, including wet and dry sclerophyll, eucalyptus forest (especially Jarrah, Eucalyptus marginata) dry woodlands and mallee remnants.	Possibly occurs as occasional transient individuals only. Large expanses of similar habitat in surrounding areas. Potential impacts anticipated to be non-existent/negligible.
Western Brush Wallaby <i>Notamacropus imma</i>	-	-	P4	Dry sclerophyll forest and woodland, including mallee areas with grassy understorey and thickets of shrubs.	Possibly occurs as occasional transient individuals only. Large expanses of similar habitat in surrounding areas. Potential impacts anticipated to be non-existent/negligible.
Heath Mouse <i>Pseudomys shortridgei</i>	EN	VU	-	Heath mouse frequently inhabits species-rich dry heathland, and open woodland and forest habitats with a heath understorey. In both the western and eastern subpopulations there appears to be a preference for a structurally complex heath. In Western Australia, the heath mouse has been trapped mostly in species-rich heath but also in mixed scrub and mallee. The species has not been located in vegetation less than 10 years post-fire and it has been known to attain high densities in heath 30 years post-fire.	Unlikely to occur. This species has not been recorded in the general area during several detailed fauna surveys over many years. Large expanses of similar habitat in surrounding areas No impact on this species conservation status will occur
Western Mouse <i>Pseudomys occidentalis</i>			P4	The western mouse shows a preference for long unburnt habitat (between 30 and 50 yrs) on sandy clay loam or sandy loam. Vegetation in suitable habitats is variable and includes sparse low shrubland, tall dense shrubland, sparse to dense shrub mallee and mid-dense woodland. All sites where the western mouse has been collected have had patches of extremely dense vegetation. On some sites, populations occur in dense vegetation surrounded by granite rocks, which may afford them protection from fire. Quandong (<i>Santalum acuminatum</i>) and sedge species are thought to be important habitat requirements in the northern part of the western mouse's range. Populations are fragmented and restricted to this type of (fragmented) habitat.	Unlikely to occur. This species has not been recorded in the general area during several detailed fauna surveys over many years. Large expanses of similar habitat in surrounding areas No impact on this species conservation status will occur

Species	Conservation Status			Habitat Description	Likelihood of Occurrence
	EPBC Act	BC Act	DBCA Priority		
Red-tailed Phascogale <i>Phascogale calura</i>	VU	CD	-	The red-tailed phascogale occurs in remnant vegetation in the southern wheatbelt of Western Australia, where annual mean rainfall is 400–500 mm. Most of the records are concentrated in an area about 150 km long in a north-south direction from Brookton to Katanning. The red-tailed phascogale is largely confined to woodlands with old-growth hollow-producing eucalypts, particularly Wandoo (<i>Eucalyptus wandoo</i>) and York gum (<i>E. loxophleba</i>), often with associated rock sheoak (<i>Allocasuarina huegeliana</i>), but has also been recorded in shrublands and various mosaics of woodland, shrubland and scrub-heath. It avoids relatively open areas and rocky ridges which are devoid of vegetation. The species prefers long unburnt (more than 50 years) patches.	Unlikely to occur. Habitat appears to be unsuitable. In addition this species has not been recorded in the general area during several detailed fauna surveys over many years. Large expanses of similar habitat in surrounding areas No impact on this species conservation status will occur.
Lake Cronin Snake <i>Paroplocephalus atriceps</i>	-	-	P3	Eucalyptus woodlands and granite outcrops.	Possibly occurs, however as the area of impact is small and there are large expanses of similar habitat in surrounding areas potential impacts are anticipated to be non-existent/negligible
Central Long-eared Bat <i>Nyctophilus major tor</i>	-	-	P4	Appears to prefer heavy eucalypt woodlands and tall woodlands with a tall shrub understorey. Less common in open woodlands.	Possibly occurs while foraging though no nearby records. No suitable refuge habitat. Large expanses of similar habitat in surrounding areas No impact on this species conservation status will occur.


4.2 Field Assessment

4.2.1 Vegetation Types

Four vegetation types were identified within the survey area. These vegetation types were identified within three landform types and comprised of three major vegetation groups according to the NVIS, Major Vegetation Group (MVG) definition (Table 4-4). These vegetation types were represented by a total of 30 Families, 75 Genera and 179 Taxa as listed in Appendix 3. A map showing the vegetation types present in the survey area is provided in Figure 4-1.

Table 4-4: Summary of vegetation types within the survey area

Major Vegetation Group	Vegetation Type	Vegetation Code	Extent within survey area	Image
Eucalypt Woodland (MVG 5)	Low woodland of <i>Eucalyptus salmonophloia</i> over low open heathland of <i>Melaleuca adnata</i> , <i>M. calyptroides</i> , <i>M. eleuterostachya</i> on clay-loam plain	CLP-EW1	8 ha (2.0%)	
Mallee Woodland and Shrubland (MVG 14)	Open mallee shrubland of <i>Eucalyptus tenera</i> / <i>E. pileata</i> over low heathland of <i>Melaleuca adnata</i> , <i>M. calyptroides</i> , <i>M. lateriflora</i> and low open shrubland of <i>Acacia deficiens</i> / <i>A. intricata</i> on sand-loam plain	SLP-MWS1	136 ha (35.5%)	

Major Vegetation Group	Vegetation Type	Vegetation Code	Extent within survey area	Image
Heathland (MVG 18)	Mid heathland of <i>Allocasuarina campestris</i> / <i>Allocasuarina comiculata</i> , <i>Acacia eremophila</i> / <i>Acacia fragilis</i> and <i>Melaleuca cordata</i> / <i>Melaleuca hamata</i> on sandplain	SP-H1	135 ha (35.3%)	
Heathland (MVG 18)	Low heathland of <i>Banksia sessilis</i> / <i>Hakea platysperma</i> and <i>Verticordia chrysantha</i> , <i>Verticordia roei</i> and <i>Grevillea incrassata</i> on sandplain	SP-H2	69 ha (18.1%)	
N/A	Cleared Vegetation (excluding exploration)	CV	35 ha (9.1%)	No image available
Total				
			383 ha	




Figure 4-1: Vegetation types within the survey area

4.2.2 Vegetation Condition

Based on the vegetation condition rating scale adapted from Keighery, 1994 and Trudgen, 1988 (Appendix 4), vegetation was rated as 'Very Good' (Figure 4-2). 'Very Good' condition depicts vegetation structure altered by obvious signs of disturbance, for example by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing. Disturbance within the survey area was a result of existing mining infrastructure and exploration gridlines.



Figure 4-2: Vegetation condition within the survey area

4.2.3 Fauna Habitat

The broad scale terrestrial fauna habitats within the survey area presented below are based on vegetation and associated landforms identified during the flora and vegetation assessment. The extent of the identified fauna habitats and a summary description of each are provided in Table 4-5 below.

Table 4-5: Main Terrestrial Fauna Habitats within the survey area



Fauna Habitat	Description	Representative Fauna Attributes	Example Image
<u>Clay-Loam Plain</u> Eucalypt Woodland approximate area = 8 ha; 2.0%	Clay-loam plain comprising Salmon Gum woodland over <i>Melaleuca</i> spp. and mixed low shrubs.	• Range of vegetation strata suitable to a variety of passerine and nonpasserine birds. • Moderate to high leaf litter due to the presence of mature Eucalypt trees. • Relatively dense shrubs providing cover for small fauna. • Ground not especially suited to burrowing species	
<u>Sand-Loam Plain</u> Heathland/ Mallee Shrubland approximate area = 341 ha; 88.8%	Sand-loam plain comprising of dense <i>Acacia/ Allocasuarina/ Banksia/ Melaleuca</i> heathland and mallee shrubland over mixed low shrubs.	• Substrate very well suited to a variety of burrowing small mammals and reptiles. • Less diverse vegetation strata supporting a less diverse avifauna assemblage.	



Figure 4-3: Main Terrestrial Fauna Habitats within the survey area

Based on the habitats present within the survey area, a list of expected vertebrate fauna species likely to occur in the survey area was compiled from information obtained during the literature review and is presented in Appendix 6. The results of some previous fauna surveys carried out in the general area are also summarised in this species listing as are the DBCA NatureMap database search results.

Not all species listed in existing databases and publications as potentially occurring within the region (i.e. *EPBC Act* Threatened Fauna and Migratory species lists, DBCA NatureMap Fauna Database and various publications) are considered likely to be present within the survey area. The list of potential fauna takes into consideration that firstly the species in question is not known to be locally/regionally extinct and secondly that suitable habitat for each species, as identified during the habitat assessment, is present within the survey area, though compiling an accurate list has limitations (see **Section 3.3 Survey limitations and constraints**).

Table 4-6 summarises the numbers of potential species based on vertebrate class considered likely to be present in the general vicinity of the survey area based on the complete list held Appendix 6. This list has been developed based on the complete list provided in Appendix 6 and using a precautionary approach adopted for the assessment. At any one time, only a subset of the listed potential species is likely to be present within the bounds of the survey area.

Table 4-6: Summary of Potential Vertebrate Fauna Species

Group	Total number of potential species	Potential number of specially protected species	Potential number of migratory species	Potential number of priority species
Amphibians	12	0	0	0
Reptiles	65	0	0	1
Birds	121	3	0	1
Non-Volant Mammals	23 ⁶	1	0	1
Volant Mammals (Bats)	9	0	0	1
Total	230⁶	4	0	4

Superscript = number of introduced species included in the total. Note: Where a species state and federal conservation status is different, the highest category is used.

4.2.4 Introduced Species

No introduced flora or fauna species were recorded during the survey.

4.2.5 Significant Flora

According to the EPA *Environmental Factor Guideline for Flora and Vegetation* (EPA, 2016a) significant flora includes:

- flora being identified as threatened or priority species;
- locally endemic flora or flora associated with a restricted habitat type (e.g. surface water or groundwater dependent ecosystems);
- new species or anomalous features that indicate a potential new species;
- flora representative of the range of a species (particularly, at the extremes of range, recently discovered range extensions, or isolated outliers of the main range);
- unusual species, including restricted subspecies, varieties or naturally occurring hybrids; and
- flora with relictual status, being representative of taxonomic groups that no longer occur widely in the broader landscape.

No Threatened flora were identified within the survey area. Three Priority flora taxa were identified within the survey area, each of which have been previously recorded by Botanica (2013) and Astron (2014); *Eutaxia hirsuta* (P2), *Microcorys* sp. Forrestania (V. English 2004) (P4) and *Rinzia triplex* (P3). Two additional Priority Flora are listed on the DBCA database as occurring within the survey area; *Rinzia torquata* (P3) and *Verticordia gracilis* (P3) however to date neither record has been able to be confirmed. These records were obtained during previous flora surveys conducted by Frost O'Connor and Associates in 2004/ 2005 and have not been identified despite numerous searches. GPS coordinates of taxa recorded within the survey area is provided in Table 4-7 and shown in Figure 4-4.

No other significant flora (as described above) was identified within the survey area. A map showing regional Threatened and Priority Flora records in relation to the survey area is provided in Appendix 2.

Table 4-7: Significant Flora recorded within the survey area

Taxon	Coordinates	Image
<i>Eutaxia hirsuta</i> (P2)	50H 751662 6409206 ³	
	50H 751274 6408561	
	50H 751276 6408603	
	50H 751276 6408606	
	50H 751277 6408606	
	50H 751277 6408605	
	50H 751279 6408604	
	50H 751281 6408623	
	50H 751425 6408949	
<i>Microcorys</i> sp. Forrestania (V. English 2004) (P4)	50H 754547 6410246 ²	
	50H 754796 6409814 ²	

Taxon	Coordinates	Image
<i>Rinzia triplex</i> (P3)	50H 753722 6409262 ^{1,2}	

¹ DBCA record (2018a); ² Botanica record (2013); ³ Astron record (2014)




Figure 4-4: Significant flora recorded within the survey area

4.2.6 Significant Vegetation

According to the EPA *Environmental Factor Guideline for Flora and Vegetation* (EPA, 2016b) significant vegetation includes:

- vegetation being identified as threatened or priority ecological communities;
- vegetation with restricted distribution;
- vegetation subject to a high degree of historical impact from threatening processes;
- vegetation which provides a role as a refuge; and
- vegetation providing an important function required to maintain ecological integrity of a significant ecosystem.

The survey area is located adjacent to the spatially mapped 500m buffer of the *Ironcap Hills Vegetation Complexes* (North Ironcap) obtained from the DBCA communities database. This Priority Ecological Community is managed by the DBCA as a Priority 3 Ecological Community. This vegetation community was first described by Newbey and Hnatiuk (1988) during the biological survey of the Lake Johnston-Hyden area as a BIF complex and was considered one of seven unique vegetation communities to the Lake Johnston-Hyden area. The description of the North Ironcap BIF complex is provided below:

The complex vegetation on banded ironstone formation had a number of species dominant at different sites. Low trees (Acacia lasiocalyx and Eucalyptus flocktoniae [Merritt]) were rare. Mallees of Eucalyptus aff. wandoo (E. livida, E. capillosa subsp. polyclada) were usually present in small areas partially lateritized, while E. eremophila occurred rarely. Tall shrubs that were occasionally present included Allocasuarina campestris ssp. campestris (also low shrub), A. corniculata, Banksia sphaerocarpa var. dolichostyla (Ironcaps), Calothamnus quadrifidus (also low shrub), Dryandra aff. cersioides, Grevillea pterosperma, Hakea subsulcata, H. scoparia, Leptospermum erubescens, Melaleuca fulgens, M. uncinata, Santalum acuminatum and Trymalium aff. ledifolium; low shrubs were Acacia sulcata var. platyphylla, Acacia sp. (KRN 5226), Chamelaucium ciliatum (south), Cryptandra miliaris, Dodonaea adenophora, D. amblyophylla (west), Dryandra sp. (KRN 5229), Melaleuca cordata, Phebalium filifolium, P. microphyllum, P. tuberculosum ssp. tuberculosum, P. aff. tuberculosum and Platysace maxwellii (west); perennial grasses of Spartochloa scirpoidea; and sedges of Lepidosperma drummondii, L. viscidum var. viscidum, Lepidosperma sp. (KRN 5232), Lepidosperma sp. (KRN 5233) and Lepidosperma sp. (KRN 6488).

No banded ironstone was identified within the survey area and the vegetation complex described above was not represented within the survey area.

According to the DPIRD (2018) Vegetation Association GIS file, the BIF complex within the Greater North Ironcap area is classified by pre-European vegetation association Forrestania 1413; Shrublands; *Acacia, Casuarina & Melaleuca* thicket which is represented in both the Southern Cross and Western Mallee subregion. This vegetation association does not occur within the survey area (see **Section 2.3**).

No other significant vegetation (as described above) was identified within the survey area. A map showing the total extent of Priority Ecological Communities in relation to the survey area is provided in Appendix 2.

4.2.7 Significant Fauna

According to the EPA *Environmental Factor Guideline for Terrestrial Fauna* (EPA, 2016d) significant fauna includes:

- Fauna being identified as a threatened or priority species;
- Fauna species with restricted distribution;
- Fauna subject to a high degree of historical impact from threatening processes; and
- Fauna providing an important function required to maintain the ecological integrity of a significant ecosystem.

No significant fauna were observed during the survey. The current status of some species on site and/or in the general area is difficult to determine, however, based on the habitats present and, in some cases, direct observations or recent nearby records, the following species of conservation significance can be regarded as possibly utilising the survey area for some purpose at times, these being:

- **Lake Cronin Snake *Paroplocephalus atriceps* - P4 (DBCA Priority Species)**
Status of this species in the survey area is difficult to determine but it has been listed as potentially present as the survey area falls within/near the species documented range. The species does however appear to be very uncommon given the paucity of documented records in the immediate vicinity. Given the small size of the survey area and the presence of large expanses of similar habitat in surrounding areas, potential impacts are anticipated to be non-existent/negligible.
- **Carnaby's Black-Cockatoo *Calyptrorhynchus latirostris* - Endangered (EPBC Act and BC Act)**
Listed as a potential species as it has infrequently been recorded in the general area. Expected to only occur very occasionally in small transient groups. The survey area contains no potential breeding habitat for this species. No impact on this species conservation status is anticipated.
- **Western Rosella (inland ssp) *Platycercus icterotis* subsp. *xanthogenys* - P4 (DBCA Priority Species)**
Listed as a potential species as it has infrequently been recorded in the general area. Expected to only occur very occasionally in small transient groups. The survey area contains no potential breeding habitat for this species. No impact on this species conservation status is anticipated.
- **Malleefowl *Leipoa ocellata* – Vulnerable (EPBC Act and BC Act)**
Listed as a potential species as it has previously been recorded in the general area. No evidence of recent malleefowl activity (active mounds, tracks, feathers or bird observations etc.) was however observed within the survey area and therefore it is expected only to occur occasionally as transient individuals. No impact on this species conservation status is anticipated.

- **Peregrine Falcon *Falco peregrinus* – Other Specially Protected (BC Act)**
This species potentially utilises some sections of the survey area as part of a much larger home range, though records in this area are uncommon. It would not breed within the survey area. No impact on this species conservation status will occur.
- **Western Brush Wallaby *Notamacropus irma* - P4 (DBCA Priority Species)**
Listed as a potential species as it has infrequently been recorded in the general area. Expected to only occur very occasionally as individuals/pairs if at all. Given the small size of the survey area and the presence of large expanses of similar habitat in surrounding areas, potential impacts anticipated to be non-existent/negligible.
- **Chuditch *Dasyurus geoffroii* - Vulnerable (EPBC Act and BC Act)**
Listed as a potential species as it has previously been recorded in the general area. Possibly occurs as occasional transient individuals only. Given the small size of the survey area and the presence of large expanses of similar habitat in surrounding areas, potential impacts anticipated to be non-existent/negligible.
- **Central Long-eared Bat *Nyctophilus major tor* – P4 (DBCA Priority Species)**
Listed as a potential species as the survey area falls within the species documented range. The species does however appear to be uncommon in the general area given the lack of documented records in the immediate vicinity. The survey area represents potential foraging habitat but lacks refuge habitat (hollow bearing trees). No impact on this species conservation status is anticipated.

It should be noted that while habitats onsite for one or more of the species listed above are considered possibly suitable, some or all may be marginal in extent/quality and therefore the fauna species considered as possibly occurring may in fact only visit the area for short periods as infrequent vagrants.

A number of other species of conservation significance, while possibly present in the general area and/or the local region are not listed as potential species due to the survey area being outside of their currently recognised range, a lack of suitable habitat or known/very likely local or regional extinction (and no subsequent recruitment from adjoining areas).

4.3 Matters of National Environmental Significance

4.3.1 *Environment Protection and Biodiversity Conservation Act 1999*

The EPBC Act protects matters of national environmental significance, and is used by the Commonwealth DAWE to list threatened taxa and ecological communities into categories based on the criteria set out in the Act (www.environment.gov.au/epbc/index.html). The Act provides a national environmental assessment and approval system for proposed developments and enforces strict penalties for unauthorised actions that may affect matters of national environmental significance. Matters of national environmental significance as defined by the Commonwealth EPBC Act include:

- Nationally threatened flora species;
- World heritage properties;
- National heritage places;
- Wetlands of international importance (often called 'Ramsar' wetlands after the international treaty under which such wetlands are listed);
- Nationally threatened ecological communities;
- Commonwealth marine area;
- The Great Barrier Reef Marine Park; and
- Nuclear actions (including uranium mining) a water resource, in relation to coal seam gas development and large coal mining development.

No matters of national environmental significance as defined by the Commonwealth EPBC Act were identified within the survey area.

4.4 Relevant State Legislation

4.4.1 *Environmental Protection Act WA 1986*

The EP Act provides for the prevention, control and abatement of pollution and environmental harm, for the conservation, preservation, protection, enhancement and management of the environment. The Act is administered by The Department of Water and Environment Regulation (DWER), which is the State Government's environmental regulatory agency.

Under Section 51C of the EP Act and the *Environmental Protection (Clearing of Native Vegetation) Regulations (Regulations) WA 2004* any clearing of native vegetation in Western Australia that is not eligible for exemption under Schedule 6 of the *EP Act 1986* or under the Regulations 2004 requires a clearing permit from the DWER or DMIRS. Under Section 51A of the *EP Act 1986* native vegetation includes aquatic and terrestrial vegetation indigenous to Western Australia, and intentionally planted vegetation declared by regulation to be native vegetation, but not vegetation planted in a plantation or planted with commercial intent. Section 51A of the *EP Act 1986* defines clearing as "the killing or destruction of; the removal of; the severing or ringbarking of trunks or stems of; or the doing of substantial damage to some or all of the native vegetation in an area, including the flooding of land, the burning of vegetation, the grazing of stock or an act or activity that results in the above". Exemptions under Schedule 6 of the EP Act and the EP Regulations do not apply in ESAs as declared under Section 51B of the EP Act or TEC listed under State and Commonwealth legislation.

No evidence of the survey area containing any TEC or Threatened Flora or Fauna was identified during the survey. Approximately 375 ha of the survey area is located within an ESA

which encompasses an approximate 10km radius of Lake Cronin as listed under the *Environmental Protection Act 1986* (EP Act).

4.4.2 Biodiversity Conservation Act 2016

This Act is used by the Western Australian DBCA for the conservation and protection of biodiversity and biodiversity components in Western Australia and to promote the ecologically sustainable use of biodiversity components in the State. Taxa are classified as ‘Threatened’ when their populations are geographically restricted or are threatened by local processes (see following sections for Threatened definitions). Under this Act all native flora and fauna are protected throughout the State. Financial penalties are enforced under this Act if threatened species are collected without an appropriate licence.

Under Section 54(1) of the BC Act, habitat is eligible for listing as critical habitat if:

- (a) it is critical to the survival of a threatened species or a threatened ecological community;*
- and*
- (b) its listing is otherwise in accordance with the ministerial guidelines.*

No threatened species or critical habitat listed under the BC Act were recorded within the survey area.

4.4.3 Conservation Reserves

The survey area is not located within a vested Conservation Reserve. The closest vested Conservation Reserve is the Lake Cronin Nature Reserve, located approximately 4.5km north-east of the survey area which is managed by DBCA as a Class A Reserve.

According to the EPA (2009) *Advice on Conservation Values and Review of Nature Reserve Proposals in the Lake Cronin Region*, the survey area occurs within a 56,750ha area within the mineralised greenstone belt in the Lake Cronin Region which is proposed to be managed under Section 33(2) of the *Conservation and Land Management Act 1984* (CALM Act) but not formally reserved. A map showing conservation areas in relation to the survey area is provided in Appendix 2.

4.5 Native Vegetation Clearing Principles

Based on the outcomes from the survey undertaken, as presented in this report, Botanica provides the following comments regarding the native vegetation clearing principles listed under Schedule 5 of the EP Act (Table 4-8).

Table 4-8: Assessment of development within the survey area against native vegetation clearing principles

Letter	Principle	Assessment	Outcome
Native vegetation should not be cleared if it:			
(a)	comprises a high level of biological diversity.	Woodlands of the Southern Cross subregion have a very high diversity of Eucalypts with as many as 170 species occurring within the subregion. The subregion also has a high diversity of Acacia species however most species (excluding Threatened and Priority Flora species) are wide ranging and usually occur in at least one, and often several, adjoining subregions (Cowan, 2001). With the exception of granite outcrops vegetation of the Western Mallee subregion is not considered to comprise of a high level of biodiversity however it does contain a high number of endemic species (Beecham & Danks, 2001). Vegetation identified within the survey area is not considered to be of high biological diversity, and is well represented outside of the survey area.	Clearing is unlikely to be at variance to this principle
(b)	comprises the whole or part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to WA.	No significant fauna were observed within the survey area. No evidence of Malleefowl activity was identified during the targeted survey. The survey area comprises of broad fauna habitats that are typical of those in the wider region. No unique fauna habitats (caves, rocky outcrops/ pools etc.) occur within the survey area. No water bodies (both perennial/ non-perennial) occur within the survey area.	Clearing is unlikely to be at variance to this principle
(c)	includes, or is necessary for the continued existence of rare flora.	No Threatened Flora taxa, pursuant to the BC Act and the EPBC Act were identified within the survey area.	Clearing is not at variance to this principle
(d)	comprises the whole or part of or is necessary for the maintenance of a threatened ecological community (TEC).	No TEC listed under the EPBC Act or by the BC Act occur within the survey area.	Clearing is not at variance to this principle
(e)	is significant as a remnant of native vegetation in an area that has been extensively cleared	The survey area occurs within the pre-European Beard vegetation association Forrestania 511 of the Southern Cross subregion and Forrestania 2048 Western Mallee subregion which retain >97% of their original pre-European vegetation extent.	Clearing is unlikely to be at variance to this principle
(f)	is growing, in, or in association with, an environment associated with a watercourse or wetland	There are no inland waters (lakes/ playas) or any perennial/ ephemeral drainage lines within the survey area.	Clearing is unlikely to be at variance to this principle
(g)	Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.	The survey area occurs within the pre-European Beard vegetation association Forrestania 511 of the Southern Cross subregion and Forrestania 2048 Western Mallee subregion which retain >97% of their original pre-European vegetation extent. Clearing within these vegetation associations	Clearing is unlikely to be at variance to this principle

Letter	Principle	Assessment	Outcome
	Native vegetation should not be cleared if it:		
		is not likely to lead to land degradation issues such as salinity, water logging or acidic soils.	
(h)	Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.	The survey area is not located within a gazetted conservation area. According to the EPA (2009) Advice on Conservation Values and Review of Nature Reserve Proposals in the Lake Cronin Region, the survey area occurs within a 56,750ha area within the mineralised greenstone belt in the Lake Cronin Region which is proposed to be managed under Section 33(2) of the CALM Act 1984 but not formally reserved. Given the small size of the survey area (54 ha) in relation to this proposed management area (56,750ha) significant impact to this land is unlikely.	Clearing is unlikely to be at variance to this principle
(i)	Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.	There are no inland waters (lakes/ playas) or any perennial/ ephemeral drainage lines within the survey area. Most rainfall is lost by evaporation or surface runoff. Only a small portion infiltrates the soil and recharges the groundwater.	Clearing is unlikely to be at variance to this principle
(j)	Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence of flooding	Rainfall is highly variable with an average annual rainfall of 339.6 mm and an evaporation rate of 2000 mm. The region is not prone to flooding and does not contain riparian vegetation.	Clearing is unlikely to be at variance to this principle

5 Recommendations

- Clearing Permit approval under the EP Act is required prior to any clearing within the survey area as the survey area is located within an ESA listed under the EP Act.
- Disturbance to Priority Flora taxa should be avoided where possible, including maintaining a 10m exclusion zone surrounding Priority Flora. Should avoidance not be possible, consultation with the DBCA Species and Communities Program is recommended prior to clearing of Priority Flora.

6 **Bibliography**

- Anstis, M. (2013). *Tadpoles and Frogs of Australia*. New Holland Publishers, Sydney.
- Aplin, K. P. and Smith, L.A. (2001). *Checklist of the frogs and reptiles of Western Australia*, Records of the Western Australian Museum Supplement No. 63, 51-74.
- ASRIS (2014). *Atlas of Australian Soils Database*. Australian Soil Resource Information System.
- Astron (2014). Forrester Nickel Operations Lounge Lizard Vegetation, Flora and Fauna Biological Assessment, prepared for Western Areas Limited, Astron Environmental Services.
- Australian Government (2018). *National Map*. Australian Government.
- Barrett, G., Silcocks, A., Barry, S., Cunningham, R. and Poulter, R. (2003). *The New Atlas of Australian Birds*. Royal Australasian Ornithologists Union, Victoria.
- Beecham, B. and Danks, A., (2001) *A Biodiversity Audit of Western Australia's 53 Biogeographical Region in 2001- Western Mallee subregion*, Department of Conservation and Land Management.
- Beard, J.S., (1990). *Plant Life of Western Australia*, Kangaroo Press Pty Ltd, NSW.
- Biota (2007a). Forrester Monitoring Survey, Flying Fox Phases I, II, III and IV. Unpublished report for Western Areas NL.
- Biota (2010). Spotted Quoll Haul Road Single Phase Fauna Survey. Unpublished report for Western Areas N.L. May 2010.
- Biota (2018), New Morning Level 1 and Targeted Terrestrial fauna Survey, prepared for Western Areas Limited, Biota Environmental Sciences.
- BoM (2020). Groundwater Dependent Ecosystems Atlas. Bureau of Meteorology
Available: <http://www.bom.gov.au/water/groundwater/gde/map.shtml>
- BoM, (2021). Hyden weather station climate data, Bureau of Meteorology.
Available: <http://www.bom.gov.au/climate>
- Botanica (2006), Flora and Vegetation Survey of the Flying Fox North East Exploration Area for Western Areas Limited, Botanica Consulting.
- Botanica (2007a), Vegetation Survey of a Proposed Extension to the Current Clearing Permit Number 691/1 at the Flying Fox mine site prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2007b), Flora and Vegetation Survey within the Greater Flying Fox mine site prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2009), Flora and Vegetation Survey within the lounge Lizard/Flying Fox area, Proposed Gravel Pit, prepared for Western Areas Limited, Botanica Consulting.
- Botanica (2013), Flora and Vegetation survey of the greater Flying Fox area, prepared for Western Areas Limited, Botanica Consulting.

Botanica (2021a), Detailed Flora and Vegetation Survey and Targeted Flora survey of the New Morning Project, prepared for Western Areas Limited, Botanica Consulting.

Botanica (2021b), Reconnaissance Flora/ Vegetation Survey & Basic Fauna Survey Lounge Lizard East Sand Pit Extension, prepared for Western Areas Limited, Botanica Consulting.

Cowan M, Graham G & McKenzie N, (2001), *A Biodiversity Audit of Western Australia's 53 Biogeographical Region in 2001- Southern Cross Subregion*, Department of Conservation and Land Management.

DAFWA (2014). *Soil Landscape System of Western Australia*, Department of Agriculture and Food Western Australia

DAWE (2021). *Protected Matters Search Tool, Environment Protection and Biodiversity Conservation Act 1999*, Department of Agriculture, Water and Environment.

DBCA (2018a). *Threatened and Priority Flora Database search results*, Department of Biodiversity, Conservation and Attractions. Results obtained May 2018.

DBCA (2018b). *Threatened Ecological Community and Priority Ecological Community database search*, Department of Biodiversity, Conservation and Attractions. Results obtained May 2018.

DBCA (2019). *2018 Statewide Vegetation Statistics (formerly the CAR Reserve Analysis)*. Department of Biodiversity, Conservation and Attractions.

DBCA (2021). *NatureMap Database search*, Department of Biodiversity, Conservation and Attractions.

DotEE (2012). *Interim Biogeographic Regionalisation for Australia (IBRA)*, Version 7, Department of the Environment and Energy.

DotEE (2017). *National Vegetation Information System (NVIS) Major Vegetation Groups, Version 4.2*, Department of the Environment and Energy.

DPIRD (2018). *Pre-European Vegetation - Western Australia (NVIS Compliant Version GIS file)*, Department of Primary Industries and Regional Development.

DPIRD (2020). *Declared Organism-database search*, Department of Primary Industries and Regional Development, Western Australia.

Available: <http://www.biosecurity.wa.gov.au/>

Duncan, Anne. & Baker, G. B. & Montgomery, Narelle. & Natural Heritage Trust (Australia) (1999). *The action plan for Australian bats* / edited by Anne Duncan, G. Barry Baker and Narelle Montgomery; with assistance from Lindy Lumsden *et al.* Natural Heritage Trust, Canberra.

EPA (2016a). *Technical Guide - Flora and Vegetation Surveys for Environmental Impact Assessment – December 2016*. Environmental Protection Authority.

EPA (2020). *Technical Guide – Terrestrial Fauna Surveys for Environmental Impact Assessment – June 2020*. Environmental Protection Authority.

Geoscience Australia (2015). *Surface Hydrology 250K Topographical Map*. Australian Government.

Gibson (2004) Flora and vegetation of the Eastern Goldfields Ranges: Part 7. Middle and South Ironcap, Digger Rock and Hatter Hill. Science Division, Department of Conservation and Land Management.

Hart, R.P. and Kitchener, D.J., (1986). *First Record of *Sminthopsis psammophila* (Marsupialia: Dasyuridae) from Western Australia*. Records of the Western Australian Museum 13(1): 139-144.

Hatton T, Evans R, (1998). *Dependence of ecosystems on groundwater and its significance to Australia*. LWRRDC Occasional Paper No 12/98.

How, R., Cooper, N.K. and Bannister, J.L. (2001). *Checklist of the mammals of Western Australia*, Records of the Western Australian Museum Supplement No. 63, 91-98.

How, R. A Newbey, K.R Dell, J. Muir, B.G & Hnatiuk, R.J, (1988), *The Biological survey of the Eastern Goldfields of Western Australia: Lake Johnston-Hyden*. Western Australian Museum Supplement No. 30.

Jackson, S. & Groves, C. (2015). *Taxonomy of Australian mammals*. CSIRO Publishing.

Johnstone, R.E. (2001). *Checklist of the birds of Western Australia*, Records of the Western Australian Museum Supplement No. 63, 75-90.

Johnstone, R.E. and Storr, G.M. (1998). *Handbook of Western Australian Birds: Volume 1 – Non-passerines (Emu to Dollarbird)*. Western Australian Museum, Perth Western Australia.

Johnstone, R.E. and Storr, G.M. (2004). *Handbook of Western Australian Birds: Volume 2 – Passerines (Blue-winged Pitta to Goldfinch)*. Western Australian Museum, Perth Western Australia.

Keighery, B. J., (1994). *Bushland Plant Survey: A guide to plant community survey for the community*. Wildflower Society of Western Australia (Inc.), Nedlands.

Morcombe, M. (2004). *Field Guide to Australian Birds*. Steve Parish Publishing, Archerfield, Queensland.

National Committee on Soil and Terrain (2009). *Australian soil and land survey field handbook* (3rd edn). CSIRO Publishing: Melbourne.

Pizzey, G & Knight, F. (2012). *The Field Guide to the Birds of Australia*. 9th Edition. Harper Collins, Sydney.

Simpson, K. and Day, N. (2010). *Field Guide to the Birds of Australia*. Penguin Books, Ringwood.

Tille, P. (2006). *Soil Landscapes of Western Australia's Rangelands and Arid Interior*, Department of Agriculture and Food Western Australia

WAHERB, (2021). *Florabase – Information on the Western Australian Flora*, Department of Biodiversity, Conservation and Attractions.

Available: <https://florabase.dpaw.wa.gov.au/>

Watson, A., Judd, S., Watson J., Lam, A. and Mackenzie, D., (2008) *The Extraordinary Nature of The Great Western Woodlands*. The Wilderness Society of WA.

Wilson, S. and Swan, G. (2017). *A Complete Guide to Reptiles of Australia*. Reed, New Holland, Sydney.

Appendix 1: Conservation Ratings BC Act and EPBC Act

Definitions of Conservation Significant Species

Code	Category
State categories of threatened and priority species	
Threatened Species (T)	
Listed by order of the Minister as Threatened in the category of critically endangered, endangered or vulnerable under section 19(1), or is a rediscovered species to be regarded as threatened species under section 26(2) of the Biodiversity Conservation Act 2016 (BC Act).	
CR	Critically Endangered Threatened species considered to be “facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with criteria set out in the ministerial guidelines”. Listed as critically endangered under section 19(1)(a) of the BC Act in accordance with the criteria set out in section 20 and the ministerial guidelines. Published under schedule 1 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018 for critically endangered fauna or the Wildlife Conservation (Rare Flora) Notice 2018 for critically endangered flora.
EN	Endangered Threatened species considered to be “facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines”. Listed as endangered under section 19(1)(b) of the BC Act in accordance with the criteria set out in section 21 and the ministerial guidelines. Published under schedule 2 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018 for endangered fauna or the Wildlife Conservation (Rare Flora) Notice 2018 for endangered flora.
VU	Vulnerable Threatened species considered to be “facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with criteria set out in the ministerial guidelines”. Listed as vulnerable under section 19(1)(c) of the BC Act in accordance with the criteria set out in section 22 and the ministerial guidelines. Published under schedule 3 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018 for vulnerable fauna or the Wildlife Conservation (Rare Flora) Notice 2018 for vulnerable flora.
Extinct species	
Listed by order of the Minister as extinct under section 23(1) of the BC Act as extinct or extinct in the wild.	
EX	Extinct Species where “<i>there is no reasonable doubt that the last member of the species has died</i>”, and listing is otherwise in accordance with the ministerial guidelines (section 24 of the BC Act). Published as presumed extinct under schedule 4 of the <i>Wildlife Conservation (Specially Protected Fauna) Notice 2018</i> for extinct fauna or the <i>Wildlife Conservation (Rare Flora) Notice 2018</i> for extinct flora.
EW	Extinct in the Wild Species that “<i>is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; and it has not been recorded in its known habitat or expected habitat, at appropriate seasons, anywhere in its past range, despite surveys over a time frame appropriate to its life cycle and form</i>”, and listing is otherwise in accordance with the ministerial guidelines (section 25 of the BC Act). Currently there are no threatened fauna or threatened flora species listed as extinct in the wild. If listing of a species as extinct in the wild occurs, then a schedule will be added to the applicable notice.
Specially protected species	
Listed by order of the Minister as specially protected under section 13(1) of the BC Act. Meeting one or more of the following categories: species of special conservation interest; migratory species; cetaceans; species subject to international agreement; or species otherwise in need of special protection.	
Species that are listed as threatened species (critically endangered, endangered or vulnerable) or extinct species under the BC Act cannot also be listed as Specially Protected species.	
IA	International Agreement/ Migratory Fauna that periodically or occasionally visit Australia or an external Territory or the exclusive economic zone; or the species is subject of an international agreement that relates to the protection of migratory species and that binds the Commonwealth; and listing is otherwise in accordance with the ministerial guidelines (section 15 of the BC Act). Includes birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and fauna subject to the <i>Convention on the Conservation of Migratory Species of Wild Animals</i> (Bonn Convention), an environmental treaty under the United Nations Environment Program. Migratory species listed under the BC Act are a subset of the migratory animals, that are known to visit Western Australia, protected under the international agreements or treaties, excluding species that are listed as Threatened species. Published as migratory birds protected under an international agreement under schedule 5 of the <i>Wildlife Conservation (Specially Protected Fauna) Notice 2018</i>.

Code	Category
CD	Species of special conservation interest Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened, and listing is otherwise in accordance with the ministerial guidelines (section 14 of the BC Act). Published as conservation dependent fauna under schedule 6 of the <i>Wildlife Conservation (Specially Protected Fauna) Notice 2018</i> .
OS	Other specially protected species Fauna otherwise in need of special protection to ensure their conservation, and listing is otherwise in accordance with the ministerial guidelines (section 18 of the BC Act). Published as other specially protected fauna under schedule 7 of the <i>Wildlife Conservation (Specially Protected Fauna) Notice 2018</i> .
Priority species Possibly threatened species that do not meet survey criteria, or are otherwise data deficient, are added to the Priority Fauna or Priority Flora Lists under Priorities 1, 2 or 3. These three categories are ranked in order of priority for survey and evaluation of conservation status so that consideration can be given to their declaration as threatened fauna or flora. Species that are adequately known, are rare but not threatened, or meet criteria for near threatened, or that have been recently removed from the threatened species or other specially protected fauna lists for other than taxonomic reasons, are placed in Priority 4. These species require regular monitoring. Assessment of Priority codes is based on the Western Australian distribution of the species, unless the distribution in WA is part of a contiguous population extending into adjacent States, as defined by the known spread of locations.	
P1	Priority 1: Poorly-known species Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.
P2	Priority 2: Poorly-known species Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.
P3	Priority 3: Poorly-known species Species that are known from several locations, and the species does not appear to be under imminent threat, or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. Such species are in need of further survey.
P4	Priority 4: Rare, Near Threatened and other species in need of monitoring (a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection but could be if present circumstances change. These species are usually represented on conservation lands. (b) Near Threatened. Species that are considered to have been adequately surveyed and that are close to qualifying for vulnerable but are not listed as Conservation Dependent. (c) Species that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.
Commonwealth categories of threatened species	
EX	Extinct Taxa where there is no reasonable doubt that the last member of the species has died.
EW	Extinct in the Wild Taxa where it is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; or it has not been recorded in its known and/or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
CR	Critically Endangered Taxa that are facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria.
EN	Endangered Taxa which are not critically endangered and is facing a very high risk of extinction in the wild in the near future, as determined in accordance with the prescribed criteria.


Code	Category
VU	Vulnerable Taxa which are not critically endangered or endangered and is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
CD	Conservation Dependent Taxa which are the focus of a specific conservation program the cessation of which would result in the species becoming vulnerable, endangered or critically endangered; or (b) the following subparagraphs are satisfied: (i) the species is a species of fish; (ii) the species is the focus of a plan of management that provides for actions necessary to stop the decline of, and support the recovery of, the species so that its chances of long term survival in nature are maximised; (iii) the plan of management is in force under a law of the Commonwealth or of a State or Territory; (iv) cessation of the plan of management would adversely affect the conservation status of the species.

Definitions of Conservation Significant Communities

Category Code	Category
State categories of Threatened Ecological Communities (TEC)	
PD	Presumed Totally Destroyed An ecological community will be listed as Presumed Totally Destroyed if there are no recent records of the community being extant and either of the following applies: - records within the last 50 years have not been confirmed despite thorough searches or known likely habitats or; - all occurrences recorded within the last 50 years have since been destroyed.
CR	Critically Endangered An ecological community will be listed as Critically Endangered when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future, meeting any one of the following criteria: The estimated geographic range and distribution has been reduced by at least 90% and is either continuing to decline with total destruction imminent, or is unlikely to be substantially rehabilitated in the immediate future due to modification; The current distribution is limited i.e. highly restricted, having very few small or isolated occurrences, or covering a small area; The ecological community is highly modified with potential of being rehabilitated in the immediate future.
EN	Endangered An ecological community will be listed as Endangered when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future. The ecological community must meet any one of the following criteria: The estimated geographic range and distribution has been reduced by at least 70% and is either continuing to decline with total destruction imminent in the short-term future, or is unlikely to be substantially rehabilitated in the short-term future due to modification; The current distribution is limited i.e. highly restricted, having very few small or isolated occurrences, or covering a small area; The ecological community is highly modified with potential of being rehabilitated in the short-term future.
VU	Vulnerable An ecological community will be listed as Vulnerable when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing high risk of total destruction in the medium to long term future. The ecological community must meet any one of the following criteria: The ecological community exists largely as modified occurrences that are likely to be able to be substantially restored or rehabilitated; The ecological community may already be modified and would be vulnerable to threatening process, and restricted in range or distribution; The ecological community may be widespread but has potential to move to a higher threat category due to existing or impending threatening processes.

Category Code	Category
Commonwealth categories of Threatened Ecological Communities (TEC)	
CE	Critically Endangered If, at that time, an ecological community is facing an extremely high risk of extinction in the wild in the immediate future (indicative timeframe being the next 10 years).
EN	Endangered If, at that time, an ecological community is not critically endangered but is facing a very high risk of extinction in the wild in the near future (indicative timeframe being the next 20 years).
VU	Vulnerable If, at that time, an ecological community is not critically endangered or endangered, but is facing a high risk of extinction in the wild in the medium-term future (indicative timeframe being the next 50 years).
Priority Ecological Communities (PEC)	
P1	Poorly-known ecological communities Ecological communities with apparently few, small occurrences, all or most not actively managed for conservation (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) and for which current threats exist.
P2	Poorly-known ecological communities Communities that are known from few small occurrences, all or most of which are actively managed for conservation (e.g. within national parks, conservation parks, nature reserves, State forest, un-allocated Crown land, water reserves, etc.) and not under imminent threat of destruction or degradation.
P3	Poorly known ecological communities Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or: Communities known from a few widespread occurrences, which are either large or within significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or; Communities made up of large, and/or widespread occurrences, that may or not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing and inappropriate fire regimes.
P4	Ecological communities that are adequately known, rare but not threatened or meet criteria for near threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.
P5	Conservation Dependent ecological communities Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

Appendix 2: Regional map of the survey area in relation to conservation areas



Appendix 3: List of species identified within each vegetation type

Red text=Priority species (WAHERB, 2021)

Family	Genus	Taxon	CLP-EW1	SLP-MWS1	SP-H1	SP-H2
Apiaceae	<i>Platysace</i>	<i>deflexa</i>			*	
Apiaceae	<i>Platysace</i>	<i>maxwellii</i>		*		
Asparagaceae	<i>Lomandra</i>	<i>effusa</i>			*	
Asparagaceae	<i>Lomandra</i>	<i>mucronata</i>	*		*	
Asteraceae	<i>Olearia</i>	<i>dampieri</i> subsp. <i>eremicola</i>			*	
Asteraceae	<i>Olearia</i>	<i>lanuginosa</i>		*		
Asteraceae	<i>Olearia</i>	<i>muelleri</i>			*	
Boraginaceae	<i>Halgania</i>	<i>andromedifolia</i>			*	
Boraginaceae	<i>Halgania</i>	<i>lavandulacea</i>	*			
Boryaceae	<i>Borya</i>	<i>constricta</i>			*	
Casuarinaceae	<i>Allocasuarina</i>	<i>acutivalvis</i>	*	*	*	
Casuarinaceae	<i>Allocasuarina</i>	<i>campestris</i>			*	
Casuarinaceae	<i>Allocasuarina</i>	<i>corniculata</i>			*	*
Casuarinaceae	<i>Allocasuarina</i>	<i>decussata</i>	*	*	*	
Casuarinaceae	<i>Allocasuarina</i>	<i>humilis</i>			*	
Casuarinaceae	<i>Allocasuarina</i>	<i>microstachya</i>				*
Celastraceae	<i>Psammomoya</i>	<i>choretroides</i>			*	
Chenopodiaceae	<i>Atriplex</i>	<i>vesicaria</i>	*			
Chenopodiaceae	<i>Chenopodium</i>	<i>curvispicatum</i>	*			
Chenopodiaceae	<i>Maireana</i>	<i>pyramidata</i>			*	
Chenopodiaceae	<i>Rhagodia</i>	<i>preissii</i> subsp. <i>preissii</i>			*	
Chenopodiaceae	<i>Sclerolaena</i>	<i>diacantha</i>	*			
Chenopodiaceae	<i>Sclerolaena</i>	<i>uniflora</i>	*			
Convolvulaceae	<i>Wilsonia</i>	<i>humilis</i>		*		
Cupressaceae	<i>Callitris</i>	<i>canescens</i>		*		
Cupressaceae	<i>Callitris</i>	<i>preissii</i>	*	*	*	*
Cyperaceae	<i>Gahnia</i>	<i>ancistrophylla</i>		*	*	
Cyperaceae	<i>Lepidosperma</i>	<i>drummondii</i>	*	*	*	
Cyperaceae	<i>Lepidosperma</i>	<i>sanguinolentum</i>				
Cyperaceae	<i>Schoenus</i>	<i>brevisetis</i>				
Dasypogonaceae	<i>Calectasia</i>	<i>valida</i>	*			
Dilleniaceae	<i>Hibbertia</i>	<i>ancistrophylla</i>			*	
Dilleniaceae	<i>Hibbertia</i>	<i>gracilipes</i>			*	*
Ericaceae	<i>Leucopogon</i>	sp. Coujinup (Burgman 1085)			*	
Ericaceae	<i>Leucopogon</i>	sp. Wheatbelt (S. Murray 257)	*			
Ericaceae	<i>Lysinema</i>	<i>ciliatum</i>	*		*	*
Ericaceae	<i>Styphelia</i>	<i>epacridis</i>			*	
Ericaceae	<i>Styphelia</i>	<i>serratifolia</i>	*		*	
Ericaceae	<i>Styphelia</i>	<i>conostephioides</i>	*			
Ericaceae	<i>Styphelia</i>	<i>dielsianus</i>			*	
Ericaceae	<i>Styphelia</i>	<i>lissanthoides</i>			*	
Euphorbiaceae	<i>Bertya</i>	<i>dimerostigma</i>			*	
Fabaceae	<i>Acacia</i>	<i>assimilis</i> subsp. <i>atroviridis</i>	*			
Fabaceae	<i>Acacia</i>	<i>binata</i>		*		
Fabaceae	<i>Acacia</i>	<i>brachyclada</i>			*	
Fabaceae	<i>Acacia</i>	<i>deficiens</i>		*	*	
Fabaceae	<i>Acacia</i>	<i>densiflora</i>			*	
Fabaceae	<i>Acacia</i>	<i>eremophila</i> var. <i>eremophila</i>			*	

Family	Genus	Taxon	CLP-EW1	SLP-MWS1	SP-H1	SP-H2
Fabaceae	<i>Acacia</i>	<i>erinacea</i>	*	*	*	
Fabaceae	<i>Acacia</i>	<i>fragilis</i>		*	*	
Fabaceae	<i>Acacia</i>	<i>hemiteles</i>		*	*	
Fabaceae	<i>Acacia</i>	<i>heteroneura</i> var. <i>jutsonii</i>			*	
Fabaceae	<i>Acacia</i>	<i>intricata</i>		*		
Fabaceae	<i>Acacia</i>	<i>leptopetala</i>		*	*	
Fabaceae	<i>Acacia</i>	<i>neuophylla</i> subsp. <i>erugata</i>			*	
Fabaceae	<i>Acacia</i>	<i>sulcata</i> var. <i>platyphylla</i>		*		
Fabaceae	<i>Aotus</i>	sp. Tortile (G.J. Keighery 3767)		*		
Fabaceae	<i>Aotus</i>	<i>tietkensis</i>		*		
Fabaceae	<i>Chorizema</i>	<i>aciculare</i> subsp. <i>aciculare</i>				*
Fabaceae	<i>Daviesia</i>	<i>benthamii</i>	*	*	*	
Fabaceae	<i>Daviesia</i>	<i>brachyphylla</i>	*			
Fabaceae	<i>Daviesia</i>	<i>lancifolia</i>	*		*	
Fabaceae	<i>Daviesia</i>	<i>nematophylla</i>			*	
Fabaceae	<i>Daviesia</i>	<i>pachyloma</i>			*	
Fabaceae	<i>Dillwynia</i>	<i>divaricata</i>		*		
Fabaceae	<i>Dillwynia</i>	<i>uncinata</i>			*	
Fabaceae	<i>Eutaxia</i>	<i>hirsuta</i> (P2)				*
Fabaceae	<i>Gastrolobium</i>	<i>melanocarpum</i>			*	
Fabaceae	<i>Gastrolobium</i>	<i>spinosum</i>	*		*	
Fabaceae	<i>Gastrolobium</i>	<i>trilobum</i>			*	*
Fabaceae	<i>Gompholobium</i>	<i>gompholobioides</i>	*			
Fabaceae	<i>Jacksonia</i>	<i>nematoclada</i>		*		
Fabaceae	<i>Senna</i>	<i>artemisioides</i> subsp. <i>filifolia</i>			*	
Fabaceae	<i>Templetonia</i>	<i>egena</i>			*	
Fabaceae	<i>Templetonia</i>	<i>sulcata</i>	*			
Goodeniaceae	<i>Coopernookia</i>	<i>strophiolata</i>		*	*	
Goodeniaceae	<i>Dampiera</i>	<i>angulata</i>			*	
Goodeniaceae	<i>Dampiera</i>	<i>eriocephala</i>			*	
Goodeniaceae	<i>Goodenia</i>	<i>pinifolia</i>	*		*	
Goodeniaceae	<i>Scaevola</i>	<i>spinescens</i>	*			
Haloragaceae	<i>Glischrocaryon</i>	<i>roei</i>	*			
Hemerocallidaceae	<i>Dianella</i>	<i>revoluta</i>			*	
Lamiaceae	<i>Dasymalla</i>	<i>terminalis</i>		*		
Lamiaceae	<i>Hemiphora</i>	<i>lanata</i>		*		
Lamiaceae	<i>Microcorys</i>	sp. Forrestania (V. English 2004) (P4)		*	*	
Lamiaceae	<i>Westringia</i>	<i>cephalantha</i>		*	*	
Lamiaceae	<i>Westringia</i>	<i>rigida</i>		*		
Lauraceae	<i>Cassytha</i>	<i>glabella</i>			*	
Lauraceae	<i>Cassytha</i>	<i>melantha</i>		*	*	
Myrtaceae	<i>Balaustion</i>	<i>pulcherrimum</i>	*		*	
Myrtaceae	<i>Beaufortia</i>	<i>interstans</i>	*		*	
Myrtaceae	<i>Beaufortia</i>	<i>schaueri</i>	*		*	
Myrtaceae	<i>Calytrix</i>	<i>breviseta</i> subsp. <i>stipulosa</i>			*	
Myrtaceae	<i>Calytrix</i>	<i>duplistipulata</i>			*	
Myrtaceae	<i>Calytrix</i>	<i>violacea</i>			*	
Myrtaceae	<i>Cyathostemon</i>	<i>ambiguus</i>		*	*	
Myrtaceae	<i>Cyathostemon</i>	<i>tenuifolius</i>			*	
Myrtaceae	<i>Darwinia</i>	<i>inconspicua</i>	*	*	*	*
Myrtaceae	<i>Eucalyptus</i>	<i>calycogona</i>		*		

Family	Genus	Taxon	CLP-EW1	SLP-MWS1	SP-H1	SP-H2
Myrtaceae	<i>Eucalyptus</i>	<i>celastroides</i> subsp. <i>celastroides</i>	*			
Myrtaceae	<i>Eucalyptus</i>	<i>cylindriflora</i>		*		
Myrtaceae	<i>Eucalyptus</i>	<i>cylindrocarpa</i>			*	
Myrtaceae	<i>Eucalyptus</i>	<i>horistes</i>	*			*
Myrtaceae	<i>Eucalyptus</i>	<i>pileata</i>		*	*	
Myrtaceae	<i>Eucalyptus</i>	<i>platycorys</i>	*		*	
Myrtaceae	<i>Eucalyptus</i>	<i>polita</i>			*	
Myrtaceae	<i>Eucalyptus</i>	<i>rigidula</i>			*	*
Myrtaceae	<i>Eucalyptus</i>	<i>salmonophloia</i>		*		
Myrtaceae	<i>Eucalyptus</i>	<i>tenera</i>		*	*	*
Myrtaceae	<i>Eucalyptus</i>	<i>transcontinentalis</i>		*		
Myrtaceae	<i>Leptospermum</i>	<i>erubescens</i>	*	*	*	
Myrtaceae	<i>Leptospermum</i>	<i>roei</i>			*	*
Myrtaceae	<i>Leptospermum</i>	<i>spinescens</i>			*	
Myrtaceae	<i>Melaleuca</i>	<i>acuminata</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>adnata</i>		*	*	
Myrtaceae	<i>Melaleuca</i>	<i>calyptroides</i>	*	*	*	
Myrtaceae	<i>Melaleuca</i>	<i>cordata</i>	*	*	*	
Myrtaceae	<i>Melaleuca</i>	<i>cucullata</i>		*	*	
Myrtaceae	<i>Melaleuca</i>	<i>eleuterostachya</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>elliptica</i>		*	*	
Myrtaceae	<i>Melaleuca</i>	<i>hamata</i>	*	*	*	
Myrtaceae	<i>Melaleuca</i>	<i>lateriflora</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>laxiflora</i>			*	
Myrtaceae	<i>Melaleuca</i>	<i>pauperiflora</i> subsp. <i>fastigiata</i>	*			
Myrtaceae	<i>Melaleuca</i>	<i>pauperiflora</i> subsp. <i>pauperiflora</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>radula</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>teuthidoides</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>thyoides</i>		*		
Myrtaceae	<i>Melaleuca</i>	<i>uncinata</i>		*	*	
Myrtaceae	<i>Micromyrtus</i>	<i>obovata</i>			*	
Myrtaceae	<i>Rinzia</i>	<i>triplex</i> (P3)		*		
Myrtaceae	<i>Thryptomene</i>	<i>kochii</i>		*	*	
Myrtaceae	<i>Verticordia</i>	<i>chrysantha</i>		*	*	*
Myrtaceae	<i>Verticordia</i>	<i>roei</i>			*	*
Olacaceae	<i>Olex</i>	<i>benthamiana</i>			*	
Pittosporaceae	<i>Marianthus</i>	<i>bicolor</i>		*		
Proteaceae	<i>Banksia</i>	<i>cirsioides</i>	*		*	*
Proteaceae	<i>Banksia</i>	<i>elderiana</i>		*		*
Proteaceae	<i>Banksia</i>	<i>erythrocephala</i>		*	*	*
Proteaceae	<i>Banksia</i>	<i>laevigata</i> subsp. <i>laevigata</i>			*	*
Proteaceae	<i>Banksia</i>	<i>sessilis</i>			*	*
Proteaceae	<i>Grevillea</i>	<i>acuaria</i>	*			
Proteaceae	<i>Grevillea</i>	<i>cagiana</i>	*		*	*
Proteaceae	<i>Grevillea</i>	<i>eristachya</i>		*	*	
Proteaceae	<i>Grevillea</i>	<i>eryngioides</i>		*		*
Proteaceae	<i>Grevillea</i>	<i>huegelii</i>		*		
Proteaceae	<i>Grevillea</i>	<i>incrassata</i>				*
Proteaceae	<i>Grevillea</i>	<i>obliquistigma</i>		*		
Proteaceae	<i>Grevillea</i>	<i>oncogyne</i>	*			
Proteaceae	<i>Grevillea</i>	<i>pterosperma</i>		*		

Family	Genus	Taxon	CLP-EW1	SLP-MWS1	SP-H1	SP-H2
Proteaceae	<i>Grevillea</i>	<i>shuttleworthiana</i> subsp. <i>obovata</i>	*			*
Proteaceae	<i>Hakea</i>	<i>corymbosa</i>	*			*
Proteaceae	<i>Hakea</i>	<i>cygna</i> subsp. <i>cygna</i>			*	
Proteaceae	<i>Hakea</i>	<i>incrassata</i>	*			
Proteaceae	<i>Hakea</i>	<i>multilineata</i>		*	*	
Proteaceae	<i>Hakea</i>	<i>newbeyana</i>			*	
Proteaceae	<i>Hakea</i>	<i>platysperma</i>				*
Proteaceae	<i>Hakea</i>	<i>scoparia</i>	*	*	*	*
Proteaceae	<i>Isopogon</i>	<i>scabriusculus</i>				*
Proteaceae	<i>Persoonia</i>	<i>coriacea</i>		*	*	
Proteaceae	<i>Persoonia</i>	<i>helix</i>	*	*	*	
Proteaceae	<i>Petrophile</i>	<i>circinata</i>				*
Proteaceae	<i>Petrophile</i>	<i>stricta</i>			*	
Proteaceae	<i>Petrophile</i>	<i>teretifolia</i>			*	
Rutaceae	<i>Boronia</i>	<i>inornata</i>		*		
Rutaceae	<i>Drummondita</i>	<i>hassellii</i>			*	
Rutaceae	<i>Microcybe</i>	<i>albiflora</i>			*	
Rutaceae	<i>Microcybe</i>	<i>ambigua</i>			*	
Rutaceae	<i>Microcybe</i>	<i>multiflora</i>	*			
Rutaceae	<i>Phebalium</i>	<i>filifolium</i>	*			
Santalaceae	<i>Exocarpos</i>	<i>aphyllus</i>	*			
Santalaceae	<i>Exocarpos</i>	<i>sparteus</i>		*	*	*
Santalaceae	<i>Leptomeria</i>	<i>preissiana</i>			*	
Santalaceae	<i>Santalum</i>	<i>acuminatum</i>	*	*	*	*
Sapindaceae	<i>Dodonaea</i>	<i>bursariifolia</i>		*		
Scrophulariaceae	<i>Eremophila</i>	<i>densiflora</i>	*			
Scrophulariaceae	<i>Eremophila</i>	<i>dichroantha</i>		*		
Scrophulariaceae	<i>Eremophila</i>	<i>drummondii</i>		*	*	
Thymelaeaceae	<i>Pimelea</i>	<i>aeruginosa</i>			*	

Appendix 4: Vegetation Condition Rating

Vegetation Condition Rating	South West and Interzone Botanical Provinces	Eremaean and Northern Botanical Provinces
Pristine	Pristine or nearly so, no obvious signs of disturbance or damage caused by human activities since European settlement.	
Excellent	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species. Damage to trees caused by fire, the presence of non-aggressive weeds and occasional vehicle tracks.	Pristine or nearly so, no obvious signs of damage caused by human activities since European settlement.
Very Good	Vegetation structure altered, obvious signs of disturbance. Disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.	Some relatively slight signs of damage caused by human activities since European settlement. For example, some signs of damage to tree trunks caused by repeated fire, the presence of some relatively non-aggressive weeds, or occasional vehicle tracks.
Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.	More obvious signs of damage caused by human activity since European settlement, including some obvious impact on the vegetation structure such as that caused by low levels of grazing or slightly aggressive weeds.
Poor		Still retains basic vegetation structure or ability to regenerate it after very obvious impacts of human activities since European settlement, such as grazing, partial clearing, frequent fires or aggressive weeds.
Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds at high density, partial clearing, dieback and grazing.	Severely impacted by grazing, very frequent fires, clearing or a combination of these activities. Scope for some regeneration but not to a state approaching good condition without intensive management. Usually with a number of weed species present including very aggressive species.
Completely Degraded	The structure of the vegetation is no longer intact and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees and shrubs.	Areas that are completely or almost completely without native species in the structure of their vegetation; i.e. areas that are cleared or 'parkland cleared' with their flora comprising weed or crop species with isolated native trees or shrubs.

Appendix 5: Conservation Significant Flora within 40km of the survey area

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Acacia asepala</i>			P2	Diffuse, much-branched shrub, 0.5-1.5 m high. Fl. yellow, Aug. Red-brown sandy loam. Undulating plains, along drainage lines.	Unlikely
<i>Acacia kerryana</i>			P2	Low, spreading, domed shrub, 0.3-1 m high. Fl. yellow, Oct to Dec or Jan to Feb. Granitic loamy sand, stony clayey loam or clayey sand. Low stony ridges, undulating plains.	Unlikely
<i>Acacia lanuginophylla</i>	EN	VU		Dense shrub, 0.5-1.2 m high. Fl. yellow, Jul to Oct. White/grey sand, clayey sand, gravelly soils. Flats, along drainage lines.	Unlikely
<i>Acacia repanda</i>			P3	Rounded to obconic, single-stemmed or much-branched shrub, 0.5-2 m high, bark 'minni-ritchi'. Fl. yellow, Jun to Aug. Loam, sandy or gravelly loam. Near granite outcrops.	Unlikely
<i>Acacia</i> sp. Mt Holland (B. Ellery BE 1147)			P1	No description available	Possible
<i>Anticoryne melanosperma</i>			P3	No description available	Possible
<i>Austrostipa</i> sp. Carlingup Road (S. Kern & R. Jasper LCH 18459)			P1	No description available	Possible
<i>Austrostipa</i> sp. Mt Holland (W.A. Thompson & J. Allen 948)			P1	No description available	Possible
<i>Baeckea</i> sp. Blue Haze Mine (P. Armstrong 06/910)			P1	No description available	Possible
<i>Baeckea</i> sp. Crossroads (B.L. Rye & M.E. Trudgen 241186)			P1	No description available	Possible
<i>Baeckea</i> sp. Forrestania (K.R. Newbey 1105)			P1	No description available	Possible
<i>Baeckea</i> sp. Hatter Hill (K.R. Newbey 3284)			P3	Narrow, open, upright shrub, to 1.3 m high. Fl. pink, Jun to Oct. Yellow-orange coarse sandy loam with laterite gravel, red-brown sandy loam with quartz pebbles. Undulating plains.	Possible
<i>Baeckea</i> sp. Koonadgin (B.L. Rye & M.E. Trudgen BLR 241137)			P3	No description available	Possible
<i>Baeckea</i> sp. Lake Cronin (K.R. Newbey 9191)			P1	Upright, spreading, moderately open shrub. Fl. white/pink, Oct. Well-drained gravelly sands. Moderately exposed, gently undulating plain.	Possible
<i>Baeckea</i> sp. North Ironcap (R.J. Cranfield 10580)			P1	Erect, open shrub, to 0.4 m high. Fl. white/pink, Oct. Red clay. Gently undulating sandplains.	Possible
<i>Baeckea</i> sp. Sheoaks Rocks (M.E. Trudgen MET5452)			P1	Open shrub, to 0.3 m high. Fl. white/pink, Nov. Yellow-brown silty sand. Mid-upper gentle slopes.	Unlikely
<i>Banksia rufa</i> subsp. <i>flavescens</i>			P3	Prostrate, ?lignotuberous shrub, to 0.45 m high. Fl. cream-yellow, Jul to Aug. Sandy loam or sand with gravel.	Possible
<i>Banksia sphaerocarpa</i> var. <i>dolichostyla</i>	VU	VU		Lignotuberous shrub, 1-3 m high. Fl. yellow-orange, Mar to May. Lateritic gravel, grey sand.	Unlikely
<i>Banksia viscida</i>			P3	Densely branched, non-lignotuberous shrub, 0.4-1 m high. Fl. yellow-orange, Jul to Oct. Gravelly soils. Lateritic rises.	Possible

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Boronia revoluta</i>	EN	VU		Shrub, 0.4-0.8 m high. Fl. pink, Jul to Aug. Stony sandy loam or sand. Plains, hillsides & summits.	Unlikely
<i>Boronia westringioides</i>			P2	Erect shrub, 0.2-0.75 m high. Fl. pink, Jul to Sep. Loamy sand. Plains.	Possible
<i>Brachyloma nguba</i>			P1	Erect, compact to spreading, mid-dense shrub, to 0.8 m high, leaves discolourous, usually 2-3 mm long; style 0.2-0.25 mm long; disc truncate. Fl. red, Apr to May. White to brown sandy clay, shallow sandy loam. Open mallee woodland, mallee scrub, flat plains.	Possible
<i>Brachyloma stenolobum</i>			P1	No description available	Possible
<i>Caladenia graniticola</i>		EN		Tuberous, perennial, herb, to 0.21 m high, plant usually single flowered. Fl. green-yellow, Oct. Gritty sandy clay, granite. Near low exposed rock outcrops.	Unlikely
<i>Caladenia hoffmanii</i>	EN	EN		Tuberous, perennial, herb, 0.13-0.3 m high. Fl. green & yellow & red, Aug to Oct. Clay, loam, laterite, granite. Rocky outcrops and hillsides, ridges, swamps and gullies.	Unlikely
<i>Calamphoreus inflatus</i>			P4	Erect, spreading shrub, 0.4-1.6 m high, to 2 m wide. Fl. blue-purple/green, Oct to Dec or Feb to Mar. Clay loam with ironstone gravel. Flats, disturbed sites.	Possible
<i>Calectasia pignattiana</i>	VU	VU		Rhizomatous, prickly herb, to 0.5 m high. Fl. blue-purple, Aug to Oct. Sand to sandy clay over granite or laterite, gravel. Plains and gentle slopes.	Unlikely
<i>Calytrix nematoclada</i>			P3	Shrub, 0.15-0.5(-1) m high. Fl. purple-pink, Sep or Nov to Dec or Jan. Yellow or grey sand. Sandplains.	Possible
<i>Chorizema circinale</i>			P3	Prostrate, scrambling, wiry shrub, to 0.4 m high. Fl. yellow & orange & red, Sep to Dec. Yellow sand, sandy clay with gravel. Flats, margin of gravel pit.	Possible
<i>Comesperma calcicola</i>			P3	Soft perennial, herb, to 0.3 m high. Fl. pink, Oct to Dec or Jan. Calcareous or semi-saline clay loams, limestone. Areas around saline water.	Unlikely
<i>Conospermum sigmoideum</i>			P2	Erect shrub, 0.2-0.5 m high. Fl. blue, Aug to Sep. Yellow sand.	Unlikely
<i>Cryptandra polyclada</i> subsp. <i>polyclada</i>			P3	Mat-forming or upright shrub, 0.1-0.7 m high. Fl. white/cream, Jan to May or Aug or Oct. Sand. Sandplains.	Possible
<i>Dampiera orchardii</i>			P2	Erect perennial, herb, 0.2-0.4 m high. Sand.	Possible
<i>Dampiera scaevolina</i>			P1	Erect to ascending perennial, herb or shrub, 0.2-0.5 m high. Fl. blue/white, Sep to Nov. Sandy & gravelly soils.	Possible
<i>Daviesia elongata</i> subsp. <i>implexa</i>			P3	Spreading or sprawling shrub, 0.4-1 m high. Fl. yellow/orange & red, Sep. Sand & laterite.	Possible
<i>Daviesia implexa</i>			P3	No description available	Possible
<i>Dicrastyliis capitellata</i>			P1	Low spreading shrub, 0.2-0.25 m high. Fl. blue-purple, May. Loamy sand, sandy loam.	Possible

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Elatine macrocalyx</i>			P3	Prostrate, glabrous, mat-forming annual, herb, sepals 2-3mm long, fruit indehiscent. Fl. white, May to Oct (probably opportunistic). Shallow sands over clay. Margins of playa lakes and clay pans.	Unlikely
<i>Eremophila biserrata</i>			P4	Prostrate shrub, to 3 m wide. Fl. green/yellow-green, Sep to Nov or Mar. Sandy or sandy clay soils. Alluvial flats, salt flats & lakes.	Unlikely
<i>Eremophila racemosa</i>			P4	Erect shrub, 0.5-1.7 m high. Fl. purple-pink-red/white, Mar or Aug to Dec. Sandy or stony loam, clay loam. Undulating plains, roadsides.	Possible
<i>Eremophila verticillata</i>	EN	CR		Low spreading shrub, up to 0.8 m high, to 1 m wide. Fl. purple-violet, Nov to Dec. Clay loam, loam over limestone.	Unlikely
<i>Eucalyptus cerasiformis</i>			P4	Mallee, 2-3.5 m high, bark smooth, grey to brown. Fl. yellow, Dec or Jan to Mar. Red loamy soils.	Possible
<i>Eucalyptus deflexa</i>			P4	(Mallee), 1-3 m high, bark smooth. Fl. pink/cream-white, Mar or May to Oct. Clay loam, sandy loam, white or yellow sand, often with gravel. Flat areas & slight rises.	Possible
<i>Eucalyptus exigua</i>			P3	Mallee, 2-5 m high, bark smooth. Fl. white-cream, Mar. Sandy loam, white sand. Sandplains.	Possible
<i>Eucalyptus georgei</i> subsp. <i>fulgida</i>			P4	Tree, 4-20 m high, bark smooth, often hanging in ribbons. Fl. cream-white. Sandy loam, clayey sand. Slight depressions.	Unlikely
<i>Eucalyptus myriadena</i> subsp. <i>parviflora</i>			P1	Mallee or tree, 3-10 m high, bark rough, coarse & flaky on trunk, smooth above. Loam. Swamps, plains.	Unlikely
<i>Eucalyptus recta</i>	EN	VU		Tree, to 15 m high, bark smooth. Sandy laterite.	Unlikely
<i>Eucalyptus retusa</i>			P1	No description available	Possible
<i>Eucalyptus steedmanii</i>	VU	VU		Tree, 2-8(-12) m high, bark smooth. Fl. white, Jan to Mar. Gravelly loam over ironstone, sand. Low hills, undulating plains.	Unlikely
<i>Eutaxia acanthoclada</i>			P3	Compact, mat-forming, prostrate shrub, to 0.3 m high. Fl. yellow/orange/red, Oct to Nov. Light brown sandy clay, shallow sandy loam, red clay over banded ironstone, gravel. Gently undulating plains.	Unlikely
<i>Eutaxia hirsuta</i>			P2	Erect, shrub, spindly shrub (broom-like). <i>Stems</i> terete, glabrous; pustules or glands absent.	Known to occur ³
<i>Eutaxia lasiocalyx</i>			P2	Low, spreading, multi-stemmed shrub, to 0.15 m high. Fl. yellow, Nov. Red sandy loam, laterite and quartz gravel. Gentle lower slopes.	Unlikely
<i>Eutaxia nanophylla</i>			P3	Straggly, rounded shrub, to 0.35 m high. Fl. Yellow & orange & red, Oct to Nov. Clayey sand, red clay, stoney clayey loam. Low-lying areas, damp flats, slopes, undulating plains, low stony ridges.	Unlikely

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Eutaxia rubricarina</i>			P3	Straggling shrub, to 0.5 m high. Fl. Orange & yellow & brown, Aug or Oct. Gravelly sand, grey to pinkish-white sandy clay, red loam. Flats, slopes, valley floors, road verges.	Unlikely
<i>Eutaxia</i> sp. North Ironcap (P. Armstrong PA 06/898)			P1	No description available	Possible
<i>Frankenia drummondii</i>			P3	Prostrate shrub. Fl. white. Sand. Lake edges.	Unlikely
<i>Gastrolobium tenue</i>			P1	Low, bushy shrub, to 0.6 m high. Fl. Orange & red & purple, Sep to Oct. Yellow sand or sandy clay. Undulating dunes, stony outcrops.	Unlikely
<i>Grevillea aneura</i>			P4	Dense, prickly shrub, 0.5-2.8 m high. Fl. red, Jun or Aug to Dec or Jan. Sand, sandy clay, gravel.	Possible
<i>Grevillea neodissecta</i>			P4	No description available	Possible
<i>Grevillea insignis</i> subsp. <i>elliottii</i>			P3	Erect, bushy, non-lignotuberous shrub, 1-2 m high. Fl. red/pink & cream & white, Oct. Gravelly sand or loam over ironstone. Hilltops or rises.	Unlikely
<i>Grevillea pilosa</i> subsp. <i>redacta</i>			P3	Spreading to prostrate, non-lignotuberous shrub, 0.4-1.2 m high. Fl. red, Feb or Oct or Dec. Sand, laterite.	Unlikely
<i>Grevillea prostrata</i>			P4	Loose, prostrate shrub, 0.04-0.1 m high, 0.8-1.2 m wide. Fl. cream-white/pink-red, Aug to Dec or Jan. White, grey or yellow sand, gravel. Sandplains.	Possible
<i>Guichenotia anota</i>			P1	Shrubs, 1 m high; branchlets hairy, not glaucous. Leaves alternate, 10-25 mm long	Possible
<i>Guichenotia asteriskos</i>			P2	Erect, compact shrub, ca 0.35 m high. Fl. white, Sep to Oct. Sandy clay or loam with gravel.	Possible
<i>Gyrostemon ditrigynus</i>			P4	Shrub, 0.4-1.5 m high. Sand, sandy clay, loam. Plains, low ironstone ridges.	Unlikely
<i>Haegiela tatei</i>			P4	Ascending to erect annual, herb, 0.02-0.08(-0.2) m high. Fl. white-yellow, Aug to Nov. Clay, sandy loam, gypsum. Saline habitats.	Unlikely
<i>Hemigenia</i> sp. Newdegate (E. Bishop 75)			P1	Spindly, erect to spreading shrub, 0.2-0.45 m high, to 0.5 m wide. Fl. blue/purple, Sep to Oct. Clay loam. Disturbed sites.	Possible
<i>Hibbertia axillibarba</i>			P1	Shrub, to 0.7 m high. Fl. yellow, Sep to Oct. Lateritic soil. Ranges.	Unlikely
<i>Hibbertia carinata</i>			P1	Shrub, to 0.4 m high. Fl. yellow, Aug to Sep. Well-drained gravelly sand, yellow sand with gravel.	Possible
<i>Hibbertia pachyphylla</i>			P3	Shrub, to 0.5 m high. Fl. yellow, Sep to Nov. White to yellow sand, brown sandy gravel, gravelly loam, laterite, granite, quartz. Undulating plains, low rises, valley floors.	Possible
<i>Hydrocotyle eichleri</i>			P3	No description available	Possible
<i>Hysterobaeckea pterocera</i>			P1	No description available	Possible
<i>Isolepis australiensis</i>			P3	Annual, grass-like or herb (sedge), 0.03-0.055 m high, glumes 0.8-1.2 mm long; stamens 1(-2); style branches 3; nut with abaxial angle acute. Fl. Jun or Sep. Silty sand, sandy clay. Lake margins, pools.	Unlikely
<i>Labichea rossii</i>			P1	No description available	Possible

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Lepidosperma amantiferrum</i>			P1	Tufted rhizomatous, herb (sedge), leaves 0.15-0.42 m high, culms and leaves distichous. Yellow sandy loam with banded ironstone gravel and rocks. Gentle lower slopes.	Unlikely
<i>Lepidosperma ferriculmen</i>			P1	Tufted rhizomatous, perennial, herb (sedge), leaves 0.16-0.38 m high, culms and leaves distichous. Well-drained orange-red sandy loam with banded ironstone gravel and rocks. Stony slopes.	Unlikely
<i>Lepidosperma lyonsii</i>			P4	Tufted rhizomatous, perennial, herb (sedge), leaves 0.31-0.53 m high, culms and leaves distichous. Pale orange skeletal sandy loam with banded ironstone gravel & rock, well-drained shallow stony loamy with quartz. Gentle hill slopes, upper slopes of large hill.	Unlikely
<i>Melaleuca macronychia</i> subsp. <i>trygonoides</i>			P3	Multi-stemmed, spreading shrub, 1-4 m high, leaves broadly elliptic. Fl. red, Feb or Jul to Aug or Oct. Sandy soils. Granite outcrops.	Unlikely
<i>Melaleuca ochroma</i>			P3	No description available	Possible
<i>Microcorys</i> sp. Forresteria (V. English 2004)			P4	Prostrate or erect shrub, 0.35-0.4 m high. Fl. white/purple, Jan or Apr. Yellow sandy clay or red-brown clay. Open woodland or cleared areas.	Known to occur ²
<i>Microcorys</i> sp. Mt Holland (D. Angus DA 2397)			P1	No description available	Possible
<i>Microseris walteri</i>			P3	No description available	Possible
<i>Mirbelia densiflora</i>			P3	Erect or straggling shrub, 0.2-1 m high. Fl. yellow-orange, Oct or Jan. Stony loam, loamy sand. Small ridges, breakaways, undulating plains.	Unlikely
<i>Mirbelia taxifolia</i>			P1	Shrub, 0.6-0.9 m high. Fl. orange-yellow, Sep. Red or yellow sand.	Possible
<i>Notisia intonsa</i>			P3	No description available	Possible
<i>Olearia laciniifolia</i>			P2	Erect, few-stemmed shrub, 0.6-1.2 m high. Fl. blue/purple & white/yellow, May to Sep. White sand. Around playa lakes.	Unlikely
<i>Orianthera exilis</i>			P2	No description available	Possible
<i>Oxymyrrhine plicata</i>			P3	No description available	Possible
<i>Paragoodia crenulata</i>	CE	VU		The flower spike has 1–3 flowers that are brown and yellow in colour. The flowering period is from July to August. This species is thought to require disturbance (DEC, 2010)	Unlikely
<i>Pityrodia scabra</i> subsp. <i>dendrotricha</i>			P3	No description available	Possible
<i>Pityrodia</i> sp. Yilgarn (A.P. Brown 2679)			P3	No description available	Possible
<i>Pterostylis echinulata</i>			P3	No description available	Possible
<i>Pultenaea daena</i>			P3	Dense, prostrate, domed shrub, to 0.07 m high. Fl. yellow, Mar. White to yellow sand or sandy loam, sandy or loamy clay, gravel, limestone, dolomite, laterite. Gently undulating plains, adjacent to salt lakes, in disturbed areas.	Unlikely
<i>Rinzia torquata</i>			P3	No description available	Known to occur ¹
<i>Rinzia triplex</i>			P3	No description available	Known to occur ^{1,2}

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Roycea pycnophylloides</i>	EN	VU		Perennial, herb, forming densely branched, silvery mats to 1 m wide. Fl. Sep. Sandy soils, clay. Saline flats.	Unlikely
<i>Scaevola tortuosa</i>			P1	Ascending perennial, herb, 0.1-0.2 m high. Fl. blue-purple/pink, Oct. Sandy clay. Margins of salt lakes.	Unlikely
<i>Seringia adenogyna</i>			P3	No description available	Possible
<i>Stenanthemum liberum</i>			P1	Dwarf shrub, ca 0.5 m high. Yellow sandy loam over laterite.	Unlikely
<i>Stylidium sejunctum</i>			P3	Caespitose perennial, herb, 0.25-0.45 m high, Leaves tufted, linear to narrowly oblanceolate, 10-30 cm long, 0.8-4 mm wide, apex acute to mucronate, margin involute, glabrous to scabrous. Membraneous scale leaves present at base of mature leaves. Scape glandular throughout. Inflorescence paniculate. Fl. white/pink-purple, Sep to Nov. Clayey sand or loam, laterite. Outcrops, upper slopes, breakaways. Mallee and Allocasuarina shrubland.	Unlikely
<i>Stylidium thylax</i>			P2	Creeping perennial, herb, 0.04-0.08 m high, Leaves adpressed to stem, ovate to lanceolate, 0.1-0.4 cm long, 0.6-1.5 mm wide, apex mucronate, margin hyaline, glabrous. Inflorescence uni-flowered, pedicels glandular. Fl. white, Oct. Sand. Gentle slopes and plains. Heath, mallee shrubland.	Possible
<i>Stylidium validum</i>			P1	Caespitose perennial, herb, 0.06-0.3 m high, Leaves tufted, oblanceolate, 1.5-10 cm long, 2.2-6 mm wide, apex acute to acuminate, margin entire, glabrous, glaucous. Scape glabrous. Inflorescence paniculate. Fl. white/pink, Sep to Oct. Clayey sand or loam, ironstone, greenstone gravel. Hillslopes and hilltops. Eucalypt woodland, mallee shrubland.	Unlikely
<i>Teucrium diabolicum</i>			P3	Compact, dwarf shrub, 0.1 m high, to 0.1 m wide. Fl. white, Apr. Hills, road verges.	Unlikely
<i>Tetradlea aphylla</i> subsp. <i>megacarpa</i>	VU	VU		Shrub, to 0.35 m high. Yellow sand, brown sandy loam, yellow-brown clay loam, gravel, laterite. Rises and ridges.	Unlikely
<i>Thysanotus lavanduliflorus</i>			P1	Caespitose perennial, herb (with tuberous roots), to 0.25 m high. Fl. purple, Nov to Dec. Sand, sandy loam.	Possible
<i>Thysanotus</i> sp. Yellowdine (A.S. George 6040)			P2	No description available	Possible
<i>Tribonanthes purpurea</i>	VU	VU		Tuberous, perennial, herb, 0.03-0.04 m high. Fl. pink-purple, Aug. Seasonally wet soils in moss swards & herbfields among granite rocks.	Unlikely
<i>Verticordia gracilis</i>			P3	Low, slender shrub, 0.15-0.6 m high. Fl. pink, Oct to Nov. Yellow sand, gravelly sand, sandy loam.	Known to occur ¹
<i>Verticordia multiflora</i> subsp. <i>solox</i>			P2	Erect to spreading shrub, 0.2-0.6 m high. Fl. yellow, Oct to Dec or Jan. Yellow sand over gravel, sand over granite.	Unlikely

Taxon	Conservation Code			Description (WAHERB, 2021)	Likelihood of Occurrence
	EPBC Act	BC Act	Priority Listing		
<i>Verticordia staminosa</i> var. <i>cyindracea</i>	EN	VU		Spreading shrub, 0.3-0.8 m high. Fl. green-yellow/yellow-brown, Jul to Oct. Soil pockets. Granite outcrops.	Unlikely
<i>Verticordia stenopetala</i>			P3	Shrub, 0.2-0.6(-1.3) m high. Fl. pink/pink-purple-red, Oct to Dec or Jan. Yellow sand, sometimes with gravel. Undulating plains.	Possible

Appendix 6: Potential Fauna Species List

Listing of Fauna Potentially Present in/near Survey Area

Lounge Lizard Sand Pits

Approximate centroid 119° 41' 36" E, 32° 25' 46" S
Compiled by Greg Harewood - February 2021
Recorded (Captured/Sighted/Heard/Signs) = X

A = How, R.A. et al. (1988). The biological survey of the eastern goldfields of Western Australia. Part 4, Lake Johnston-Hyden Study Area, Records of the WAM, supplement No. 30. (Lake Cronin)
B = Biota (2006a/2007a). Forrestania Monitoring Survey, Flying Fox Phases I, II, III and IV. Unpublished report for Western Areas NL.
Biota (2006b). Forrestania Water Disposal Pipeline Survey – Fauna and Faunal Assemblages Report. Unpublished report for Western Areas NL.
Biota (2007b). Diggers South Fauna Survey – Phase I. Unpublished report for Western Areas NL.
Biota (2010). Spotted Quoll Haul Road Single Phase Fauna Survey. Unpublished report for Western Areas N.L. May 2010.
Biota (2018). New Morning Level 1 and Targeted Terrestrial Fauna Survey Unpublished report for Western Areas N.L. Nov 2018.
C = KLA (2010). Cazaly Resources Ltd. Parker Range Iron Ore Project. Fauna Assessment. Unpublished report for Cazaly Resources Limited, August 2011.
D = DBCA (2021). NatureMap Database search. "By Circle"119° 41' 36" E, 32° 25' 56" S – Survey Area (plus 40 km buffer). 20 February 2021.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Amphibia						
Myobatrachidae Ground or Burrowing Frogs						
<i>Crinia pseudinsignifera</i>	Bleating Froglet	LC		X		X
<i>Heleioporus albopunctatus</i>	Western Spotted Frog	LC	X	X		X
<i>Limnodynastes dorsalis</i>	Western Banjo Frog	LC	X			X
<i>Myobatrachus gouldii</i>	Turtle Frog	LC				
<i>Neobatrachus albigipes</i>	White-footed Trilling Frog	LC				X
<i>Neobatrachus centralis</i>	Trilling Frog	LC	X			

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
<i>Neobatrachus kunapalari</i>	Kunapalari Frog	LC				X
<i>Neobatrachus pelobatoides</i>	Humming Frog	LC	X			X
<i>Neobatrachus</i> sp.	Unidentified Burrowing Frog	LC				
<i>Neobatrachus sutor</i>	Shoemaker Frog	LC	X			X
<i>Pseudophryne guentheri</i>	Crawling Toadlet	LC				X
<i>Pseudophryne occidentalis</i>	Western Toadlet	LC	X	X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Reptilia						
Diplodactylidae Geckoes						
<i>Crenadactylus ocellatus</i>	Clawless Gecko	LC	X	X		X
<i>Diplodactylus granariensis granariensis</i>	Wheatbelt Stone Gecko	LC	X	X	X	
<i>Diplodactylus pulcher</i>	Western Saddled Ground Gecko	LC		X	X	X
<i>Lucasium maini</i>	Main's Ground Gecko	LC	X	X	X	X
<i>Oedura reticulata</i>	Reticulated Velvet Gecko	LC	X	X	X	
<i>Strophurus intermedius</i>	Southern Spiny-tailed Gecko	LC				
<i>Strophurus spinigerus inornatus</i>	Orange-eyed Southwestern Spiny-tailed Gecko	LC	X	X		
Gekkonidae Geckoes						
<i>Christinus marmoratus</i>	Marbled Gecko	LC			X	
<i>Gehyra variegata</i>	Variegated Dittella	LC	X	X		X
<i>Heteronotia binoei</i>	Bynoe's Gecko	LC	X			
<i>Underwoodisaurus milii</i>	Barking Gecko	LC	X	X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Pygopodidae Legless Lizards						
<i>Delma australis</i>	Marble-faced Delma	LC		X		X
<i>Delma butleri</i>	Unbanded Delma	LC				
<i>Delma fraseri</i>	Fraser's Legless Lizard	LC	X	X		X
<i>Lialis burtonis</i>	Burton's Legless Lizard	LC	X	X		X
<i>Pygopus lepidopodus</i>	Common Scaly Foot	LC		X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Agamidae Dragon Lizards						
<i>Ctenophorus cristatus</i>	Crested Bicycle Dragon	LC	X	X	X	X
<i>Ctenophorus isolepis</i>	Goldfields Military Sand Dragon	LC				
<i>Ctenophorus maculatus</i>	Spotted Military Dragon	LC	X	X		X
<i>Ctenophorus ornatus</i>	Ornate Crevice Dragon	LC				X
<i>Ctenophorus salinarum</i>	Salt Pan Dragon	LC	X			X
<i>Moloch horridus</i>	Thorny Devil	LC	X	X		X
<i>Pogona minor minor</i>	Western Bearded Dragon	LC	X	X	X	
<i>Rankinia adalaidensis chapmani</i>	Eastern Heath Dragon	LC	X	X		
Varanidae Monitor's or Goanna's						
<i>Varanus gouldii</i>	Gould's Sand Monitor	LC	X	X		X
<i>Varanus rosenbergi</i>	Heath Monitor	LC		X		X
<i>Varanus tristis</i>	Black-headed Monitor	LC				

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Scincidae Skinks						
<i>Cryptoblepharus buchananii</i>	Buchanan's Snake-eyed Skink	LC	X	X	X	X
<i>Ctenotus atlas</i>	Southern Malle Ctenotus	LC	X			X
<i>Ctenotus impar</i>	Odd-striped Ctenotus	LC	X	X		X
<i>Ctenotus schomburgkii</i>	Barred Wedge-snout Ctenotus	LC	X	X		X
<i>Ctenotus xenopleura</i>	Wide-striped Sandplain Ctenotus	LC				
<i>Cyclodomorphus melanops elongatus</i>	Eastern Slender Blue-tongue	LC	X			
<i>Egernia formosa</i>	Goldfields Crevice Skink	LC				X
<i>Egernia inornata</i>	Desert Skink	LC			X	
<i>Egernia richardi</i>	Woodland Crevice Skink	LC	X	X	X	X
<i>Eremiascincus richardsonii</i>	Broad-banded Sand Swimmer	LC				
<i>Hemiergis initialis initialis</i>	Sth Five-toed Mulch Skink	LC		X		
<i>Hemiergis peronii peronii</i>	Four-toed Earless Skink	LC				

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
<i>Lerista distinguenda</i>	SW Four-toed Lerista	LC	X	X		X
<i>Lerista dorsalis</i>	Southern Four-toed Lerista	LC		X		
<i>Lerista kingi</i>	King's Three-toed Slider	LC				
<i>Lerista picturata</i>	Goldfields Robust Lerista	LC		X		X
<i>Lerista timida</i>	Dwarf Three-toed Slider	LC				
<i>Liopholis multiscutata</i>	Bull Skink	LC	X	X	X	X
<i>Menetia greyii</i>	Dwarf Skink	LC	X	X	X	X
<i>Morethia butleri</i>	Woodland Dark-flecked Morethia	LC	X		X	X
<i>Morethia obscura</i>	Shrubland Pale-flecked Morethia	LC	X	X		X
<i>Tiliqua occipitalis</i>	Western Bluetongue	LC	X	X		X
<i>Tiliqua rugosa</i>	Bobtail	LC		X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Typhlopidae Blind Snakes						
<i>Ramphotyphlops australis</i>	Southern Blind Snake	LC	X	X	X	
<i>Ramphotyphlops bicolor</i>	Dark-spined Blind Snake	LC				
<i>Ramphotyphlops bituberculatus</i>	Prong-snouted Blind Snake	LC				
<i>Ramphotyphlops hamatus</i>	Northern Hook-snouted Blind Snake	LC				
Boidae Pythons, Boas						
<i>Morelia imbricata</i>	Southern Carpet Python	LC				

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class		Common Name		Conservation Status			
Family	Species			A	B	C	D

Elapidae Elapid Snakes							
<i>Brachyuropsis semifasciata</i>		Southern Shovel-nosed Snake	LC				
<i>Echiopsis curta</i>		Bardick	LC		X		X
<i>Neelaps bimaculatus</i>		Black-naped Snake	LC	X			X
<i>Parasuta gouldii</i>		Gould's Hooded Snake	LC	X	X		X
<i>Parasuta nigriceps</i>		Black-backed Snake	LC		X		X
<i>Paroplocephalus atriceps</i>		Lake Cronin Snake	P3	X	X		X
<i>Pseudechis australis</i>		Mulga Snake	LC	X			X
<i>Pseudonaja affinis</i>		Dugite	LC	X	X		X
<i>Pseudonaja modesta</i>		Ringed Brown Snake	LC			X	
<i>Simoselaps bertholdi</i>		Jan's Banded Snake	LC		X		X

Aves

Casuariidae Emus, Cassowaries							
<i>Dromaius novaehollandiae</i>		Emu	LC	X	X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Megapodiidae Moundbuilders						
<i>Leipoa ocellata</i>	Malleefowl	S3 VU VU	X	X	X	X
Phasianidae Quails, Pheasants						
<i>Coturnix pectoralis</i>	Stubble Quail	LC				
Accipitridae Kites, Goshawks, Eagles, Harriers						
<i>Accipiter cirrocephalus</i>	Collared Sparrowhawk	LC	X	X		X
<i>Accipiter fasciatus</i>	Brown Goshawk	LC				X
<i>Aquila audax</i>	Wedge-tailed Eagle	LC		X	X	X
<i>Aquila morphnoides</i>	Little Eagle	LC	X	X		
<i>Circus assimilis</i>	Spotted Harrier	LC				X
<i>Elanus caeruleus</i>	Black-shouldered Kite	LC				
<i>Haliastur sphenurus</i>	Whistling Kite	LC				X
<i>Hamirostra isura</i>	Square-tailed Kite	LC	X	X		

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Falconidae Falcons						
<i>Falco berigora</i>	Brown Falcon	LC	X	X	X	X
<i>Falco cenchroides</i>	Australian Kestrel	LC	X	X		X
<i>Falco longipennis</i>	Australian Hobby	LC	X			X
<i>Falco peregrinus</i>	Peregrine Falcon	S7 LC		X		X
Otididae Bustards						
<i>Ardeotis australis</i>	Australian Bustard	LC	X			X
Turnicidae Button-quails						
<i>Turnix velox</i>	Little Button-quail	LC		X		X
Charadriidae Lapwings, Plovers, Dotterels						
<i>Vanellus tricolor</i>	Banded Lapwing	LC				

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Columbidae Pigeons, Doves						
<i>Ocyphaps lophotes</i>	Crested Pigeon	LC				X
<i>Phaps chalcoptera</i>	Common Bronzewing	LC	X	X	X	X
<i>Phaps elegans</i>	Brush Bronzewing	LC		X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Psittacidae Parrots						
<i>Cacatua roseicapilla</i>	Galah	LC	X	X	X	X
<i>Calyptorhynchus latirostris</i>	Camaby's Black-Cockatoo	S2 EN EN	X	X		X
<i>Glossopsitta porphyrocephala</i>	Purple-crowned Lorikeet	LC	X	X	X	
<i>Melopsittacus undulatus</i>	Budgerigar	LC				X
<i>Neophema elegans</i>	Elegant Parrot	LC	X	X	X	X
<i>Nymphicus hollandicus</i>	Cockatiel	LC				X
<i>Platycercus icterotis xanthogenys</i>	Western Rosella (inland ssp)	P4	X	X	X	
<i>Platycercus varius</i>	Mulga Parrot	LC		X		X
<i>Platycercus zonarius</i>	Australian Ringneck	LC	X	X	X	X
<i>Polytelis anthopeplus</i>	Regent Parrot	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Cuculidae Parasitic Cuckoos						
<i>Cacomantis flabelliformis</i>	Fan-tailed Cuckoo	LC	X	X		X
<i>Chrysococcyx basalis</i>	Horsfield's Bronze Cuckoo	LC	X		X	X
<i>Chrysococcyx lucidus</i>	Shining Bronze Cuckoo	LC		X		
<i>Chrysococcyx osculans</i>	Black-eared Cuckoo	LC				
<i>Cuculus pallidus</i>	Pallid Cuckoo	LC	X	X		
Strigidae Hawk Owls						
<i>Ninox novaeseelandiae</i>	Boobook Owl	LC	X		X	
Tytonidae Barn Owls						
<i>Tyto alba</i>	Barn Owl	LC				
Podargidae Frogmouths						
<i>Podargus strigoides</i>	Tawny Frogmouth	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Caprimulgidae Nightjars						
<i>Eurostopodus argus</i>	Spotted Nightjar	LC	X	X	X	X
Aegothelidae Owlet-nightjars						
<i>Aegotheles cristatus</i>	Australian Owlet-nightjar	LC	X	X		X
Halcyonidae Tree Kingfishers						
<i>Todiramphus pyrrhopygia</i>	Red-backed Kingfisher	LC				
<i>Todiramphus sanctus</i>	Sacred Kingfisher	LC				X
Meropidae Bee-eaters						
<i>Merops ornatus</i>	Rainbow Bee-eater	JA LC	X	X	X	X
Climacteridae Treecreepers						
<i>Climacteris rufa</i>	Rufous Treecreeper	LC	X	X		

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Maluridae Fairy Wrens, GrassWrens						
<i>Malurus lamberti</i>	Variegated Fairy-wren	LC				
<i>Malurus leucopterus</i>	White-winged Fairy-wren	LC				X
<i>Malurus pulcherrimus</i>	Blue-breasted Fairy-wren	LC	X	X	X	X
<i>Malurus splendens</i>	Splendid Fairy-wren	LC				X
<i>Stipiturus malachurus</i>	Southern Emu-wren	LC		X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Acanthizidae Thornbills, Geryones, Fieldwrens & Whitefaces						
<i>Acanthiza apicalis</i>	Broad-tailed Thornbill	LC	X	X	X	X
<i>Acanthiza chrysorrhoa</i>	Yellow-rumped Thornbill	LC	X	X		X
<i>Acanthiza uropygialis</i>	Chestnut-rumped Thornbill	LC	X	X		X
<i>Aphelocephala leucopsis</i>	Southern Whiteface	LC				
<i>Calamanthus campestris</i>	Rufous Fieldwren	LC	X	X		X
<i>Gerygone fusca</i>	Western Gerygone	LC	X	X		X
<i>Hylacola cauta whitlocki</i>	Shy Heath-wren (western)	LC	X	X		
<i>Pyrholaemus brunneus</i>	Redthroat	LC	X	X	X	X
<i>Sericornis frontalis</i>	White-browed Scrubwren	LC	X			X
<i>Smicromis brevirostris</i>	Weebill	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Pardalotidae Pardalotes						
<i>Pardalotus punctatus</i>	Spotted Pardalote	LC	X	X	X	X
<i>Pardalotus striatus</i>	Striated Pardalote	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmis - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Meliphagidae Honeyeaters, Chats						
<i>Acanthagenys rufogularis</i>	Spiny-cheeked Honeyeater	LC		X		X
<i>Anthochaera carunculata</i>	Red Wattlebird	LC	X	X	X	X
<i>Certhionyx niger</i>	Black Honeyeater	LC			X	
<i>Certhionyx variegatus</i>	Pied Honeyeater	LC				X
<i>Epthianura albifrons</i>	White-fronted Chat	LC	X			X
<i>Epthianura tricolor</i>	Crimson Chat	LC				
<i>Lichenostomus cratitius</i>	Purple-gaped Honeyeater	LC	X	X		X
<i>Lichenostomus leucotis</i>	White-eared Honeyeater	LC	X	X	X	X
<i>Lichenostomus ornatus</i>	Yellow-plumed Honeyeater	LC	X	X	X	
<i>Lichenostomus plumulus</i>	Grey-fronted Honeyeater	LC		X		
<i>Lichenostomus virescens</i>	Singing Honeyeater	LC		X	X	
<i>Lichmera indistincta</i>	Brown Honeyeater	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
<i>Manorina flavigula</i>	Yellow-throated Miner	LC	X	X	X	X
<i>Melithreptus brevirostris</i>	Brown-headed Honeyeater	LC	X	X	X	X
<i>Phylidonyris albigrons</i>	White-fronted Honeyeater	LC	X	X	X	
<i>Phylidonyris melanops</i>	Tawny-crowned Honeyeater	LC	X	X		
<i>Phylidonyris nigra</i>	White-cheeked Honeyeater	LC		X		
<i>Phylidonyris novaehollandiae</i>	New Holland Honeyeater	LC		X		X
Petroicidae Australian Robins						
<i>Drymodes brunneopygia</i>	Southern Scrub-robin	LC	X	X	X	X
<i>Eopsaltria griseogularis</i>	Western Yellow Robin	LC	X	X	X	
<i>Microeca fascians</i>	Jacky Winter	LC	X	X		X
<i>Petroica cucullata</i>	Hooded Robin	LC	X	X		
<i>Petroica goodenovii</i>	Red-capped Robin	LC	X	X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Pomatostomidae Babblers						
<i>Pomatostomus superciliosus</i>	White-browed Babbler	LC	X	X		X
Cinclosomatidae Whipbirds, Wedgebills, Quail Thrushes						
<i>Cinclosoma castanotus</i>	Chestnut Quail-thrush	LC	X	X		
Neosittidae Sittellas						
<i>Daphoenositta chrysoptera</i>	Varied Sittella	LC	X	X	X	X
Pachycephalidae Crested Shrike-tit, Crested Bellbird, Shrike Thrushes, Whistlers						
<i>Colluricincla harmonica</i>	Grey Shrike-thrush	LC	X	X	X	X
<i>Falcunculus frontatus</i>	Crested Shrike-tit	LC				X
<i>Oreoica gutturalis</i>	Crested Bellbird	LC	X	X	X	X
<i>Pachycephala inornata</i>	Gilbert's Whistler	LC				X
<i>Pachycephala pectoralis</i>	Golden Whistler	LC	X	X	X	
<i>Pachycephala rufiventris</i>	Rufous Whistler	LC		X	X	X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status			
		A	B	C	D
Dicruridae Monarchs, Magpie Lark, Flycatchers, Fantails, Drongo					
<i>Grallina cyanoleuca</i>	Magpie-lark	LC	X		X
<i>Myiagra inquieta</i>	Restless Flycatcher	LC			X
<i>Rhipidura fuliginosa</i>	Grey Fantail	LC	X	X	X
<i>Rhipidura leucophrys</i>	Willie Wagtail	LC	X	X	X
Campephagidae Cuckoo-shrikes, Trillers					
<i>Coracina novaehollandiae</i>	Black-faced Cuckoo-shrike	LC	X	X	X
<i>Lalage tricolor</i>	White-winged Triller	LC	X	X	X
Artamidae Woodswallows, Butcherbirds, Currawongs					
<i>Artamus cinereus</i>	Black-faced Woodswallow	LC	X	X	X
<i>Artamus cyanopterus</i>	Dusky Woodswallow	LC	X	X	X
<i>Artamus personatus</i>	Masked Woodswallow	LC			X
<i>Artamus superciliosus</i>	White-browed Woodswallow	LC			

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Cracticidae Currawongs, Magpies & Butcherbirds						
<i>Cracticus nigrogularis</i>	Pied Butcherbird	LC		X	X	X
<i>Cracticus tibicen</i>	Australian Magpie	LC		X		X
<i>Cracticus torquatus</i>	Grey Butcherbird	LC	X	X		X
<i>Strepera versicolor</i>	Grey Currawong	LC		X	X	X
Corvidae Ravens, Crows						
<i>Corvus bennetti</i>	Little Crow	LC				X
<i>Corvus coronoides</i>	Australian Raven	LC		X	X	X
<i>Corvus sp.</i>	Unidentified corvid	LC				
Motacillidae Old World Pipits, Wagtails						
<i>Anthus australis</i>	Australian Pipit	LC	X	X		X
Estrilidae Grass Finches & Mannikins						
<i>Taeniopygia guttata</i>	Zebra Finch	LC				X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Dicaeidae Flowerpeckers						
<i>Dicaeum hirundinaceum</i>	Mistletoebird	LC	X			X
Hirundinidae Swallows, Martins						
<i>Cheramoeca leucosternus</i>	White-backed Swallow	LC				
<i>Hirundo ariel</i>	Fairy Martin	LC				
<i>Hirundo neoxena</i>	Welcome Swallow	LC				X
<i>Hirundo nigricans</i>	Tree Martin	LC	X	X		
Sylviidae Old World Warblers						
<i>Cinclooramphus cruralis</i>	Brown Songlark	LC				
<i>Cinclooramphus mathewsi</i>	Rufous Songlark	LC				
Zosteropidae White-eyes						
<i>Zosterops lateralis</i>	Silvereye	LC	X	X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Mammalia						
Tachyglossidae Echidnas						
<i>Tachyglossus aculeatus</i>	Echidna	LC	X	X		X
Dasyuridae Carnivorous Marsupials						
<i>Antechinomys laniger</i>	Kultarr	LC				
<i>Dasyurus geoffroii</i>	Chuditch	S3 VU VU		X		X
<i>Ningauai yvonneae</i>	Southern Ningauai	LC	X			X
<i>Sminthopsis crassicaudata</i>	Fat-tailed Dunnart	LC		X		X
<i>Sminthopsis dolichura</i>	Little long-tailed Dunnart	LC			X	X
<i>Sminthopsis gilberti</i>	Gilbert's Dunnart	LC	X	X		X
<i>Sminthopsis granulipes</i>	White-tailed Dunnart	LC	X	X		X
<i>Sminthopsis griseoventer</i>	Grey-bellied Dunnart	LC		X		
<i>Sminthopsis ooldea</i>	Ooldea Dunnart	LC				X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Burramyidae Pygmy Possums						
<i>Cercartetus concinnus</i>	Western Pygmy-possum	LC	X	X	X	X
Tarsipedidae Honey Possum						
<i>Tarsipes rostratus</i>	Honey Possum	LC		X		
Macropodidae Kangaroos, Wallabies						
<i>Macropus fuliginosus</i>	Western Grey Kangaroo	LC	X	X		X
<i>Macropus ima</i>	Western Brush Wallaby	P4 LC		X		
<i>Macropus robustus</i>	Euro	LC	X			
Molossidae Freetail Bats						
<i>Ozimops kitcheneri</i>	Western Freetail-bat	LC	X	X	X	
<i>Tadarida australis</i>	White-striped Freetail-bat	LC	X	X	X	

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Vespertilionidae Ordinary Bats						
<i>Chalinolobus gouldii</i>	Gould's Wattled Bat	LC	X	X	X	X
<i>Chalinolobus morio</i>	Chocolate Wattled Bat	LC		X	X	X
<i>Nyctophilus geoffroyi</i>	Lesser Long-eared Bat	LC	X	X	X	X
<i>Nyctophilus major tor</i>	Central Long-eared Bat	P4				
<i>Scotorepens balstoni</i>	Inland Broad-nosed Bat	LC			X	X
<i>Vespadelus baverstocki</i>	Inland Forest Bat	LC				X
<i>Vespadelus regulus</i>	Southern Forest Bat	LC	X	X	X	X
Muridae Rats, Mice						
<i>Mus musculus</i>	House Mouse	Introduced	X	X	X	X
<i>Notomys mitchellii</i>	Mitchell's Hopping-mouse	LC	X	X	X	X
<i>Pseudomys albocinereus</i>	Ash-grey Mouse	LC	X	X		X
<i>Pseudomys bolami</i>	Bolam's Mouse	LC	X			

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.

Class Family Species	Common Name	Conservation Status	A	B	C	D
Canidae Dogs, Foxes						
<i>Canis lupus</i>	Dingo/Dog	LC/Introduced	X	X		
<i>Vulpes vulpes</i>	Red Fox	Introduced	X	X		X
Felidae Cats						
<i>Felis catus</i>	Cat	Introduced		X	X	X
Camelidae Camels						
<i>Camelus dromedarius</i>	Camel	Introduced				
Leporidae Rabbits, Hares						
<i>Oryctolagus cuniculus</i>	Rabbit	Introduced	X	X		X

BC Act Status - S1 to S7, EPBC Act Status - CR = Critically Endangered, EN = Endangered, VU = Vulnerable, EX = Extinct, DPaW Priority Status - P1 to P4, Int. Agmts - CA = CAMBA, JA = JAMBA, RK = ROKAMBA, IUCN Red List Category Definitions LC = Least Concern - see Appendix A and <https://www.iucnredlist.org/resources/categories-and-criteria> for others.