



**Reconnaissance Flora and Vegetation  
Survey of the Reedy Central Project  
Area- June 2026**

Prepared for



**Valiant Gold**

**Valiant Gold Ltd**

FINAL V2.0  
July 2026

Prepared by:  
Native Vegetation Solutions  
PO Box 41  
KALGOORLIE  
Ph: (08) 9021 5818  
Mob: 0407 998 953  
Email: [eren@nativevegsolutions.com.au](mailto:eren@nativevegsolutions.com.au)

|           |                                                                                  |           |
|-----------|----------------------------------------------------------------------------------|-----------|
| <b>1</b>  | <b>INTRODUCTION .....</b>                                                        | <b>1</b>  |
| 1.1       | PURPOSE AND SCOPE .....                                                          | 3         |
| 1.2       | STATUTORY FRAMEWORK AND GUIDANCE .....                                           | 3         |
| 1.2.1     | Western Australian Biodiversity Conservation Act 2016 .....                      | 3         |
| 1.2.2     | Environmental Protection Act 1986 .....                                          | 4         |
| 1.2.3     | Environment Protection and Biodiversity Conservation Act 1999 .....              | 4         |
| 1.2.4     | Flora .....                                                                      | 4         |
| 1.2.5     | Ecological Communities and Vegetation .....                                      | 5         |
| <b>2.</b> | <b>EXISTING ENVIRONMENT .....</b>                                                | <b>7</b>  |
| 2.1       | GEOLOGY AND VEGETATION.....                                                      | 7         |
| 2.2       | CLIMATE .....                                                                    | 7         |
| 2.2.1     | Temperature.....                                                                 | 7         |
| 2.2.2     | Rainfall .....                                                                   | 7         |
| <b>3.</b> | <b>ASSESSMENT METHODOLOGY .....</b>                                              | <b>9</b>  |
| 3.1       | PERSONNEL AND REPORTING .....                                                    | 9         |
| 3.2       | PRELIMINARY DESKTOP STUDY .....                                                  | 9         |
| 3.2.1     | Known Previous Flora and Vegetation Surveys.....                                 | 9         |
| 3.2.2     | Environment Protection and Biodiversity Conservation Act Protected Matters ..... | 11        |
| 3.2.3     | Threatened Flora and Communities .....                                           | 11        |
| 3.2.4     | Environmentally Sensitive Areas (ESAs) and Conservation Reserves .....           | 11        |
| 3.2.5     | Vegetation Type, Extent and Status .....                                         | 11        |
| 3.2.6     | Wetlands.....                                                                    | 11        |
| 3.2.7     | Dieback.....                                                                     | 11        |
| 3.3       | SITE INVESTIGATION .....                                                         | 12        |
| 3.3.1     | Licenses.....                                                                    | 12        |
| 3.3.2     | Field Methods.....                                                               | 12        |
| 3.3.3     | Post-Field Methods.....                                                          | 13        |
| 3.3.4     | Mapping.....                                                                     | 13        |
| 3.3.5     | IBSA Data Package .....                                                          | 13        |
| 3.4       | NOMENCLATURE AND TAXONOMY .....                                                  | 14        |
| 3.5       | LIMITATIONS .....                                                                | 14        |
| <b>4.</b> | <b>RESULTS.....</b>                                                              | <b>15</b> |
| 4.1       | PRELIMINARY DESKTOP ASSESSMENT .....                                             | 15        |
| 4.1.1     | EPBC Act Protected Matters .....                                                 | 15        |
| 4.1.2     | Threatened Flora and Communities .....                                           | 15        |
| 4.1.3     | Environmentally Sensitive Areas and Conservation Reserves .....                  | 15        |
| 4.1.4     | Land Systems .....                                                               | 16        |
| 4.1.5     | Vegetation Type, Extent and Status .....                                         | 16        |
| 4.1.6     | Wetlands.....                                                                    | 18        |
| 4.1.7     | Dieback.....                                                                     | 18        |
| 4.2       | FIELD ASSESSMENT .....                                                           | 19        |
| 4.2.1     | Threatened Flora.....                                                            | 19        |
| 4.2.2     | Vegetation Type, Extent and Status .....                                         | 20        |
| 4.2.3     | Weeds .....                                                                      | 36        |
| 4.2.4     | Vegetation Condition .....                                                       | 37        |
| 4.2.5     | Clearing Principals .....                                                        | 38        |

|                            |           |
|----------------------------|-----------|
| <b>5. DISCUSSION .....</b> | <b>39</b> |
| <b>6. REFERENCES .....</b> | <b>40</b> |
| <b>7. GLOSSARY .....</b>   | <b>42</b> |

### Appendices

|                                                              |    |
|--------------------------------------------------------------|----|
| Appendix 1: Relevant Government Database Search Results..... | 47 |
| Appendix 2: Threatened Flora Databases Search Results .....  | 58 |
| Appendix 3: Vegetation Definitions.....                      | 60 |
| Appendix 4: Vegetation Mapping .....                         | 63 |
| Appendix 5: Species List.....                                | 72 |

### Figures

|                                                                              |    |
|------------------------------------------------------------------------------|----|
| Figure 1: Regional map of survey location .....                              | 2  |
| Figure 2: Mean temperature ranges for Cue weather station.....               | 7  |
| Figure 3: Monthly and mean rainfall for Cue weather station .....            | 8  |
| Figure 4: Central Reedy survey area in comparison to past survey areas ..... | 10 |
| Figure 5: <i>Dodonaea amplisemina</i> (P4) .....                             | 19 |
| Figure 6: Vegetation Group A within the survey area .....                    | 21 |
| Figure 7: Vegetation Group B within the survey area.....                     | 22 |
| Figure 8: Vegetation Group C within the survey area.....                     | 23 |
| Figure 9: Vegetation Group D within the survey area .....                    | 24 |
| Figure 10: Vegetation Group E within the survey area .....                   | 25 |
| Figure 11: Vegetation Group F within the survey area .....                   | 26 |
| Figure 12: Vegetation Group G within the survey area .....                   | 27 |
| Figure 13: Vegetation Group H within the survey area .....                   | 28 |
| Figure 14: Vegetation Group I within the survey area .....                   | 29 |
| Figure 15: Vegetation Group J within the survey area.....                    | 30 |
| Figure 16: Vegetation Group K within the survey area .....                   | 31 |
| Figure 17: Vegetation Group L within the survey area .....                   | 32 |
| Figure 18: Vegetation Group M within the survey area.....                    | 33 |
| Figure 19: Vegetation Group N within the survey area .....                   | 34 |
| Figure 20: Existing disturbance within the survey area .....                 | 35 |

### Tables

|                                                                                                                                               |    |
|-----------------------------------------------------------------------------------------------------------------------------------------------|----|
| Table 1: Previous Flora Surveys .....                                                                                                         | 9  |
| Table 2: List of potential survey limitations.....                                                                                            | 14 |
| Table 3: Land Systems occurring within the survey area (DPIRD, 2017) .....                                                                    | 16 |
| Table 4: Extent of Beard Associations within the survey area.....                                                                             | 17 |
| Table 5: Summary of information regarding pre-European and current vegetation extent of Vegetation Association 39 within the survey area..... | 17 |
| Table 6: Summary of information regarding pre-European and current vegetation extent of Vegetation Association 18 within the survey area..... | 18 |
| Table 7: Priority Flora recorded in the survey area .....                                                                                     | 19 |
| Table 8: Vegetation Group Summary .....                                                                                                       | 20 |
| Table 9: Weed Abundance recorded in the survey area .....                                                                                     | 36 |
| Table 10: Assessment of impact using the native vegetation clearing principles .....                                                          | 38 |

## 1 INTRODUCTION

Valiant Gold Ltd (ASX:VAL) are proposing to develop their Reedy Central project area located in Western Australia's Murchison Region (DCCEEW, 2026).

VAL require a reconnaissance flora and vegetation survey to determine any impact to flora and vegetation within the Reedy Central project area.

Native Vegetation Solutions (NVS) was supplied with a survey area located approximately 55 km south of Meekatharra, in the Murchison Region (MUR) of Western Australia (Figure 1).

The total survey area received from VAL covered approximately 971.21 ha. The survey area intersects Mining Tenements M20/12, M20/45, M20/68, M20/77, M20/214, and M20/421. A NVCP was historically approved for the same area for Westgold Resources Ltd, however due to a change in Tenement ownership, a new NVCP application is required by VAL. The new NVCP application requires a more recent flora and vegetation survey to support the application.

This report will encompass results of the reconnaissance flora and vegetation survey within the Reedy Central project survey area.



Figure 1: Regional map of survey location

## 1.1 Purpose and Scope

The objective of this report is to document the results of the flora and vegetation component of a reconnaissance assessment conducted in accordance with:

- Environmental Factor Guideline- Flora and Vegetation (EPA, 2016); and
- Technical Guidance- Flora and Vegetation Surveys for Environmental Impact Assessment (EPA, 2016a).

A reconnaissance assessment has two components:

- 1). Desktop study which includes a literature review and a search of the relevant databases;
- 2). Reconnaissance survey of the survey area to verify the desktop survey, to define vegetation units present in the area, search for species of conservation significance and to determine potential sensitivity to impact.

As part of the reporting for the reconnaissance assessment, NVS has conducted a flora and vegetation survey which includes broad-scale vegetation mapping and vegetation condition mapping of the survey area.

The scope of work for the reconnaissance flora and vegetation survey was to:

- conduct a desktop study that includes a literature review and search of the relevant databases;
- describe the vegetation associations in the survey area;
- prepare an inventory of species occurring in the survey area;
- identify any vegetation communities or flora species of conservation significance;
- map broad-scale vegetation groups found within the survey area, including vegetation condition; and
- provide recommendations, including the management of perceived impacts to flora and vegetation within the survey area.

## 1.2 Statutory Framework and Guidance

This assessment took into account relevant sections of Commonwealth and State legislation and guidelines:

- *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act)
- *Environmental Protection Act 1986* (EP Act)
- *Biodiversity Conservation Act 2016* (BC Act)
- *Biosecurity and Agriculture Management Act 2007* (BAM Act)

The Minister for the Environment publishes lists of flora species in need of special protection because they are considered rare, likely to become extinct, or are presumed extinct. The current listings were published in the Government Gazette on 30 April 2024 (Lawn, 2024) and were taken into account.

As well as those listed above, the assessment took into account relevant sections of:

- EPA (2023) *Statement of environmental principles, factors, objectives and aims of EIA*;
- EPA (2016) *Environmental Factor Guideline: Flora and Vegetation*; and
- EPA (2016a) *Technical Guidance - Flora and Vegetation Surveys for Environmental Impact Assessment*, known as *Flora and Vegetation Technical Guidance*

### 1.2.1 Western Australian Biodiversity Conservation Act 2016

The Western Australian *Biodiversity Conservation Act 2016* (BC Act) provides for the conservation, protection and ecologically sustainable use of biodiversity and biodiversity components in Western Australia. The BC Act replaced the *Wildlife Conservation Act 1950*.

Threatened species (both flora and fauna) that meet the categories listed within the Act are highly protected and require authorisation by the Ministerial to take or disturb. These are known as Threatened Flora and Threatened Fauna. The conservation categories of Critically Endangered, Endangered and Vulnerable have been aligned with those detailed in the EPBC Act, as below.

Flora and fauna species may be listed as being of special conservation interest if they have a naturally low population, restricted natural range, are subject to or recovering from a significant population decline or reduction of range or are of special interest, and the Minister considers that taking may result in depletion of the species. Migratory species and those subject to international agreement are also listed under the BC Act. These are known as specially protected species in the BC Act.

Threatened Ecological Communities (TECs) are also protected under BC Act and are categorised using the same criteria as threatened species.

### **1.2.2 Environmental Protection Act 1986**

The *EP Act 1986* was created to provide for an Environmental Protection Authority (the EPA) that has the responsibility for:

- prevention, control and abatement of pollution and environmental harm
- conservation, preservation, protection, enhancement and management of the environment
- matters incidental to or connected with the above.

The EPA is responsible for providing the guidance and policy under which environmental assessments are conducted. It conducts environmental impact assessments (based on the information included in environmental assessments and provided by the proponent), initiates measures to protect the environment and provides advice to the Minister responsible for environmental matters.

### **1.2.3 Environment Protection and Biodiversity Conservation Act 1999**

At a Commonwealth level, Threatened taxa are protected under the EPBC Act, which lists species and ecological communities that are considered Critically Endangered, Endangered, Vulnerable, Conservation Dependent, Extinct, or Extinct in the Wild (Section 6 below).

### **1.2.4 Flora**

#### **1.2.4.1 Threatened and Priority Flora**

Conservation significant flora species are those that are listed as TF (Threatened Flora) and (within Western Australia) as PF (Priority Flora). TF species are listed as threatened by the Western Australian Department of Biodiversity Conservation and Attractions (DBCA) and protected under the provisions of the BC Act. Some State-listed TF are provided with additional protection as they are also listed under the Commonwealth EPBC Act. Species can also be listed under the EPBC Act without being listed under the BC Act.

Flora are listed as PF where populations are geographically restricted or threatened by local processes, or where there is insufficient information to formally assign them to TF categories. Whilst PF are not specifically listed in the BC Act, some may qualify as being of special conservation interest and these may require a greater level of protection than unlisted species. Generally though, PF have no statutory protection. They are generally considered in environmental impact assessments under the state approval processes by Department of Mines, Industry Regulation and Safety (DEMIRS) under the *Mining Act 1978* and DBCA under the EP Act. Under this approval process measures are usually taken to protect and avoid PF.

There are seven categories covering State-listed TF and PF species (DBCA, 2023) which are defined in Section 7 below. PF for Western Australia are regularly reviewed by DBCA whenever new information becomes available, with species status altered or removed from the list (Lawn, 2024) when data indicates that they no longer meet the requirements outlined in Section 7 below.

#### **1.2.4.2 Other Significant Flora**

According to the Flora and Vegetation Technical Guidance (EPA 2016a) other than being listed as Threatened or Priority Flora, a species can be considered as significant if it is considered to be:

- locally endemic or association with a restricted habitat type (e.g., surface water or groundwater dependent ecosystems)
- a new species or has anomalous features that indicate a potential new species

- at the extremes of range, recently discovered range extensions (generally considered greater than 100 km or in a different bioregion), or isolated outliers of the main range
- unusual species, including restricted subspecies, varieties or naturally occurring hybrids and
- relictual status, being representative of taxonomic groups that no longer occur widely in the broader landscape.

## **1.2.5 Ecological Communities and Vegetation**

### **1.2.5.1 Threatened and Priority Ecological Communities**

#### **Nationally Listed Threatened Ecological Communities**

An ecological community is a naturally occurring group of plants, animals and other organisms interacting in a unique habitat. The complex range of interactions between the component species provides an important level of biological diversity in addition to genetics and species. At Commonwealth level, Threatened Flora and TECs are protected under the Commonwealth EPBC Act. An ecological community may be categorised into one of the three subcategories:

- Critically Endangered, if it is facing an extremely high risk of extinction in the wild in the immediate future
- Endangered, if it is not critically endangered and is facing a very high risk of extinction in the wild in the near future and
- Vulnerable, if it is not critically endangered or endangered, and is facing a high risk of extinction in the wild in the medium-term future.

#### **State Listed Threatened Ecological Communities**

The Western Australian DBCA also maintains a list of TECs, protected under the BC Act, which are further categorised into three subcategories much like those of the EPBC Act.

#### **State Listed Priority Ecological Communities**

DBCA maintains a list of Priority Ecological Communities (PECs). PECs include potential TECs that do not meet survey criteria, or that are not adequately defined.

### **1.2.5.2 Other Significant Vegetation**

According to the Flora and Vegetation Technical Guidance (EPA 2016a), other than being listed as a TEC or PEC, vegetation can be considered as significant if it is considered to have:

- restricted distribution
- a degree of historical impact from threatening processes
- a role as a refuge; and/or
- provides an important function required to maintain ecological integrity of a significant ecosystem.

### **1.2.5.3 Declared Pest Plants**

The Western Australian Organism List (WAOL) details organisms listed as Declared Pests under the BAM Act). Under the BAM Act, Declared Pests are listed as one of the three categories, or exempt:

- C1 (exclusion), that applies to pests not established in Western Australia; control measures are to be taken to prevent their entry and establishment
- C2 (eradication), that applies to pests that are present in Western Australia but in low numbers or in limited areas where eradication is still a possibility
- C3 (management), that applies to established pests where it is not feasible or desirable to manage them in order to limit their damage; or
- Exempt (no category).

## 2. EXISTING ENVIRONMENT

### 2.1 Geology and Vegetation

The survey area lies in the Murchison (MUR) bioregion, more specifically the Eastern Murchison (MUR01) subregion. The Eastern Murchison subregion covers over 7.8 million hectares and lies on the northern parts of the 'Southern Cross' and 'Eastern Goldfields' Terrains of the Yilgarn Craton. Characterised by its internal drainage, and extensive areas of elevated red desert sandplains with minimal dune development. Salt lake systems are associated with the occluded Paleodrainage system, broad plains of red-brown soils and breakaway complexes as well as red sandplains (CALM, 2002).

The vegetation is dominated by Mulga Woodlands often rich in ephemerals; hummock grasslands, saltbush shrublands and *Tecticornia* shrublands. (CALM, 2002)

### 2.2 Climate

The climate of the Murchison Region is classified as arid with 200 mm of rainfall, sometimes in summer but usually in winter (CALM, 2002). The nearest official meteorological weather station with the most complete and up to date temperature information is Cue (station number 007017), which is located approximately 48.5 km south of the survey area.

#### 2.2.1 Temperature

Mean annual minimum temperature at Cue is 14.7°C and mean annual maximum temperature is 28.4°C (BOM, 2026). The coldest temperatures are attained in July (mean minimum temperature 6.9°C), the hottest is January (mean maximum temperature 37.8°C) and diurnal temperature variations are relatively consistent throughout the year (Figure 2).




Figure 2: Mean temperature ranges for Cue weather station

#### 2.2.2 Rainfall

The annual average rainfall at Cue is 230.8 mm, which falls (>1 mm) on an average of 28.3 rain-days (BOM, 2026). Larger rainfall events occur from January to July (Figure 3). Prior to the survey in 2026, rainfall in March and June exceeded the monthly averages, whilst February was on par with its monthly average. Rainfall for all other months remained below monthly averages (BOM, 2026).




**Figure 3: Monthly and mean rainfall for Cue weather station**

### 3. ASSESSMENT METHODOLOGY

#### 3.1 Personnel and Reporting

The following personnel were involved in the Reconnaissance flora and vegetation survey:

- Mr Eren Reid (*BSc- Biological Science*), Principal Botanist, Native Vegetation Solutions, undertook the survey, vegetation mapping, data collation, field identification of flora, preparation and review of the report. Mr Eren Reid has over 22 years' experience in botanical surveys throughout the Murchison Region and over a variety of environments across Western Australia.

#### 3.2 Preliminary Desktop Study

A preliminary assessment of the survey area and its potential constraints was undertaken by reviewing relevant government agency managed databases (Sections 3.2.1 to 3.2.6, and Appendices 1 & 2) and consulting with government agencies where necessary. The following sections provide a summary of desktop searches undertaken for the project.

##### 3.2.1 Known Previous Flora and Vegetation Surveys

Three known Reconnaissance Flora and Vegetation Surveys have previously been completed near the Reedy Central Project in April 2015, April 2016 and November 2021. The results of these are summarised in Table 1 below. These reports were referenced for locations of conservation significant flora.

**Table 1: Previous Flora Surveys**

| Reference      | Date          | Survey Type                           | Location relevant to current survey area | Threatened Flora recorded                                                                                                |
|----------------|---------------|---------------------------------------|------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| MWH, (2015)    | April 2015    | Level 1 Flora and Fauna Survey        | Encompassed existing area                | <ul style="list-style-type: none"><li>• <i>Ptilotus beardii</i> (P3)- locations outside of current survey area</li></ul> |
| MWH (2016)     | April 2016    | Level 1 Flora and Fauna Survey        | 4km east of current survey area          | <ul style="list-style-type: none"><li>• <i>None</i></li></ul>                                                            |
| Stantec (2022) | November 2021 | Reconnaissance Flora and Fauna Survey | 6.35km north of current survey area      | <ul style="list-style-type: none"><li>• <i>Tecticornia cymbiformis</i> (P3)</li></ul>                                    |




Figure 4: Central Reedy survey area in comparison to past survey areas

### **3.2.2 Environment Protection and Biodiversity Conservation Act Protected Matters**

The *EPBC Act* Protected Matters Search tool was utilised to provide results for matters of National Environmental Significance within the survey area using the survey area as the search criteria with a 20 km buffer (DCCEEW, 2026a).

### **3.2.3 Threatened Flora and Communities**

The Threatened and Priority Flora Database managed by the Department of Biodiversity, Conservation and Attractions (DBCA) was searched for threatened and priority flora within a 20km radius of the survey area (DBCA, 2026a).

The TEC and PEC database was searched to determine the presence of PECs or TECs (DBCA, 2026), with Geographic Information System (GIS) data supplied for assessment, within a 20 km radius of the survey area.

### **3.2.4 Environmentally Sensitive Areas (ESAs) and Conservation Reserves**

The Department of Water and Environmental Regulation (DWER, 2026) Clearing Permit System Map Viewer was used to determine the location of any ESAs and Conservation Reserves.

### **3.2.5 Vegetation Type, Extent and Status**

Vegetation extent and status data was sourced from the Department of Agriculture and Food (DAFWA) report and its associated GIS file (Shepherd *et al*, 2002). This data comprises Beard's Pre-European vegetation groups.

DBCA's Statewide Vegetation Statistics (DBCA, 2019) was also referenced for the current extent of Beard's Vegetation Groups. The purpose of examining this information is to determine if the survey area lies within any vegetation groups defined by Beard that may have been subjected to widescale clearing for European settlement. The national objectives and targets for biodiversity conservation recognise that the retention of 30% or more of the pre-clearing extent of a Beard vegetation association is necessary if Australia's biological diversity is to be protected.

### **3.2.6 Wetlands**

The potential of wetlands within the project area was determined by examining DWER's Clearing Permit System Map Viewer (DWER, 2026).

### **3.2.7 Dieback**

Under normal circumstances Dieback is only considered a potential issue for any project if the project area lies within the Southwest Land Division and the mean annual rainfall of the area is greater than 400 mm. There is no record of *Phytophthora cinnamomi* (Dieback) establishing in natural ecosystems in regions receiving <400mm rainfall per annum (CALM, 2003).

However, as indicated within the more recent Dieback guidelines (DBCA, 2020), other species of *Phytophthora* may persist east of the 400mm isohyet in unusually wet conditions. It is therefore recommended to conduct a risk assessment as per these guidelines.

### **3.3 Site Investigation**

A site visit of the survey area was carried out by Botanist Eren Reid from Native Vegetation Solutions on the 25<sup>th</sup> and 26<sup>th</sup> of 2026 to examine the flora and vegetation groups contained within the survey area. A total of 24 hours was spent on site traversing the survey area, by Yamaha Viking All-Terrain Vehicle (ATV) and on foot.

The survey was conducted in accordance with relevant Environmental Protection Authority's (EPA's) Statements and Technical Guidance (Section 1.1).

The EPA uses the Interim Biogeographic Regionalisation of Australia (IBRA) as the largest unit for Environmental Impact Assessment (EIA) decision making in relation to the conservation of biodiversity. Given the scale and nature of the proposed disturbance as well as the existing disturbance, and that the survey area is located within the Murchison (MUR) IBRA region, a reconnaissance flora and vegetation survey was deemed adequate.

#### **3.3.1 Licenses**

A Scientific License was not required for the field work as no samples were collected for identification. All taxa were able to be identified in the field.

#### **3.3.2 Field Methods**

Prior to the field work, the aerial photography was examined and representative sample sites for relevés were chosen to provide coverage over all potential vegetation types.

In the field, 20m x 20m non-permanent relevé sites were established at these sites, taking into account representation of surrounding vegetation and vegetation boundaries. Relevé sites are represented in Appendix 4.

Each relevé site was captured on a TwoNav Aventura GPS at ±4m accuracy, using Universal Transverse Mercator location on GDA94 datum. Digital photographs were taken of each representative vegetation group present in the survey area.

Data collected at each relevé included:

- Photograph of representative vegetation group;
- GPS Location;
- Species Present;
- Population Count/Estimate of Conservation Significant Flora (if present);
- Disturbance Level; and
- Vegetation Condition

The vegetation structure was assessed using the method developed by Muir (1977). Definitions of the vegetation structure are presented in Appendix 3.

The condition of each relevé was assessed using the method developed by Keighery (1994). Definitions of the condition scale are presented in Appendix 3.

Vegetation groups were mapped using the methods listed in Section 3.3.4 below.

Opportunistic recording of plant taxa and vegetation group mapping was also utilised in the survey area between relevé sampling points, via wandering traverses. Smaller singular relevé sites were also utilised as opportunistic sample sites to record taxa and assist in mapping vegetation groups.

All relevé sample sites and GPS tracks are included in Appendix 4.

### 3.3.3 Post-Field Methods

Taxa were identified with the use of information published on Florabase (WAHERB, 2026). Threatened flora range extensions and new locations were submitted to the Western Australian Herbarium (WAHERB) as per the EPA Technical Guidelines (EPA 2016a).

Species information was transferred into Microsoft Excel<sup>®</sup> worksheets representing presence/absence of species per vegetation group.

### 3.3.4 Mapping

Vegetation mapping was produced via GPS recorded information in the field, cross-referenced with vegetation descriptions made in the field, overlaid on aerial imagery of the survey area. The GPS utilised (TwoNav Aventura GPS) displayed aerial imagery, hence real-time mapping of vegetation groups was available during field work.

Vegetation Health Condition was assessed in the field with reference to Keighery (1994).

GPS tracks and waypoints recorded during field work are presented in Appendix 4.

### 3.3.5 IBSA Data Package

The Environmental Protection Authority (EPA), Department of Water and Environmental Regulation (DWER) and Department of Mines, Petroleum and Exploration (DMPE) require Index of Biodiversity Surveys for Assessments (IBSA) Data Packages to be submitted to support assessment and compliance under the *Environmental Protection Act 1986*.

An IBSA data package is a single file in .zip format, containing:

- one survey report in .pdf format;
- one plain-text survey report in .txt format; and
- a set of electronic data files, comprising:
  - one survey details spatial dataset in shapefile (.shp, etc.) or MapInfo (.tab, etc.) format; and
  - one or more survey data spatial datasets, as required, in shapefile (.shp, etc.) or MapInfo (.tab, etc.) format.

The IBSA Data package for this survey will be submitted via the DWER IBSA Submission Portal.

### 3.4 Nomenclature And Taxonomy

Nomenclature follows that used by the WAHERB.

The WAHERB has updated its sequence and arrangement of collections to conform to the systematic sequence of the Angiosperm Phylogeny Group (APGIV), with the result that many Families and Genera have been moved or renamed. This report attempts to follow those changes in relation to species recorded during this survey.

### 3.5 Limitations

Table 2 lists potential limitations that may have affected the survey.

**Table 2: List of potential survey limitations**

| Potential Limitations                                               | Constraint (Y/N) | Comment                                                                                                                                                                                                                                                                                                                                                                                           |
|---------------------------------------------------------------------|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Competency and experience of the consultants undertaking the survey | N                | Experienced and competent personnel conducted the survey. Eren Reid ( <i>BSc</i> ) has over 22 years' experience in botanical surveys throughout the Murchison Region and over a variety of environments across Western Australia.                                                                                                                                                                |
| Scope                                                               | N                | The Scope of work was adequately defined. Vascular flora species were the focus of the survey and were thoroughly sampled.                                                                                                                                                                                                                                                                        |
| Proportion of flora identified during survey                        | N                | As the survey was planned to target species of conservation significance and flora within a defined survey area, a complete census of the species present was attempted (Approx. 95%). Sufficient identifications were made to allow vegetation descriptions to be made.                                                                                                                          |
| Sources of information                                              | N                | Threatened and Priority Flora GIS information was available from DBCA.                                                                                                                                                                                                                                                                                                                            |
| Proportion of the task achieved                                     | N                | All tasks completed.                                                                                                                                                                                                                                                                                                                                                                              |
| Timing/Season                                                       | N                | The reconnaissance flora and vegetation survey was conducted in June 2026. Flowering annual species were present within the survey area, suggesting recent above average rainfall in March, and normal average rainfall in February 2026 was sufficient for the period of survey. The most recent rainfall received in the area was from 12 <sup>th</sup> to 16 <sup>th</sup> June 2026 (34.2mm). |
| Disturbance in survey area                                          | N                | Minor disturbance (historical mining access tracks and exploration) was observed within the survey area, however, did not compromise the results of the survey as these areas were avoided whilst collecting data.                                                                                                                                                                                |
| Intensity of survey effort                                          | N                | The survey intensity is considered to have been sufficient for a reconnaissance survey according to EPA (2016) guidelines. Areas most likely to contain threatened and priority species were targeted. Vegetation mapping sites were selected to provide adequate coverage of the survey area.                                                                                                    |
| Resources                                                           | N                | Resources, in terms of time, equipment, support and personnel were adequate to undertake and complete the reconnaissance survey.                                                                                                                                                                                                                                                                  |
| Access problems                                                     | N                | All the areas in need of survey were easily accessible from existing tracks, or by foot.                                                                                                                                                                                                                                                                                                          |
| Availability of contextual information on the region                | N                | Contextual information regarding vegetation and flora of the Murchison bioregion is readily available. Adequate information was able to be accessed from available databases.                                                                                                                                                                                                                     |

## **4. RESULTS**

### **4.1 Preliminary Desktop Assessment**

#### **4.1.1 EPBC Act Protected Matters**

Results of the EPBC Protected Matters search tool are included in Appendix 1.

The EPBC Protected Matters report indicated no TECs or Commonwealth Reserves within the requested survey area.

The search results indicated that no Threatened plant species or habitat occurs in the buffer of the requested survey area.

#### **4.1.2 Threatened Flora and Communities**

The DBCA database searches revealed a potential for no Threatened and eight Priority Flora species to occur within a 20km radius of the survey area (DBCA, 2026a). No known locations of Threatened or Priority Flora occur within the survey area, with the closest Priority Flora located approximately 2.39 km east of the survey area, as shown on Map 3 of Appendix 4.

Priority Flora species within a 20 km radius of the survey area are displayed in Map 3 of Appendix 4.

Results of the threatened flora database search are included in Appendix 2 which includes the likelihood of each species to occur within the survey area.

The PEC/TEC search (DBCA, 2026a) revealed that no TECs or PECs fall within the survey area, or within 10km of the survey area.

#### **4.1.3 Environmentally Sensitive Areas and Conservation Reserves**

No ESA's or Conservation Reserves intersect with the survey area (DWER, 2026).

#### 4.1.4 Land Systems

As part of the Rangeland resource surveys, the Department of Agriculture mapped the Land Systems of Western Australia (DPIRD, 2017). The Land Systems occurring within the survey area are listed in Table 3 below and displayed in Appendix 4.

**Table 3: Land Systems occurring within the survey area (DPIRD, 2017)**

| Land System       | Description                                                                                                                                                                                                                  | Extent of Survey Area (ha) | % Of Survey Area (%) |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|----------------------|
| Violet system     | Gently undulating gravelly plains on greenstone, laterite and hardpan, with low stony rises and minor saline plains; supporting groved mulga and bowgada shrublands and occasionally chenopod shrublands                     | 401.21                     | 41.38%               |
| Millex system     | Plains on granite, with irregularly distributed low sandy banks and saline alluvial plains lightly strewn with quartz mantles; supporting mulga shrublands and low halophytic shrublands                                     | 160.02                     | 16.50%               |
| Mindura system    | Low hills, ridges and outcrops of granite, gneiss and quartz above convex, quartz-strewn interfluvies and lower plains supporting sparse acacia shrublands becoming denser in drainage floors                                | 128.59                     | 13.26%               |
| Gabanintha system | Greenstone ridges, hills and footslopes supporting sparse acacia and other mainly non-halophytic shrublands                                                                                                                  | 127.83                     | 13.18%               |
| Wiluna system     | Low greenstone hills with occasional lateritic breakaways and broad stony slopes, lower saline stony plains and broad drainage tracts; supporting sparse mulga and other acacia shrublands with patches of halophytic shrubs | 83.58                      | 8.62%                |
| Jundee system     | Hardpan plains with variable gravelly mantles and minor sandy banks supporting weakly groved mulga shrublands                                                                                                                | 55.47                      | 5.72%                |
| Sherwood system   | Breakaways, kaolinised footslopes and extensive gently sloping plains on granite supporting mulga shrublands and minor halophytic shrublands                                                                                 | 12.89                      | 1.33%                |

#### 4.1.5 Vegetation Type, Extent and Status

Two vegetation units defined by Beard (1990) were identified as part of the desktop assessment. The vegetation units identify the pre-European extent of vegetation, as mapped by Beard (1990). The national objectives and targets for biodiversity conservation recognise that the retention of 30% or more of the pre-clearing extent of Beard's vegetation associations is necessary if Australia's biological diversity is to be protected.

Information relating to known Beard (1990) vegetation units within the survey area has been summarised in Table 4, Table 5 and Table 6 below. This information has been compiled through both desktop assessments and the site visit.

The extent of the Beard vegetation unit within the survey area at all scales is less than 1% of the total area at each scale (Table 4). The single Beard vegetation unit is above the 30% threshold at a State, bioregional and subregional scale.

**Table 4: Extent of Beard Associations within the survey area**

| Beard Vegetation Association | Extent within survey area (ha) | % of survey area (%) | By Association WA | By IBRA Region (MUR) | By IBRA Sub-region (MUR01) | By Shire (Shire of Cue) |
|------------------------------|--------------------------------|----------------------|-------------------|----------------------|----------------------------|-------------------------|
| 39                           | 715.68                         | 73.81%               | <1%               | <1%                  | <1%                        | <1%                     |
| 18                           | 253.89                         | 26.19%               | <1%               | <1%                  | <1%                        | <1%                     |

**Table 5: Summary of information regarding pre-European and current vegetation extent of Vegetation Association 39 within the survey area**

| Factor                              | Value                                            |                                 |                                             |                                                         |                                  |
|-------------------------------------|--------------------------------------------------|---------------------------------|---------------------------------------------|---------------------------------------------------------|----------------------------------|
| Beard Vegetation Association*       | 39                                               |                                 |                                             |                                                         |                                  |
| Vegetation Association Description* | Shrublands; mulga scrub                          |                                 |                                             |                                                         |                                  |
| Pre-European Extent (ha)            | Scale                                            |                                 |                                             |                                                         |                                  |
|                                     | <b>By Association*</b><br>(WA)                   | <b>By Association**</b><br>(WA) | <b>By IBRA Region**</b><br><i>Murchison</i> | <b>By IBRA Sub-region**</b><br><i>Eastern Murchison</i> | <b>By LGA**</b><br>CUE, SHIRE OF |
|                                     | 4,856,768                                        | 6,613,567                       | 1,148,400                                   | 711,329                                                 | 135,321                          |
| % Pre-European Extent Remaining**   | 100.00%                                          | 99.83%                          | 99.10%                                      | 98.68%                                                  | 99.05%                           |
| Surrounding Land Use***             | Mining, Exploration, Prospecting, Pastoral Lease |                                 |                                             |                                                         |                                  |
| Weed prevalence***                  | Low                                              |                                 |                                             |                                                         |                                  |

\* Source: Shepherd *et al.* (2002) Appendix 2

\*\*Source: DBCA, (2019)

\*\*\*Source: Field Assessment

**Table 6: Summary of information regarding pre-European and current vegetation extent of Vegetation Association 18 within the survey area**

| Factor                              | Value                                            |                                 |                                      |                                                  |                                  |
|-------------------------------------|--------------------------------------------------|---------------------------------|--------------------------------------|--------------------------------------------------|----------------------------------|
| Beard Vegetation Association*       | 18                                               |                                 |                                      |                                                  |                                  |
| Vegetation Association Description* | Low woodland; mulga ( <i>Acacia aneura</i> )     |                                 |                                      |                                                  |                                  |
| Pre-European Extent (ha)            | Scale                                            |                                 |                                      |                                                  |                                  |
|                                     | <b>By Association*</b><br>(WA)                   | <b>By Association**</b><br>(WA) | <b>By IBRA Region**</b><br>Murchison | <b>By IBRA Sub-region**</b><br>Eastern Murchison | <b>By LGA**</b><br>CUE, SHIRE OF |
|                                     | 22,029,557                                       | 19,892,306                      | 12,403,172                           | 10,269,896                                       | 881,736                          |
| % Pre-European Extent Remaining**   | 100.00%                                          | 99.75%                          | 99.68%                               | 99.66%                                           | 99.67%                           |
| Surrounding Land Use***             | Mining, Exploration, Prospecting, Pastoral Lease |                                 |                                      |                                                  |                                  |
| Weed prevalence***                  | Low                                              |                                 |                                      |                                                  |                                  |

\* Source: Shepherd *et al.* (2002) Appendix 2

\*\*Source: DBCA, (2019)

\*\*\*Source: Field Assessment

#### 4.1.6 Wetlands

The DWER Clearing Permit System Map Viewer revealed no waterbodies within the survey area (DWER, 2026).

#### 4.1.7 Dieback

The survey area lies south of the 26<sup>th</sup> parallel, however receives average annual rainfall of 230.8 mm. There is no record of *Phytophthora cinnamomi* establishing in natural ecosystems in regions receiving less than 400mm rainfall per annum (CALM, 2003).

However, as indicated within the more recent Dieback guidelines (DBCA, 2020), other species of *Phytophthora* may persist east of the 400mm isohyet in unusually wet conditions. It is therefore recommended to conduct a risk assessment as per these guidelines.

Additionally, all measures should be taken to prevent any possible soil contamination (including seeds of non-native species *etc.*) which poses a risk in the survey area during seasonally favourable conditions.

## 4.2 Field Assessment

### 4.2.1 Threatened Flora

No Threatened Flora were recorded in the survey area.

One Priority Flora taxon, *Dodonaea amplisemina* (P4), was recorded at one location, with a total abundance of 30 plants.

**Table 7: Priority Flora recorded in the survey area**

| Taxon                            | Abundance | Vegetation Group Code | Observation Date | GDA2020 Longitude | GDA2020 Latitude |
|----------------------------------|-----------|-----------------------|------------------|-------------------|------------------|
| <i>Dodonaea amplisemina</i> (P4) | 30        | N                     | 26/06/2026       | -27.137863        | 118.25912        |

FloraBase returned 39 records of this taxon within a 300km radius of the survey area (WAHERB, 2026). The closest FloraBase record is located 45.3km southwest of the survey area. These records estimate the regional population to be approximately 10,579 from only 10 of these records. The remaining 29 records do not provide an estimate of abundance at their location.



**Figure 5: *Dodonaea amplisemina* (P4)**

#### 4.2.2 Vegetation Type, Extent and Status

A total of 31 Families, 58 Genera and 128 Species were recorded within the survey area. Fourteen major vegetation groups were recorded in the survey area and range from Completely Degraded to Very Good condition (using the scale of Keighery 1994, see Appendix 3). Existing disturbance within the survey area is comprised of historic exploration and mining activities, and access roads.

No unique or restricted vegetation communities were identified, and all vegetation types/communities are common, widespread and well represented in the Eastern Murchison subregion.

The summary of vegetation groups contained within the survey area is summarised in Table 8 below. Maps of the survey area can be seen in Appendix 4.

**Table 8: Vegetation Group Summary**

| <b>Vegetation Group</b>                                                                                            | <b>Veg Group Code</b> | <b>Families</b> | <b>Genera</b> | <b>Species</b> | <b>Area (ha)</b> | <b>Percentage of survey area (%)</b> |
|--------------------------------------------------------------------------------------------------------------------|-----------------------|-----------------|---------------|----------------|------------------|--------------------------------------|
| Mulga over low shrubland and tussock grass                                                                         | A                     | 14              | 21            | 45             | 384.39           | 39.58%                               |
| Mulga over low shrubland and tussock grass over rocky outcrops                                                     | B                     | 14              | 22            | 32             | 5.12             | 0.53%                                |
| Open Mulga Drainage Vegetation                                                                                     | C                     | 14              | 22            | 34             | 67.08            | 6.91%                                |
| Mulga thicket                                                                                                      | D                     | 11              | 16            | 24             | 5.05             | 0.52%                                |
| Mulga over <i>Eremophila forrestii</i> shrubland                                                                   | E                     | 9               | 9             | 22             | 17.14            | 1.76%                                |
| Mulga Creekline Vegetation                                                                                         | F                     | 9               | 16            | 27             | 54.86            | 5.65%                                |
| Mulga over <i>Acacia ramulosa</i> on rocky flats                                                                   | G                     | 13              | 16            | 26             | 99.83            | 10.28%                               |
| <i>Acacia quadrimarginea</i> shrubland                                                                             | H                     | 6               | 8             | 18             | 6.15             | 0.63%                                |
| Mulga over chenopod shrubland                                                                                      | I                     | 13              | 20            | 30             | 6.33             | 0.65%                                |
| Mulga over <i>Senna sp. Meekatharra</i> and <i>Eremophila forrestii</i> over low shrubland on rocky laterite hills | J                     | 13              | 15            | 32             | 100.60           | 10.36%                               |
| <i>Acacia quadrimarginea</i> shrubland on ironstone ridge                                                          | K                     | 8               | 11            | 21             | 31.38            | 3.23%                                |
| Mulga shrubland on laterite breakaways                                                                             | L                     | 8               | 9             | 11             | 2.22             | 0.23%                                |
| <i>Acacia</i> shrubland on granite outcropping                                                                     | M                     | 4               | 4             | 9              | 2.69             | 0.28%                                |
| Mulga over quartz outcrop                                                                                          | N                     | 11              | 12            | 24             | 17.10            | 1.76%                                |
| Existing Disturbance                                                                                               |                       | N/A             | N/A           | N/A            | 171.29           | 17.66%                               |
| <b>Total survey area</b>                                                                                           |                       | <b>31*</b>      | <b>58*</b>    | <b>128*</b>    | <b>971.21#</b>   | <b>100.00%#</b>                      |

Note: \* Within total survey area (not sum of column)

# Sum of column

The vegetation groups within the survey area are described in more detail below.

#### **4.2.2.1 Mulga over low shrubland and tussock grass (A)**

This Tree Mallee (Muir, 1977) consisted of 14 Families, 21 Genera and 45 Species. The vegetation group was approximately 384.39 ha which makes up 39.58% of the survey area.



**Figure 6: Vegetation Group A within the survey area**

#### **4.2.2.2 Mulga over low shrubland and tussock grass over rocky outcrops (B)**

This Tree Mallee (Muir, 1977) consisted of 14 Families, 22 Genera and 32 Species. The vegetation group was approximately 5.12 ha which makes up 0.53% of the survey area.



**Figure 7: Vegetation Group B within the survey area**

#### 4.2.2.3 Open Mulga Drainage Vegetation (C)

This Tree Mallee (Muir, 1977) consisted of 14 Families, 22 Genera and 34 Species. The vegetation group was approximately 67.08 ha which makes up 6.91% of the survey area.



**Figure 8: Vegetation Group C within the survey area**

#### **4.2.2.4 Mulga thicket (D)**

This Tree Mallee (Muir, 1977) consisted of 11 Families, 16 Genera and 24 Species. The vegetation group was approximately 5.05 ha which makes up 0.52% of the survey area.



**Figure 9: Vegetation Group D within the survey area**

#### 4.2.2.5 Mulga over *Eremophila forrestii* shrubland (E)

This Tree Mallee (Muir, 1977) consisted of nine Families, nine Genera and 22 Species. The vegetation group was approximately 17.14 ha which makes up 1.76% of the survey area.



**Figure 10: Vegetation Group E within the survey area**

#### 4.2.2.6 Mulga Creekline Vegetation (F)

This Tree Mallee (Muir, 1977) consisted of nine Families, 16 Genera and 27 Species. The vegetation group was approximately 54.86 ha which makes up 5.65% of the survey area.



**Figure 11: Vegetation Group F within the survey area**

#### 4.2.2.7 Mulga over *Acacia ramulosa* on rocky flats (G)

This Tree Mallee (Muir, 1977) consisted of 13 Families, 16 Genera and 26 Species. The vegetation group was approximately 99.83 ha which makes up 10.28% of the survey area.



Figure 12: Vegetation Group G within the survey area

#### 4.2.2.8 *Acacia quadrimarginea* shrubland (H)

This Tree Mallee (Muir, 1977) consisted of six Families, eight Genera and 18 Species. The vegetation group was approximately 6.15 ha which makes up 0.63% of the survey area.



**Figure 13: Vegetation Group H within the survey area**

#### **4.2.2.9 Mulga over chenopod shrubland (I)**

This Tree Mallee (Muir, 1977) consisted of 13 Families, 20 Genera and 30 Species. The vegetation group was approximately 6.33 ha which makes up 0.65% of the survey area.



**Figure 14: Vegetation Group I within the survey area**

#### 4.2.2.10 Mulga over *Senna* sp. Meekatharra and *Eremophila forrestii* over low shrubland on rocky laterite hills (J)

This Tree Mallee (Muir, 1977) consisted of 13 Families, 15 Genera and 32 Species. The vegetation group was approximately 100.60 ha which makes up 10.36% of the survey area.



Figure 15: Vegetation Group J within the survey area

#### 4.2.2.11 *Acacia quadrimarginea* shrubland on ironstone ridge (K)

This Tree Mallee (Muir, 1977) consisted of eight Families, 11 Genera and 21 Species. The vegetation group was approximately 31.38 ha which makes up 3.23% of the survey area.



**Figure 16: Vegetation Group K within the survey area**

#### **4.2.2.12 Mulga shrubland on laterite breakaways (L)**

This Tree Mallee (Muir, 1977) consisted of eight Families, nine Genera and 11 Species. The vegetation group was approximately 2.22 ha which makes up 0.23% of the survey area.



**Figure 17: Vegetation Group L within the survey area**

#### 4.2.2.13 *Acacia* shrubland on granite outcropping (M)

This Tree Mallee (Muir, 1977) consisted of four Families, four Genera and nine Species. The vegetation group was approximately 2.69 ha which makes up 0.28% of the survey area.



**Figure 18: Vegetation Group M within the survey area**

#### **4.2.2.14 Mulga over quartz outcrop (N)**

This Tree Mallee (Muir, 1977) consisted of 11 Families, 12 Genera and 24 Species. The vegetation group was approximately 17.10 ha which makes up 1.76% of the survey area.



**Figure 19: Vegetation Group N within the survey area**

#### 4.2.2.15 Existing Disturbance

Existing disturbance within the survey area consisted of historic exploration/mining, access roads and was approximately 171.29 ha which makes up 17.64% of the survey area.



**Figure 20: Existing disturbance within the survey area**

### 4.2.3 Weeds

Three weed species were recorded within the survey area, *Carrichtera annua*, *Cucumis myriocarpus* and *Rumex vesicarius*. None of these species are considered a Declared Plant under the *BAM Act* (DPIRD, 2026). Locations and abundances of weeds recorded in the survey area are listed in Table 9 below.

**Table 9: Weed Abundance recorded in the survey area**

| <b>Taxon</b>                 | <b>Waypoint</b> | <b>Veg Code</b> | <b>Abundance</b> | <b>GDA2020 Longitude</b> | <b>GDA2020 Latitude</b> |
|------------------------------|-----------------|-----------------|------------------|--------------------------|-------------------------|
| <i>Carrichtera annua</i> *   | wpt002          | B               | 20               | 118.298                  | -27.084                 |
| <i>Carrichtera annua</i> *   | wpt004          | B               | 20               | 118.297307               | -27.0861                |
| <i>Carrichtera annua</i> *   | wpt006          | B               | 20               | 118.296392               | -27.084                 |
| <i>Carrichtera annua</i> *   | wpt007          | C               | 20               | 118.294693               | -27.0902                |
| <i>Cucumis myriocarpus</i> * | wpt007          | C               | 10               | 118.294693               | -27.0902                |
| <i>Carrichtera annua</i> *   | wpt009          | C               | 20               | 118.293867               | -27.0954                |
| <i>Cucumis myriocarpus</i> * | wpt009          | C               | 10               | 118.293867               | -27.0954                |
| <i>Carrichtera annua</i> *   | wpt010          | C               | 20               | 118.29327                | -27.0897                |
| <i>Carrichtera annua</i> *   | wpt012          | C               | 20               | 118.292715               | -27.0816                |
| <i>Carrichtera annua</i> *   | wpt013          | C               | 20               | 118.291615               | -27.0896                |
| <i>Carrichtera annua</i> *   | wpt014          | C               | 20               | 118.291645               | -27.0986                |
| <i>Carrichtera annua</i> *   | wpt016          | C               | 20               | 118.290208               | -27.0977                |
| <i>Carrichtera annua</i> *   | wpt018          | C               | 20               | 118.290112               | -27.0881                |
| <i>Carrichtera annua</i> *   | wpt020          | C               | 20               | 118.2899                 | -27.0819                |
| <i>Carrichtera annua</i> *   | wpt021          | D               | 20               | 118.28805                | -27.0835                |
| <i>Carrichtera annua</i> *   | wpt022          | C               | 20               | 118.288676               | -27.096                 |
| <i>Carrichtera annua</i> *   | wpt028          | E               | 20               | 118.285647               | -27.1099                |
| <i>Carrichtera annua</i> *   | wpt031          | E               | 20               | 118.285411               | -27.079                 |
| <i>Rumex vesicarius</i> *    | wpt035          | A               | 50               | 118.281357               | -27.0923                |
| <i>Rumex vesicarius</i> *    | wpt039          | F               | 20               | 118.276153               | -27.0967                |
| <i>Rumex vesicarius</i> *    | wpt042          | F               | 20               | 118.278642               | -27.1088                |
| <i>Rumex vesicarius</i> *    | wpt044          | A               | 200              | 118.280728               | -27.1075                |
| <i>Carrichtera annua</i> *   | wpt047          | G               | 20               | 118.28124                | -27.119                 |
| <i>Rumex vesicarius</i> *    | wpt047          | G               | 2                | 118.28124                | -27.119                 |
| <i>Carrichtera annua</i> *   | wpt048          | D               | 20               | 118.282905               | -27.1179                |
| <i>Carrichtera annua</i> *   | wpt050          | D               | 20               | 118.282832               | -27.11                  |
| <i>Carrichtera annua</i> *   | wpt055          | C               | 20               | 118.271975               | -27.1138                |
| <i>Carrichtera annua</i> *   | wpt056          | G               | 20               | 118.272242               | -27.1081                |
| <i>Carrichtera annua</i> *   | wpt058          | G               | 20               | 118.271737               | -27.1034                |
| <i>Cucumis myriocarpus</i> * | wpt062          | I               | 6                | 118.269313               | -27.1135                |
| <i>Carrichtera annua</i> *   | wpt067          | G               | 20               | 118.279863               | -27.1188                |
| <i>Carrichtera annua</i> *   | wpt068          | E               | 20               | 118.27651                | -27.128                 |
| <i>Carrichtera annua</i> *   | wpt070          | G               | 20               | 118.276412               | -27.1306                |
| <i>Rumex vesicarius</i> *    | wpt076          | G               | 150              | 118.273902               | -27.1271                |
| <i>Carrichtera annua</i> *   | wpt076          | G               | 20               | 118.273902               | -27.1271                |

#### **4.2.4 Vegetation Condition**

Evidence of historic exploration and access tracks was observed during the field assessment.

Overall, the condition of the vegetation was determined to range from “Completely Degraded” to “Very Good” with most of the area falling into the “Good” Category. Areas which were affected by historic exploration and clearing were deemed in “Degraded” or “Completely Degraded” condition. A map of the vegetation condition within the survey is depicted in Appendix 4.

#### 4.2.5 Clearing Principals

The *Environmental Protection Act (1986)* outlines 10 principles that are to be used in the assessment of native vegetation clearing permit applications which are also applicable for other assessments and approvals (Table 10). Where possible, native vegetation should not be cleared if any of the following principles are compromised.

**Table 10: Assessment of impact using the native vegetation clearing principles**

| Principle                                                                                                                                     | Response                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| It comprises a high level of biological diversity.                                                                                            | A total of 31 Families, 58 Genera and 128 Flora Taxon were recorded within the survey area. Fourteen major vegetation groups were recorded in the survey area and range from Completely Degraded to Very Good condition.<br><br>No unique or restricted vegetation communities were identified, and all vegetation types/communities are common, widespread and well represented in the Eastern Murchison subregion and adjoining bioregions.                                              |
| It comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia. | N/A- not assessed in this report                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| It includes, or is necessary for the continued existence of, rare flora.                                                                      | No Threatened Flora were recorded in the survey area. One Priority Flora taxon, <i>Dodonaea amplisemina</i> (P4), was recorded at one location, with a total abundance of 30 plants. Wider searches within this vegetation group did not record any further populations. The recorded 30 plants in the survey area represents 0.28% of the regional population. Removal of this taxon from the survey area is not considered a significant impact to the conservation value of this taxon. |
| It comprises the whole or a part of, or is necessary for the maintenance of, a threatened ecological community.                               | The area does not contain any threatened ecological communities.                                                                                                                                                                                                                                                                                                                                                                                                                           |
| It is significant as a remnant of native vegetation in an area that has been extensively cleared.                                             | The area is not a remnant.                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| It is growing in, or in association with, an environment associated with a watercourses or wetland.                                           | The area does not contain a natural wetland or salt lake. Broad ephemeral creeklines are present in the survey area, which provide short term runoff drainage from adjacent areas.                                                                                                                                                                                                                                                                                                         |
| The clearing of the vegetation is likely to cause appreciable land degradation.                                                               | N/A- not assessed in this report                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| The clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.           | Clearing of vegetation is unlikely to impact on the environmental values of the bioregion, or any nearby conservation areas.                                                                                                                                                                                                                                                                                                                                                               |
| The clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.                               | Elevations within the project development area range from 423 to 536mAHD. There are no major water bodies such as rivers or lakes within the Project area, with relatively minor creek/ drainage lines occurring in parts. All creeks and drainages within the survey area are ephemeral in nature, only carrying runoff following heavy rainfall events.                                                                                                                                  |
| The clearing of the vegetation is likely to cause or exacerbate the incidence of flooding.                                                    | N/A- not assessed in this report                                                                                                                                                                                                                                                                                                                                                                                                                                                           |

## 5. DISCUSSION

The field assessment established that the condition of the vegetation in the proposed disturbance area ranged from “Completely Degraded” to “Very Good” with most of the area falling into the “Good” category. Areas which were affected by historic exploration were deemed in “Degraded” or “Completely Degraded” condition. No areas of vegetation were assessed to be in “Pristine” condition.

Three weed species were recorded within the survey area, *Carrichtera annua*, *Cucumis myriocarpus* and *Rumex vesicarius*. None of these species are considered a Declared Plant under the *BAM Act* (DPIRD, 2026).

No Threatened Flora were recorded in the survey area.

One Priority Flora taxon, *Dodonaea amplisemina* (P4), was recorded at one location, with a total abundance of 30 plants. Wider searches within this vegetation group did not record any further populations. FloraBase returned 39 records of this taxon within a 300km radius of the survey area (WAHERB, 2026). These records estimate the regional population to be approximately 10,579 from only 10 of these records.

The removal of 30 plants from the survey area represents 0.28% of the regional population and is not considered a significant impact to the conservation value of this taxon.

No PECs or TECs were recorded in the survey area.

No unique or restricted vegetation communities were identified, and all vegetation types/communities are common, widespread and well represented in the Eastern Murchison subregion.

Any proposed disturbance/clearing of vegetation will result in a loss of some plants and vegetation. However, given the size of the area and the extent of the Beard (1990) vegetation association elsewhere, the impact on the vegetation and its component flora will not affect the conservation values of either or create fragmentation or patches of remnant vegetation.

The following recommendations arise from the reconnaissance flora survey:

- Locations of Priority Flora should be avoided, where practicable, when planning activities of disturbance;
- Weed control measures should be implemented during and following earthworks; and
- Dust control measures should be implemented during earthworks.

## 6. REFERENCES

Beard, J.S., (1990), *Plant Life of Western Australia*, Kangaroo Press Pty Ltd, NSW

BOM, (2026), *Climate Data Online*, Bureau of Meteorology,

<http://www.bom.gov.au/climate/averages/>

Accessed: 25/07/2026

CALM, (2002), *A Biodiversity Audit of Western Australia's 53 Biogeographical Subregions in 2002-Murchison (MUR01 Eastern Murchison synopses)*, Department of Conservation and Land Management

CALM, (2003), *Phytophthora cinnamomi and Diseases Caused by It, Volume 1-Management Guidelines*, Department of Conservation and Land Management

<https://library.dbca.wa.gov.au/FullTextFiles/021873.pdf>

Accessed: 25/07/2026

DBCA, (2019), *2018 Statewide Vegetation Statistics incorporating the CAR Reserve Analysis (Full Report)- Current as of March 2019*, WA Department of Biodiversity, Conservation and Attractions, Perth

<https://catalogue.data.wa.gov.au/dataset/dbca-statewide-vegetation-statistics>

Accessed: 25/07/2026

DBCA, (2020), *Phytophthora Dieback Management Manual, October 2020*, Department of Biodiversity, Conservation and Attractions

DBCA, (2026), *TEC/PEC Database Results Ref:60\_0726EC*, Department of Biodiversity, Conservation and Attractions

DBCA, (2026a), *Threatened Flora Database Results Ref: 61\_0726FL*, Department of Biodiversity, Conservation and Attractions

DBCA, (2023) *Conservation Codes for Western Australian Flora and Fauna*. Department of Biodiversity, Conservation and Attractions, Western Australia, August 2023

DCCEEW, (2026), *Interim Biogeographic Regionalisation for Australia (IBRA) Version 7.1 (Subregions)*, Updated 26 February 2026, Department of Climate Change, Energy, the Environment and Water, Australian Government

<https://www.environment.gov.au/land/nrs/science/ibra>

Accessed: 25/07/2026

DCCEEW (2026a), *Protected Matters Search Tool*, Department of Climate Change, Energy, the Environment and Water, Australian Government

<http://www.environment.gov.au/epbc/protected-matters-search-tool>

Accessed: 25/07/2026

DPIRD, (2017), *NRInfo Digital Mapping*, Department of Primary Industries and Regional Development

<https://maps.agric.wa.gov.au/nrm-info/>

Accessed: 25/07/2026

DPIRD, (2026), *Declared Plants Database*, Department of Primary Industries and Regional Development, Western Australia

<https://www.agric.wa.gov.au/pests-weeds-diseases/weeds/declared-plants>

Accessed: 25/07/2026

DWER, (2026), *Clearing Permit System Map Viewer*, Department of Water and Environmental Regulation  
<https://cps.dwer.wa.gov.au/main.html>  
Accessed: 25/07/2026

EPA, (2016), *Environmental Factor Guideline: Flora and Vegetation*, Environmental Protection Authority, Western Australia

EPA, (2016a), *Technical Guidance- Flora and Vegetation Surveys for Environmental Impact Assessment*, Environmental Protection Authority, Western Australia

EPA, (2023), *Statement of environmental principles, factors, objectives and aims of EIA*, Environmental Protection Authority, Western Australia

Keighery, B.J., (1994), *Bushland Plant Survey; A guide to plant community survey for the Community*, Wildflower Society of Western Australia (Inc.) Nedlands

Lawn, G., (2024), *Biodiversity Conservation (Listing of Native Species) (Flora) Order 2024*, published in the Gazette on 30 April 2024 at pages 1135 to 1140, Western Australian Government

Muir, B.G., (1977), *Biological Survey of the Western Australian Wheatbelt. Pt. 2. Vegetation and habitat of the Bendering Reserve*. Records of the Western Australian Museum Supplement 3

MWH, (2015), *CMGP Reedy Project Dewatering Program - Level 1 Flora and Fauna Survey- April 2015*, Unpublished Report Prepared for Metals X Ltd

MWH, (2016), *Culculli and Paddy Italiano Level 1 Flora, Vegetation and Fauna Assessment- April 2016*, Unpublished Report Prepared for Metals X Ltd

Shepherd, D.P., Beeston, G.R. and Hopkins, A.J.M., (2002), *Land-Use and Vegetation in Western Australia- National Land and Water Resources Audit Report*, Technical Report 250, Department of Agriculture Western Australia

Stantec, (2022), *Boomerang Open Pit Dewatering Discharge Pipeline Level 1 Flora and Fauna Study- November 2021*, Unpublished Report Prepared for Westgold Resources Ltd

WAHERB, (2026), *Florabase- the Western Australian Flora*, Department of Parks and Wildlife  
<http://florabase.dpaw.wa.gov.au/>  
Accessed 25/07/2026

## 7. GLOSSARY

### Acronyms:

|                 |                                                                                                                           |
|-----------------|---------------------------------------------------------------------------------------------------------------------------|
| <b>BOM</b>      | Bureau of Meteorology, Australian Government                                                                              |
| <b>BSc</b>      | Bachelor of Science                                                                                                       |
| <b>CALM</b>     | Department of Conservation and Land Management (now DBCA)                                                                 |
| <b>CPS</b>      | Clearing Permit System (DWER)                                                                                             |
| <b>DBCA</b>     | Department of Biodiversity, Conservation and Attractions, Western Australia                                               |
| <b>DCCEEW</b>   | Department of Climate Control, Energy, the Environment and Water, Australian Government                                   |
| <b>DMPE</b>     | Department of Mines Petroleum and Exploration, Western Australia                                                          |
| <b>DPAW</b>     | Department of Parks and Wildlife, Western Australia (now DBCA)                                                            |
| <b>DPIRD</b>    | Department of Primary Industries and Regional Development, Western Australia                                              |
| <b>DRF</b>      | Declared Rare Flora (now classed as Threatened Flora)                                                                     |
| <b>DWER</b>     | Department of Water and Environmental Regulation, Western Australia                                                       |
| <b>EPA</b>      | Environmental Protection Authority, Western Australia                                                                     |
| <b>EP Act</b>   | Environmental Protection Act 1986, Western Australia                                                                      |
| <b>EPBC Act</b> | Environment Protection and Biodiversity Conservation Act 1999 (Commonwealth Act)                                          |
| <b>ESA</b>      | Environmentally Sensitive Area                                                                                            |
| <b>GIS</b>      | Geographical Information System                                                                                           |
| <b>ha</b>       | Hectare (10,000 square metres)                                                                                            |
| <b>IBRA</b>     | Interim Biogeographic Regionalisation for Australia, DCCEEW                                                               |
| <b>IUCN</b>     | International Union for the Conservation of Nature and Natural Resources – commonly known as the World Conservation Union |
| <b>km</b>       | Kilometres                                                                                                                |
| <b>m</b>        | Metres                                                                                                                    |
| <b>MUR</b>      | Murchison Bioregion (IBRA)                                                                                                |
| <b>MUR01</b>    | Eastern Murchison Subregion (IBRA)                                                                                        |
| <b>NVS</b>      | Native Vegetation Solutions                                                                                               |
| <b>PEC</b>      | Priority Ecological Community, Western Australia                                                                          |
| <b>PF</b>       | Priority Flora                                                                                                            |
| <b>Ramsar</b>   | A wetland site designated of international importance under the Ramsar Convention (UNESCO)                                |
| <b>TEC</b>      | Threatened Ecological Community                                                                                           |
| <b>TF</b>       | Threatened Flora                                                                                                          |
| <b>UNESCO</b>   | United Nations Educational, Scientific and Cultural Organization                                                          |
| <b>WA</b>       | Western Australia                                                                                                         |
| <b>WAHERB</b>   | Western Australian Herbarium (DBCA)                                                                                       |
| <b>WAOL</b>     | Western Australian Organism List (DPIRD)                                                                                  |

## Definitions:

**DBCA (2023) Conservation Codes for Western Australian Flora and Fauna. Department of Biodiversity, Conservation and Attractions, Western Australia, August 2023: -**

### **T Threatened species:**

Listed by order of the Minister as Threatened in the category of critically endangered, endangered or vulnerable under section 19(1), or is a rediscovered species to be regarded as threatened species under section 26(2) of the *Biodiversity Conservation Act 2016* (BC Act).

**Threatened fauna** is that subset of ‘Specially Protected Fauna’ listed under schedules 1 to 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for Threatened Fauna.

**Threatened flora** is that subset of ‘Rare Flora’ listed under schedules 1 to 3 of the *Wildlife Conservation (Rare Flora) Notice 2018* for Threatened Flora.

The assessment of the conservation status of these species is based on their national extent and ranked according to their level of threat using IUCN Red List categories and criteria as detailed below.

### **CR Critically endangered species**

Threatened species considered to be “facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with criteria set out in the ministerial guidelines”.

Listed as critically endangered under section 19(1)(a) of the BC Act in accordance with the criteria set out in section 20 and the ministerial guidelines. Published under schedule 1 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for critically endangered fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for critically endangered flora.

### **EN Endangered species**

Threatened species considered to be “facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines”.

Listed as endangered under section 19(1)(b) of the BC Act in accordance with the criteria set out in section 21 and the ministerial guidelines. Published under schedule 2 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for endangered fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for endangered flora.

### **VU Vulnerable species**

Threatened species considered to be “facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with criteria set out in the ministerial guidelines”.

Listed as vulnerable under section 19(1)(c) of the BC Act in accordance with the criteria set out in section 22 and the ministerial guidelines. Published under schedule 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for vulnerable fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for vulnerable flora.

**Extinct species:**

Listed by order of the Minister as extinct under section 23(1) of the BC Act as extinct or extinct in the wild.

**EX Extinct species**

Species where “*there is no reasonable doubt that the last member of the species has died*”, and listing is otherwise in accordance with the ministerial guidelines (section 24 of the BC Act).

Published as presumed extinct under schedule 4 of the Wildlife Conservation (Specially Protected Fauna) Notice 2018 for extinct fauna or the Wildlife Conservation (Rare Flora) Notice 2018 for extinct flora.

**EW Extinct in the wild species**

Species that “is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; and it has not been recorded in its known habitat or expected habitat, at appropriate seasons, anywhere in its past range, despite surveys over a time frame appropriate to its life cycle and form”, and listing is otherwise in accordance with the ministerial guidelines (section 25 of the BC Act).

Currently there are no threatened fauna or threatened flora species listed as extinct in the wild. If listing of a species as extinct in the wild occurs, then a schedule will be added to the applicable notice.

**Specially protected species**

Listed by order of the Minister as specially protected under section 13(1) of the BC Act. Meeting one or more of the following categories: species of special conservation interest; migratory species; cetaceans; species subject to international agreement; or species otherwise in need of special protection.

Species that are listed as threatened species (critically endangered, endangered or vulnerable) or extinct species under the BC Act cannot also be listed as Specially Protected species.

**MI Migratory species**

Fauna that periodically or occasionally visit Australia or an external Territory or the exclusive economic zone; or the species is subject of an international agreement that relates to the protection of migratory species and that binds the Commonwealth; and listing is otherwise in accordance with the ministerial guidelines (section 15 of the BC Act).

Includes birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and fauna subject to the *Convention on the Conservation of Migratory Species of Wild Animals* (Bonn Convention), an environmental treaty under the United Nations Environment Program. Migratory species listed under the BC Act are a subset of the migratory animals, that are known to visit Western Australia, protected under the international agreements or treaties, excluding species that are listed as Threatened species.

Published as migratory birds protected under an international agreement under schedule 5 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

### **CD Species of special conservation interest (conservation dependent fauna)**

Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened, and listing is otherwise in accordance with the ministerial guidelines (section 14 of the BC Act).

Published as conservation dependent fauna under schedule 6 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

### **OS Other specially protected species**

Fauna otherwise in need of special protection to ensure their conservation, and listing is otherwise in accordance with the ministerial guidelines (section 18 of the BC Act).

Published as other specially protected fauna under schedule 7 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

### **P Priority Species**

Priority is not a listing category under the BC Act. The Priority Flora and Fauna lists are maintained by the department and are published on the department's website.

All fauna and flora are protected in WA following the provisions in Part 10 of the BC Act. The protection applies even when a species is not listed as threatened or specially protected, and regardless of land tenure (State managed land (Crown land), private land, or Commonwealth land).

Species that may possibly be threatened species that do not meet the criteria for listing under the BC Act because of insufficient survey or are otherwise data deficient, are added to the Priority Fauna or Priority Flora Lists under Priorities 1, 2 or 3. These three categories are ranked in order of prioritisation for survey and evaluation of conservation status so that consideration can be given to potential listing as threatened.

Species that are adequately known, meet criteria for near threatened, or are rare but not threatened, or that have been recently removed from the threatened species list or conservation dependent or other specially protected fauna lists for other than taxonomic reasons, are placed in Priority 4. These species require regular monitoring.

Assessment of priority status is based on the Western Australian distribution of the species, unless the distribution in WA is part of a contiguous population extending into adjacent States, as defined by the known spread of locations.

**Priority 1: Poorly-known species**

Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.

**Priority 2: Poorly-known species**

Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.

**Priority 3: Poorly-known species**

Species that are known from several locations and the species does not appear to be under imminent threat or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. These species need further survey.

**Priority 4: Rare, Near Threatened and other species in need of monitoring**

- a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection but could be if present circumstances change. These species are usually represented on conservation lands.
- b) Near Threatened. Species that are considered to have been adequately surveyed and that are close to qualifying for vulnerable but are not listed as a conservation dependent specially protected species.
- c) Species that have been removed from the list of threatened species or lists of conservation dependent or other specially protected species, during the past five years for reasons other than taxonomy.
- d) Other species in need of monitoring.

## **Appendix 1: Relevant Government Database Search Results**



Australian Government

Department of Climate Change, Energy,  
the Environment and Water

## EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected. Please see the caveat for interpretation of information provided here.

Report created: 25-Jul-2026

[Summary](#)

[Details](#)

[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



## Summary

### Matters of National Environment Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

|                                                               |      |
|---------------------------------------------------------------|------|
| <a href="#">World Heritage Properties:</a>                    | None |
| <a href="#">National Heritage Places:</a>                     | None |
| <a href="#">Wetlands of International Importance (Ramsar)</a> | None |
| <a href="#">Great Barrier Reef Marine Park:</a>               | None |
| <a href="#">Commonwealth Marine Area:</a>                     | None |
| <a href="#">Listed Threatened Ecological Communities:</a>     | None |
| <a href="#">Listed Threatened Species:</a>                    | 9    |
| <a href="#">Listed Migratory Species:</a>                     | 8    |

### Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <https://www.dcceew.gov.au/parks-heritage/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

|                                                                     |      |
|---------------------------------------------------------------------|------|
| <a href="#">Commonwealth Lands:</a>                                 | None |
| <a href="#">Commonwealth Heritage Places:</a>                       | None |
| <a href="#">Listed Marine Species:</a>                              | 10   |
| <a href="#">Whales and Other Cetaceans:</a>                         | None |
| <a href="#">Critical Habitats:</a>                                  | None |
| <a href="#">Commonwealth Reserves Terrestrial:</a>                  | None |
| <a href="#">Australian Marine Parks:</a>                            | None |
| <a href="#">Habitat Critical to the Survival of Marine Turtles:</a> | None |

### Extra Information

This part of the report provides information that may also be relevant to the area you have

|                                                         |      |
|---------------------------------------------------------|------|
| <a href="#">State and Territory Reserves:</a>           | None |
| <a href="#">Regional Forest Agreements:</a>             | None |
| <a href="#">Nationally Important Wetlands:</a>          | None |
| <a href="#">EPBC Act Referrals:</a>                     | 1    |
| <a href="#">Key Ecological Features (Marine):</a>       | None |
| <a href="#">Biologically Important Areas:</a>           | None |
| <a href="#">Bioregional Assessments:</a>                | None |
| <a href="#">Geological and Bioregional Assessments:</a> | None |

## Details

### Matters of National Environmental Significance

| Listed Threatened Species                                                                                       |                       |                                                        | [ Resource Information ] |
|-----------------------------------------------------------------------------------------------------------------|-----------------------|--------------------------------------------------------|--------------------------|
| Status of Conservation Dependent and Extinct are not MNES under the EPBC Act.<br>Number is the current name ID. |                       |                                                        |                          |
| Scientific Name                                                                                                 | Threatened Category   | Presence Text                                          | Buffer Status            |
| <b>BIRD</b>                                                                                                     |                       |                                                        |                          |
| <a href="#">Aphelocephala leucopsis</a><br>Southern Whiteface [529]                                             | Vulnerable            | Species or species habitat likely to occur within area | In feature area          |
| <a href="#">Calidris acuminata</a><br>Sharp-tailed Sandpiper [874]                                              | Vulnerable            | Species or species habitat may occur within area       | In feature area          |
| <a href="#">Calidris ferruginea</a><br>Curlew Sandpiper [856]                                                   | Critically Endangered | Species or species habitat likely to occur within area | In feature area          |
| <a href="#">Falco hypoleucos</a><br>Grey Falcon [929]                                                           | Vulnerable            | Species or species habitat may occur within area       | In feature area          |
| <a href="#">Leipoa ocellata</a><br>Malleefowl [934]                                                             | Vulnerable            | Species or species habitat likely to occur within area | In feature area          |
| <a href="#">Pezoporus occidentalis</a><br>Night Parrot [59350]                                                  | Critically Endangered | Species or species habitat may occur within area       | In feature area          |
| <a href="#">Tringa nebularia</a><br>Common Greenshank, Greenshank [832]                                         | Endangered            | Species or species habitat may occur within area       | In buffer area only      |
| <b>REPTILE</b>                                                                                                  |                       |                                                        |                          |
| <a href="#">Egernia stokesii badia</a><br>Western Spiny-tailed Skink, Baudin Island Spiny-tailed Skink [64483]  | Endangered            | Species or species habitat may occur within area       | In buffer area only      |

| Scientific Name                                                        | Threatened Category | Presence Text                                                | Buffer Status   |
|------------------------------------------------------------------------|---------------------|--------------------------------------------------------------|-----------------|
| <b>SPIDER</b>                                                          |                     |                                                              |                 |
| <a href="#">Idiosoma nigrum</a>                                        |                     |                                                              |                 |
| Shield-backed Trapdoor Spider, Black<br>Rugose Trapdoor Spider [66798] | Vulnerable          | Species or species<br>habitat likely to occur<br>within area | In feature area |

#### Listed Migratory Species [ Resource Information ]

| Scientific Name                | Threatened Category | Presence Text                                                | Buffer Status   |
|--------------------------------|---------------------|--------------------------------------------------------------|-----------------|
| <b>Migratory Marine Birds</b>  |                     |                                                              |                 |
| <a href="#">Apus pacificus</a> |                     |                                                              |                 |
| Fork-tailed Swift [678]        |                     | Species or species<br>habitat likely to occur<br>within area | In feature area |

#### Migratory Terrestrial Species

|                                   |  |                                                        |                 |
|-----------------------------------|--|--------------------------------------------------------|-----------------|
| <a href="#">Motacilla cinerea</a> |  |                                                        |                 |
| Grey Wagtail [642]                |  | Species or species<br>habitat may occur<br>within area | In feature area |
| <a href="#">Motacilla flava</a>   |  |                                                        |                 |
| Yellow Wagtail [644]              |  | Species or species<br>habitat may occur<br>within area | In feature area |

#### Migratory Wetlands Species

|                                        |                       |                                                              |                     |
|----------------------------------------|-----------------------|--------------------------------------------------------------|---------------------|
| <a href="#">Actitis hypoleucos</a>     |                       |                                                              |                     |
| Common Sandpiper [59309]               |                       | Species or species<br>habitat may occur<br>within area       | In feature area     |
| <a href="#">Calidris acuminata</a>     |                       |                                                              |                     |
| Sharp-tailed Sandpiper [874]           | Vulnerable            | Species or species<br>habitat may occur<br>within area       | In feature area     |
| <a href="#">Calidris ferruginea</a>    |                       |                                                              |                     |
| Curlew Sandpiper [856]                 | Critically Endangered | Species or species<br>habitat likely to occur<br>within area | In feature area     |
| <a href="#">Calidris melanotos</a>     |                       |                                                              |                     |
| Pectoral Sandpiper [858]               |                       | Species or species<br>habitat may occur<br>within area       | In feature area     |
| <a href="#">Tringa nebularia</a>       |                       |                                                              |                     |
| Common Greenshank, Greenshank<br>[832] | Endangered            | Species or species<br>habitat may occur<br>within area       | In buffer area only |

## Other Matters Protected by the EPBC Act

| Listed Marine Species                                                                     |                       | [ Resource Information ]                                                   |                 |
|-------------------------------------------------------------------------------------------|-----------------------|----------------------------------------------------------------------------|-----------------|
| Scientific Name                                                                           | Threatened Category   | Presence Text                                                              | Buffer Status   |
| <b>Bird</b>                                                                               |                       |                                                                            |                 |
| <a href="#">Actitis hypoleucos</a><br>Common Sandpiper [59309]                            |                       | Species or species habitat may occur within area                           | In feature area |
| <a href="#">Apus pacificus</a><br>Fork-tailed Swift [678]                                 |                       | Species or species habitat likely to occur within area overfly marine area | In feature area |
| <a href="#">Calidris acuminata</a><br>Sharp-tailed Sandpiper [874]                        | Vulnerable            | Species or species habitat may occur within area                           | In feature area |
| <a href="#">Calidris ferruginea</a><br>Curlew Sandpiper [856]                             | Critically Endangered | Species or species habitat likely to occur within area overfly marine area | In feature area |
| <a href="#">Calidris melanotos</a><br>Pectoral Sandpiper [858]                            |                       | Species or species habitat may occur within area overfly marine area       | In feature area |
| <a href="#">Chalcites osculans as Chrysococcyx osculans</a><br>Black-eared Cuckoo [83425] |                       | Species or species habitat likely to occur within area overfly marine area | In feature area |
| <a href="#">Merops ornatus</a><br>Rainbow Bee-eater [670]                                 |                       | Species or species habitat may occur within area overfly marine area       | In feature area |
| <a href="#">Motacilla cinerea</a><br>Grey Wagtail [642]                                   |                       | Species or species habitat may occur within area overfly marine area       | In feature area |

| Scientific Name                                                         | Threatened Category | Presence Text                                                        | Buffer Status       |
|-------------------------------------------------------------------------|---------------------|----------------------------------------------------------------------|---------------------|
| <a href="#">Motacilla flava</a><br>Yellow Wagtail [644]                 |                     | Species or species habitat may occur within area overfly marine area | In feature area     |
| <a href="#">Tringa nebularia</a><br>Common Greenshank, Greenshank [832] | Endangered          | Species or species habitat may occur within area overfly marine area | In buffer area only |

## Extra Information

| EPBC Act Referrals                                                                                           |           |                       | [ Resource Information ] |                 |
|--------------------------------------------------------------------------------------------------------------|-----------|-----------------------|--------------------------|-----------------|
| Title of referral                                                                                            | Reference | Referral Outcome      | Assessment Status        | Buffer Status   |
| Not controlled action                                                                                        |           |                       |                          |                 |
| <a href="#">Improving rabbit biocontrol: releasing another strain of RHDV, sthrn two thirds of Australia</a> | 2015/7522 | Not Controlled Action | Completed                | In feature area |

# Caveat

## 1 PURPOSE

This report is designed to assist in identifying the location of matters of national environmental significance (MNES) and other matters protected by the Environment Protection and Biodiversity Conservation Act 1999 (Cth) (EPBC Act) which may be relevant in determining obligations and requirements under the EPBC Act.

The report contains the mapped locations of:

- World and National Heritage properties;
- Wetlands of International and National Importance;
- Commonwealth and State/Territory reserves;
- distribution of listed threatened, migratory and marine species;
- listed threatened ecological communities; and
- other information that may be useful as an indicator of potential habitat value.

## 2 DISCLAIMER

This report is not intended to be exhaustive and should only be relied upon as a general guide as mapped data is not available for all species or ecological communities listed under the EPBC Act (see below). Persons seeking to use the information contained in this report to inform the referral of a proposed action under the EPBC Act should consider the limitations noted below and whether additional information is required to determine the existence and location of MNES and other protected matters.

Where data is available to inform the mapping of protected species, the presence type (e.g. known, likely or may occur) that can be determined from the data is indicated in general terms. It is the responsibility of any person using or relying on the information in this report to ensure that it is suitable for the circumstances of any proposed use. The Commonwealth cannot accept responsibility for the consequences of any use of the report or any part thereof. To the maximum extent allowed under governing law, the Commonwealth will not be liable for any loss or damage that may be occasioned directly or indirectly through the use of, or reliance on the contents of this report.

## 3 DATA SOURCES

### Threatened ecological communities

For threatened ecological communities where the distribution is well known, maps are generated based on information contained in recovery plans, State vegetation maps and remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

### Threatened, migratory and marine species

Threatened, migratory and marine species distributions have been discerned through a variety of methods. Where distributions are well known and if time permits, distributions are inferred from either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc.) together with point locations and described habitat; or modelled (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where little information is available for a species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc.).

In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More detailed distribution mapping methods are used to update these distributions when time permits.

## 4 LIMITATIONS

The following species and ecological communities have not been mapped and do not appear in this report:

- threatened species listed as extinct or considered vagrants;
- some recently listed species and ecological communities;
- some listed migratory and listed marine species, which are not listed as threatened species; and
- migratory species that are very widespread, vagrant, or only occur in Australia in small numbers.

The following groups have been mapped, but may not cover the complete distribution of the species:

- listed migratory and/or listed marine seabirds, which are not listed as threatened, have only been mapped for recorded breeding sites; and
- seals which have only been mapped for breeding sites near the Australian continent

The breeding sites may be important for the protection of the Commonwealth Marine environment.

Refer to the metadata for the feature group (using the Resource Information link) for the currency of the information.

## Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [-Office of Environment and Heritage, New South Wales](#)
- [-Department of Environment and Primary Industries, Victoria](#)
- [-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [-Department of Environment, Water and Natural Resources, South Australia](#)
- [-Department of Land and Resource Management, Northern Territory](#)
- [-Department of Environmental and Heritage Protection, Queensland](#)
- [-Department of Parks and Wildlife, Western Australia](#)
- [-Environment and Planning Directorate, ACT](#)
- [-Birdlife Australia](#)
- [-Australian Bird and Bat Banding Scheme](#)
- [-Australian National Wildlife Collection](#)
- [-Natural history museums of Australia](#)
- [-Museum Victoria](#)
- [-Australian Museum](#)
- [-South Australian Museum](#)
- [-Queensland Museum](#)
- [-Online Zoological Collections of Australian Museums](#)
- [-Queensland Herbarium](#)
- [-National Herbarium of NSW](#)
- [-Royal Botanic Gardens and National Herbarium of Victoria](#)
- [-Tasmanian Herbarium](#)
- [-State Herbarium of South Australia](#)
- [-Northern Territory Herbarium](#)
- [-Western Australian Herbarium](#)
- [-Australian National Herbarium, Canberra](#)
- [-University of New England](#)
- [-Ocean Biogeographic Information System](#)
- [-Australian Government, Department of Defence](#)
- [Forestry Corporation, NSW](#)
- [-Geoscience Australia](#)
- [-CSIRO](#)
- [-Australian Tropical Herbarium, Cairns](#)
- [-eBird Australia](#)
- [-Australian Government – Australian Antarctic Data Centre](#)
- [-Museum and Art Gallery of the Northern Territory](#)
- [-Australian Government National Environmental Science Program](#)
- [-Australian Institute of Marine Science](#)
- [-Reef Life Survey Australia](#)
- [-American Museum of Natural History](#)
- [-Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [-Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact us](#) page.


[@Commonwealth of Australia](#)

[Department of Climate Change, Energy, the Environment and Water](#)


GPO Box 3000

Canberra ACT 2601 Australia

+61 2 6274 1111



DWER's Clearing Permit System Map Viewer showing no ESA's (dark green shaded areas) within the survey area (DWER, 2026)



DWER Clearing Permit System Map Viewer showing no waterbodies within the survey area (DWER, 2026)

## **Appendix 2: Threatened Flora Databases Search Results**

GIS information provided in the Search results listed the following species within a 20km radius of the survey area (DBCA, 2026a):

| <b>Taxon</b>                   | <b>Conservation Code</b> | <b>Comment (Post field work)</b>                          |
|--------------------------------|--------------------------|-----------------------------------------------------------|
| <i>Acacia speckii</i>          | P4                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Calytrix verruculosa</i>    | P3                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Maireana prosthochaeta</i>  | P3                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Prostanthera petrophila</i> | P3                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Ptilotus beardii</i>        | P3                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Ptilotus luteolus</i>       | P3                       | Unlikely- possible suitable habitat, extensively searched |
| <i>Sida picklesiana</i>        | P3                       | Unlikely- No suitable habitat                             |
| <i>Tecticornia cymbiformis</i> | P3                       | Unlikely- No suitable habitat                             |

### **Appendix 3: Vegetation Definitions**

## Vegetation Condition Definitions (Keighery, 1994)

**Pristine (1).** Pristine or nearly so, no obvious signs of disturbance.

**Excellent (2).** Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species.

**Very Good (3).** Vegetation structure altered, obvious signs of disturbance.  
For example, disturbance to vegetation structure caused by repeating fires, the presence of some more aggressive weeds, dieback, logging and grazing.

**Good (4).** Vegetation structure significantly altered by very obvious signs of multiple disturbance.

Retains basic vegetation structure or ability to regenerate it.

For example, disturbance to vegetation structure caused by frequent fires, the presence of some very aggressive weeds at high density, partial clearing, dieback and grazing.

**Degraded (5).** Basic vegetation structure severely impacted by disturbance.

Scope for regeneration but not to a state approaching good condition without intensive management.

For example, disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.


**Completely Degraded (6).** The structure of the vegetation is no longer intact and the area is completely or almost completely without native species.

These areas are often described as 'parkland cleared' with the flora compromising weed or crop species with isolated trees or shrubs.


## Vegetation Structure Definitions (Muir, 1977)

| Life Form/Height Class | Canopy Cover          |                          |                       |                           |
|------------------------|-----------------------|--------------------------|-----------------------|---------------------------|
|                        | Dense<br>70-100%<br>d | Mid-Dense<br>30-70%<br>c | Sparse<br>10-30%<br>i | Very Sparse<br>2-10%<br>r |
| T Trees>30m            | Dense Tall Forest     | Tall Forest              | Tall Woodland         | Open Tall Woodland        |
| M Trees 15-30m         | Dense Forest          | Forest                   | Woodland              | Open Woodland             |
| LA Trees 5-15m         | Dense Low Forest A    | Low Forest A             | Low Woodland A        | Open Low Woodland A       |
| LB Trees<5m            | Dense Low Forest B    | Low Forest B             | Low Woodland B        | Open Low Woodland B       |
| KT Mallee tree form    | Dense Tree Mallee     | Tree Mallee              | Open Tree Mallee      | Very Open Tree Mallee     |
| KS Mallee shrub form   | Dense Shrub Mallee    | Shrub Mallee             | Open Shrub Mallee     | Very Open Shrub Mallee    |
| S Shrubs>2m            | Dense Thicket         | Thicket                  | Scrub                 | Open Scrub                |
| SA Shrubs 1.5-2.0m     | Dense Heath A         | Heath A                  | Low Scrub A           | Open Low Scrub A          |
| SB Shrubs 1.0-1.5m     | Dense Heath B         | Heath B                  | Low Scrub B           | Open Low Scrub B          |
| SC Shrubs 0.5-1.0m     | Dense Low Heath C     | Low Heath C              | Dwarf Scrub C         | Open Dwarf Scrub C        |
| SD Shrubs 0.0-0.5m     | Dense Low Heath D     | Low Heath D              | Dwarf Scrub D         | Open Dwarf Scrub D        |
| P Mat plants           | Dense Mat Plants      | Mat Plants               | Open Mat Plants       | Very Open Mat Plants      |
| H Hummock Grass        | Dense Hummock Grass   | Mid-Dense Hummock Grass  | Hummock Grass         | Open Hummock Grass        |
| GT Bunch grass >0.5m   | Dense Tall Grass      | Tall Grass               | Open Tall Grass       | Very Open Tall Grass      |
| GL Bunch grass <0.5m   | Dense Low Grass       | Low Grass                | Open Low Grass        | Very Open Low Grass       |
| J Herbaceous spp.      | Dense Herbs           | Herbs                    | Open Herbs            | Very Open Herbs           |
| VT Sedges >0.5m        | Dense Tall Sedges     | Tall Sedges              | Open Tall Sedges      | Very Open Tall Sedges     |
| VL Sedges <0.5m        | Dense Low Sedges      | Low Sedges               | Open Low Sedges       | Very Open Low Sedges      |
| X Ferns                | Dense Ferns           | Ferns                    | Open Ferns            | Very Open Ferns           |
| Mosses, liverwort      | Dense Mosses          | Mosses                   | Open Mosses           | Very Open Mosses          |


## **Appendix 4: Vegetation Mapping**




**Map 1: Survey Area**




**Map 2: NVS GPS Data**




**Map 3: DBCA Databases Search Results**




**Map 4: Land Systems**




**Map 5: Beard Pre-European Vegetation Groups**



Map 6: Vegetation Groups



**Map 7: Vegetation Condition**



**Map 8: Priority Flora Locations**

**Appendix 5: Species List**

## Species List per Vegetation Group

| Family         | Genus                | Taxon                                                      | A, P, NN | A | B | C | D | E | F | G | H | I | J | K | L | M | N |
|----------------|----------------------|------------------------------------------------------------|----------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Aizoaceae      | <i>Tetragonia</i>    | <i>Tetragonia eremaea</i>                                  | A        |   |   | * | * |   |   | * |   |   |   |   |   |   |   |
| Aizoaceae      | <i>Trianthema</i>    | <i>Trianthema triquetrum</i>                               | A        |   |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Amaranthaceae  | <i>Ptilotus</i>      | <i>Ptilotus helipteroides</i>                              | A        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Amaranthaceae  | <i>Ptilotus</i>      | <i>Ptilotus obovatus</i>                                   | P        | * | * | * | * | * | * | * |   | * | * |   | * | * | * |
| Amaranthaceae  | <i>Ptilotus</i>      | <i>Ptilotus roei</i>                                       | P        | * |   | * |   | * |   |   |   |   |   |   |   |   |   |
| Amaranthaceae  | <i>Ptilotus</i>      | <i>Ptilotus rotundifolius</i>                              | P        | * | * |   |   |   |   | * | * |   | * | * |   |   | * |
| Amaranthaceae  | <i>Ptilotus</i>      | <i>Ptilotus schwartzii</i>                                 | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Apocynaceae    | <i>Leichhardtia</i>  | <i>Leichhardtia australis</i>                              | P        |   | * | * | * | * |   |   |   | * |   |   |   |   | * |
| Asteraceae     | <i>Calotis</i>       | <i>Calotis plumulifera</i>                                 | A        |   |   | * |   |   | * |   |   |   |   |   |   |   |   |
| Asteraceae     | <i>Helipterum</i>    | <i>Helipterum craspedioides</i>                            | A        |   |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Asteraceae     | <i>Myriocephalus</i> | <i>Myriocephalus oldfieldii</i>                            | A        |   | * | * |   |   |   | * |   |   |   |   |   |   |   |
| Asteraceae     | <i>Siemssenia</i>    | <i>Siemssenia capillaris</i>                               | A        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Asteraceae     | <i>Vittadinia</i>    | <i>Vittadinia sulcata</i>                                  | A        |   |   |   |   |   | * |   |   |   |   |   |   |   |   |
| Brassicaceae   | <i>Carrichtera</i>   | <i>Carrichtera annua</i> *                                 | A,NN     |   | * | * | * | * |   | * |   |   |   |   |   |   |   |
| Brassicaceae   | <i>Lepidium</i>      | <i>Lepidium platypetalum</i>                               | P        |   | * |   |   |   |   |   |   |   |   |   | * |   |   |
| Campanulaceae  | <i>Lithotoma</i>     | <i>Lithotoma petraea</i>                                   | P        |   |   |   |   |   |   |   |   |   |   |   |   |   | * |
| Chenopodiaceae | <i>Atriplex</i>      | <i>Atriplex codonocarpa</i>                                | A        |   |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Atriplex</i>      | <i>Atriplex holocarpa</i>                                  | A        |   |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Atriplex</i>      | <i>Atriplex quinii</i>                                     | P        |   |   |   |   |   | * |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Atriplex</i>      | <i>Atriplex semibaccata</i>                                | A        |   |   |   |   |   | * |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Atriplex</i>      | <i>Atriplex vesicaria</i>                                  | P        |   |   |   | * |   |   |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Dysphania</i>     | <i>Dysphania melanocarpa</i>                               | A        | * | * | * |   |   | * | * |   | * |   |   |   |   |   |
| Chenopodiaceae | <i>Enchylaena</i>    | <i>Enchylaena tomentosa</i> var. <i>tomentosa</i>          | P        |   |   |   | * |   | * |   | * | * |   | * |   |   |   |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana georgei</i>                                    | P        |   |   |   |   |   |   | * |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana planifolia</i>                                 | P        |   | * |   |   |   | * |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana pyramidata</i>                                 | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana thesioides</i>                                 | P        |   |   |   |   |   | * | * |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana triptera</i>                                   | P        | * | * | * |   | * |   | * | * | * | * | * | * | * | * |
| Chenopodiaceae | <i>Maireana</i>      | <i>Maireana villosa</i>                                    | P        |   |   |   |   |   |   | * |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Rhagodia</i>      | <i>Rhagodia eremaea</i>                                    | P        |   |   |   | * |   | * |   |   | * |   |   |   |   |   |
| Chenopodiaceae | <i>Sclerolaena</i>   | <i>Sclerolaena diacantha</i>                               | P        | * |   |   |   |   | * |   |   | * |   |   | * |   |   |
| Chenopodiaceae | <i>Sclerolaena</i>   | <i>Sclerolaena ericantha</i>                               | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Sclerolaena</i>   | <i>Sclerolaena patenticuspis</i>                           | P        |   |   | * | * |   |   |   |   |   |   |   |   |   |   |
| Chenopodiaceae | <i>Tecticornia</i>   | <i>Tecticornia pterygosperma</i> subsp. <i>denticulata</i> | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Combretaceae   | <i>Alectryon</i>     | <i>Alectryon oleifolius</i> subsp. <i>canescens</i>        | P        |   |   |   | * |   |   |   |   |   |   |   |   |   |   |
| Cucurbitaceae  | <i>Cucumis</i>       | <i>Cucumis myriocarpus</i> *                               | A,NN     |   |   | * |   |   |   |   |   | * |   |   |   |   |   |
| Euphorbiaceae  | <i>Euphorbia</i>     | <i>Euphorbia tannensis</i> subsp. <i>eremophila</i>        | A        | * | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia ?nigripilosa</i> subsp. <i>nigripilosa</i>       | P        |   |   |   |   |   |   |   |   |   |   |   |   | * | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia</i> aff. <i>quadriramarginea</i>                 | P        |   |   |   |   |   |   |   |   |   |   |   |   |   | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia aneura</i>                                       | P        | * |   |   | * | * | * | * | * | * | * | * | * | * | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia aptaneura</i>                                    | P        | * | * | * |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia ayersiana</i>                                    | P        |   |   |   |   | * |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia balsamea</i>                                     | P        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia craspedocarpa</i>                                | P        | * |   | * |   | * | * |   |   | * | * |   |   |   | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia fuscaneura</i>                                   | P        |   |   | * |   |   | * |   |   | * |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia grasbyi</i>                                      | P        | * | * |   |   |   |   |   | * |   | * | * |   |   | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia incurvaneura</i>                                 | P        | * | * | * | * | * | * | * | * | * | * | * | * |   | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia kempeana</i>                                     | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia mulganeura</i>                                   | P        |   |   |   |   | * |   |   |   |   | * |   |   | * | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia oswaldii</i>                                     | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia pruinocarpa</i>                                  | P        | * |   |   | * |   |   | * | * |   | * | * | * |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia pteraneura</i>                                   | P        | * |   | * |   | * | * | * | * | * | * | * | * |   | * |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia quadriramarginea</i>                             | P        |   |   |   |   |   |   |   | * |   |   | * |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia ramulosa</i> var. <i>linophylla</i>              | P        |   |   |   |   | * |   | * |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia sclerosperma</i> subsp. <i>sclerosperma</i>      | P        |   |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia synchronicia</i>                                 | P        | * |   | * |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Acacia</i>        | <i>Acacia tetragonophylla</i>                              | P        | * | * | * | * | * | * | * | * | * | * | * | * |   |   |
| Fabaceae       | <i>Indigofera</i>    | <i>Indigofera monophylla</i>                               | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Fabaceae       | <i>Senna</i>         | <i>Senna artemisioides</i> subsp. <i>xsturtii</i>          | P        | * | * |   |   |   |   |   | * |   |   | * |   |   |   |
| Fabaceae       | <i>Senna</i>         | <i>Senna artemisioides</i> subsp. <i>helmsii</i>           | P        | * | * |   |   |   |   |   |   | * |   |   |   |   | * |
| Fabaceae       | <i>Senna</i>         | <i>Senna</i> sp. Austin                                    | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Fabaceae       | <i>Senna</i>         | <i>Senna</i> sp. Meekatharra                               | P        | * | * |   |   |   |   |   | * |   | * | * |   |   | * |
| Geraniaceae    | <i>Erodium</i>       | <i>Erodium crinitum</i>                                    | P        | * | * |   | * |   |   |   |   |   |   |   |   |   |   |
| Goodeniaceae   | <i>Goodenia</i>      | <i>Goodenia mimuloides</i>                                 | A        | * | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Goodeniaceae   | <i>Scaevola</i>      | <i>Scaevola spinescens</i>                                 | P        | * |   |   |   |   |   |   |   |   | * |   | * |   |   |
| Lamiaceae      | <i>Prostanthera</i>  | <i>Prostanthera althoferi</i> subsp. <i>althoferi</i>      | P        |   |   |   |   |   |   |   |   |   |   |   |   |   | * |

| Family           | Genus              | Taxon                                                      | A, P, NN | A | B | C | D | E | F | G | H | I | J | K | L | M | N |
|------------------|--------------------|------------------------------------------------------------|----------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Lamiaceae        | <i>Teucrium</i>    | <i>Teucrium teucriflorum</i>                               | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Malvaceae        | <i>Abutilon</i>    | <i>Abutilon cryptopetalum</i>                              | P        |   |   |   | * |   |   |   |   |   |   |   |   |   |   |
| Malvaceae        | <i>Abutilon</i>    | <i>Abutilon oxycarpum</i>                                  | P        |   |   |   | * | * |   |   |   | * |   |   |   |   |   |
| Malvaceae        | <i>Sida</i>        | <i>Sida calyxhymenia</i>                                   | P        | * |   |   | * |   |   |   |   |   |   |   |   |   |   |
| Malvaceae        | <i>Sida</i>        | <i>Sida ectogama</i>                                       | P        | * |   |   |   |   |   |   | * |   | * | * |   |   | * |
| Malvaceae        | <i>Sida</i>        | <i>Sida</i> sp. dark green fruits                          | P        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Malvaceae        | <i>Sida</i>        | <i>Sida</i> sp. Excedentifolia                             | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Malvaceae        | <i>Sida</i>        | <i>Sida</i> sp. Golden calyces glabrous                    | P        |   |   |   |   |   |   |   | * |   | * | * |   |   |   |
| Montiaceae       | <i>Calandrinia</i> | <i>Calandrinia baccata</i>                                 | A        |   |   | * |   |   | * |   |   |   |   |   |   |   |   |
| Montiaceae       | <i>Calandrinia</i> | <i>Calandrinia creethae</i>                                | A        |   |   | * |   |   | * |   |   |   |   |   |   |   |   |
| Montiaceae       | <i>Calandrinia</i> | <i>Calandrinia eremaea</i>                                 | A        | * |   | * |   |   | * |   |   |   |   |   |   |   |   |
| Montiaceae       | <i>Calandrinia</i> | <i>Calandrinia polyandra</i>                               | A        |   |   | * |   |   | * | * |   |   |   |   |   |   |   |
| Myrtaceae        | <i>Aluta</i>       | <i>Aluta maisonneuvei</i> subsp. <i>auriculata</i>         | P        |   |   |   |   |   |   |   |   |   |   | * |   |   |   |
| Myrtaceae        | <i>Calytrix</i>    | <i>Calytrix desolata</i>                                   | P        |   | * |   |   |   |   |   |   |   |   |   |   | * |   |
| Myrtaceae        | <i>Micromyrtus</i> | <i>Micromyrtus sulphurea</i>                               | P        |   |   |   |   |   |   |   |   |   |   | * |   |   |   |
| Pittosporaceae   | <i>Pittosporum</i> | <i>Pittosporum angustifolium</i>                           | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Plantaginaceae   | <i>Plantago</i>    | <i>Plantago turrifera</i>                                  | A        |   |   | * |   |   |   | * |   |   |   |   |   |   |   |
| Poaceae          | <i>Aristida</i>    | <i>Aristida holathera</i> var. <i>holathera</i>            | A        | * | * |   |   | * |   | * |   |   | * |   |   |   |   |
| Poaceae          | <i>Enneapogon</i>  | <i>Enneapogon caeruleus</i>                                | A        | * | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Poaceae          | <i>Eragrostis</i>  | <i>Eragrostis eriopoda</i>                                 | P        |   |   |   |   |   |   | * |   |   | * |   |   |   |   |
| Poaceae          | <i>Eriachne</i>    | <i>Eriachne pulchella</i> subsp. <i>pulchella</i>          | A        | * | * |   |   |   | * |   |   |   |   |   |   |   |   |
| Poaceae          | <i>Monachather</i> | <i>Monachather paradoxus</i>                               | P        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Polygonaceae     | <i>Rumex</i>       | <i>Rumex vesicarius</i> *                                  | A, NN    | * |   |   |   |   | * | * |   |   |   |   |   |   |   |
| Proteaceae       | <i>Grevillea</i>   | <i>Grevillea berryana</i>                                  | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Proteaceae       | <i>Hakea</i>       | <i>Hakea lorea</i> subsp. <i>lorea</i>                     | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Proteaceae       | <i>Hakea</i>       | <i>Hakea preissii</i>                                      | P        |   |   | * |   |   |   |   |   | * |   |   |   |   | * |
| Proteaceae       | <i>Hakea</i>       | <i>Hakea recurva</i> subsp. <i>recurva</i>                 | P        | * |   |   |   |   |   |   |   | * |   |   | * |   |   |
| Pteridaceae      | <i>Cheilanthes</i> | <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>           | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Rubiaceae        | <i>Psyrax</i>      | <i>Psyrax latifolia</i>                                    | P        |   |   |   |   | * |   |   |   |   | * |   |   |   |   |
| Rubiaceae        | <i>Psyrax</i>      | <i>Psyrax rigidula</i>                                     | P        | * |   |   |   | * |   |   |   |   |   |   |   |   |   |
| Rubiaceae        | <i>Psyrax</i>      | <i>Psyrax suaveolens</i>                                   | P        | * |   |   |   |   |   | * | * |   |   | * |   |   |   |
| Santalaceae      | <i>Exocarpos</i>   | <i>Exocarpos aphyllus</i>                                  | P        |   |   |   |   |   |   |   |   |   |   |   | * |   |   |
| Santalaceae      | <i>Santalum</i>    | <i>Santalum spicatum</i>                                   | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Sapindaceae      | <i>Dodonaea</i>    | <i>Dodonaea amplisemina</i> (P4)                           | P        |   |   |   |   |   |   |   |   |   |   |   |   |   | * |
| Sapindaceae      | <i>Dodonaea</i>    | <i>Dodonaea petiolaris</i>                                 | P        |   |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila exilifolia</i>                               | P        | * |   |   |   |   |   |   |   |   | * |   |   | * | * |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila forrestii</i> subsp. <i>forrestii</i>        | P        | * |   |   |   | * |   |   |   |   | * |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila galeata</i>                                  | P        | * |   |   |   | * | * | * | * |   | * | * |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila georgei</i>                                  | P        |   |   |   |   | * |   |   |   |   | * |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila glutinosa</i>                                | P        |   |   |   |   |   | * |   |   |   |   |   |   |   | * |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila jucunda</i> subsp. <i>jucunda</i>            | P        | * |   |   |   |   |   |   |   |   | * |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila lachnocalyx</i>                              | P        | * |   |   |   |   |   |   |   |   |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila latrobei</i> subsp. <i>latrobei</i>          | P        |   |   |   |   | * |   |   | * |   |   | * |   | * | * |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila linearis</i>                                 | P        | * |   | * | * | * |   |   | * | * |   | * |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila longifolia</i>                               | P        |   |   |   | * |   |   |   |   |   |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila macmillaniana</i>                            | P        | * | * |   | * |   |   |   |   |   | * |   |   | * | * |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>    | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i> | P        |   |   |   |   |   |   |   |   |   |   |   | * |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila pantonii</i>                                 | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila platycalyx</i> subsp. <i>Granites</i>        | P        |   | * |   |   |   |   |   |   |   |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila platycalyx</i> subsp. <i>Leonora</i>         | P        |   |   |   | * |   | * |   |   |   |   |   |   |   |   |
| Scrophulariaceae | <i>Eremophila</i>  | <i>Eremophila youngii</i> subsp. <i>youngii</i>            | P        |   |   |   | * |   |   |   |   |   | * |   |   | * |   |
| Solanaceae       | <i>Nicotiana</i>   | <i>Nicotiana obliqua</i>                                   | A        |   |   |   |   |   |   | * |   |   |   |   |   |   |   |
| Solanaceae       | <i>Nicotiana</i>   | <i>Nicotiana rosulata</i>                                  | A        |   |   | * | * |   | * |   |   |   |   |   |   |   |   |
| Solanaceae       | <i>Solanum</i>     | <i>Solanum lasiophyllum</i>                                | P        |   | * | * | * |   | * | * |   | * | * |   |   |   |   |
| Stylidiaceae     | <i>Stylidium</i>   | <i>Stylidium longibracteatum</i>                           | P        |   |   |   |   |   |   |   |   |   |   | * |   |   |   |
| Thymelaeaceae    | <i>Pimelea</i>     | <i>Pimelea microcephala</i> subsp. <i>microcephala</i>     | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Thymelaeaceae    | <i>Pimelea</i>     | <i>Pimelea spiculigera</i> var. <i>spiculigera</i>         | P        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Zygophyllaceae   | <i>Roepera</i>     | <i>Roepera apiculata</i>                                   | A        |   |   |   |   |   |   |   |   | * |   |   |   |   |   |
| Zygophyllaceae   | <i>Roepera</i>     | <i>Roepera ovata</i>                                       | A        | * |   | * |   |   |   |   |   | * |   |   |   |   |   |
| Zygophyllaceae   | <i>Tribulus</i>    | <i>Tribulus astrocarpus</i>                                | A        | * |   | * |   |   |   |   |   |   |   |   |   |   |   |