



1. Application details

1.1. Permit application details

Permit application No.: 2745/1
Permit type: Purpose Permit

1.2. Proponent details

Proponent's name: Pioneer Nickel Limited

1.3. Property details

Property: M74/163
Local Government Area: Shire of Ravensthorpe
Colloquial name: Exploration Drilling

1.4. Application

Clearing Area (ha)	No. Trees	Method of Clearing	For the purpose of:
0.935		Mechanical Removal	Mineral Exploration

2. Site Information

2.1. Existing environment and information

2.1.1. Description of the native vegetation under application

Vegetation Description Beard Vegetation Associations have been mapped at a 1:250,000 scale for the whole of Western Australia. One Beard vegetation Association has been mapped within the application area (GIS Database; Shepherd *et al.*, 2001).

691: Shrublands; *Dryandra quercifolia* & *Eucalyptus* spp. thicket.

The application area was surveyed by McQuoid in August 2008 (McQuoid, 2008). The following four vegetation units were identified within the application area according to the descriptions of Craig *et al.* (2007):

1. *Eucalyptus flocktoniae*/E. phenax (Eflo/Ephe): A tall mallee community with a sparse shrub understorey. Common species found within this vegetation unit include:

Mallees: *Eucalyptus flocktoniae* subsp. *flocktoniae*, *E. phenax* subsp. *phenax*, *E. calycogona* subsp. *calycogona* and *E. suggrandis* subsp. *suggrandis*;

Tall Shrubs: *Melaleuca hamata* and *M. pauperiflora* subsp. *pauperiflora*;

Mid Shrubs: *Exocarpos aphyllus*, *Beyeria brevifolia* var. *brevifolia*, *Daviesia nematophylla*, *Dodonaea concinna*, *Gastrolobium parviflorum* forma 'broad', *Hakea verrucosa*, *Melaleuca* sp. Gorse, *Phebalium tuberculatum* and *Senna artemisioides* subsp. *x artemisioides*;

Low Shrubs: *Boronia inornata*, *Dodonaea pinifolia*, *Lasiopetalum compactum* and *Platysace maxwellii*; and

Dwarf Shrubs: *Acacia erinacea*, *A. glaucoptera* forma 'spreading', *A. ingrata*, *A. pusilla*, *Eremophila densifolia* and *Grevillea huegelii*.

2. *Eucalyptus falcata*/E. pleurocarpa (Efal/Eple): This heterogeneous mallee heath complex dominates the landscape on the laterites and is characterised by patches of mallee, interspersed with a diversity of shrubs, many being proteaceous. Common species found within this vegetation unit include:

Mallees: *Eucalyptus pleurocarpa*, *E. falcata* subsp. *falcata*, *E. uncinata*, *E. incrassata* and *E. phaenophylla* subsp. *phaenophylla*;

Tall Shrubs: *Banksia lemmaniana*, *B. laevigata* subsp. *laevigata*, *Beaufortia orbifolia*, *Grevillea coccinea* subsp. *coccinea*, *Hakea pandanica* subsp. *crassifolia* and *Melaleuca hamata*;

Mid Shrubs: *Acacia fragilis*, *A. heterochroa* subsp. *heterochroa*, *Allocasuarina humilis*, *Beyeria brevifolia* var. *brevifolia*, *Boronia ternate* var. *elongate*, *Calothamnus quadrifidus*, *Dryandra cirsioides*, *D. foliosissima*, *D. pallida*, *Gastrolobium parviflorum* forma 'broad', *Hakea cygna* subsp. *cygna*, *H. lissocarpa*, *H. obtuse*, *H. subsulcata*, *Isopogon polycephalus*, *Jacksonia viscosa*, *Kunzea cincinnata*, *Leptospermum spinescens*, *Melaleuca rigidifolia*, *M. subtrigona*, *Petrophile seminuda*, *Rhadinothamnus rudis* subsp. *amblycarpus* and *Taxandria spathulata*;

Low Shrubs: *Beaufortia schaueri*, *Hakea marginata*, *Hibbertia mucronata*, *Lasiopetalum compactum*, *Leucopogon conostephioides*, *Petrophile glauca*, *Philothea gardneri* subsp. *Ravensthorpe*; and

Dwarf Shrubs: *Rinzia communis*.

3. *Eucalyptus sporadica* (Espo): Generally a diverse thicket community with an overstorey of mallees that include *Eucalyptus sporadica*; and are generally restricted to drainage lines. Common species found within this vegetation unit include:

Mallees: *Eucalyptus falcata* subsp. *falcata*, *E. flocktoniae* subsp. *flocktoniae*, *E. incrassata*, *E. phenax* subsp.

phenax. *E. pluricaulis* ssp. *pluricaulis* and *E. sporadica*;

Tall Shrubs: *Exocarpus sparteus*, *Hakea laurina* and *Melaleuca hamata*;

Mid Shrubs: *Acacia durabilis*, *A. fragilis*, *A. sulcata* var. *platyphylla*, *Calothamnus quadrifidus*, *Gastrolobium parviflorum* forma 'broad', *Grevillea patentiloba* subsp. *platypoda*, *Phebalium tuberosum*, *Rhadinothamnus rudis* subsp. *amblycarpus* and *Trymalium elachophyllum*;

Low Shrubs: *Dodonaea pinifolia*, *Lasiopetalum compactum* and *Platysace maxwellii*;

Dwarf Shrubs: *Marianthus mollis*;

Grasses/Herbs: *Cassytha melantha*; and

Sedges: *Lepidosperma* sp. *Ravensthorpe* and *Lepidosperma* sp. GFC 8278.

4. *Allocasuarina campestris* (Alca): Shrub community dominated by *Allocasuarina campestris* usually associated with outcrops or granite. Common species found within this vegetation unit include:

Tall Shrubs: *Allocasuarina campestris*, *Leptospermum maxwellii*, *Melaleuca hamata* and *Santalum acuminatum*;

Mid Shrubs: *Calothamnus quadrifidus*, *Hakea verrucosa*, *Kunzea cincinnata*, *Melaleuca rigidifolia* and *Petrophile seminuda*;

Low Shrubs: *Dodonaea pinifolia* and *Platysace maxwellii*;

Sedges/Sedge-Like: *Lepidosperma brunonianum*; and

Grasses: *Spartochloa scirpoidea* and *Neurachne alopecuroidea*.

One species of introduced flora was recorded within the application area: Bridal Creeper (*Asparagus asparagoides*) (Pioneer Nickel Ltd, 2006).

Clearing Description	Pioneer Nickel Ltd (from this point referred to as Pioneer) has applied to clear up to 0.935 hectares of native vegetation for the purposes of mineral exploration and access tracks. The area has been previously disturbed by historical mining and exploration activities.
Vegetation Condition	Excellent: Vegetation structure intact; disturbance affecting individual species, weeds non-aggressive (Keighery, 1994) to Good: Structure significantly altered by multiple disturbance; retains basic structure/ability to regenerate (Keighery, 1994).
Comment	The vegetation condition was derived from a vegetation survey conducted by Craig <i>et al.</i> (2007).

3. Assessment of application against clearing principles

(a) Native vegetation should not be cleared if it comprises a high level of biological diversity.

Comments **Proposal may be at variance to this Principle**

The application area occurs within the Fitzgerald (ESP1) sub-region of the Esperance Bioregion of the Interim Biogeographic Regionalisation of Australia (IBRA) (GIS Database). This sub-region is characterised by myrtaceous and proteaceous scrub and mallee heaths on sand plains overlying Eocene sediments. Herb fields and heaths occur on abrupt granite tors and quartzite ranges that rise from the plain, while Eucalypt woodlands occur in the gullies and alluvial foot-slopes (CALM, 2002). The vegetation described within the application area is typical of the bioregion (McQuoid, 2008).

The application area is located within the Elverdton-Desmond area. The Elverdton-Desmond area and surrounds are one of the three areas of highest flora endemism in Western Australia, with more than 60 endemic species with a range of less than 30 kilometres (Pioneer, 2006). The application area is within the Ravensthorpe Range (Red Book System 3.8). This area is recognised as an Environmentally Sensitive Area (ESA), and forms part of the Fitzgerald Biosphere, providing an important linkage between the Fitzgerald River National Park and crown land east of the vermin proof fence (GIS Database; Craig *et al.*, 2007). The Ravensthorpe Range area has been listed by the Australian Heritage Council on the Register of National Estate as it contains many rare and endemic plant species and is the only natural locality of several eucalypts (DEWHA, 2008).

The Ravensthorpe Range - Fitzgerald River National Park contains 75 endemic flora species and makes a significant contribution to the South Coast Region's biodiversity values (Danks, 2004). The plants of the South Coast Region belong to 756 genera in 146 families. Myrtaceae, Proteaceae, Papilionaceae, Epacridaceae, Mimosaceae and Orchidaceae families are particularly species rich and diverse within the area (Danks, 2004).

One introduced species was recorded during the survey (McQuoid, 2008). Bridal Creeper (*Asparagus asparagoides*) is a 'Declared Plant' under the Agriculture and Related Resources Protection Act, 1976. Bridal Creeper is listed as P1, therefore should a clearing permit be granted, it is recommended that a condition be imposed for the purposes of weed management.

An area search of the Western Australian Museum's Faunabase conducted by the assessing officer suggests that the application area is diverse in avian and reptile species, particularly Skinks (20) (Western Australian Museum, 2008). The database search found 73 avian species from 28 families and 47 reptile species from 7 families as potentially occurring within the application area, or within a 50 kilometre radius of the application area.

Based on the above, the proposed clearing may be at variance to this Principle. However, the Ravensthorpe area has had a long history of exploration and mining (approximately 100 years) and as a result there has been considerable disturbance in the Elverdton-Desmond area (Pioneer, 2006). Considering the relatively small size and disturbed nature of the application area, it is unlikely that the proposed clearing will compromise the diversity of the vegetation within the application area in the long term.

Methodology CALM (2002)
Craig *et al.* (2007)
Danks (2004)
DEWHA (2008)
McQuoid (2008)
Pioneer (2006)
Western Australian Museum (2008)
GIS Database
- Interim Biogeographic Regionalisation of Australia
- Clearing Regulations - Environmentally Sensitive Areas

(b) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

Comments Proposal is not likely to be at variance to this Principle

The assessing officer has conducted a search of the Western Australian Museum's online fauna database between the co-ordinates 120.6859°E, 33.1546°S and 119.5898°E, 34.0604°S, representing a 50 kilometre radius around the application area.

This search identified 12 Amphibian, 73 Avian, 33 Mammalian, 47 Reptilian and 25 Fish species that may occur within the application area (Western Australian Museum, 2008). Of these, the following species of conservation significance have previously been recorded within the search area: Southern Death Adder (*Acanthophis antarcticus*), *Ctenotus gemmula*, Red Tailed Phascogale (*Phascogale calura*), Quenda (*Isoodon obesulus fusciventer*), Rufous Fieldwren (*Calamanthus campestris*), Recherche Cape Barren Goose (*Cereopsis novaehollandiae grisea*), Hooded Plover (*Charadrius rubricollis*), Brush Bronzewing (*Phaps elegans*), Crested Shrike Tit (*Falcunculus frontatus*) and the Western Rosella (*Platycercus icterotis*).

Pioneer (2006) conducted a desktop search of the Department of Environment and Conservation (DEC) threatened fauna database and the Department of the Environment and Heritage (DEH) fauna of conservation significance database to identify species of conservation significance that had been recorded within the area specified. The co-ordinates used were similar to those used by the assessing officer above. In addition to those species listed above, the following fauna species of conservation significance were identified through this database search: *Lerista viduata*, Chuditch (*Dasyurus geoffroii*), Numbat (*Myrmecobius fasciatus*), Dibbler (*Parantechinus apicalis*), Heath mouse (*Pseudomys shorridgei*), Western Brush Wallaby (*Macropus irma*), Western Mouse (*Pseudomys occidentalis*), Tammar wallaby (*Macropus eugenii derbianus*), Malleefowl (*Leipoa ocellata*), Carnaby's Black Cockatoo (*Calyptorhynchus latirostris*), White-Tailed Black Cockatoo (*Calyptorhynchus* sp.), Western Ground Parrot (*Pezoporus wallicus flaviventris*), Western Bristlebird (*Dasyornis longirostris*), Shy Heathwren (*Hylacola cauta whitlocki*), Crested Bellbird (*Oreoica gutturalis gutturalis*), White-Browed Babbler (*Pomatostomus superciliosus ashbyi*) and the Western Whipbird (*Psophodes nigrogularis oberon*).

Based on habitat requirements, the following species are most likely to occur within the application area:

The Chuditch (Schedule 1 - Fauna that is rare or likely to become extinct, *Wildlife Conservation (Specially Protected Fauna) Notice, 2008*) is known to have occupied a wide range of habitats from woodlands, dry sclerophyll forests, riparian vegetation, beaches and deserts (DEC, 2008a). The chuditch now has a contracted range in south-western Australia, in areas dominated by sclerophyll forest or drier woodland and mallee scrub land (Serena & Soderquist, 2004). Given the large home ranges of the chuditch, it is possible that the vegetation within the application area may provide suitable habitat for this species. However, given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

The Dibbler (Schedule 1 - Fauna that is rare or likely to become extinct, *Wildlife Conservation (Specially Protected Fauna) Notice, 2008*) typically occupies heath and mallee-heath vegetation communities, preferring dense, long unburnt coastal heath (DEC, 2008b). The vegetation within the application area may provide suitable habitat for this species, however given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

Malleefowl (Schedule 1 - Fauna that is rare or likely to become extinct, *Wildlife Conservation (Specially Protected Fauna) Notice, 2008*) are largely confined to arid and semi-arid woodland that is dominated by mallee eucalypts on sandy soils, with less than 430 millimetres of rainfall annually (DEC, 2008c). However, they can also occur in habitats of acacia, paperbark, sheoak and other scrubs as well as eucalypt woodland and coastal heaths with an abundant layer of leaf litter for use in nest mounds (Garnett & Crowley, 2000). It is unlikely that

the application area would provide significant habitat for this species given the vegetation types are well represented within the bioregion and the area proposed to clear is small (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares).

The Western Ground Parrot (Schedule 1 - Fauna that is rare or likely to become extinct, *Wildlife Conservation (Specially Protected Fauna) Notice, 2008*) lives in floristically diverse heath lands where it feeds on fruit, seeds and leaves (Environment, 2008b). The application may provide suitable habitat for this species, however the last record of the Western Ground Parrot from the Ravensthorpe area was in 1987. Therefore, it is unlikely that the application area provides significant habitat for this species.

The wheat belt species of Western Rosella (Schedule 1 - Fauna that is rare or likely to become extinct, *Wildlife Conservation (Specially Protected Fauna) Notice, 2008*) is described as utilising woodland habitat. Threats include clearing for agriculture affecting availability of food and nesting hollows (Garnett & Crowley, 2000). This species has vast amounts of suitable habitat in the bioregion and due to the small nature of the application area (0.935 hectares) in relation to the larger sub-region (1,572,329 hectares) it is unlikely that it would provide significant habitat for this species.

Lerista viduata (P4 - DEC Priority Fauna List) is a species of skink that shelters at the base of trees and shrubs in Eucalypt woodlands amongst the leaf litter or on loamy clay soils (Wilson & Swan, 2004). Few populations of this species have been recorded; however one population was located at nearby Kundip. There is little information regarding the habitat requirements of this species, therefore the assessing officer is unable to determine with any certainty whether the vegetation within the application area is significant habitat for this species. The vegetation types within the application area are well represented throughout the bioregion and given the small size of the area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area provides significant habitat for this species.

The Western Brush Wallaby (P4 - DEC Priority Fauna List) was very common in the early days of settlement, however their range has been severely reduced and fragmented due to clearing for agriculture (DEC, 2008d). The optimum habitat for this species is open forest or woodland, particularly open seasonally wet flats with low grasses and open scrubby thickets (DEC, 2008d). The vegetation within the application area may provide suitable habitat for this species, however the habitat of mallee over heath may not be preferred habitat and so it is unlikely that the application area provides significant habitat for this species.

The Western Mouse (P4 - DEC Priority Fauna List) prefers long unburnt vegetation on sandy clay loam or sandy loam (DEC, 2008e). Suitable habitat can be varied and includes sparse low shrub land, tall dense shrub land, sparse to dense shrub mallee and mid-dense woodlands, with all preferred habitats having patches of extremely dense vegetation (DEC, 2008e). The soils of the application area are generally sandy and clayey and so may provide suitable habitat for this species. Given the small area proposed to clear (0.935 hectares) in relation to the size of the bioregion (1,572,329 hectares) it is unlikely that it would provide significant habitat for the Western Mouse.

Crested Bellbirds (P4 - DEC Priority Fauna List) inhabit the shrub layer of Eucalypt woodland, mallee, acacia shrubland, Triodia hummock grassland, saltbush and heath, where they feed on insects and seeds (Environment, 2008a). The Crested Bellbird range has contracted towards the inland in south-western Australia, and it is possible that the mallees present in the application area may provide suitable habitat for this species. Given that the vegetation types are well represented within the bioregion and the area proposed to clear is small (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area would provide significant habitat for this species.

The White-Browed Babbler (P4 - DEC Priority Fauna List) lives in Eucalypt forests and woodlands, and forages on and near the ground for insects and seeds (Environment, 2008b). The vegetation within the application area may provide suitable habitat for this species, however given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

The Western Whipbird (P4 - DEC Priority Fauna List) occurs from the Stirling Range east to Munglinup and north to Lake Grace (Environment, 2008c). This species inhabits mallee and heath, with recent observations being recorded from Kundip (Pioneer, 2006). The vegetation within the application area may provide suitable habitat for this species, however given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

The Crested Shrike-Tit (P4 - DEC Priority Fauna List) is endemic to Western Australia, south-west of a line between Geraldton and Point Culver (Serventy & Whittell, 2000). The Crested Shrike-Tit inhabits eucalypt forest and woodland, favouring smooth barked *Eucalyptus diversicolor*, *E. wandoo*, *E. salmonophloia*, *E. rudis* and *Acacia acuminata* over rough barked species (Serventy & Whittell, 2000). The vegetation within the application area provides suitable habitat for this species, however given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

The Tammar Wallaby (P5 - DEC Priority Fauna List) inhabits coastal scrub, heath, dry sclerophyll forest and

thickets in mallee and woodland (DEC, 2008f). The vegetation within the application area may provide suitable habitat for this species, however given that the vegetation types are well represented throughout the bioregion and the small area proposed to clear (0.935 hectares) in relation to the size of the sub-region (1,572,329 hectares) it is unlikely that the application area contains significant habitat for this species.

Based on the above the proposed clearing is not likely to be at variance to this Principle despite the possible presence of suitable habitat for various fauna species. The application area is not likely to provide significant habitat for the above species, particularly as parts of the application area have been degraded by previous disturbance from mining activities and given the low impact and localised nature of the proposed clearing activities, the proposed clearing is unlikely to significantly impact on fauna habitats in the area, or cause significant habitat fragmentation in the local area.

Methodology	DEC (2008a) DEC (2008b) DEC (2008c) DEC (2008d) DEC (2008e) DEC (2008f) Environment (2008a) Environment (2008b) Environment (2008c) Environment (2008d) Garnett & Crowley (2000) Pioneer (2006) Serena & Soderquist (2004) Serventy & Whittell (2000) Western Australian Museum (2008) Wilson & Swan (2004)
--------------------	--

(c) Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, rare flora.

Comments Proposal may be at variance to this Principle

According to available databases, no Declared Rare Flora (DRF) species occur within the application area (GIS Database). One population of *Marianthus villosus* (DRF) has been recorded approximately 0.34 kilometres north-east of the application area (GIS Database). Four populations of *Daviesia megacalyx* (DRF) have been recorded approximately 1.46 - 2.23 kilometres south-east of the application area (GIS Database).

A botanical consultant conducted a floristic survey of the application area in August 2008 (McQuoid, 2008). No species of Declared Rare Flora were recorded during the flora survey. Eight Priority Flora species were recorded during the flora survey, namely *Astartea* sp. Jerdacuttup (P1); *Guichenotia anota* ms (P1); *Guichenotia apetala* (P1); *Acacia durabilis* (P3); *Boronia oxyantha* subsp. *brevicalyx* (P3); *Leptospermum* sp. Bandalup Hill (P3); *Siegfriedia darwinioides* (P4) and *Spyridium glaucum* (P4) (McQuoid, 2008).

Guichenotia anota ms (P1) is a shrub 0.3-1 metres high found on sandy, loamy gravel and undulating land (Western Australian Herbarium, 2008). Two individuals were recorded from the application area (McQuoid, 2008). *Guichenotia anota* ms has previously been recorded only nine times and appears to be localised within the Ravensthorpe area (Western Australian Herbarium, 2008).

Guichenotia apetala (P1) is a compact, much-branched shrub 0.15-0.4 metres high found on gravel and laterite (Western Australian Herbarium, 2008). One population consisting of two individuals was recorded from the application area (McQuoid, 2008). Previous records indicate that *Guichenotia apetala* is locally abundant within the Ravensthorpe area, with it being recorded 25 times previously (Western Australian Herbarium, 2008).

Acacia durabilis (P3) is a slender, moderately open, spinescent shrub 0.7-2 metres high. It is found on rocky or lateritic clay, sandy clay and on stony ridges and hills (Western Australian Herbarium, 2008). One population consisting of two individuals was recorded from the application area (McQuoid, 2008). *Acacia durabilis* has previously been recorded along the coast from Albany to Ravensthorpe with records indicating that 36 populations have been recorded, some with 1000+ individuals (Western Australian Herbarium, 2008).

Siegfriedia darwinioides (P4) is a much branched spreading shrub 0.2-1 metres high. It is found on gravelly loam or sandy soils and clay and flats and ridges (Western Australian Herbarium, 2008). Three populations were recorded from the application area with population sizes ranging from 1 - 6 individuals (McQuoid, 2008). *Siegfriedia darwinioides* has been recorded across the lower south-west coast from Albany to Esperance (Western Australian Herbarium, 2008).

Spyridium glaucum (P4) is an erect or spreading shrub 0.5-1 metres high and is found on clay (Western Australian Herbarium, 2008). Three populations were recorded from the application area with population sizes ranging from 4 - 50+ plants (McQuoid, 2008). Previous records indicate that this species is localised around Ravensthorpe, with populations ranging from 10-1000+ plants (Western Australian Herbarium, 2008).

The above Priority Flora species were located within the application area and are likely to be impacted by the proposed clearing activities; however *Acacia durabilis* and *Spyridium glaucum* are known to regenerate from soil stored seed, while *Siegfriedia darwinioides*, *Guichenotia apetala* and *Guichenotia anota* ms may also respond to disturbance (McQuoid, 2008).

The Assessing Officer carried out a search for *Astartea* sp. Jerdacuttup, *Boronia oxyantha* subsp. *brevicalyx* and *Leptospermum* sp. Bandalup Hill on FloraBase on 3 October 2008 which noted that the conservation status of these species has been altered to Not Threatened (Western Australian Herbarium, 2008).

Based on the above, the proposed clearing may be at variance to this Principle. It is recommended that should a permit be granted, a condition be imposed on the permit with regard to flora management and stockpiling of all cleared topsoil and vegetation.

Methodology McQuoid (2008)
Western Australian Herbarium (2008)
GIS Database
- Declared Rare and Priority Flora List

(d) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of a threatened ecological community.

Comments Proposal is not likely to be at variance to this Principle

A search of available databases reveals that there are no Threatened Ecological Communities (TEC's) within the application area, although the Ravensthorpe System is planned for review (GIS Database; Craig, 2006).

The vegetation communities of the Elverdton-Desmond area lie within the Ravensthorpe System of Beard, which has been proposed for inclusion in CALM's (now Department of Environment and Conservation) Threatened Ecological Community database (Craig, 2006). Within the Ravensthorpe System five vegetation communities are recognised as being ecosystems at risk;

- Proteaceous heath thickets of the Ravensthorpe Range laterite upland;
- Mallee woodlands of breakaway slopes of Ravensthorpe Range;
- *Eucalyptus* spp. on red loams on lower foothills of the eastern Ravensthorpe Range;
- Pale grey sand ridge magnesite - *Eucalyptus purpurata* low forest on ridgetops and upper slopes; and
- Unique mallee on magnesite ridges west of Bandalup Hill (Craig, 2006).

A plant community of interest, approximately 1.7 kilometres south-west from the proposed clearing area on top of Mt Desmond, is being surveyed for occurrence and possible nomination as a TEC (DEC, 2006; DEC, 2007). The proposed TEC is a *Banksia* community, while the application area is within an *Allocasuarina*, *Melaleuca* and *Eucalyptus* vegetation association.

Due to the distance from the application area, these ecosystems are unlikely to be affected by the proposed clearing.

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology Craig (2006)
DEC (2006)
DEC (2007)
GIS Database
- Threatened Ecological Communities

(e) Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.

Comments Proposal is not at variance to this Principle

The application area falls within the IBRA Esperance Bioregion. Shepherd *et al.* (2001) report that approximately 51.1% of the pre-European vegetation still exists in this Bioregion. The vegetation in the application area is recorded as Beard Vegetation Association 691: Shrublands; *Dryandra quercifolia* & *Eucalyptus* spp. thicket (GIS Database; Shepherd *et al.*, 2001). According to Shepherd *et al.*, (2001) there is approximately 96.6% of this vegetation type remaining within the Bioregion (see table below).

Therefore the vegetation within the application area is not a significant remnant of native vegetation within an area that has been extensively cleared.

	Pre-European area (ha)*	Current extent (ha)*	Remaining %*	Conservation Status**	Pre-European % in IUCN Class I-IV Reserves (and post clearing %)
IBRA Bioregion – Esperance	2,899,944	1,483,240	~51.1%	Least Concern	~28.4% (~54%)
IBRA Subregion – Fitzgerald	1,570,670	844,885	~53.8%	Least Concern	~27.7% (~50.5%)
Local Government – Ravensthorpe	1,355,762	865,382	~63.8%	Least Concern	N/A
Beard veg assoc. – State					
691	45,589	34,951	~76.7%	Least Concern	~65.8% (~82.7%)
Beard veg assoc. – Bioregion					
691	35,491	34,274	~96.6%	Least Concern	~84.5% (~84.3%)
Beard veg assoc. - subregion					
691	35,491	34,274	~96.6%	Least Concern	~84.5% (~84.3%)

* Shepherd *et al.* (2001) updated 2005

** Department of Natural Resources and Environment (2002)

Based on the above, the proposed clearing is not at variance to this Principle.

Methodology Department of Natural Resources and Environment (2002)
Shepherd *et al.* (2001)
GIS Database
- Pre-European Vegetation

(f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.

Comments Proposal is at variance to this Principle

According to known GIS datasets, there are no known perennial watercourses or water bodies within the application area. There are numerous minor, non-perennial watercourses running in a westerly direction through the application area, which would occasionally flow during heavy rainfall events (GIS Database).

The application area experiences a rainfall of approximately 426 millimetres/year according to the nearest recording station at Ravensthorpe, located approximately 8.5 kilometres to the north-west (BOM, 2008). The application area also experiences a pan evaporation rate of approximately 1987 mm/year (Luke *et al.*, 1987).

It is the proponent's responsibility to liaise with the Department of Water to determine whether a Bed and Banks permit is required.

Based on the above, the proposed clearing is at variance to this Principle.

However, it is necessary to consider the proposed clearing in context:

- the watercourses have previously been disturbed by mineral exploration activity,
- the watercourses flow very rarely, with a very limited flow duration,
- vegetation downstream of the watercourses is likely to benefit from occasional flows, but is not likely to be dependent on them for survival,
- the vegetation surrounding the watercourses is not riparian vegetation.

On this basis, it is unlikely that the proposed clearing will impact on any watercourses or wetlands of significant environmental value.

Methodology BOM (2008)
Luke *et al.* (1987)
GIS Database
- Hydrography -Linear

(g) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

Comments Proposal is not likely to be at variance to this Principle

Pioneer describes the soils of the application area as generally sandy and clayey with some areas of ironstone gravels (Pioneer, 2006).

According to available GIS Databases, there is one soil type (MM15) within the application area (GIS Database). This soil type is described as;

- (i) on rolling to undulating terrain, brown and grey cracking clays
- (ii) on rolling areas, similar shallow soils, with a complex association of soils often containing some ironstone gravels (DAFF, 2008).

The Department of Agriculture and Food Western Australia (DAFWA, 2006) describes the landscape and soils of the 'Ravensthorpe Zone' as rolling low hills on greenstone (mafic and ultramafic) and soils as alkaline sandy duplex soils with some clays, sands and gravels.

Sandy earths have a moderate to high risk of wind erodibility and seasonal water logging may occur over the sandy topsoil and clays (Schoknecht, 2002), whilst the Ravensthorpe area is described as having a moderate to high salinity risk (DAFWA, 2006). However, the linear nature of the clearing suggests that the potential for wind erosion is low, and provided the disturbed areas are rehabilitated after drilling is completed there would be minimal risk of increased salinity and/or water logging.

There is no known occurrence of *Phytophthora cinnamomi* (PC Dieback) within the proposed clearing area (DEC, 2007). The Ravensthorpe area is recognised as being at risk to the introduction of PC dieback. The proposed exploration activities should be undertaken during dry soil conditions to minimise the introduction of the plant pathogen (DEC, 2006).

Based on the above, the proposed clearing is not likely to be at variance to this Principle. It is recommended that should a permit be granted, a condition be imposed on the permit with regard to hygiene to prevent the introduction and/or spread of PC dieback and weeds. In addition, a condition be imposed on the permit with regard to the stockpiling of all cleared topsoil and vegetation for rehabilitation purposes.

Methodology DAFF (2008)
DAFWA (2006)
DEC (2006)
DEC (2007)
Pioneer (2006)
Schoknecht (2002)
GIS Database
- Soils - Statewide

(h) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.

Comments Proposal may be at variance to this Principle

The application area is located within the Elverdton-Desmond area. The Elverdton-Desmond area and surrounding areas are one of the three areas of highest flora endemism in Western Australia, with more than 60 endemic species with a range of less than 30 kilometres (Pioneer, 2006). The application area forms part of the Ravensthorpe Range Area, recognised as an Environmentally Sensitive Area (ESA), and has been listed by the Australian Heritage Council on the Register of National Estate (GIS Database; DEWHA, 2008).

The Ravensthorpe Ranges (Redbook System 3.8) are classified as a C class nature reserve for its high floral diversity (EPA, 1974). The Ravensthorpe Range Area is rich in floral species and contains many rare and endemic plant species, including many species that are not known from the surrounding country. Some of these species are endemic, with some being common to the area but rare elsewhere (EPA, 1974). The Ravensthorpe Ranges are also a focal point for species of Eucalyptus, with over 20 taxa occurring in the area and it is the only natural locality of several eucalypts (DEWHA, 2008).

The application area lies within the eastern section Fitzgerald Biosphere which is a part-tenured management concept recognised by United Nations Educational, Scientific and Cultural Organisation (UNESCO) as well as by State and Commonwealth Governments (Craig *et al.*, 2007). The concept includes a core area (Fitzgerald River National Park), buffer zone (Crown land and invested reserves) and a zone of co-operation (private freehold farmland: cleared and uncleared) (Craig *et al.*, 2007).

The Ravensthorpe Range vegetation is recognised as an important linkage between the Fitzgerald River National Park and crown land east of the vermin proof fence, allowing for the continuity of biological processes (Craig *et al.*, 2007). The long term sustainability and viability of this corridor will largely depend on maintaining the vegetation in excellent condition. Secondary impacts such as spread of PC dieback, weeds, erosion and

drainage effects can impact an area far in excess of the immediate project area. However, given the small area proposed to be cleared and prior disturbances that have taken place, it is unlikely that the proposed clearing will significantly impede the use of this corridor by fauna.

Based on the above, the proposed clearing may be at variance to this Principle. However, the small area proposed to be cleared is unlikely to significantly impact on the environmental values of the surrounding areas. It is recommended that should a permit be granted, a condition be imposed on the permit with regard to hygiene. In addition, a condition should be imposed on the permit with regard to the stockpiling of all cleared topsoil and vegetation for rehabilitation purposes.

Methodology Craig *et al.* (2007)
DEWHA (2008)
EPA (1974)
Pioneer (2006)
GIS Database
- Environmentally Sensitive Areas

(i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.

Comments Proposal is not likely to be at variance to this Principle

According to available databases, the application area is not located within a Public Drinking Water Source Area (PDWSA) (GIS Database).

There are no permanent water bodies or watercourses within the application area (GIS Database). The application area is located in a temperate Mediterranean region and experiences an average annual rainfall of approximately 426 millimetres (CALM, 2002; BOM, 2008).

The application area is located within the Yilgarn South West Groundwater Province (GIS Database). The groundwater salinity within the application area is approximately 7,000 - 14,000 milligrams/Litre Total Dissolved Solids (TDS) (GIS Database). Given the size of the area to be cleared (0.935 hectares) compared to the size of the Yilgarn South West Groundwater Province (24,601,260 hectares) (GIS Database), the proposed clearing is not likely to cause salinity levels within the application area to alter significantly.

There are no known Groundwater Dependent Ecosystems within the application area (GIS Database).

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology BOM (2008)
CALM (2002)
GIS Database
- Public Drinking Water Source Area
- Hydrography - Linear
- Groundwater - Provinces
- Groundwater Salinity
- Potential Groundwater Dependent Ecosystems

(j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence or intensity of flooding.

Comments Proposal is not likely to be at variance to this Principle

The application area is located on the Jerdacuttup River Catchment area, with the Cordingup Creek to the north and Jerdacuttup River to the north and east (GIS Database). The low rises of shallow soils are intersected by weak drainage lines that eventually drain into Cordingup Creek and the Jerdacuttup River (Pioneer, 2006). The topography of the application area is gently undulating, suggesting that the area is not likely to be subject to flooding (GIS Database).

The small area to be cleared (0.935 hectares) in relation to the size of the Jerdacuttup River Catchment area (173,928 hectares) is not likely to lead to an increase in flood height or duration (GIS Database).

Based on the above the proposed clearing is not likely to be at variance to this Principle.

Methodology Pioneer (2006)
GIS Database
- Hydrographic Catchments - Catchments
- Topographic Contours - Statewide
- Geographic Names

Planning instrument, Native Title, Previous EPA decision or other matter.

Comments

There are two native title claims (WC98_070 and WC96_109) over the area under application. This claim has been registered with the National Native Title Tribunal on behalf of the claimant group. However, the tenement has been granted in accordance with the future act regime of the *Native Title Act 1993* and the nature of the act (i.e. the proposed clearing activity) has been provided for in that process, therefore the granting of a clearing permit is not a future act under the *Native Title Act 1993*.

There are several known Aboriginal Sites of Significance within the vicinity of the application area (GIS Database). It is the proponent's responsibility to comply with the *Aboriginal Heritage Act 1972* and ensure that no Sites of Aboriginal Significance are damaged through the clearing process.

It is the proponent's responsibility to liaise with the Department of Environment and Conservation and the Department of Water, to determine whether a Works Approval, Water Licence, Bed and Banks Permit, or any other licences or approvals are required for the proposed works.

The applicant has been granted three other clearing permits in the vicinity. One of these was referred to the Environmental Protection Authority, and subsequently it was determined that the proposal could be managed under the provisions of Part V of the *Environmental Protection Act 1986*.

No public submissions were received in regard to this Permit application.

Methodology GIS Database
- Aboriginal Sites of Significance
- Native Title Claims

4. Assessor's comments

Comment

The proposal has been assessed against the Clearing Principles, and the proposal is not at variance to Principle (e), is not likely to be at variance to Principles (b), (d), (g), (i) and (j) and may be at variance to Principles (a), (c) and (h), and is at variance to Principle (f).

It is recommended that should a permit be granted, conditions be imposed on the permit with regards to flora management, weed management and hygiene, rehabilitation, recording the areas cleared and reporting.

5. References

- Bureau of Meteorology (2008) BOM Website - Climate Averages by Number, Averages for RAVENSTHORPE.
www.bom.gov.au/climate/averages/tables/cw_010633.shtml (Accessed 10 October 2008).
- CALM (2002) A Biodiversity Audit of Western Australia's 53 Biogeographic Subregions in 2002. Department of Conservation and Land Management, Western Australia.
- Craig G.F., Sandiford E.M., Hickman E.J., Rick A.M. and Newell J. (2007) The Vegetation of the Ravensthorpe Range, Western Australia: I. Mt Short to South Coast Highway. WA Department of Environment and Conservation, South Coast Natural Resource Management Inc.
- Craig, G.F. (2006) Elverdton - Desmond MLA74/163 Declared Rare and Priority Flora Survey, Part II. Prepared for Pioneer Nickel Ltd, Western Australia.
- DAFF (2008) Department of Agriculture, Fisheries and Forestry - Digital Atlas of Australian Soils.
<http://www.daff.gov.au/brs/data-tools/daas-download> (Accessed 15 October 2008).
- DAFWA (2006) Department of Agriculture and Food Western Australia: Ravensthorpe Area Catchment Appraisal. Resource Management Technical Report 311.
- Danks A. (2004) South Coast Biodiversity: An Overview of Biodiversity Values, Threats and Conservation in the South Coast Region. Department of Conservation and Land Management, Albany.
- DEC (2006) Biodiversity Coordination Section Biodiversity Advice for Land Clearing application 1575/1 Pioneer Nickel. Email advice to the Department of Industry and Resources, dated 22 November 2006.
- DEC (2007) Biodiversity Advice for Land Clearing Application. Advice to Assessing Officer, Native Vegetation Assessment Branch, Department of Industry and Resources (DoIR) and the Environmental Protection Authority, received 19 April 2007, Ravensthorpe DEC Branch.
- DEC (2008a) Fauna Species Profiles - Chuditch (*Dasyurus geoffroii*)
www.dec.wa.gov.au/component/option,com_docman/task,doc_download/gid,125/Itemid,1288/ (Accessed 20 October 2008).
- DEC (2008b) Fauna Species Profiles - Dibbler (*Parantechinus apicalis*)
www.dec.wa.gov.au/component/option,com_docman/Itemid,/gid,126/task,doc_download/ - (Accessed 16 October 2008).
- DEC (2008c) Fauna Species Profiles - Malleefowl (*Leipoa ocellata*)
www.dec.wa.gov.au/component/option,com_docman/Itemid,/gid,118/task,doc_download/ - (Accessed 16 October 2008).
- DEC (2008d) Fauna Species Profiles - Western Brush Wallaby (*Macropus irma*)
www.dec.wa.gov.au/component/option,com_docman/Itemid,/gid,135/task,doc_download/ - (Accessed 15 October 2008).
- DEC (2008e) Fauna Species Profiles - Western Mouse (*Pseudomys occidentalis*)

- www.dec.wa.gov.au/component/option,com_docman/Itemid,/gid,148/task,doc_download/ - (Accessed 20 October 2008).
- DEC (2008f) Fauna Species Profiles - Tammar Wallaby (*Macropus eugenii*)
www.dec.wa.gov.au/component/option,com_docman/Itemid,/gid,133/task,doc_download/ - (Accessed 16 October 2008).
- Department of Natural Resources and Environment (2002) Biodiversity Action Planning. Action planning for native biodiversity at multiple scales; catchment bioregional, landscape, local. Department of Natural Resources and Environment, Victoria.
- DEWHA (Department of the Environment, Water, Heritage and the Arts) (2008) Australian Heritage Database.
<http://www.environment.wa.gov.au> (accessed 28 October 2008).
- Environment (2008a) www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/pubs/crested-bellbird.pdf
 - (Accessed 15 October 2008).
- Environment (2008b) www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/pubs/ground-parrot-w.pdf
 (Accessed 20 October 2008).
- Environment (2008c) www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/pubs/wbrowed-babbler.pdf - (Accessed 15 October 2008).
- Environment (2008d) www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/pubs/w-whipbird-wm.pdf - (Accessed 15 October 2008).
- EPA (1974) Conservation Reserves in Western Australia - Report of the Conservation Through Reserves Committee to the Environmental Protection Authority: Section 1, Systems 1-5 "CTRC Green Book". Environmental protection Authority, Perth.
- Garnett, S.T. and Crowley, G.M. (2000) Action Plan for Australian Birds 2000. Environment Australia, Canberra.
- Keighery, B.J. (1994) Bushland Plant Survey: A Guide to Plant Community Survey for the Community. Wildflower Society of WA (Inc). Nedlands, Western Australia.
- Luke, G.J., Burke, K.L. and O'Brien, T.M. (1987) Evaporation Data for Western Australia. Resource Management Technical Report No. 65. Department of Agriculture, Western Australia.
- McQuoid N.K. (2008) Survey for Declared Rare and Priority Flora of Manganese Deposit Drill Pads and Access Tracks, Mt Chester Area Ravensthorpe Range. Prepared for Pioneer Nickel Ltd. Unpublished Report dated July 2008.
- Pioneer Nickel Ltd (2006) Additional information to accompany the application for a clearing permit (purpose permit): Native Vegetation Management Plan. Pioneer Nickel Limited, Perth.
- Schoknecht, N. (2002) Soil Groups of Western Australia. A Simple Guide to the Main Soils of Western Australia. Resource Management Technical Report 246. Edition 3.
- Serena M. and Soderquist T.R. (2004) Western Quoll. In The Mammals of Australia. Strahm R. (E.d). Reed, New Holland.
- Serventy, D.L. and Whittell, H.M. (2000) Action Plan for Australian Birds 2000.
<http://www.environment.gov.au/biodiversity/threatened/publications/action/birds2000/recovery.html> (Accessed 15 July 2008).
- Shepherd, D.P., Beeston, G.R. and Hopkins, A.J.M. (2001) Native Vegetation in Western Australia, Extent, Type and Status. Resource Management Technical Report 249. Department of Agriculture, Western Australia.
- Western Australian Herbarium (2008) - FloraBase - The Western Australian Flora. Department of Environment and Conservation. <http://florabase.calm.wa.gov.au/> (Accessed 3 October 2008).
- Western Australian Museum (2008) Faunabase - Western Australian Museum, Queensland Museum and Museum and Art Gallery of NT Collections Databases. http://www.museum.wa.gov.au/faunabase/_asp_bin/AreaSearchcx.asp?d (Accessed 7 October 2008).
- Wilson, S. and Swan, G. (2003) A Complete Guide to Reptiles of Australia. Reeds New Holland.

6. Glossary

Acronyms:

BoM	Bureau of Meteorology, Australian Government.
CALM	Department of Conservation and Land Management, Western Australia.
DAFWA	Department of Agriculture and Food, Western Australia.
DA	Department of Agriculture, Western Australia.
DEC	Department of Environment and Conservation
DEH	Department of Environment and Heritage (federal based in Canberra) previously Environment Australia
DEP	Department of Environment Protection (now DoE), Western Australia.
DIA	Department of Indigenous Affairs
DLI	Department of Land Information, Western Australia.
DoE	Department of Environment, Western Australia.
DoIR	Department of Industry and Resources, Western Australia.
DOLA	Department of Land Administration, Western Australia.
DoW	Department of Water
EP Act	Environment Protection Act 1986, Western Australia.
EPBC Act	Environment Protection and Biodiversity Conservation Act 1999 (Federal Act)
GIS	Geographical Information System.
IBRA	Interim Biogeographic Regionalisation for Australia.
IUCN	International Union for the Conservation of Nature and Natural Resources – commonly known as the World Conservation Union
RIWI	Rights in Water and Irrigation Act 1914, Western Australia.

s.17 Section 17 of the Environment Protection Act 1986, Western Australia.
TECs Threatened Ecological Communities.

Definitions:

{Atkins, K (2005). *Declared rare and priority flora list for Western Australia, 22 February 2005. Department of Conservation and Land Management, Como, Western Australia* } :-

- P1 Priority One - Poorly Known taxa:** taxa which are known from one or a few (generally <5) populations which are under threat, either due to small population size, or being on lands under immediate threat, e.g. road verges, urban areas, farmland, active mineral leases, etc., or the plants are under threat, e.g. from disease, grazing by feral animals, etc. May include taxa with threatened populations on protected lands. Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
- P2 Priority Two - Poorly Known taxa:** taxa which are known from one or a few (generally <5) populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
- P3 Priority Three - Poorly Known taxa:** taxa which are known from several populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but are in need of further survey.
- P4 Priority Four – Rare taxa:** taxa which are considered to have been adequately surveyed and which, whilst being rare (in Australia), are not currently threatened by any identifiable factors. These taxa require monitoring every 5–10 years.
- R Declared Rare Flora – Extant taxa (= Threatened Flora = Endangered + Vulnerable):** taxa which have been adequately searched for, and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection, and have been gazetted as such, following approval by the Minister for the Environment, after recommendation by the State's Endangered Flora Consultative Committee.
- X Declared Rare Flora - Presumed Extinct taxa:** taxa which have not been collected, or otherwise verified, over the past 50 years despite thorough searching, or of which all known wild populations have been destroyed more recently, and have been gazetted as such, following approval by the Minister for the Environment, after recommendation by the State's Endangered Flora Consultative Committee.

{Wildlife Conservation (Specially Protected Fauna) Notice 2005} [Wildlife Conservation Act 1950] :-

- Schedule 1 Schedule 1 – Fauna that is rare or likely to become extinct:** being fauna that is rare or likely to become extinct, are declared to be fauna that is need of special protection.
- Schedule 2 Schedule 2 – Fauna that is presumed to be extinct:** being fauna that is presumed to be extinct, are declared to be fauna that is need of special protection.
- Schedule 3 Schedule 3 – Birds protected under an international agreement:** being birds that are subject to an agreement between the governments of Australia and Japan relating to the protection of migratory birds and birds in danger of extinction, are declared to be fauna that is need of special protection.
- Schedule 4 Schedule 4 – Other specially protected fauna:** being fauna that is declared to be fauna that is in need of special protection, otherwise than for the reasons mentioned in Schedules 1, 2 or 3.

{CALM (2005). *Priority Codes for Fauna. Department of Conservation and Land Management, Como, Western Australia* } :-

- P1 Priority One: Taxa with few, poorly known populations on threatened lands:** Taxa which are known from few specimens or sight records from one or a few localities on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, active mineral leases. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P2 Priority Two: Taxa with few, poorly known populations on conservation lands:** Taxa which are known from few specimens or sight records from one or a few localities on lands not under immediate threat of habitat destruction or degradation, e.g. national parks, conservation parks, nature reserves, State forest, vacant Crown land, water reserves, etc. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P3 Priority Three: Taxa with several, poorly known populations, some on conservation lands:** Taxa which are known from few specimens or sight records from several localities, some of which are on lands not under immediate threat of habitat destruction or degradation. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P4 Priority Four: Taxa in need of monitoring:** Taxa which are considered to have been adequately surveyed, or for which sufficient knowledge is available, and which are considered not currently threatened or in need of special protection, but could be if present circumstances change. These taxa are usually represented on conservation lands.
- P5 Priority Five: Taxa in need of monitoring:** Taxa which are not considered threatened but are subject to a specific conservation program, the cessation of which would result in the species becoming threatened within five years.

Categories of threatened species (*Environment Protection and Biodiversity Conservation Act 1999*)

- EX** **Extinct:** A native species for which there is no reasonable doubt that the last member of the species has died.
- EX(W)** **Extinct in the wild:** A native species which:
(a) is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; or
(b) has not been recorded in its known and/or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
- CR** **Critically Endangered:** A native species which is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria.
- EN** **Endangered:** A native species which:
(a) is not critically endangered; and
(b) is facing a very high risk of extinction in the wild in the near future, as determined in accordance with the prescribed criteria.
- VU** **Vulnerable:** A native species which:
(a) is not critically endangered or endangered; and
(b) is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
- CD** **Conservation Dependent:** A native species which is the focus of a specific conservation program, the cessation of which would result in the species becoming vulnerable, endangered or critically endangered within a period of 5 years.