



1. Application details

1.1. Permit application details

Permit application No.: 2912/1
Permit type: Purpose Permit

1.2. Proponent details

Proponent's name: Pilbara Manganese Pty Ltd

1.3. Property details

Property: Mining Lease 45/430
Mining Lease 45/431
Local Government Area: Shire Of East Pilbara
Colloquial name: Big Mack West Rock Stockpile Project

1.4. Application

Clearing Area (ha)	No. Trees	Method of Clearing	For the purpose of:
70		Mechanical Removal	Mineral Production

2. Site Information

2.1. Existing environment and information

2.1.1. Description of the native vegetation under application

Vegetation Description	Clearing Description	Vegetation Condition	Comment
<p>The area applied to clear has been broadly mapped at a scale of 1:250,000 as (Shepherd et al, 2001):</p> <ul style="list-style-type: none"> Beard Vegetation Association 173: hummock grasslands, shrub steppe; kanji over soft spinifex and <i>Triodia wiseana</i> on basalt; and Beard Vegetation Association 177: hummock grasslands, sparse shrub steppe; <i>Acacia bivenosa</i> over hard spinifex <i>Triodia brizoides</i>. <p>Mattiske Consulting (2007) undertook a flora and vegetation survey for the mining tenements (M45/430 and M45/431) within which the application area is located. The survey was conducted in May 2007 and consisted of surveying numerous sampling sites that were chosen to represent the different plant communities within the survey area (Mattiske Consulting, 2007).</p> <p>MBS Environmental (2008) have mapped the vegetation communities of the proposed project area. Fourteen vegetation units were recorded within the survey area with the following seven occurring within the application area (MBS Environmental, 2008):</p> <ol style="list-style-type: none"> 1) Woodland of <i>Eucalyptus camaldulensis</i> var. <i>obtusa</i> over <i>Acacia trachycarpa</i>, <i>Acacia inaequilatera</i> and <i>Ficus brachypoda</i> over <i>Cenchrus ciliaris</i> along major watercourses. 2) Scrub or Thicket of <i>Carissa lanceolata</i>, <i>Petalostylis labicheoides</i>, <i>Acacia bivenosa</i> and <i>Acacia ancistrocarpa</i> over <i>Triodia pungens</i>, <i>Triodia basedowii</i>, <i>Cenchrus ciliaris</i> and <i>Chrysopogon fallax</i> along minor watercourses. 3) Scrub or Low Shrubland of <i>Acacia ancistrocarpa</i>, <i>Acacia arida</i>, <i>Acacia acradenia</i>, <i>Petalostylis labicheoides</i>, <i>Gossypium australe</i>, <i>Acacia synchronicia</i> and <i>Acacia inaequilatera</i> over <i>Triodia longiceps</i> and <i>Triodia wiseana</i> with patches of <i>Cenchrus ciliaris</i> on flats, often associated with major watercourses. 4) Low Shrubland of <i>Acacia arida</i> and <i>Acacia hilliana</i> over <i>Triodia wiseana</i> and <i>Dampiera candidans</i> on slopes and hilltops. 5) Hummock Grassland of <i>Triodia longiceps</i> with scattered <i>Acacia bivenosa</i>, <i>Acacia synchronicia</i> and <i>Acacia ptychophylla</i> on flats and lower slopes. 6) Hummock Grassland of <i>Triodia longiceps</i> and <i>Triodia wiseana</i> with 	<p>This clearing permit application is for a purpose permit to clear up to 70ha of native vegetation within an area of approximately 152ha (GIS Database). The proposed clearing is located approximately 100km east of Nullagine and 400km south-east of Port Hedland (GIS Database).</p> <p>The purpose of the proposed clearing is for a waste rock stockpile and associated haul road (Pilbara Manganese, 2008). Vegetation clearing will be undertaken via mechanical means using a bulldozer and vegetation and topsoil will be stockpiled for rehabilitation purposes (Pilbara Manganese, 2008).</p>	<p>Very Good: Vegetation structure altered; obvious signs of disturbance (Keighery, 1994).</p>	<p>The vegetation condition rating was based on the flora and vegetation survey of the proposed clearing area which was conducted by Mattiske Consulting in May 2007.</p> <p>Apart from localised disturbances around roads and drill sites, mainly weed and fire has impacted on the condition of the tenement vegetation (Mattiske Consulting, 2007).</p>

occasional *Grevillea wickhamii* subsp. *hispidula* on flats and lower slopes.

7) Hummock Grassland of *Triodia basedowii*, *Triodia pungens* and *Triodia wiseana* with *Acacia bivenosa*, *Acacia pyrifolia* var. *morrisonii*, *Acacia synchronicia*, *Hakea lorea* subsp. *lorea* and emergent *Corymbia hamersleyana* and *Corymbia aspera* on undulating plains and slopes.

3. Assessment of application against clearing principles

(a) Native vegetation should not be cleared if it comprises a high level of biological diversity.

Comments **Proposal is not likely to be at variance to this Principle**

The application area is located within the Chichester subregion of the Pilbara Interim Biogeographic Regionalisation for Australia (IBRA) bioregion (GIS Database). The Chichester subregion is characterised by undulating Archaean granite and basalt plains with significant areas of basaltic ranges (CALM, 2002). At a broad scale, vegetation can be described as shrub steppe characterised by *Acacia inaequilatera* over *Triodia wiseana* hummock grasslands on plains, while *Eucalyptus leucophloia* tree steppes occur on ranges (CALM, 2002).

The geomorphology of the application area consists of low hills with steep slopes (MBS Environmental, 2008). Small chert and sandstone outcrops occur on the steeper slopes, primarily in the north-west part of the application area (MBS Environmental, 2008). Within the application area weakly incised drainage systems exist that flow westwards into the Oakover River, located approximately 6km from the application area (MBS Environmental, 2008). The flows within these creeks are ephemeral with no nearby permanent pools or waterholes (MBS Environmental, 2008). The vegetation within the application area is described broadly as varying from *Triodia* hummock grassland dominated plains, slopes and hills to *Acacia* shrubland over tussock grass in minor drainage lines (MBS Environmental, 2008). The proposed clearing area is close to other mining operations and has suffered disturbance from fire and weed invasion (MBS Environmental, 2008).

Several flora and vegetation surveys have been conducted over the Woodie Woodie tenements (MBS Environmental, 2008). A flora and vegetation survey was conducted by Mattiske Consulting in 2007 which identified a total of 127 taxa from 33 families and 71 genera within the application area. The most common families within the application area were the Grass family (*Poaceae*), Amaranth family (*Amaranthaceae*) and Acacia family (*Mimosaceae*) (MBS Environmental, 2008).

Mattiske Consulting (2007) identified four weed species within the survey area: Kapok Bush (*Aerva javanica*), Buffel Grass (*Cenchrus ciliaris*), Common Purslane (*Portulaca oleracea*) and Thornapple (*Datura leichhardtii*). The presence of these introduced weed species lowers the biodiversity value of the proposed clearing area. Care must be taken to ensure that the proposed clearing activities do not spread or introduce weed species to non-infested areas. Should a clearing permit be granted, it is recommended that a condition be imposed for the purposes of weed management.

Western Wildlife has undertaken two fauna surveys in 2006 and 2006/2007, over numerous tenements within the Woodie Woodie region, including the application area. They have identified four broad habitat types that will be affected by the proposed clearing, however, MBS Environmental (2008) has concluded that none of the landforms or habitat types are unique at the local or regional scale. Table 1 below shows the number of vertebrate fauna species recorded during the 2006/2007 survey in comparison to the number of species with the potential to occur:

Table 1: Vertebrate fauna species with the potential to occur within the Woodie Woodie tenements:

	Amphibians	Reptiles	Birds	Mammals	Total
Potential to occur	7	79	135	50	268
Recorded during survey	5	59	84	20	155

(Davis and Wilcox, 2007)

This table indicates that the application area is potentially high in bird and reptile species in particular, however, the landforms, vegetation types and fauna habitats in the application area are well represented in the Pilbara region (MBS Environmental, 2008). Furthermore, the application area is located immediately adjacent to an established mining area and pit, therefore, it is unlikely that the application area has greater biological diversity than other, undisturbed areas nearby.

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology CALM (2002)
Davis and Wilcox (2007)
Mattiske Consulting (2007)

(b) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.

Comments Proposal may be at variance to this Principle

Western Wildlife was commissioned by Consolidated Minerals Ltd to undertake vertebrate fauna surveys in the Woodie Woodie Project Area in 2006 and 2006/2007. These surveys were conducted in accordance with the Environmental Protection Authority (EPA) Position Statement No.3 and Guidance Statement 56: *Guidance for the Assessment for Environmental Factors - Terrestrial Fauna for Environmental Impact Assessment in Western Australia* (EPA 2002; 2004). The 2006/2007 reconnaissance fauna survey examined ten prospective areas within the Woodie Woodie tenements, including an area adjacent to the application area, while the 2006 reconnaissance fauna survey used opportunistic sampling methods to survey a further eight prospective areas, including the application area (Davis and Wilcox, 2006; 2007).

From these surveys, Davis and Wilcox (2007) identified four broad habitat types within the Woodie Woodie tenements:

1. *Cenchrus ciliaris* dominated plains and minor creek lines with emergent *Acacia*;
2. *Triodia* hummock grassland dominated plains;
3. Scrub / *Triodia* hummock grassland on low rocky hills and mesas;
4. Tall shrubland of *Acacia*.

The proposed clearing is likely to result in the following impacts to fauna:

- mortality of vertebrate and invertebrate species in the clearing footprint area. Sedentary species and young animals are particularly susceptible;
- displacement of mobile species in the proposed clearing area into surrounding habitats;
- temporary loss of habitat for foraging and shelter; and
- localised disturbance from noise and dust pollution.

Vegetation Community 1: Woodland of *Eucalyptus camaldulensis* var. *obtusa* over *Acacia trachycarpa*, *Acacia inaequilatera* and *Ficus brachypoda* over *Cenchrus ciliaris* along major watercourses, occurs within the southernmost portion of the application area (MBS Environmental, 2008). MBS Environmental (2008), report that this habitat unit is common throughout the Woodie Woodie tenements. However, this vegetation unit may still represent significant habitat for some fauna species, in particular the River Red Gum trees which may be of importance for bird species, including migratory birds.

The Woodie Woodie tenements are known to get visits from various migratory bird species including the Rainbow Bee-eater (*Merops ornatus*) and the Wood Sandpiper (*Tringa glareola*) which have been sighted within the application area on numerous occasions (MBS Environmental, 2008). The clearing area is quite large and migratory bird species may visit the Woodie Woodie area as it may be significant habitat for migratory bird species; in particular Vegetation Community 1. It is recommended that should a permit be granted, a condition be imposed for the protection of this vegetation community.

In addition to migratory bird species there are numerous other fauna species of conservation significance that have the potential to occur, with several having been recorded within the Woodie Woodie tenements during the fauna surveys. These were:

- Australian Bustard (*Ardeotis australis*) – Priority 4 on the Department of Environment and Conservation's (DEC) Threatened and Priority Fauna list;
- Star Finch (*Neochima ruficauda subclarescens*) – Priority 4 on the DEC's Threatened and Priority Fauna list;
- Princess Parrot (*Polytelis alexandrae*) – Priority 4 on the DEC's Threatened and Priority Fauna list;
- Western Pebble-mound Mouse (*Pseudomys chapmani*) – Priority 4 on the DEC's Threatened and Priority Fauna list.

The Australian Bustard is a dispersive species with widespread movements over long distances (DECC, 2005). The Australian Bustard is known to inhabit grasslands, low shrublands, grassy woodlands as well as altered environments such as croplands and airfields (DECC, 2005). The species usually breeds on bare ground, on low sandy ridges or stony rises (DECC, 2005). This species is slow to take flight and is therefore vulnerable to being killed by vehicles (Davis and Wilcox, 2007). This species has been recorded on numerous occasions throughout the Woodie Woodie tenements (Davis and Wilcox, 2007) and therefore, would be likely to occur within the application area. However, given the widespread distribution of this species it is unlikely that the vegetation within the application area would represent significant habitat for this species.

The Star Finch has a patchy distribution within the Pilbara, and can be found at low densities where it occurs (Garnett and Crowley, 2000). The Star Finch inhabits the dense vegetation around swamps, rivers and permanent waterholes in the larger watercourses of the north-west of Western Australia (Johnstone and Storr,

2004, as cited in Davis and Wilcox, 2007). The Star Finch is likely to occur seasonally in small numbers around waterholes in the area, and six individuals have previously been recorded on the Woodie Woodie tenements. Based on the above, it is possible that this species may occur near the ephemeral creekline within the application area, following rainfall, however, it is more likely that this species could be found in areas where permanent sources of water occur. Therefore, it is unlikely that the vegetation within the application area represents significant habitat for this species.

The Princess Parrot may occur in the area based on its known distribution. Swales between sand dunes are the preferred habitat of the Princess Parrot and nests have been recorded in hollows of River Red Gum (*Eucalyptus camaldulensis*) and Desert Oak (*Allocasuarina decaisneana*) (Higgins, 1999). It is possible that the Princess Parrot may sporadically inhabit the Woodie Woodie area, in particular the creek line habitat comprising River Red Gums. Approximately 1.61ha of the River Red Gum habitat occurs within the area applied to clear.

The Western Pebble-mound Mouse generally occurs on gentler slopes of rocky ranges where the ground is covered by a stony mulch and vegetated by hard Spinifex, often with an overstorey of eucalypts and scattered shrubs (Van Dyck and Strahan, 2008). Mounds are often sited close to narrow ribbons of *Acacia*-dominated scrub that grow along incised drainage lines (Van Dyck and Strahan, 2008). Numerous mounds of this species have been found throughout the Woodie Woodie tenements, on low rocky hills, however, all mounds found have been old with no evidence of recently active mounds (Davis and Wilcox, 2007). Suitable habitat for this species is present within the application area, although, it is considered more likely to occur in areas with larger hills (Davis and Wilcox, 2007). Pilbara Manganese should make all contractors aware that Western Pebble-mound Mouse mounds may be present in rocky places within the application area, and that these should be avoided wherever possible.

Based on the above, the proposed clearing may be at variance to this Principle.

Methodology Davis and Wilcox (2006)
Davis and Wilcox (2007)
DECC (2005)
EPA (2002)
EPA (2004)
Garnett and Crowley (2000)
Higgins (1999)
MBS Environmental (2008)
Van Dyck and Strahan (2008)

(c) Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, rare flora.

Comments Proposal is not likely to be at variance to this Principle

Mattiske Consulting undertook a flora and vegetation survey of the application area in May 2007. The survey involved choosing sampling sites within representative vegetation types and then describing and systematically sampling the vegetation within each site (Mattiske Consulting, 2007). In addition MBS Environmental (2008) undertook a desktop literature review of previous flora surveys from the Woodie Woodie tenements and a search of the DEC and Environmental Protection and Biodiversity Conservation (EPBC) databases to compile a potential Declared Rare Flora (DRF) and Priority Flora species list for the proposed clearing area.

According to available databases, there are no known records of DRF within 100km of the application area (GIS Database). In addition no DRF has previously been recorded within the Woodie Woodie tenements during previous flora and vegetation surveys (MBS Environmental, 2008).

According to available databases, there are no known records of Priority Flora within the proposed clearing area (GIS Database). Following a search of DEC and EPBC databases, MBS Environmental (2008) have identified 16 Priority species that could potentially occur in the region based on known distributions. In addition, a search by MBS Environmental (2008) of the Western Australian Herbarium specimen database indicates that two Priority species may occur within the Woodie Woodie tenement area: *Lepidium amelum* (Priority 1) and *Dampiera atriplicina* (Priority 2). These species were not found within the application area during the flora and vegetation survey conducted by Mattiske Consulting (2007).

The survey conducted by Mattiske Consulting (2007) identified two Priority species within Mining Lease 45/431: *Goodenia* sp. East Pilbara (Priority 1) and *Tephrosia* sp. Cathedral Gorge (Priority 3). These species are located approximately 100m from the application area and are therefore unlikely to be impacted by the proposed clearing.

The vegetation communities present within the proposed clearing area are typical of those found within the Hamersley IBRA subregion (MBS Environmental, 2008). It is not expected that the proposed clearing will result in the loss of habitat necessary for the continued existence of any DRF or Priority Flora species.

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology Matiske Consulting (2007)
MBS Environmental (2008)
GIS Database
- Declared Rare and Priority Flora list

(d) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of a threatened ecological community.

Comments Proposal is not likely to be at variance to this Principle

There are no known Threatened Ecological Communities (TEC's) within the area applied to clear or within 100km of the application area (GIS Database).

MBS Environmental (2008) report that no TEC's were identified during the flora survey of the application area.

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology MBS Environmental (2008)
GIS Database
- Threatened Ecological Communities

(e) Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.

Comments Proposal is not at variance to this Principle

The application area falls within the Pilbara IBRA Bioregion (GIS Database). Shepherd et al. (2001) report that approximately 99.9% of the pre-European vegetation still exists in this Bioregion (see table below).

The vegetation in the application area is recorded as (Shepherd et al, 2001):

- **Beard Vegetation Association 173:** Hummock grasslands, shrub steppe, Kanji over soft spinifex and *Triodia wiseana* on basalt; and
- **Beard Vegetation Association 177:** Hummock grasslands, sparse shrub steppe; *Acacia bivenosa* over hard spinifex *Triodia brizoides*.

According to Shepherd et al. (2001) approximately 100% of both of these vegetation associations remain within the Bioregion (see table below).

Therefore, the vegetation within the application area is not a significant remnant of native vegetation within an area that has been extensively cleared.

	Pre-European area (ha)*	Current extent (ha)*	Remaining %*	Conservation Status**	Pre-European % in IUCN Class I-IV Reserves
IBRA Bioregion – Pilbara	17,804,164	17,794,651	~99.9	Least Concern	6.3
Beard veg assoc. – State					
173	1,753,116	1,753,116	~100	Least Concern	7.5
177	169,446	169,446	~100	Least Concern	0.0
Beard veg assoc. – Bioregion					
173	1,752,533	1,752,533	~100	Least Concern	7.5
177	169,446	169,446	~100	Least Concern	0.0

* Shepherd et al. (2001) updated 2005

** Department of Natural Resources and Environment (2002)

Based on the above, the proposed clearing is not at variance to this Principle.

Methodology Shepherd et al. (2001)
Department of Natural Resources and Environment (2002)

GIS Database
- Interim Biogeographic Regionalisation for Australia
- Pre-European Vegetation

(f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.

Comments Proposal is at variance to this Principle

The application area contains several ephemeral drainage lines (GIS Database). MBS Environmental (2008) have reported three vegetation units within the application area that are generally associated with watercourses:

- 1) Woodland of *Eucalyptus camaldulensis* var. *obtusa* over *Acacia trachycarpa*, *Acacia inaequilatera* and *Ficus brachypoda* over *Cenchrus ciliaris* along major watercourses.
- 2) Scrub or Thicket of *Carissa lanceolata*, *Petalostylis labicheoides*, *Acacia bivenosa* and *Acacia ancistrocarpa* over *Triodia pungens*, *Triodia basedowii*, *Cenchrus ciliaris* and *Chrysopogon fallax* along minor watercourses.
- 3) Scrub or Low Shrubland of *Acacia ancistrocarpa*, *Acacia arida*, *Acacia acradenia*, *Petalostylis labicheoides*, *Gossypium australe*, *Acacia synchronicia* and *Acacia inaequilatera* over *Triodia longiceps* and *Triodia wiseana* with patches of *Cenchrus ciliaris* on flats, often associated with major watercourses.

These vegetation units are distributed within the application area as follows:

- Vegetation Unit 1 = 1.61ha;
- Vegetation Unit 2 = 20.74ha; and
- Vegetation Unit 3 = 36.69ha

MBS Environmental (2008) report that Vegetation Unit 3 is a common vegetation unit outside of the application area and vegetation maps provided by MBS Environmental (2008) confirm that this vegetation unit is well represented in surrounding areas.

MBS Environmental (2008) report that based on preliminary designs approximately 9ha of Vegetation Unit 2 will be impacted for the proposed project, and at this stage no clearing of Vegetation Unit 1 is deemed necessary. Vegetation Unit 2 is well represented throughout the Woodie Woodie tenements (MBS Environmental, 2008), and therefore the proposed clearing of 9ha of this vegetation type is unlikely to have a significant impact on the overall representation of the vegetation unit in the local area.

Should a permit be granted, it is recommended that if any watercourses are to be disturbed the proponent should liaise with the Department of Water to determine whether a Bed and Banks permit is necessary for the proposed works.

Based on the above, the proposed clearing is at variance to this Principle. It is recommended that should a permit be granted, a condition be imposed for the protection of Vegetation Unit 1.

Methodology MBS Environmental (2008)
GIS Database
- Hydrography - linear

(g) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

Comments Proposal is not likely to be at variance to this Principle

The majority of the application area has been mapped as occurring within the Coonigmah Land System (GIS Database). The Coonigmah Land System consists of plateau surfaces, low hills with steep slopes and undulating uplands supporting hard Spinifex grasslands (Van Vreeswyk et al., 2004). This land system is considered to have a very low erosion risk and the vegetation is not susceptible to degradation (Van Vreeswyk et al., 2004).

MBS Environmental (2008) have listed potential sources of land degradation from the proposed clearing:

- Wind erosion from topsoil stripping;
- Wind and water erosion of topsoil stockpiles;
- Wind and water erosion of rehabilitated surfaces e.g. waste rock stockpiles;
- Water erosion due to changes to the surface flow;
- Soil compaction;
- Soil contamination;
- Introduction and/or spread of weeds.

MBS Environmental (2008) report that Pilbara Manganese will implement management strategies in order to minimise land degradation, which include:

- Minimising the area requiring vegetation removal;
- Confining vehicle movements to defined haul roads and tracks;
- Conducting topsoil-stripping activities during periods of low winds;
- Establishing vegetation on bare surfaces on completion of activities;
- Stockpiling topsoil for use in rehabilitation;
- Implementation of a weed management program;
- Progressive rehabilitation of completed surfaces to minimise active areas exposed;
- Scarifying of compacted tracks prior to rehabilitation of the site.

Based on the above, the proposed clearing is not likely to be at variance to this Principle. It is recommended that should a permit be granted, conditions be imposed for the purposes of weed management and rehabilitation.

Methodology MBS Environmental (2008)
Van Vreeswyk et al. (2004)
GIS Database
- Rangelands System Mapping

(h) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.

Comments Proposal is not likely to be at variance to this Principle

The proposed clearing is not located within close proximity to any conservation areas (GIS Database). The nearest DEC managed land is the Rudall River National Park located approximately 90km south-east of the application area (GIS Database).

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology GIS Database
- CALM Managed Land and Waters

(i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.

Comments Proposal is not likely to be at variance to this Principle

The application area is located in an arid region with an average annual rainfall of approximately 327mm falling mainly during the summer months (MBS Environmental, 2008). Based on an average annual evaporation rate of approximately 3,800mm (MBS Environmental, 2008), any surface water resulting from rain events is relatively short-lived.

The application area has several ephemeral drainage lines running through it (GIS Database). Based on the climate of the region these creeks are expected to be dry except following significant rainfall events which are typically associated with tropical cyclones.

The groundwater and surface water of the Woodie Woodie region is well documented with over ten years of monitoring data (MBS Environmental, 2008). The groundwater and surface water within the Woodie Woodie region has pH ranging between 7.2 and 8.5 and is generally fresh to brackish (MBS Environmental, 2008).

The natural water table is more than 20 metres below natural ground level (MBS Environmental, 2008). Therefore, the impact of vegetation removal on groundwater levels is not likely to be significant. In addition, due to the arid climate, surface water runoff is expected to be minimal except following significant rainfall. Hence, the proposed clearing of the native vegetation is unlikely to have any significant impact on surface water flows or groundwater level or quality.

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology MBS Environmental (2008)
GIS Database
- Hydrography, linear

(j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence or intensity of flooding.

Comments Proposal is not likely to be at variance to this Principle

The application area is located in an arid region where the average annual evaporation rate greatly exceeds the average annual rainfall (MBS Environmental, 2008). There are no permanent watercourses within the

application area, however, several ephemeral drainage lines dissect the proposed clearing area (GIS Database). These drainage lines are expected to be dry for most of the year, and would likely only flow briefly immediately following significant rainfall.

Natural flood events do occur in the Pilbara following cyclonic activity. However, the proposed clearing is not expected to increase the incidence or intensity of such events given the size of the area to be cleared (70ha) in relation to the Oakover River catchment area (2,001,756ha) (GIS Database).

Based on the above, the proposed clearing is not likely to be at variance to this Principle.

Methodology MBS Environmental (2008)
GIS Database
- Hydrographic Catchments - catchments

Planning instrument, Native Title, Previous EPA decision or other matter.

Comments

There is one Native Title claim (WC99/008) over the area under application (GIS Database). This claim has been registered with the National Native Title Tribunal on behalf of the claimant group. However, the tenements have been granted in accordance with the future act regime of the *Native Title Act 1993* and the nature of the act (i.e. the proposed clearing activity) has been provided for in that process, therefore the granting of a clearing permit is not a future act under the *Native Title Act 1993*.

According to available databases there is one Aboriginal Site of Significance (site ID: 6329) within the application area (GIS Database). It is the proponent's responsibility to comply with the *Aboriginal Heritage Act 1972* and ensure that no Aboriginal Sites of Significance are damaged through the clearing process.

It is the proponent's responsibility to liaise with the Department of Environment and Conservation and the Department of Water, to determine whether a Works Approval, Water Licence, Bed and Banks permit, or any other licences or approvals are required for the proposed works.

There were no public submissions received during the public comments period.

Methodology GIS Database
- Aboriginal Sites of Significance
- Native Title Claims

4. Assessor's comments

Comment

The proposal has been assessed against the Clearing Principles, and is not at variance to Principle (e), is not likely to be at variance to Principles (a), (c), (d), (g), (h), (i) and (j), may be at variance to Principle (b) and is at variance to Principle (f).

Should the permit be granted it is recommended that conditions be imposed on the permit for the purposes of weed management, rehabilitation, protection of riparian vegetation, record keeping and permit reporting.

5. References

- CALM (2002) A Biodiversity Audit of Western Australia's 53 Biogeographic Subregions in 2002. Department of Conservation and Land Management, Western Australia.
- Davis, R.A. and Wilcox, J.A. (2006) Greensnake and Radio Hill, Woodie Woodie: Targeted Threatened Fauna Survey. Western Wildlife, Western Australia.
- Davis, R.A. and Wilcox, J.A. (2007) Woodie Woodie Project Area: Baseline Fauna Survey 2006/2007. Western Wildlife, Western Australia.
- DECC (2005) Australian Bustard - profile. Department of Environment and Climate Change. Available online from: <http://www.threatenedspecies.environment.nsw.gov.au/tsprofile/profile.aspx?id=10063>. Accessed 24 September, 2008.
- Department of Natural Resources and Environment (2002) Biodiversity Action Planning. Action planning for native biodiversity at multiple scales; catchment bioregional, landscape, local. Department of Natural Resources and Environment, Victoria.
- EPA (2002) Terrestrial Biological Surveys as an element of biodiversity protection. Position Statement No. 3. March 2002. Environmental Protection Authority
- EPA (2004) Guidance for the Assessment of Environmental Factors - terrestrial flora and vegetation surveys for Environmental Impact Assessment in Western Australia. Report by the EPA under the Environmental Protection Act 1986. No. 51 WA.
- Garnett, S.T. and Crowley, G.M. (2000) Action Plan for Australian Birds 2000. Environment Australia, Canberra.
- Higgins, P.J. (ed) (1999) Handbook of Australian, New Zealand and Antarctic Birds, Vol 4. Parrots to Dollarbird, Oxford University Press, Melbourne.
- Keighery, B.J. (1994) Bushland Plant Survey: A Guide to Plant Community Survey for the Community. Wildflower Society of WA (Inc). Nedlands, Western Australia.

- Mattiske Consulting (2007) Flora and Vegetation Survey of the Eat Tenement, Woodie Woodie. Mattiske Consulting Pty Ltd, Western Australia.
- MBS Environmental (2008) Purpose Permit Application, Big Mack West Waste Rock Stockpile: Native Vegetation Management Plan and Assessment of Clearing Principle. Martinick Bosch Sell Pty Ltd, Western Australia
- Pilbara Manganese (2008) EP Act - Pilbara Manganese Pty Ltd - Clearing Permit 2912/1. Pilbara Manganese Pty Ltd, Western Australia.
- Shepherd, D.P., Beeston, G.R. and Hopkins, A.J.M. (2001) Native Vegetation in Western Australia, Extent, Type and Status. Resource Management Technical Report 249. Department of Agriculture, Western Australia.
- Van Dyck, S. and Strahan, R. (eds.) (2008) The Mammals of Australia. Third Edition. New Holland Publisher (Australia) Pty Ltd, Australia.
- Van Vreeswyk, A.M.E., Payne, A.L., Hennig, P. and Leighton, K.A. (2004) An Inventory and Condition Survey of the Pilbara Region, Western Australia. Department of Agriculture, Western Australia.

6. Glossary

Acronyms:

BoM	Bureau of Meteorology, Australian Government.
CALM	Department of Conservation and Land Management, Western Australia.
DAFWA	Department of Agriculture and Food, Western Australia.
DA	Department of Agriculture, Western Australia.
DEC	Department of Environment and Conservation
DEH	Department of Environment and Heritage (federal based in Canberra) previously Environment Australia
DEP	Department of Environment Protection (now DoE), Western Australia.
DIA	Department of Indigenous Affairs
DLI	Department of Land Information, Western Australia.
DoE	Department of Environment, Western Australia.
DMP	Department of Mines and Petroleum, Western Australia.
DOLA	Department of Land Administration, Western Australia.
DoW	Department of Water
EP Act	Environment Protection Act 1986, Western Australia.
EPBC Act	Environment Protection and Biodiversity Conservation Act 1999 (Federal Act)
GIS	Geographical Information System.
IBRA	Interim Biogeographic Regionalisation for Australia.
IUCN	International Union for the Conservation of Nature and Natural Resources – commonly known as the World Conservation Union
RIWI	Rights in Water and Irrigation Act 1914, Western Australia.
s.17	Section 17 of the Environment Protection Act 1986, Western Australia.
TECs	Threatened Ecological Communities.

Definitions:

{Atkins, K (2005). *Declared rare and priority flora list for Western Australia, 22 February 2005*. Department of Conservation and Land Management, Como, Western Australia} :-

- P1** **Priority One - Poorly Known taxa:** taxa which are known from one or a few (generally <5) populations which are under threat, either due to small population size, or being on lands under immediate threat, e.g. road verges, urban areas, farmland, active mineral leases, etc., or the plants are under threat, e.g. from disease, grazing by feral animals, etc. May include taxa with threatened populations on protected lands. Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
- P2** **Priority Two - Poorly Known taxa:** taxa which are known from one or a few (generally <5) populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
- P3** **Priority Three - Poorly Known taxa:** taxa which are known from several populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but are in need of further survey.
- P4** **Priority Four – Rare taxa:** taxa which are considered to have been adequately surveyed and which, whilst being rare (in Australia), are not currently threatened by any identifiable factors. These taxa require monitoring every 5–10 years.
- R** **Declared Rare Flora – Extant taxa (= Threatened Flora = Endangered + Vulnerable):** taxa which have been adequately searched for, and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection, and have been gazetted as such, following approval by the Minister for the Environment, after recommendation by the State's Endangered Flora Consultative Committee.
- X** **Declared Rare Flora - Presumed Extinct taxa:** taxa which have not been collected, or otherwise verified, over the past 50 years despite thorough searching, or of which all known wild populations have been

destroyed more recently, and have been gazetted as such, following approval by the Minister for the Environment, after recommendation by the State's Endangered Flora Consultative Committee.

{Wildlife Conservation (Specially Protected Fauna) Notice 2005} [Wildlife Conservation Act 1950] :-

- Schedule 1** **Schedule 1 – Fauna that is rare or likely to become extinct:** being fauna that is rare or likely to become extinct, are declared to be fauna that is need of special protection.
- Schedule 2** **Schedule 2 – Fauna that is presumed to be extinct:** being fauna that is presumed to be extinct, are declared to be fauna that is need of special protection.
- Schedule 3** **Schedule 3 – Birds protected under an international agreement:** being birds that are subject to an agreement between the governments of Australia and Japan relating to the protection of migratory birds and birds in danger of extinction, are declared to be fauna that is need of special protection.
- Schedule 4** **Schedule 4 – Other specially protected fauna:** being fauna that is declared to be fauna that is in need of special protection, otherwise than for the reasons mentioned in Schedules 1, 2 or 3.

{CALM (2005). Priority Codes for Fauna. Department of Conservation and Land Management, Como, Western Australia} :-

- P1** **Priority One: Taxa with few, poorly known populations on threatened lands:** Taxa which are known from few specimens or sight records from one or a few localities on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, active mineral leases. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P2** **Priority Two: Taxa with few, poorly known populations on conservation lands:** Taxa which are known from few specimens or sight records from one or a few localities on lands not under immediate threat of habitat destruction or degradation, e.g. national parks, conservation parks, nature reserves, State forest, vacant Crown land, water reserves, etc. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P3** **Priority Three: Taxa with several, poorly known populations, some on conservation lands:** Taxa which are known from few specimens or sight records from several localities, some of which are on lands not under immediate threat of habitat destruction or degradation. The taxon needs urgent survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna.
- P4** **Priority Four: Taxa in need of monitoring:** Taxa which are considered to have been adequately surveyed, or for which sufficient knowledge is available, and which are considered not currently threatened or in need of special protection, but could be if present circumstances change. These taxa are usually represented on conservation lands.
- P5** **Priority Five: Taxa in need of monitoring:** Taxa which are not considered threatened but are subject to a specific conservation program, the cessation of which would result in the species becoming threatened within five years.

Categories of threatened species (Environment Protection and Biodiversity Conservation Act 1999)

- EX** **Extinct:** A native species for which there is no reasonable doubt that the last member of the species has died.
- EX(W)** **Extinct in the wild:** A native species which:
(a) is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; or
(b) has not been recorded in its known and/or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
- CR** **Critically Endangered:** A native species which is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria.
- EN** **Endangered:** A native species which:
(a) is not critically endangered; and
(b) is facing a very high risk of extinction in the wild in the near future, as determined in accordance with the prescribed criteria.
- VU** **Vulnerable:** A native species which:
(a) is not critically endangered or endangered; and
(b) is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
- CD** **Conservation Dependent:** A native species which is the focus of a specific conservation program, the cessation of which would result in the species becoming vulnerable, endangered or critically endangered within a period of 5 years.