



➤ Supporting Document for Amendment to CPS 6323/1

Christmas Island Phosphates
August 2024

Prepared by: SK, JK, RG
Version No: V001
Version date: 15/08/2024

E info@cirp.com
P +61 8 6250 4900
cirp.com.au



CHRISTMAS ISLAND PHOSPHATES





Table of Contents

Introduction	3
Background.....	3
Location.....	3
Tenure.....	4
Conservation Levy	4
Ownership	5
Proposed Amendment	5
Extension of Permit	5
Conditions	6
Clearing to Date	6
Relinquished Areas	6
Existing Environment	9
Climate	9
Hydrology	10
Hydrogeology	10
Geology, Soils and Landforms	10
Land Use.....	11
Land Systems	12
Environmental Sensitive Areas	12
Flora and Vegetation	15
Vegetation Types.....	15
Vegetation Condition	23
Significant Changes.....	28
Flora of Conservation Significance	28
Fauna and Habitat.....	39
Heritage.....	51
Environmental Management.....	52
Clearing Activities.....	52
Flora Management	52
Fauna Management	52
Weed Management.....	53
Mine Relinquishment.....	53
Ten Clearing Principals.....	53
Conclusion	58
References	59
Appendices	62



Introduction

Christmas Island is an Indian Ocean territory of Australia, located approximately 2 600 km north-west of Perth. Phosphate Resources Limited (PRL) operates phosphate mining, processing and shipping operations from Christmas Island under the trading name of Christmas Island Phosphates (CIP). PRL were provided with a mining lease and environmental approval to mine in 1997 and operated under this basis until the application of Western Australian legislation required the company to apply for clearing permits under the *Environmental Protection Act 1986* (WA) (CI) (EP Act) for all future clearing. The original lease granted was for 10 years, with a second lease provided in 1997 for a further 21 years, and a lease amendment in 2013 taking the current lease term until 2034. PRL have progressively applied for and obtained approval for clearing in target sites over mining lease MCI 70/1A.

Background

In June 2015, clearing permit CPS 6323/1 was granted under section 51E of the EP Act 1986 (WA)(CI), which is valid for the duration of 23 July 2015 to 31 December 2024. Areas with a high risk of environmental impacts were excluded from the application which was reduced in size from the initially proposed 132.53 ha to the approved 128.53 ha due to the proximity of Abbott’s Booby nests.

The clearing permit, CPS 6323/1, permits up to 128.53 hectares (ha) of clearing for the purpose of stockpile recovery and insitu phosphate mining.

Location

Christmas Island is located in the Indian Ocean, 10°30’ South, 105°40’ East, and approximately 2 600 km north-west of Perth, Western Australia. Christmas Island is 500 km south of the Indonesian capital Jakarta (Figure 1).




Figure 1: Location of Christmas Island (PRL 2014)



Legislative Framework

The legislative framework for Christmas Island is complex. The *Christmas Island Act 1958* outlines the governance arrangements for the island. Sections 8 and 8E of the Act make provision for the laws of Western Australia and the Commonwealth to apply on Christmas Island. The Minister lists selected Western Australian laws to be applied in the *Applied Laws (Implementation) Ordinance 1992*. The island is governed under Commonwealth legislation and administered by the Department of Infrastructure, Transport, Regional Development, Communication and the Arts (DITRDCA). Applied Western Australian laws are administered by the relevant Commonwealth Minister, by Commonwealth officers acting under ministerial delegations, or by State officers exercising delegated power and acting pursuant to inter-government service agreements under Section 8H of the act.

CIP operations (mining, processing, transport, and shiploading) are regulated through the Licence for Prescribed Premises issued under the *Environmental Protection Act 1986* (WA) (CI) (EP Act). The licence specifies monitoring and reporting requirements.

Clearing of native vegetation is regulated under provisions in the EP Act (Section 51) and under current governance arrangements any clearing within MCI 70/1A must be assessed and approved under this legislation.

Tenure

CIP was granted approval to re-mine and/or remove existing stockpiles of low-grade ore over approximately 2,000 hectares (14% of the Island) of previously cleared areas outside the National Park in 1990 by a lease with the Commonwealth issued under the *Lands Ordinance 1987*. Mining Lease 70/1 was granted to CIP by the Commonwealth in 1997 under the *Mining Act 1978* (WA) (CI) following approval under the *Environment Protection (Impact of Proposals) Act 1974* (Commonwealth) (EPIP Act). The approval for disturbance was carried over to the *Environment Protection and Biodiversity Conservation Act 1999* (Commonwealth) (EPBC Act) when the EPBC Act came into effect in 2000. In June 2013, MCI 70/1A was renewed until 2034 as part of a rewriting of the mining lease undertaken largely to clarify and update the royalty and conservation levy provisions and rehabilitation obligations.

MCI 70/1A is the principal lease for the Christmas Island operation. CPS 6323/1 resides completely within MCI 70/1A.

The lease outlines conditions that must be met by Phosphate Resources Ltd and includes the payment of royalties, rainforest protection, mine management and road maintenance and lease Covenant 8.3, which requires that the Lessee 'shall undertake to comply with all the requirements of the Environmental Management Plan' (EMP). The EMP is revised every five years, and the most recent EMP covers the 2024 – 2029 period (currently under assessment). The lease is administered by DITRDCA and expires in 2034.

Conservation Levy

The rehabilitation of former mining areas is undertaken as part of the Christmas Island Minesite to Forest Rehabilitation Program. The program is funded by a conservation levy paid to the Territory Administration by Phosphate Resources Limited as a provision of Mining Lease MCI 70/1A. The program is operated by Parks Australia under a Memorandum of Understanding between the Director of National Parks and the Territory Administration. The conservation levy is paid on all lease areas for all rock and dust exported.

PRL must pay the Conservation Levy quarterly to the Department for the purpose of rehabilitating cleared areas and other high priority conservation activities on Christmas Island as the Commonwealth sees fit.



Ownership

Phosphate Resources Limited (PRL) was established in 1990 to reopen the Christmas Island phosphate mine, first established in 1891, which had been shut down in 1987. PRL trades as Christmas Island Phosphates which is one of PRL's subsidiaries (Figure 2). CIP are the current operator of the phosphate mine which exports phosphate rock throughout the South East Asian market.




Figure 2: Corporate Structure

Proposed Amendment

On 23 June 2015, Christmas Island Phosphates was granted Clearing Permit 6323/1 which proposed to clear a total of 128.53 ha, over 23 separate areas for the purposes of phosphate mining. The existing CPS outlines the approved areas (shown in the plans Plan 6323/1a, Plan 6323/1b, Plan 6323/1c, Plan 6323/1d, and Plan 6323/1e). CIP does not seek to make any changes to the amount of native vegetation to be cleared in this application. All the proposed clearing within CPS 6323 will be undertaken in historically cleared areas within MCI 70/1A. No primary rainforest will be cleared for mining operations.

Areas within the Permit fall into the following categories:

- Stockpiles built by CIP;
- Areas dominated with fern vegetation which was previously considered a weed due to the low environmental values of these areas;
- Areas dominated by weed species;
- Areas with secondary regrowth;
- Areas which have been cleared in the past ten years but have had vegetation regeneration; and
- Areas with high value vegetation which will be protected.

Extension of Permit

Currently CPS 6323/1 expires 31 December 2024. PRL request the permit expiry aligns with MCI 70/1A tenure expiry 26 June 2034 to ensure continuation of operations.



Conditions

PRL request to remove Condition 6(c); *At least once in each 3-month period prior to the Weed Management Plan required in condition 6(b) being approved, the Permit Holder must remove or kill any weeds growing within areas cleared under this permit.* As the CIP Weed Management Plan was approved in May 2017 by the then Department of Infrastructure and Regional Development this sub condition is no longer current.

PRL also request to remove Condition 7; *Fauna Management - The Permit Holder must implement and adhere to the document 'A Management Plan for Christmas Island Pipistrelle in relation to Vegetation Clearing on Mining Leases, May 2015.'* The Christmas Island Pipistrelle Bat which was considered 'as possibly become extinct' in the 2014 application (2009) was transferred from Critically Endangered category to Extinct in 2021 - [Amendments to the EPBC Act list of threatened species - March 2021](#) (DCCEEW, 2021).

Clearing to Date

Approximately, 9.17 hectares has been cleared under CPS 6323/1 (Table 1; Figure 3; Figure 4).

Table 1: Clearing under CPS 6323/1

CPS 6323/1 Area	2015	2016	2017	2018	2019	2020	2021	2022	2023	Grand Total
101-MB4	0.0152	0.0002	0.0131					0.1996		0.2283
101-MB5	0.1829	0.0381								0.2209
101-STP17A	0.6090			0.1307						0.7398
106-MB10		0.0753		0.0520	0.0943					0.2217
106-MB13		0.0236								0.0236
133B-MB7			5.2253				2.3404	0.0824	0.0852	7.7335
Total	0.8072	0.1373	5.2385	0.1827	0.0943		2.3404	0.2821	0.0851	9.1680

Relinquished Areas


No MCI 70/1A areas within CPS 6323/1 have been relinquished by PRL to the underlying landholder to date.



**Phosphate Resources Limited
Site Plan - CIP Clearing To Dec 2023
CPS6323**



- Legend**
- CIP-Clearing_CPS6323_upto_240101
 - CIP-Clearing_CPS6323_upto_240101
 - Approved Clearing Permits
 - CPS_6323
 - Tenements
 - MCI_70_10
 - MCI_70_17
 - MCI_70_18
 - MCI_70_19
 - MCI70_1A_230315
 - GCI_701_Mar2021
 - GCI_702_Mar2021
 - RAMSAR_SITES_CI
 - National Park




Coordinate System: GDA 1994 MGA Zone 48
Projection: Transverse Mercator
Datum: GDA 1994
False Easting: 500,000.0000
False Northing: 10,000,000.0000
Central Meridian: 105.0000
Scale Factor: 0.9996
Latitude Of Origin: 0.0000
Units: Meter

Scale: 1:4,267


Figure 3: Clearing conducted under CPS 6323/1 (Map 1)



Phosphate Resources Limited
Site Plan - CIP Clearing To Dec 2023
CPS6323



- Legend**
- CIP-Clearing_CPS6323_upto_240101
 - CIP-Clearing_CPS6323_upto_240101
 - Approved Clearing Permits
 - CPS_6323
 - Tenements
 - MCI_70_10
 - MCI_70_17
 - MCI_70_18
 - MCI_70_19
 - MCI70_1A_230315
 - GCL_701_Mar2021
 - GCL_702_Mar2021
 - RAMSAR_SITES_CI
 - National Park



Coordinate System: GDA 1994 MGA Zone 48
Projection: Transverse Mercator
Datum: GDA 1994
False Easting: 500,000,000
False Northing: 10,000,000,000
Central Meridian: 105,0000
Scale Factor: 0.9996
Latitude Of Origin: 0.0000
Units: Meter

Scale: 1:2,731

Figure 4: Clearing conducted under CPS 6323/1 (Map 2)



Existing Environment

Climate

Christmas Island lies on the southern edge of the inter-tropical convergence zone and the climate is dominated by a low-pressure trough that seasonally circles the equator. The Island has a tropical monsoonal climate with distinct wet and dry seasons and little seasonal variation in temperature. The dry season (May to November) is dominated by low and sporadic rainfall with consistent south-east trade winds. The wet season generally occurs from December to April with the island receiving most of its rainfall during this period. Christmas Island's average annual rainfall over 30 years is around 2 000mm but may vary considerably from year to year; with the island's lowest recorded annual rainfall being 1 067mm in 1987, and the highest 5 121mm in 2016 (BOM, 2023). Temperatures remain relatively uniform throughout the year, with an average daily maximum of 28°C in March/ April and average daily minimum of 22°C in August/ September. The Island has high humidity (80-90%) throughout the year.

During the wet season, Christmas Island is subject to the influence of north-west monsoons, which typically cause high swells and high winds, with gusts of over 100 km/h. Many cyclones have passed nearby the Island, resulting in strong winds and heavy rainfall on the Island. The last cyclone to cause any real damage to the island was Cyclone Gillian, which caused significant damage to rainforest communities in 2014, with trees blown over and foliage stripped from many areas.

The island has distinct microclimates due to the island's geography, with rainfall measurements indicating that there are significant variations in rainfall (Falkland, 1999) across the island. Average rainfall at Rocky Point (Settlement), Jedda Cave and South Point (1 931, 2 375, & 1 907mm respectively) for example shows the significant variation across the Island. Daily evaporation generally exceeds rainfall between July and October. In the six-month period preceding and during the survey, covering the dry season (May 2023 to November 2023), Christmas Island received significantly less than long-term average for rainfall, with temperatures slightly above the mean average (Figure 5).

The Christmas Island Aero weather station (Station Number 200790) has operated from 1972 to the current date and is situated on the plateau near the airport at an elevation on 261m.




Figure 5: Rainfall and Temperature 2023 at Christmas Island Aero (BoM 2023)



Hydrology

A major feature of the Christmas Island geomorphology is the lack of surface drainage. Rainfall mostly infiltrates the land surface and is utilised by plants, contributes to soil water stores or recharges to groundwater. There is therefore no significant surface drainage network except down gradient of springs that arise at the interface between limestone and basalt formations (Hollingsworth, 2003).

Christmas Island's soils are generally highly permeable and there is consequently little runoff or erosion. In the Wet Season when the soils are saturated, runoff can occur during heavy rainfall providing some risk of erosion and sedimentation. However, given the high natural infiltration rates the risk of erosion and sedimentation is generally localised to compacted areas such as roads and stockpile pads. Infiltration tests by Puhlovich et al. (2003) indicate that soil infiltration rates are typically substantially higher than hourly rainfall intensities.

Perennial (permanent) surface aquatic habitats (freshwater) on Christmas Island are limited to a number of spring-fed streams found along coastal or sloping areas of the Island. Hosnie's Spring and The Dales are both listed as a Wetland of International Importance under the Convention on Wetlands of International Importance, Water Fowl Habitat 1971 (known as the 'Ramsar Convention', with the sites referred to as Ramsar sites within internal documents) and are listed in the Directory of Important Wetlands in Australia.

Hydrogeology

There are three key hydrogeological units on the Island; shallow, residual soils, which overlie fractured, unconfined – semi-confined aquifers within the karstic limestone rocks, which in turn overlie relatively impermeable volcanic basement rocks (Puhlovich et al., 2003). Groundwater levels on the Island are reflected by the location of the unconfined water table within the karst limestone aquifers. Limestone aquifers can be recharged when rainfall permeates through the soil zone into the underlying aquifers or by direct runoff of rainfall into karst features such as dolines and sinkholes that occur across the Island (Puhlovich et al., 2003). Assessments suggest that approximately half of all incident rainfall passes through the soil zone and recharges the underlying limestone aquifers (Hollingsworth, 2003; Falkland, 1999). Groundwater discharge occurs at surface springs such as Hosnie's Spring and offshore springs such as those found at Flying Fish Cove (Puhlovich et al., 2003). The complex behaviour and extent of weathered/fractured rock aquifers on the Island are not well understood.

Geology, Soils and Landforms

Christmas Island is one of a series of submarine seamounts that rise above the 5 500 m deep abyssal areas of the West Australian Basin. At the core of the island are volcanic rocks, mainly composed of basalt with a layer of limestone generally covering these volcanic rocks, with occasionally outcrops, particularly along the present coastline. A series of geological uplifts and successive layering of coral reefs over the basalt core of the island have led to the development of new cliffs and terraces from the ocean, forming stepped terraces and inland cliffs.

The oldest (Upper Eocene) limestones are found near the island summit, 330 m above sea level and the youngest (Pliocene) near present sea level. The limestone is mixed with dolomite sediments, basalts, and tuffs. A layer of phosphate-rich soil material covers the limestone over about half of the island. The island is characterised by sea cliffs that rise via a series of terraces to a central plateau. The shoreline is dominated by cliffs and extensive shore platforms with a few small beaches and Flying Fish Cove which has a relatively large beach and shallow platform being the only safe harbour for much of the year. The island's natural landscape is dominated by karstic surface landforms and cave systems (Grimes, 2001). Figure 6 shows a cross section of Christmas Island (exaggerated vertical scale) showing volcanic core, limestone capping, water flow lines and typical cave development (source Grimes 2001).

Terrace areas on the Island reflect the height above sea level in different geologic periods. Renewed vulcanism and a series of geological uplifts at different periods have resulted in a tiered effect. The oldest limestones



near the peak of the Island formed during the Eocene period (Grimes, 2001). Most of the Island's limestone deposits were formed during the Tertiary (late Oligocene to mid Miocene age), with the youngest limestones deposited on the lowest terrace in the late Quaternary (Grimes, 2001).

The limestone is mixed with dolomite sediments, basalts, and tuffs. A layer of phosphate-rich soil material covers the limestone over about half of the Island. Marine sediments and guano deposition have formed the Island's phosphatic soils.




Figure 6: Diagrammatic cross-section of Christmas Island (Grimes 2001)

Land Use

There are a range of current land uses on Christmas Island with the largest land use by area being nature conservation in the form of the Christmas Island National Park (CINP), which occupies 63% of the island. Other land uses include residential housing, tourism, recreation, transport and the provision of utility services (Figure 7). Numerous businesses and services have been developed on the island to support the phosphate mine, detention centre, tourism industry and the island population, including construction, retail/hospitality, training/education, Government services, maritime services, agriculture, sport/recreation, maintenance, arts, and the airport.




Figure 7: Land Tenure and Uses (Environment Australia 2014, Geoscience Australia 2014)




Figure 8: Land Tenure Areas and Assigned Regions

Land Systems

The national and regional planning framework for the systematic development of a comprehensive, adequate and representative 'CAR' National Reserve System is provided by the Interim Biogeographic Regionalisation for Australia (IBRA). IBRA was developed in 1993-94 and is endorsed by all levels of government as a key tool for identifying land for conservation under Australia's Strategy for the National Reserve System 2009-2030. The nationally agreed regionalisation was published in Thackway and Cresswell 1995, An Interim Biogeographic Regionalisation for Australia: a framework for establishing the national system of reserves.

As per the Bioregion map for IBRA, managed by DCCEW, Christmas Island is in the Interim Biogeographic Regionalisation of Australia (IBRA7) as part of ITI (Indian Tropical Islands, ITI01) (Australia's Bioregions - DCCEWc 2023).

Environmental Sensitive Areas

An Environmentally Sensitive Area (ESA) means an area declared in Regulation 6 of the *Environmental Protection (Clearing of Native Vegetation) Regulations 2004* to be an Environmentally Sensitive Area. These areas are protected under legislation (EP Act).

The ESA on Christmas Island covers areas including:

- The National Heritage Listed "Christmas Island Natural Areas"
- The RAMSAR listed nationally important wetlands Hosnie's Springs and The Dales

The "Christmas Island Natural Areas" ESA on Christmas Island was put in place under the EPBC Act but has subsequently been repealed; however, the gazetted *Environmental Protection (Clearing of Native Vegetation)*



Regulations 2004 ruling is still in place. The “Christmas Island Natural Areas” is applicable across a large part of the island ruled to be “natural”, including previously cleared areas, and areas immediately adjacent to CIP infrastructure. This prohibits CIP conducting any works which would ordinarily be allowed as exemptions for clearing native vegetation for certain reasons (i.e. maintenance requirements). A small portion of the CPS 6323/1 area falls outside of the ESA in the Dogs Head Region (*Figure 9*).

CPS 6323/1 does not intersect the RAMSAR listed wetlands however the Dales area is adjacent to the sub-region in North West Point (*Figure 9*).

A Protected Matters Search using the DCCEEW tool did not identify any additional areas requiring specific management (DCCEEWc, 2024).

Asplenium listeri, *Tectaria devexa* var. *minor* and *Pneumatopteris truncata* are unlikely to be found in the proposed clearing areas (see Flora of Conservation Significance).



**Phosphate Resources Limited
Environmentally Sensitive Area
Christmas Island**




Figure 9: Map of ESA, RAMSAR and CPS 6323/1



Flora and Vegetation

The flora of Christmas Island has very strong taxonomic relationships with those of the Indo-Malayan and Malesian regions. Some species (e.g. *Terminalia catappa*, *Inocarpus fagifer* and *Gyrocarpus americanus*) are more widely distributed, and largely unremarkable, tropical Indo-Pacific littoral flora that extends as far east as the islands of Polynesia. Despite its isolated location, estimated 10-million-year history above sea level, and diverse 'high-island' landscape, only eighteen species are known to be endemic (denoted by §) to Christmas Island.

The flora comprises about 420 species of vascular plants with approximately 242 species indigenous to the island and a further 177 species introduced since human occupation (Director of National Parks, 2014b). The general taxonomic status and affinities of the flora are relatively well known but detailed records of population distributions across the island (and its range of habitats) are limited for many species. Some additional species of ferns, herbs and shrubs continue to be added to the flora from further collections, especially from poorly accessible terrace areas on the island.

Christmas Island National Park (CINP), which covers some 63% of the island, is assigned to IUCN reserve category II, most of which is uncleared primary rainforest; approximately 25% of the island's original vegetation has been cleared for mining and infrastructure.

Three features of the ecology of the island's native vegetation are notable:

- The occurrence of many of the widespread Indo-Malesian species in habitats that would be considered extremely atypical elsewhere in their natural ranges, and associated with this, the exceptionally large stature of some of these species.
- The low diversity of canopy and sub-canopy species and the lack of structural complexity (e.g. relatively poor development of robust woody vines and rattans, the absence of aroids and of gingers in the understorey) in the Island's rainforests; and the very low diversity and lack of speciation amongst plant genera that elsewhere in the region is characteristic of early successional, and frequently disturbed, rainforest environments (e.g. *Macaranga*, *Claoxylon* and *Pipturus*).
- The geology, geomorphology and climate on Christmas Island create the biophysical environment and constraints for the vegetation communities. These factors determine the soil nutrient status, the seasonal availability of moisture and the degree of exposure to wind, which in turn control the distribution, structure and functioning of the natural vegetation.

Vegetation Types

Vegetation mapping of the Island was initially undertaken by Mitchell (1985) for the Australian Nature Conservation Agency. This mapping had limited use due to its broad scale and spatial inaccuracy. Flora of Australia Volume 50 (Du Puy, 1993a) listed eight vegetation types for the Island.

An island wide vegetation mapping project was undertaken from 2011 to 2014 to attempt to map vegetation with better spatial accuracy, to determine height categories and to apply these consistently across the Island. The process included a Light Detection and Ranging (LiDAR) survey, review of historic aerial photography which captured past clearing, and ground truthing. Additional categories were added to include wetland vegetation and regrowth in cleared areas. The Christmas Island Vegetation and Clearing Map was developed through a collaborative project by Geoscience Australia, CIP, CINP and the Commonwealth Department of the Environment (Geoscience Australia, 2014).



The map classified the full extent of Christmas Island into vegetation and land cover classes (Figure 10 - Figure 13; Table 2), though boundaries are not always perfectly geographically accurate. Flora of Australia's vegetation types, which are still referred to in some documents, were recategorized to the following:

- 'Primary rainforest' - closed canopy evergreen forest,
- 'Marginal rainforest' - semi-deciduous forest,
- 'Areas with surface water' - perennial wetland forest,
- 'Open forest, scrubby forest and vine forest' and 'inland cliffs' - semi-deciduous scrub,
- 'Coastal fringe' and 'shore cliffs and spray zone' - coastal fringe vegetation, and
- 'Mined areas' - rehabilitation, regrowth and weed dominated veg and pioneer regrowth.

There are seven major structural types of secondary vegetation (i.e. vegetation that has established in areas that have been cleared of the original natural forest) on Christmas Island. The occurrence of these secondary vegetation types in specific disturbed areas on Christmas Island reflects the influence of four main factors:

- The areal extent and the severity /intensity of the original disturbance (especially whether the soil profile has been disrupted or removed),
- Proximity to the nearest forest boundaries and/ or to weed-infested areas,
- The subsequent disturbance or management history of the area (especially the frequency and intensity of further disturbances including fire), and
- The time since the last major disturbance occurred.

Whether or not the soil profile has been removed or disturbed, the proximity to forest and/or to weed infested areas and the occurrence of further disturbance are the major environmental determinants of secondary vegetation type and successional pathways. There are four main successional pathways, as defined by Bennett in 2010:

- Arrested successions dominated by ferns,
- Stagnant successions dominated by thickets of exotic shrubs and vines,
- Reconstructive successions dominated by *Macaranga tanarius*, and
- Retrogressive successions leading to fernlands.

There are two vegetation types within CPS 6323/1:

- **Regrowth; and**
- **Weed dominated vegetation and pioneer regrowth.**



Table 2: Vegetation Types of Christmas Island (Du Puy 1993, and Geoscience Australia 2014)

Vegetation Type	Geoscience Australia			Indicator Species
	Level 1	Level 2	Description	
Primary Rainforest	Closed canopy evergreen forest	Closed canopy evergreen forest (tall or moderate)	Generally found on the plateau and terraces, with a closed uneven canopy up to 40 m in height. Some trees emerge up to 10 m above the canopy. Often supports ferns and orchids, young palms and lilies in the understory.	<i>Bolbitis heteroclita</i> , <i>Syzygium nervosum</i> , <i>Hernandia ovigera</i> , <i>Planchonella nitida</i> , <i>Pisonia umbellifera</i> , <i>Corymborkis veratrifolia</i> , <i>Ehretia javanica</i>
Marginal Rainforest	Semi-deciduous forest	Semi-deciduous forest	Generally found on the slopes and terraces down to the coast - and some plateau areas. Higher occurrence of semi-deciduous trees compared to Closed Canopy Evergreen, which lose a portion of leaves during the dry season. Tree height generally 10-25m	<i>Terminalia</i> , <i>Gyrocarpus</i> , <i>Erythrina variegata</i> , <i>Premna serratifolia</i> , <i>Pisonia grandis</i> , <i>Ochrosia ackeringae</i>
Inland Cliffs and Open Forest, scrubby forest and vine forest	Semi-deciduous scrub	Semi-deciduous scrub	Found on the terraces, steep slopes and inland cliffs. Semi-deciduous canopy with vines and shrub understory Tree height generally <10m.	<i>Colubrina pedunculata</i> ^S <i>Canavalia cathartica</i> , <i>Carmona retusa</i> , <i>Cycads</i>
Areas with Surface Water	Perennial wetland forest	<i>Inocarpus fagifer</i> dominant	Areas of fresh water runoff on the lower terraces dominated by <i>Inocarpus fagifer</i> .	<i>Inocarpus fagifer</i>
		<i>Hibiscus tiliaceus</i> dominant	Areas of fresh water runoff on the shore terrace dominated by <i>Hibiscus tiliaceus</i> .	<i>Hibiscus tiliaceus</i>
		<i>Bruguiera</i> dominant	A single patch of vegetation dominated by <i>Bruguiera</i> at Hosnie's Spring. Occurring in an area of fresh water runoff on the shore terrace.	<i>Bruguiera gymnorhiza</i>
Coastal Fringe OR Shore cliffs and spray zone, and Mined Areas	Coastal fringe vegetation	Coastal herbland	Found between the coastal scrub and coastal cliffs in exposed areas. Class is dominated by low-lying herbs, sedges and grasses.	<i>Portulaca tuberosa</i> , <i>Ischaemum nativitatis</i> ^S , <i>Oplismenus compositus</i> , <i>Sporobolus virginicus</i>
		Coastal shrubland	Dense salt-tolerant vegetation growing between the coastal herbland and the terrace cliffs.	<i>Pandanus christmatensis</i> ^S , <i>Scaevola</i> , <i>Pemphis</i> , <i>Argusia</i>



Vegetation Type	Geoscience Australia			Indicator Species	
	Level 1	Level 2	Description		
Coastal Fringe OR Shore cliffs and spray zone, and Mined Areas				<i>argentea</i> , <i>Cordia cordata</i> , <i>Guettarda</i>	
	Rehabilitation	Rehabilitation	Areas where forest rehabilitation has taken place. The standard of forest varies depending on the type of rehabilitation completed, species planted and management regime.	A mix of up to 30 native tree species when initially planted, dependent on characteristic of the site and year of rehabilitation. <i>Macaranga</i> , <i>Dysoxylum</i> , <i>Calophyllum</i> , <i>Tristiropsis</i>	
	Regrowth	Regrowth	Generally, well developed regrowth vegetation over 5 m mean tree height. May include some introduced or weed species.	Various species – dependent on adjacent vegetation	
	Weed dominated vegetation and pioneer regrowth		* <i>Leucaena leucocephala</i>	Monoculture of * <i>Leucaena leucocephala</i> . Often occurring as regrowth in previously cleared areas.	* <i>Leucaena leucocephala</i>
			Fern field	Expanse of low-lying ferns often growing on limestone pinnacles.	<i>Nephrolepis biserrata</i> , <i>Microsorium scolopendria</i> , <i>Psilotum nudum</i>
			Mixed weed and pioneer regrowth	Regrowth vegetation with a mean tree height of <5m. Can vary between native and introduced species depending on the location and time since clearing. Tends to have a higher occurrence of weed species compared to the 'Regrowth' category.	* <i>Muntingia calabura</i> , * <i>Psidium</i> sp. (<i>Guava</i>), * <i>Mimosa</i> , <i>Passionfruit</i> , <i>Macaranga</i>

Note: Endemic flora taxa are denoted by §, foreign introduced weed species are denoted by*




Figure 10 Geoscience 2014 Classified Vegetation within Northern Region




Figure 11 Geoscience 2014 Classified Vegetation within Central Region




Figure 12 Geoscience 2014 Classified Vegetation within Southern Region




Figure 13 Geoscience 2014 Classified Vegetation within Western Region



Vegetation Condition

A field reconnaissance survey was undertaken by Mr Leonard Stapp (BSc Environmental Management – major in Conservation Biology) of Christmas Island Environmental Services to determine if there was significant change to information provided for the original CPS application. Leonard has extensive experience in botanical surveying, as well as considerable local knowledge of the flora and fauna of Christmas Island, as he has been providing environmental services such as botanical field surveys and assessments, species identification, conduct of monitoring operations and weed management programs for Parks Australia on Christmas Island, working as a Field Supervisor/ Technical Coordinator; and as sole proprietor at Christmas Island Environmental Services. Leonard also has widespread experience working in Environmental Advisor and Bushland Regeneration roles in mainland WA.

The reconnaissance survey was conducted ranking sites vegetation conditions scores based on the Keighery Condition Scale (Keighery, B.J. 1994), with some allowances and modifications made to accommodate the unique tropical environment on Christmas Island (Table 3).

Table 3: Modified Vegetation Condition Score (Keighery, B.J. 1994)

Condition Rating	Keighery Definition	Christmas Island Definition
Pristine (1)	Pristine or nearly so, no obvious signs of disturbance	Mature, undisturbed rainforest or very advanced secondary regrowth. Disturbance is limited to cyclone damage. Climax species dominate and full structural complexity is present with epiphytic orchids and ferns, terrestrial orchids and ferns, § <i>Pandanus</i> , pals, buttressing and woody lianas
Excellent (2)	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species	Little to no weed species and a largely intact landform with few signs of disturbance. The vegetation structure is beginning to develop the full complexity and diversity expected in rainforest, with climax species present and a range of features included such as buttressing, palms, ferns, epiphytes and orchids, but the canopy is likely to be more open than in undisturbed rainforest
Very Good (3)	Vegetation structure altered, obvious signs of disturbance. For example, disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing	Low diversity vegetation with few signs of disturbance or moderately diverse vegetation on a disturbed landform, a canopy dominated by pioneer species through recruitment of climax species may be evident, an open canopy or poor structural complexity. Few weeds present or weeds are limited to less aggressive species.
Good (4)	Vegetation structure significantly altered by very obvious signs of multiple disturbance. Retains basic vegetation structure or ability to regenerate it. For example, disturbance to vegetation structure caused by very frequent fires,	Multiple signs of disturbance or with low diversity and no structural complexity. Typically displays little to no recruitment of climax tree species. Numerous weeds present or a dense



Condition Rating	Keighery Definition	Christmas Island Definition
	the presence of some very aggressive weeds at high density, partial clearing, dieback and grazing	understory dominated by one or two native fern species
Degraded (5)	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. For example, disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing	Vegetation at very early successional stages with a canopy of one to two species and an understory dominated by ferns or weeds, typically adjacent to completely degraded land. May have the potential to develop into better quality vegetation as the canopy develops if weeds are managed
Completely degraded (6)	The structure of the vegetation is no longer intact, and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees or shrubs	The structures of primary rainforest are no longer present and native species are limited to only a few species, for example weed dominated shrubland with sparse emergent <i>Macaranga</i> or a near monoculture <i>Nephrolepis bierrata</i> fernland. The landscape may be highly disturbed, often with significantly reduced soil and pinnacles exposed. Predominantly lacking a native canopy and with little to no potential to develop one due to dense groundcover of shallow soil

Data collected was compared against the vegetation data collated for the original application to identify any areas of significant change. No significant changes in vegetation type or condition category were identified.



Figure 14 (left) Example of Classified Category 6 Vegetation (Field 133B -MB6); (right) Example of Classified Category 4 Vegetation (Field 133B - MB6)



Table 4: Vegetation Condition and Comments

Location	Vegetation Condition Rating/s	Total Ha	2023 Ground-truthing Assessment	Key Issues in 2014 application
100-SPWMB2	6	2.87	Pinnacles Fern fields Old rehab field	Near to known Tectaria <i>devexa</i> population
100-SPWMB3	5	1.33	Pinnacle field	Near to known Tectaria <i>devexa</i> population
101-MB4	5, 6	1.42	Fern field Existing cleared areas	Small section adjacent to National Park, medium-high Red Crab densities
101-MB5	6	0.32	Abbotts Booby nest out of clearing permit area by approx. 50m Existing cleared areas	Adjacent to National Park
101-MB6	6	0.67	Fern fields	Adjacent to National Park
101-STP17A	5, 6	1.55	Primary regen Pinnacles Ravine Fern field	Nil
106-MB10	4, 5	1.38	Edge to existing cleared areas, Tracks Narrow strip of good forest	Tall regrowth. Abbott's nest site to the south-west of the site so no likely turbulence issues
106-MB11	5, 6	2.01	Fern fields Drainage sump	Visual amenity, Abbott's nest site 24m to the east, across the road so no turbulence issues
106-MB13	4, 5	2.41	Rocky ground, some areas weed free Edge to existing cleared areas in some parts	Adjacent to National Park
106-MB14a	3, 4, 5, 6	8.51	Fern fields Young trees Edge to fern fields Edge to existing cleared areas Narrow strip of good forest	Adjacent to National Park. Abbott's nest sites < 300m to the north-west but the change in
106-MB14b	3, 4, 5, 6	5.68	Areas with no weeds but open canopy	vegetation height is



Location	Vegetation Condition Rating/s	Total Ha	2023 Ground-truthing Assessment	Key Issues in 2014 application
			Was assessed as one location 106-MB14 in original application, but approved as two locations	unlikely to be significant
116-STP23D	4, 5, 6	2.85	Red crabs in stockpiles, but impenetrable in areas due to fern fields Fern fields	Adjacent to National Park. Abbott's nest site 63m to the east so no turbulence issues
133A-MB8	5, 6	1.43	Fern fields Cordia thickets Red crabs	Near Spleenwort population
133B-MB2	4, 5	1.84	Red crabs, Robber crab Abbotts Booby nest out of clearing permit area by approx. 20m	Adjacent to National Park, medium Red Crab densities
133B-MB3	1, 3, 4, 5	2.1	Red crabs	Medium-high Red Crab densities. Some tall regrowth
133B-MB4	3, 5, 6	1.97	Red crabs Weedy edges, fern fields	Some tall regrowth
133B-MB5	4, 5, 6	8.88	Soil stockpiles Edge of stockpile Edge of fern fields Road buffers Red crabs	Visual amenity. Adjacent to Primary Rainforest
133B-MB6	3, 4, 5, 6	14.11	Abbotts Booby nest out of clearing permit area by approx. 50m Fern fields Edge of low – medium density regen Edge of medium density to fern field Soil stockpiles Red crabs	Adjacent to Primary rainforest. Near <i>Tectaria devexa</i> population
133B-MB7	3, 4, 5, 6	31.5	Codia thickets Edge of existing cleared areas Soil stockpiles Edge of stockpiles looking over fern fields Open ground Fern fields Existing cleared areas	Visual amenity



Location	Vegetation Condition Rating/s	Total Ha	2023 Ground-truthing Assessment	Key Issues in 2014 application
			Soil stockpile to cleared area Red crabs Flying foxes	
133B-MB8	3, 4, 5, 6	19.28	Fern fields	Visual amenity
133B-MB9	3, 4, 5, 6	9.85	Sparse Cordia Thicket Edge of fern field	Adjacent to National Park
139-STP25B	6	1.28	Fern field	Adjacent to National Park. Near Jewel Orchid population
140-MB3	5, 6	5.45		Adjacent to National Park, and The Dales RAMSAR site boundary, but not in a wetland area. High densities of Red Crabs and Pipistrelle records
Total		128.53		



Significant Changes

No locations were identified as having significant changes to the quality of vegetation against the 2014 application. Where minor changes were identified in vegetation it was generally accepted that changes had been to the degradation of the vegetation or incursion of weeds.

Flora of Conservation Significance

A desktop assessment of the likelihood of EPBC listed fauna species occurring in the Survey Area, either resident or transient, was made on the basis of their breeding ecology, habitat preferences and known location records held by Parks Australia, and consultants who have previously conducted surveys for clearing permits including Range to Reef Environmental and JBS&G. An assessment of the impact of the proposed clearing has been made against the clearing principles (Table 7) in later in this report, together with other notes on impact and proposed mitigation.

Likelihood of presence of conservation significant species within proposed clearing areas was estimated and the potential for significant impact to the species was examined. The assessment suggested that the risk posed to conservation significant species from the proposed mining activity would be low and not significant in the context of species conservation. The general absence of natural habitat and low biodiversity value of the area suggested limited habitat important for the lifecycle of listed species.

The Draft Christmas Island Biodiversity Conservation Plan (2014) lists five criteria for determining the significance of fauna on the island and identified 27 species as significant. For the purposes of the plan, a significant species was defined as a native terrestrial species which meets one or more of the following criteria:

- A species listed (or under consideration for listing) as threatened under the EPBC Act,
- A species with an important or 'keystone' role in maintaining the island's ecology or which characterises a significant ecosystem,
- Species which are of conservation concern (those which have a substantial decline on Christmas Island) but not listed as threatened,
- An endemic vertebrate, and/ or
- A species of international conservation significance with strong community support for its conservation

Table 6 includes the species listed in the Significant Christmas Island Species, listed in the Conservation Plan draft, and is updated with the current list of Christmas Island species found on the DCCEEW EPBC Act List of Threatened Species Databases (Flora and Fauna) (DCCEEWc, 2023).

Three species which occur on Christmas Island are listed as threatened species under the *Environmental Protection and Biodiversity Conservation Act 1999* (Table 5).



Table 5: Conservation Significant Flora Species

Genus (Family)	Conservation Status (listed under EPBC Act)	Habitat	Description	Distribution
<i>Asplenium listeri</i> (Aspleniaceae)	CR	Limestone rock crevices in dry, exposed areas	A lithophytic fern with short erect fronds, 3.5 – 9 cm long, which grow in a crown	Christmas Island
<i>Tectaria devexa</i> <i>var. minor</i> (Dryopteridaceae)	EN	Primary rainforest (tall and largely undisturbed), above 80 metres elevation; both in deeper soils and as a lithophyte (on mossy pinnacles at the base of a slope, a wet site).	A small, tufted, terrestrial fern with pale green fronds.	Christmas Island and Sri Lanka.
<i>Pneumatopteris truncata</i> (Thelypteridaceae)	CR	Permanently moist sites in semi-deciduous closed forest.	A large terrestrial fern with an erect rhizome and fronds growing in a crown to 120 cm long. The fronds have aerophores (respiratory structures) at the base of the pinnae.	Fragmented distribution over Asia and Malaysia and two sites on Christmas Island.



See Figures 15 – 22 for the known locations of listed flora near the boundaries of CPS6323/1.

No flora species of conservation significance is known to occur within CPS 6323/1.




PHOSPHATE
RESOURCES

Phosphate Resources Limited
Northern Part 1
CPS6323






Phosphate Resources Limited
Northern Part 1
CPS6323






PHOSPHATE
RESOURCES

Phosphate Resources Limited
Central Region Part 1
CPS6323






Phosphate Resources Limited
Central Region Part 2
CPS6323






**Phosphate Resources Limited
Central Region Part 3
CPS6323**






Phosphate Resources Limited
Southern Region Part 1
CPS6323





PHOSPHATE
RESOURCES


Phosphate Resources Limited
Southern Part 2
CPS6323





PHOSPHATE
RESOURCES

Phosphate Resources Limited
North West Point Region
CPS6323





***Asplenium listeri* (Christmas Island Spleenwort)**

The Christmas Island Spleenwort occurs in limestone rock crevices in dry, exposed areas on Christmas Island and hence is not associated with proposed clearing sites. The likelihood of Christmas Island Spleenwort being impacted by the proposed clearing is negligible due to an absence of suitable habitat in proposed clearing areas.



Figure 23: Christmas Island Spleenwort (*Asplenium listeri*⁸)

Tectaria devexa* var. *minor

Tectaria devexa var. *minor* grows colonially, mainly on the plateau, in primary rainforest (tall and largely undisturbed), above 80 metres elevation; both in deeper soils and as a lithophyte (on mossy pinnacles at the base of a slope, a wet site). Ten populations of *Tectaria devexa* have been identified at Christmas Island, with the majority of these occurring in the National Park. No known populations of *Tectaria devexa* have been identified at proposed clearing locations from previous surveys or the 2023 ground truthing, though additional risk-based management measures may be proposed to mitigate against the risk of *Tectaria devexa* being disturbed.



Figure 24: *Tectaria devexa* var. *minor*

Pneumatopteris truncata

Pneumatopteris truncata is known from two sites at Christmas Island (Hugh's Dale and Blowholes Ravine), and has a specialised habitat associated with groundwater seepage (Reddell et al, 2020). Neither location is within or close to the proposed clearing areas, which all occur within regrowth closed canopy evergreen forest, excepting one site which is on the boundary of evergreen and semi-deciduous forest. *Pneumatopteris truncata* grows in permanently moist sites in semi-deciduous closed forest. Due to the small number of known



populations, its habitat requirements are not known. The likelihood of *Pneumatopteris truncata* being impacted by the proposed clearing is negligible due to the proposed clearing areas occurring on the plateau in mining regrowth.



Figure 25: *Pneumatopteris truncata*

Fauna and Habitat

EPBC Act listed species, red crabs and robber crabs were identified as those prioritised for assessment of potential impact. Evidence which was used to identify potential habitats included assessment against known breeding habitats and locations, and preferred habitats. This was compared against findings from the original application. In addition, whilst conducting the flora surveys, the areas were also surveyed for evidence of fauna habitat, particularly Abbotts's Booby. No locations were identified as being affected by the proposed clearing areas.



Table 6: Fauna of Conservation Significance

Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
Birds				
<i>Papasula abbotti</i> Abbott's Booby	EN	A large sea bird about 80 cm from beak to tail, with off-white plumage; black panda-like eye patches; black wings, flank marks and tails; and black outer ends to its blue webbed feet. Males have pale grey bills, tipped black and females have black-tipped pink bills (Pizzey & Knight 1997).	<p>Endemic to Christmas Island. Most nests for this species are situated on the central and western areas of the Island (Department of National Parks, 2016). This species prefers to nest on the lee side of slopes and gullies, with a clear area below and immediately downwind to facilitate take-off and landing. The original application in 2014 had 4 ha removed from the initial application due to proximity to Abbotts Booby breeding areas.</p> <p>Tall rainforest of the island is the only remaining breeding habitat of the Abbott's Booby (<i>Papasula abbotti</i>) which is listed as Endangered under the EPBC Act 1999. Abbott's Booby is a marine species which spends much of its time at sea but needs to come ashore to breed. It nests in tall rainforest trees in the western, central and northern portions of Christmas Island. Most nest trees are associated with uneven terrain created by gullies, hillsides or cliffs and are predominantly in uneven canopy containing emergent trees. It prefers nest sites with a clear area below and immediately downwind to facilitate take-off and landing. South-east trade winds prevail between April and November, thus emergent trees which can be approached from the north-west are most often used as nest sites (Director of National Parks, 2014).</p> <p>Nest sites for Abbott's Booby have been mapped in island wide surveys, and ground truthing activities also noted any Abbotts bobby nest locations. Three sites outside of the clearing permit area have been identified</p>	Possible; however critical nesting habitat unlikely to be impacted



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
			as containing nesting sites for the Abbott's booby, but as no clearing of primary rainforest is proposed it is unlikely that the Abbott's booby will be present in areas to be cleared.	
<i>Chalcophaps indica natalis</i> Christmas Island Emerald Dove	EN	A small pigeon about 23 to 26 cm long, with a mass of about 80 to 135 g. It is mostly purplish-brown or red-brown with iridescent dark-green wings, dark brown irises, an orange bill (with a darker base) and red or purple-red legs and feet. The male differs from the female in having a grey crown, white forehead, white stripe above each eye, narrow white line across each shoulder, and dark grey (rather than red-brown) lower back, rump and tail (Department of the Environment, 2014).	<p>Whilst confined to Christmas Island, this species is widespread and common in areas of rainforest (Director of National Parks, 2014b).</p> <p>The Christmas Island Emerald Dove is endemic to Christmas Island. It is estimated that more than 60% of the population occurs within the boundaries of the National Park. It is most common in tall closed evergreen rainforest and open semi-deciduous rainforest, especially on the terraces that surround the central plateau of the island, but is also regularly observed in deciduous scrub, disturbed vegetation such as thickets of weeds and secondary regrowth (including areas dominated by <i>Muntingia calabura</i>), and settled areas (on lawns, in gardens and around houses), and on forest tracks (Department of the Environment, 2014).</p> <p>It is likely that the Christmas Island Emerald Dove will be present in areas proposed for clearing, however, the species is highly mobile, capable of relocating during the clearing process, and with better than 60% of the population estimated to be within the National Park, the proposed clearing is not considered a significant threat to the species.</p>	Likely; however critical nesting habitat unlikely to be impacted
<i>Fregata andrewsi</i> Christmas Island frigatebird	EN	A very large seabird with a mainly black body, a glossy green sheen to the feathers of its head and	Endemic to Christmas Island. Known to fly vast distances to feed, and can be seen in Indonesia, however	Unlikely



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
		<p>back, and varying patches of white on the underbody. It has slender, long wings, a deeply forked tail and a long bill with a hooked tip. Its legs are dull pink, and its feet are black above with pale soles. Adult males have a large, red gular (throat) pouch which can be inflated (Department of the Environment, 2014). It is the rarest endemic seabird on Christmas Island. (DCCEE, 2024a).</p>	<p>Christmas Island is the only place in the world where they breed (Department of National Parks, 2016),</p> <p>The Christmas Island Frigatebird is a pelagic, marine species that frequents tropical waters of the Indian Ocean but breeds only on Christmas Island. Breeding colonies are currently confined to the terrace forests of Christmas Island. The three recorded breeding colonies are known as the Golf Course, Cemetery and Margaret Beaches colonies. Tree species used for nesting include Beach Almond (<i>Terminalia catappa</i>), Stinkwood (<i>Celtis timorensis</i>), Strangler Fig (<i>Ficus species</i>), Propeller tree (<i>Gyrocarpus americanus</i>) and Pongamia (<i>Pongamia pinnata</i>). Multiple pairs will nest in the same tree, sometimes less than 1 m apart. These nest tree species occur right around the island, and yet the species nests only in a small area of the island (DCCEE, 2024a).</p> <p>It is unlikely that the Christmas Island Frigatebird will be present in areas proposed for clearing which are confined to the upper terraces.</p>	
<p><i>Accipiter hiogaster natalis</i> Christmas Island Goshawk</p>	<p>EN</p>	<p>A large hawk with a dark grey head and upperparts, mostly rufous underparts with fine and sometimes almost imperceptible white barring, yellow irises, yellow eyelids, greenish-yellow cere, yellow legs and feet and black talons. Females are larger than males and the throat of the female has a rufous-brown wash while</p>	<p>Whilst confined to Christmas Island, this species is widespread and has been recorded in all major Island habitats from primary and marginal rainforests and areas of regrowth vegetation (Hill, 2004b).</p> <p>The Christmas Island Goshawk is an endemic, territorial resident of Christmas Island. It occurs in all forest types on the island, but is commonly seen in secondary forest, settlements or rehabilitated habitat. The Christmas Island Goshawk nests in tall trees in forest patches of more than</p>	<p>Likely; however critical nesting habitat unlikely to be impacted</p>



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
		<p>the throat of the male is white with grey spots. Adults weigh from 200-490 g and are around 40 cm length. The Christmas Island Goshawk feeds on large insects, small birds, mammals and reptiles (Department of the Environment, 2014).</p>	<p>1.5 ha in area. Its nest sites are often near cleared land, possibly because these sites provide better visibility and access to food (Department of the Environment, 2014). Primary and Marginal Rainforest have been identified as critical habitat for the species and it is understood that regrowth vegetation may also be critical habitat for the species, though the quality of regrowth necessary has not yet been identified (Hill, 2004).</p> <p>It is likely that the Christmas Island Goshawk will be present in areas proposed for clearing but proposed clearing areas are not used as nesting habitat.</p>	
<p><i>Ninox natalis</i> Christmas Island Hawk-Owl</p>	<p>VU</p>	<p>A small owl is about the size of a common pigeon (26 to 29 cm and weighing approximately 140 to 210 g) but with a barred breast. It has an unmistakable boo-book call. Unlike other raptors, including many owls, there is little difference in size between males and females with females being only slightly larger than the male. The Hawk-Owl feeds mainly on large insects and sometimes on small vertebrates. It is nocturnal, feeding mainly in the understorey of primary forest, or in secondary vegetation along roadsides (Department of the Environment, 2014).</p>	<p>This species is confined to Christmas Island. This species occupies permanent territories in all forest types on the island, with highest densities in primary forest and lowest in post-mining regrowth. Breeding does not occur in secondary growth because trees there are not old enough to have developed hollows. Habitat critical to the survival of the Christmas Island Hawk-Owl is defined as all Primary and Marginal rainforest, and all secondary growth rainforest that provides suitable habitat. Suitable secondary growth forest has not been identified (Hill, 2004).</p> <p>It is likely that the Christmas Island Hawk-Owl will forage in areas proposed for clearing but proposed clearing areas are not used as nesting habitat due to the predominance of secondary growth in these areas.</p>	<p>Likely; however critical nesting habitat unlikely to be impacted</p>



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
<i>Turdus poliocephalus erythropleurus</i> Christmas Island Thrush	EN	A subspecies of Island Thrush, 21 to 22 cm long, with a mass of 44 to 68 g. It has brown to dark-grey upperparts, a white chin and throat, a grey brown to grey breast, dull orange flanks, and a belly that is dull orange on the sides and white in the centre. It has dark brown irises, a prominent ring of yellow to orange-yellow skin around each eye, a yellow to orange bill, and yellow to orange-yellow legs and feet (Department of the Environment, 2014).	<p>Whilst confined to Christmas Island, this species is widespread on over the island (Director of National Parks, 2014b).</p> <p>The Island Thrush (Christmas Island) is endemic to, and common in most habitats on Christmas Island, including tall, closed evergreen rainforest, open semi-deciduous rainforest, secondary regrowth, thickets of weeds and semi-deciduous vines, settled areas (where it forages on lawns and nests on buildings), and on the Christmas Island golf course. It is most common in the evergreen rainforest and open rainforest on the coastal and higher terraces and plateau of Christmas Island. It is least common in disturbed habitats (such as urban areas, regrowth and post-mining wasteland) and in suboptimal endemic vegetation such as thickets of Pandanus and patches of low vegetation in coastal areas (Department of Environment, 2014). No habitat has been identified as critical to the subspecies (Director of National Parks, 2014).</p> <p>It is likely that the Christmas Island Thrush will be present in areas proposed for clearing as it is a habitat generalist, but the species is highly mobile and will self-relocate during any clearing activities and re-establish in the disturbed habitat.</p>	Likely; however highly mobile
<i>Phaethon lepturus fulvus</i> Christmas Island White-tailed Tropicbird, Golden Bosunbird	EN	The white-tailed tropicbird (Christmas Island) is a medium-sized, whitish seabird with long central tail-feathers ('streamers'), a yellow bill and black wing-bars	Although this migratory species historically was not known to breed on Christmas Island, it has been found using nest boxes installed for the Christmas Island Hawk-owl as part of a collaboration between Christmas Island Phosphates and Christmas Island District High School	Possible; however critical breeding habitat unlikely to be impacted



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
		<p>on its upper wings. It is distinguished from the white-tailed tropicbird (Indian Ocean) by having a golden or apricot wash to its white plumage (Pizzey & Knight 1997; Dunlop et al., 2001). However, there is considerable variation in the extent and intensity of the apricot wash (Department of the Environment, 2014).</p>	<p>(pers. comm. Andrew Hill). The Golden Bosun has been regarded as a subspecies on the White-tailed tropicbird, however one that is confined to Christmas Island. There has been an observed/ inferred and projected to continue decline in numbers (DCCEEW, 2023b) however the rate and cause of decline has yet to be established. Feral cat and rat predation is presumed to be significant in the rate of decline.</p> <p>The white-tailed tropicbird (Christmas Island) is endemic to Christmas Island, which is its only known breeding location. It is widely distributed across the island (Christmas Island National Park, 2013) and roosts and forages over the Indian Ocean. Both adults and juveniles appear to disperse widely and have been recorded south and south-east of Christmas Island (Marchant & Higgins, 1990). It can utilise a range of nest-sites, including hollows in rainforest trees and crevices on rock faces, cliffs and quarries (Dunlop et al., 1988b). The threat from clearing has been assessed as passed. Potential nest sites are plentiful, the species is adaptable to new nest sites, and some breeding appears to occur in most parts of Christmas Island (Dunlop et al., 1988b). No habitat has been identified as critical to the subspecies (Director of National Parks, 2014).</p> <p>It is possible that the White-Tailed Tropicbird (Golden bosun) will be present in areas proposed for clearing as it is a habitat generalist but the species is highly mobile and will self-relocate during any clearing activities, and re-establish in the disturbed habitat.</p>	



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
<p><i>Phaethon rubricauda westralis</i> Red-tailed Tropicbird, Indian Ocean Tropicbird</p>	EN	<p>The Red-tailed tropicbird is a medium-sized seabird with a body length of around 100cm, with a wingspan of around 115 cm, and weight between 600g to 1kg (Marchant & Higgins 1990; Menkhorst et al. 2017). Adult birds are mainly white with a bright red bill and very long red tail streamers. The only dark markings are a short black eye-stripe with comma-like hook; dark outer primary feathers shafts can also be observed, mostly in birds in flight. Both sexes have similar plumage and show no seasonal variation. Juveniles are heavily barred and scaled with black across most of their upper parts, lack tail streamers, and have a black beak (DEECC, 2023)</p>	<p>This species is endemic to Christmas Island, which contains one of the largest breeding populations in Australia. Cat and rat predation on the island are considered the source behind the decline (DCCEEW, 2023a). Preferred nesting sites on Christmas Island includes rock crevices, and under vegetation (R Willacy et al. 2021).</p>	Unlikely
Mammals				
<p><i>Crociodura attenuata trichura</i> Christmas Island Shrew</p>	CR	<p>The Christmas Island shrew is a small insectivorous mammal that varies from light-brown or reddish-brown to dark-slate grey in colouration, and weighs 4.5–6 g (Schulz 2004). All members of the <i>Crociodura</i> genus are small, with a head and body length of 40–180 mm and tail length of 40–110 mm,</p>	<p>Listed as Critically Endangered but its current distribution on island is unknown. The last two individuals were captured in 1984 and 1985 (Woinarski <i>et al.</i> 2014). The most recent records (in 1984 and 1985) are of single individuals from tall plateau rainforest in deep soils (Tranter pers. comm., cited in Schulz 2004) and terrace rainforest with shallow soils (Goh pers. comm., cited in Schulz 2004), with one recorded from a fallen bird's nest fern (Meek 2000). The majority of remaining primary</p>	Unlikely



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
		and a distinctly pointed muzzle (Schulz 2004)	plateau and terrace rainforest (about 75 percent of the original native rainforest vegetation remains), including the locations of the 1984 and 1985 sightings, is now protected within the Christmas Island National Park (Woinarski et al. 2014; Director of National Parks 2014). It is unknown whether the shrew occurs in post-mined secondary regrowth, and there are no records from areas of human habitation on the island.	
<i>Pipistrellus murrayi</i> Christmas Island Pipistrelle	EX	-	-	N/A
<i>Pteropus melanotus natalis</i> Christmas Island Flying Fox	CR	The Christmas Island Flying Fox is small, compared to most Australian Pteropus species, weighing on average 350 g when fully grown (Department of the Environment, 2014). The subspecies is described as having uniformly long, black fur, giving a 'chubby' appearance. A faint reddish collar of fur exists on some individuals. The subspecies forages during the day as well as at night (Department of the Environment, 2014).	The Christmas Island Flying Fox is endemic to Christmas Island. Three breeding colonies are known (DCCEEWe. 2024); and the entire island is used for foraging, with suitable fruit and nectar being found in most vegetation types including rainforests, residential gardens, and post-mine revegetation. They disperse seeds as pollen as they go, making them an important part of the rainforest system (Parks Australia, 2024). All recorded roosts have been located on the coastal terrace or around the first land cliff and semi-deciduous forest. All previously uncleared vegetation is considered critical foraging habitat. Mined areas are not considered critical habitat for this species. It is possible that the Christmas Island Flying Fox will be present in areas proposed for clearing but these areas are not known roost sites.	Possible; however individuals and roosting habitat unlikely to be affected
<i>Rattus macleari</i> Maclear's rat	EX	-	-	N/A



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
<i>Rattus nativitatis</i> Bulldog rat	EX	-	-	N/A
Reptiles				
<i>Ramphotyphlops exocoeti</i> Christmas Island Blind Snake	VU	A brown snake growing to 35 cm, with a pale brown belly, rounded snout, scales in 20 rows at mid-body and an oval scale at the tip of the snout (Department of the Environment, 2014). The Christmas Island Blind Snake almost certainly preys on the eggs, larvae and pupae of ants and termites.	The Christmas Island Blind Snake is endemic to Christmas Island. The few records with information on locality or habitat suggest that this species occurs primarily where the deeper soils and primary rainforests occur on the island's central plateau. The Christmas Island Blind Snake is a fossorial snake found in the sub-surface soil and surface litter of the forest floor. It is usually active on the surface only at night (Cogger, 2006). No specimens have been found since 2009, and it is assumed that the species occurs only in parts of the island where dense vegetation is abundant (DCCEEWD, 2024) so it is considered unlikely that the species will occur in proposed clearing areas.	Unlikely
<i>Lepidodactylus listeri</i> Lister's Gecko	EW	A brown reptile growing to 5 cm with a broad, pale fawn/grey vertebral stripe which expands to cover the top of the head and matches the colour and pattern of the tail, scattered darker-brown flecks/blotches on the back, whitish belly and body covered with small, smooth scales. Eats a broad spectrum of small invertebrates, with termites, small beetles, bugs and slaters predominating (Department of the Environment, 2014).	Endemic to Christmas Island, where was previously found to have been most abundant on the plateau area in primary rainforest (Cogger, 2006). Lister's Gecko is considered extinct in the wild (Parks Australia, 2024) so it is unlikely that the species will occur in proposed clearing areas.	Would Not Occur



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
<i>Cyrtodactylus sadleiri</i> Giant Gecko	EN	A large gecko, around 8-10 cm from head to vent, with an average body weight of 15 g. The species is dark grey, brown or blackish in colour, flecked with a mixture of darker and lighter patches, with obscure paler bands across the dorsal side. It has a long slender tail with alternating cream and grey-brown bands (Director of National Parks, 2012a). The Giant Gecko is a nocturnal predator of small invertebrates. (Threatened Species Scientific Committee, 2013).	<p>The giant gecko is endemic to Christmas Island and is likely to occur in small populations scattered across the Island, occurring in all habitats except areas lacking in tree or shrub cover (Cogger, et al., 1983; TSSC, 2013).</p> <p>Found in all island habitats, except for areas lacking trees and shrubs, including formerly mined areas with dense regrowth. Evergreen tall, closed forest is considered critical habitat for this species (Director of National Parks, 2014). Numbers of Giant Geckos recorded during targeted surveys have markedly declined due to predation from wolf snakes, giant centipedes, and rats but given that the species utilises revegetated mining areas, it is considered possible that the species may be present in areas proposed for clearing.</p>	Possible; however highly mobile
<i>Emoia nativitatis</i> (Christmas Island Forest Skink)	EX	-	-	N/A
Invertebrates - Crustaceans				
<i>Gecarcoidea natalis</i> Red Crab	N/A	A land crab with distinctive red colouring and an adult carapace measuring up to 11.6 cm across. Males are larger than females, but females have a much broader abdomen and usually have smaller claws than males. Red crabs are diurnal, and their diet consists mainly of fallen leaves, fruits, flowers and seedlings. They	Endemic to Christmas Island, the red crab is abundant across most parts of the island and critical to the ecology of the island (Director of National Parks, 2014). They are a keystone species in the forest ecology, influencing the structure and function of the rainforest where the majority of them live, by selectively consuming seeds and seedlings and leaf litter, turning over the soil and spreading seeds by burrowing, and fertilising soil with their droppings (Department of the Environment, 2014). Habitat critical to their survival occurs throughout the	Likely; however the crab burrow density in these areas is less than in the high density coastal terraces (Parks Australia 2023 Crab Burrow Density).



Species	Status	Description	Distribution and Habitat	Likelihood of Occurrence
		are not solely vegetarian and will eat other dead crabs, birds, the introduced giant African snail and palatable human refuse (Department of the Environment, 2014).	whole island (Director of National Parks, 2014). It is likely that the Red Crab will be present in areas proposed for clearing.	
<i>Birgus latro</i> Robber crab (Coconut Crab)	N/A	The largest terrestrial crustacean in the world with a weight of up to 4 kg and a leg span up to 1 m. The robber crab is related to the hermit crab but does not carry a shell. They are semi-nomadic and omnivorous, feeding on fruit, coconut, carrion and other crabs, including the Red Crab (Krieger et al., 2012).	The robber crab is abundant but critical to the ecology of the island (Director of National Parks, 2014). Christmas Island has the world's largest and best protected population of the world's biggest land crustacean (Parks Australia 2024). They are found in most parts of the island but are also a keystone species in the forest ecology, influencing the structure and function of the rainforest by foraging the forest floor, feeding mainly on seeds, fruits and the pith of fallen trees. They also scavenge the carcasses of dead animals. It is likely that the Robber Crab will be present in areas proposed for clearing. The main threat to Robber Crabs are vehicles.	Likely; however, the successful physical relocation of robber crabs will be a suitable mitigation
<i>Discoplax celeste</i> Blue Crab	N/A	A large crab with a deep and rounded carapace with a width of 45 mm or more. Adults are overall blue to bluish white (Ng and Davie, 2012).	Endemic to Christmas Island, with a restricted distribution in perennially wet/moist areas and seepages of the coastal terraces and below, though they may range further during the wet season. The species is of conservation significance because it characterises a significant ecosystem. All spring and wetland areas are considered critical habitat (Director of National Parks, 2014). The species will be absent from proposed clearing areas which are confined to the upper terraces.	N/A



Heritage

There are no Aboriginal Sites of significance or Native Title Claims over the area however there are sites on the National Heritage List (EPBC Act). Christmas Island has been mined for phosphate since the 1890s and contains a unique collection of structures and sites of heritage significance associated with this mining and blended cultural past.

Under the National Heritage List, comprising of natural and historical places, nine sites were identified on Christmas Island for its outstanding heritage significance to the Australian nation (DoE, 2004a, b, c, d, e, f, g, h, i). Three of which are located within PRL lease areas:

- Drumsite Industrial Area,
- Industrial and Administrative Group (Phosphate Rock Storage Bins, Phosphate Dust Storage Building and Downhill Conveyor System (from Drumsite to Dry Storage Bins), and Phosphate Loading Cantilevers), and
- Phosphate Hill Historic Area (Phosphate Hill Mine Workings).

National heritage sites are protected and thus would require approval for an action that will or is likely to have a significant impact on matters of National Environmental Significance (NES). **There are no heritage places within CPS 6323/1.**




Figure 26: National Heritage List Sites



Environmental Management

Clearing Activities

Key management commitments from the Christmas Island Phosphate Environmental Management Plan 2024 – 2029 (ENV-Plan-001) which will reduce the impact of clearing upon the biodiversity values of the island are listed below:

- Undertake works in accordance with procedure ENV-SOP-002: Vegetation Clearing Procedure.
- Ensure a Mine Works Clearing Permit is completed, and any restrictions/conditions highlighted and understood by all involved prior to starting any clearing activities.
- Comply with all conditions of clearing permits.
- Provide an annual report to DWER for clearing permits in accordance with permit conditions.
- Undertake mine site planning for all clearing operations to identify potential erosion potential and mitigation strategies in liaison with neighbouring land managers (where applicable).
- Ensure appropriate pre-clearing survey and flagging is undertaken to identify areas to be cleared, boundaries of National Park and other areas to be protected (heritage and/or rainforest areas). CIP will include standard Buffer Zones in all areas to be cleared (against roadway and site boundaries) and will comply with any additional buffer zones as imposed by the Permit Conditions.
- Utilise an appropriate onsite audit and reporting program for mine site operations to thoroughly risk assess sites prior to clearing and assess compliance with clearing conditions and boundary impacts to surrounding vegetation and habitats.
- Identify any primary habitat within MCI 70/1A and ensure appropriate management of mining activities to protect these areas
- Ensure no primary rainforest within MCI 70/1A is disturbed or cleared; and minimise the amount of native vegetation to be cleared.

Flora Management

- Undertake appropriate assessments in clearing permit approvals processes to ensure no threatened flora species are located within proposed mining areas.
- Risk-based pre-clearing ground truthing will be undertaken where identified as required.
- Should threatened species be identified, CIP will contact DCCEEW & Parks Australia to determine appropriate measures to protect the populations located.
- Undertake appropriate assessments of areas proposed for clearing to assess potential presence of other special flora species and assess management measures to minimise impacts.

Fauna Management

Red Crabs

- Liaise with Parks Australia to determine the most appropriate timing of clearing if areas are identified to have high crab burrow density to minimise impacts on the Red Crab population.

Robber Crabs

- Relocate Robber Crabs before areas are cleared for mining operations.

Abbott's Booby

- Assess potential impacts on Abbotts Booby and other threatened species in the clearing permit approval process to ensure no significant impacts on habitat or abundance because of clearing.
- Conduct risk-based pre-clearing ground truthing; with particular care paid prior to clearing the sites identified previously as being adjacent to or nearby known Abbott's Booby nesting areas.
- Ensure all minimum buffers and boundaries are enforced utilising the Mines Works Permit.



Weed Management

- Undertake weed management in accordance with the Weed Management Plan (ENV-Plan-003).
- Undertake weed control on a prioritised basis based on meeting regulatory requirements, minimising the establishment of new populations of weeds into 'clean' areas, and protecting important habitat communities from priority weed species.
- Undertake monitoring of priority weed locations and follow up controls.
- Ensure vehicles are cleaned regularly (e.g. high-pressure hose at designated wash down bays) to avoid movement and spread of weed species through transfer of soil and plant material.
- Work collaboratively with Island partners to address weed management of priority Island wide weed species.
- Provide training and awareness programs for the workforce on weed and pest identification, reporting and management.

Mine Relinquishment

No areas within CPS 6323/1 have been relinquished since the commencement of CPS 6323/1. Where areas are identified as part of relinquishment planning all relinquishment tenement conditions shall be complied with prior to submission of the partial surrender application. Rehabilitation is undertaken by Parks Australia.

Ten Clearing Principals

An assessment carried out against the ten clearing principles provided under Part V Division 2 of the EP Act is shown in Table 7.



Table 7: Assessment of ten clearing principles

Clearing Principle	Is the Proposal at Variance?	Justification for Variance
<p>Principle (a) Native vegetation should not be cleared if it comprises a high level of biological diversity.</p>	<p>Proposal is not likely to be at variance to Principle (a)</p>	<p>Christmas Island is home to approximately 420 species of vascular plants, including 177 introduced species and 242 species thought to be indigenous to the island. Eighteen species are known to be endemic to Christmas Island, with the other flora being more widely distributed through the Indo-Malayan and Malesian regions, or throughout the tropical Indo-Pacific. No Priority Flora species are listed for Christmas Island. Areas of the island which are previously uncleared and retain high biodiversity have been reserved as National Park, including the island's two Ramsar wetlands. Areas which have been previously cleared have been allocated to the Shire of Christmas Island for future development, or to Mining Lease for future mining.</p> <p>A total of 75 flora species from 52 families have been recorded in the proposed clearing areas, this included 53 native species (three endemic) and 22 introduced species. The native flora taxa recorded included the endemic species: <i>Hoya aldrichii</i>[§], <i>Arenga listeri</i>[§], <i>Pandanus elatus</i>[§].</p> <p><i>A.listeri</i>[§] and <i>P.elatus</i>[§] were common throughout the surveyed areas. <i>H.aldrichii</i>[§] was found at few sites during the field survey but is not common in closed canopy evergreen forest, preferring the coastal terraces where it is common in the shrublands.</p> <p>All areas proposed for clearing have been previously cleared, though some of these have regrowth vegetation that is up to 40 years old. Condition of regrowth varies, with some sites being weed dominated. Sites were assessed on a case-by-case basis to determine whether they are representative of natural vegetation in the ground truthing conducted. Given that all areas proposed for clearing have been previously cleared, the proposal not likely to be at variance to this principle</p>
<p>Principle (b) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna.</p>	<p>Proposal is not likely to be at variance to Principle (b)</p>	<p>Christmas Island provides habitat for several fauna indigenous to the island including fourteen native bird species and nine species of seabird which use the island for breeding. Three seabird taxa and nine land bird taxa are endemic to the island. A further 104 migratory or vagrant bird species have been recorded on the island. Eight of the island's endemic birds are listed as threatened under the EPBC Act 1999 (DCCEEW, 2024).</p> <p>Five endemic native mammals were historically recorded on Christmas Island with only one, the Christmas Island flying-fox, now known to remain. Christmas Island has six species of native terrestrial reptiles, five of them endemic. Mammal and reptile species indigenous to the island have undergone a catastrophic decline since human</p>



Clearing Principle	Is the Proposal at Variance?	Justification for Variance
		<p>settlement, largely because of the introduction of invasive species. Consequently, it is considered unlikely that any mammal or reptile species of conservation significance rely upon habitat in areas proposed to be cleared.</p> <p>Crustaceans are the most conspicuous invertebrate fauna of Christmas Island, with the island supporting over 20 terrestrial and intertidal crab species of which 14 are regarded as true land crabs, depending on the ocean only for their larval development. Three land-crab species have been identified as conservation significant: the Red Crab, Blue Crab and Robber Crab. Blue Crabs have a restricted distribution and do not occur within the areas proposed for clearing. Red crabs are common throughout the island but are a keystone species in the island's ecology. Robber crabs have a wide distribution across many Indian and Pacific oceanic islands. Although abundant on Christmas Island their exact conservation status is unknown. Robber crabs are present in proposed clearing areas in much lower numbers than red crabs.</p> <p>The two species of conservation significance thought to be at greatest risk of impact from the proposed clearing are the Abbott's Booby and the Red Crab. The 2014 CPS 6323/1 application removed approximately 4ha from the original application (Field 106 - MB14, which was then split into Fields 106 – MB14a and b) which was the area most likely to present a risk of disturbance to Abbott's nest sites.</p> <p>The clearing application is of previously cleared (lower) regrowth vegetation and few sites are near known Abbott's Booby nest sites, the nearest Abbott's Booby nest site identified during ground truthing was approximately 50m from boundary of proposed clearing. Red crabs are abundant throughout the island and their preferred habitat is in undisturbed rainforest, most of which is protected in the Christmas Island National Park. For these reasons, it is considered that the proposal is not likely to be at variance to this principle.</p>
<p>Principle (c) Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, threatened flora.</p>	<p>Proposal is not likely to be at variance to Principle (c)</p>	<p>There are three species listed as Threatened under the EPBC Act 1999. These are <i>Asplenium listeri</i>^s (Christmas Island Spleenwort), listed as Critically Endangered, <i>Tectaria devexa var. minor</i>, listed as Endangered and <i>Pneumatopteris truncata</i>, listed as Critically Endangered.</p> <p><i>A. listeri</i>^s occurs in limestone rock crevices in dry, exposed areas on Christmas Island and hence is not associated with proposed clearing sites.</p> <p><i>T. devexa</i> grows colonially, mainly on the plateau, in primary rainforest (tall and largely undisturbed), above 80 metres elevation; both in deeper soils and as a lithophyte (on mossy pinnacles at the base of a slope, a wet site).</p>



Clearing Principle	Is the Proposal at Variance?	Justification for Variance
		<p>Ten populations of <i>T. devexa</i> have been identified at Christmas Island, with the majority of these occurring in the National Park. No known populations of <i>T. devexa</i> have been identified at proposed clearing locations.</p> <p><i>P. truncata</i> is known from two sites at Christmas Island. Neither location is within the proposed clearing areas. <i>P. truncata</i> grows in permanently moist sites in semi-deciduous closed forest and is known to be found in two sub-populations (one at Hugh's Dale, the other at Blowholes Ravine, neither of which are associated with any of the proposed clearing sites. No known populations of <i>P. truncata</i> have been identified at proposed clearing locations.</p>
<p>Principle (d) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of a threatened ecological community.</p>	<p>Proposal is not at variance to Principle (d)</p>	<p>There are no listed Threatened Ecological Communities on Christmas Island. Therefore, the clearing as proposed is not at variance to this principle.</p>
<p>Principle (e) Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.</p>	<p>Proposal is not likely to be at variance to Principle (e)</p>	<p>Approximately 75% of Christmas Island is covered with native vegetation and 84% of this (63% of total island area) is protected within National Park. The proposed clearing occurs on land that have previously been cleared for phosphate mining and consists of predominantly regrowth vegetation and weed species, together with some lower value native species common on the island. The vegetation under application is not part of the island's original forests and the vegetation for many of the application areas is in a degraded condition and no longer representative of rainforest structural values. The vegetation within the application area is not considered to be significant remnant vegetation as it is in an extensively cleared landscape and therefore is not at variance to this principle.</p>
<p>Principle (f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.</p>	<p>Proposal is not likely to be at variance to Principle (f).</p>	<p>None of the proposed clearing is associated with or adjacent to a watercourse or wetland. Perennial surface water features on Christmas Island are limited to spring fed streams on coastal or sloping areas of the island.</p> <p>All the area under application is situated on the plateau and not within wetland areas. This proposal is not at variance to this principle.</p>



Clearing Principle	Is the Proposal at Variance?	Justification for Variance
<p>Principle (g) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.</p>	<p>Proposal is not likely to be at variance to Principle (g)</p>	<p>The interior of the island is slightly undulating plateau, from about 160-360m above sea level. The area under application is situated on the plateau with relatively little relief, and above the terraces. Due to the nature of phosphate mining topsoil will be removed in areas for in situ mining and all other areas will be mined to ground level. All areas that are mined in situ will be left as limestone boulders. The land is currently approved for mining so its land capability will not be impacted by clearing. No wind erosion, water erosion, salinity, eutrophication, or waterlogging is expected because of the clearing, so this proposal is not likely to be at variance to this principle.</p>
<p>Principle (h) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.</p>	<p>Proposal may be at variance to Principle (h)</p>	<p>There are four noteworthy conservation areas at Christmas Island. These are the Christmas Island National Park, the Ramsar wetlands Hosnie's Springs and The Dales, and the Commonwealth Heritage Site "Christmas Island Natural Areas". One area proposed for clearing is adjacent to The Dales Ramsar wetlands and several areas are adjacent to the Christmas Island National Park.</p> <p>Most areas under application are highly disturbed and unlikely to be acting as an effective buffer to the adjacent conservation areas. The clearing of some sites as proposed may be at variance to this principle as clearing of the applied area could result in adjacent conservation areas being exposed to weed invasion.</p>
<p>Principle (i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.</p>	<p>Proposal is not likely to be at variance to Principle (i)</p>	<p>Due to the high natural rate of infiltration at Christmas Island, erosion and sedimentation is generally localised to compacted areas such as roads and stockpiles. There is no potential for deterioration of underground water because of clearing. Due to the location of the areas proposed to be cleared, it is unlikely that the clearing of native vegetation for phosphate mining will cause deterioration in the quality of surface water or groundwater within the local area. Therefore, this proposal is not likely to be at variance to this principle.</p>
<p>Principle (j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence or intensity of flooding.</p>	<p>Proposal is not likely to be at variance to Principle (j)</p>	<p>Christmas Island's soils and karstic limestone rocks are generally highly permeable so clearing on the island does not cause or exacerbate flooding. As the clearing as proposed is not likely to cause or exacerbate waterlogging or flooding and as the water features on Christmas Island are not close to the applied area, the clearing as proposed is not likely to be at variance to this principle.</p>



Conclusion

This proposed clearing permit amendment is a request to extend Clearing Permit CPS 6323/1 (Purpose Permit) approved on 23 June 2015, for the clearing of native vegetation by Christmas Island Phosphates for the areas expressly covered in the application for the purpose of stockpile recovery and insitu phosphate mining.

CPS 6323/1, as approved June 2015, required in Condition 5, that CIP (the Permit Holder) have regard for the following principles (in order of preference): 5(a) avoid the clearing of native vegetation, 5(b) minimise the amount of native vegetation to be cleared, and 5(c) reduce the impact of clearing on any environmental value. All clearing conducted to date has been in accordance with these principles.

All other conditions of CPS 6323/1 have been complied with; however, several changes are being requested as part of this application for extension, including:

- Extension of the duration of the Permit from December 2024, to align with the tenement expiry MCI 70/ 1A in June 2034, to allow the remaining clearing and mining activity to be completed;
- Removal of Condition 6(c) from CPS 6323/1 approval dated June 2015 ‘At least once in each 3-month period prior to the Weed Management Plan required in Condition 6(b) being approved, the Permit Holder must remove or kill any weeds growing within areas cleared under this permit’ as the Weed Management Plan required under Condition 6(b) was approved in May 2017 by the then Department of Infrastructure and Regional Development, and will continue to form weed management over CPS 6323/1; and
- Removal of Condition 7 from CPS 6323/1 approval dated June 2015 ‘The Permit Holder must implement and adhere to the document “A Management Plan for the Christmas Island Pipistrelle in relation to vegetation Clearing on Mining Leases, May 2015”’ as the Pipistrelle Bat had its status changed to extinct on the EPBC Act List of Threatened Species in 2021.

Thorough ground truthing has been conducted which has identified no significant changes in the condition of the proposed clearing areas within CPS 6323/1 from the assessments completed in 2014, and the CPS approved in June 2015. Changes to classification of several fauna species on the EPBC Act have been recorded. Risks assessment of impacts on flora and fauna have been improved with further mapping since the 2014 application which has been utilised during this assessment.

Abbott’s Booby nests were identified as located at least 50 metres distance to the areas of clearing during the ground truthing undertaken for this assessment; however, CIP are confident that the measures in place are sufficient to minimise any disturbance.

CIP has re-assessed the proposed clearing of up to 128.53 ha within CPS 6323/1 against the ten clearing principles and the clearing been conducted to date. The assessment has found that the clearing may be at variance to clearing principle (h) and not at variance with any of the other clearing principles. No significant changes against the assessment report submitted in 2014 have been identified, other than the reduction of field 133B – M14 to avoid proximity to Abbott’s Booby nesting sites.



References

- Andrews, C. (1900). A monograph of Christmas Island (Indian Ocean). London: British Museum (Natural History).
- ABS (2023). 2021 Christmas Island, Census All persons QuickStats. Australian Bureau of Statistics (ABS), Canberra.
- ANZECC. 2000. Review of the National Strategy for the Conservation of Australia's Biological Diversity, Australian and New Zealand Environment and Conservation Council, Environment Australia, Canberra, Australian Capital Territory.
- Beard, J.S. 1990. Plant Life of Western Australia. Kangaroo Press, Perth, Western Australia.
- Bennett, M. (2010). Native vegetation clearing application CIP MCI 70/1 Mining Clearing Permit Local Permit ID: ML105 MB1. Unpublished report prepared for Christmas Island Phosphates, Indian Ocean.
- BOM. 2023. Climatic averages for Australian Sites, publicly available data prepared by the Bureau of Meteorology, Commonwealth of Australia, URL: <http://www.bom.gov.au/climate/dwo/IDCJDW6026.latest.shtml>
- Butz, M. 2004. National Recovery Plan for *Tectaria devexa*. Department of the Environment and Heritage. Canberra, Australia.
- Claussen, J. (2005). Native Plants of Christmas Island (Vol. Flora of Australia Supplementary Series Number 22). Canberra: Australian Biological Resources Study and Christmas Island Natural History Association.
- Commonwealth of Australia. 2013. Mining Lease No. MCI 70/1. Signed by the Honorable Catherine King MP, Minister for Regional Services, Local Communities and Territories on Behalf of the Commonwealth of Australia, 27 June 2013.
- DBCA 2023 Florabase – the Western Australian Flora. Descriptions by the Western Australian Herbarium. Department of Biodiversity, Conservation and Attractions. Perth. Western Australia.
- DCCEEWa. 2023. Conservation Advice for *Phaethon rubricauda Westralia* (Indian Ocean Red-tailed tropicbird). Accessed December 2023. URL: <https://www.environment.gov.au/biodiversity/threatened/species/pubs/91824-conservation-advice-21122023.pdf>
- DCCEEWb. 2023. Conservation Advice *Phaethon lepturus fulvus* (White Tailed Tropic Bird Christmas Island). Accessed December 2023. URL: <https://www.environment.gov.au/biodiversity/threatened/species/pubs/26021-conservation-advice.pdf>
- DCCEEWc. 2023. EPBC Act List of Threatened Fauna. Accessed December 2023. URL: <https://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl>
- DCCEEWd 2023. Australia's Bioregions (IBRA). Accessed December 2023. URL: <https://www.dcceew.gov.au/environment/land/nrs/science/ibra>
- DCCEEW. 2024 Amendments to the EPBC Act list of threatened species. Accessed May 2024. <https://www.dcceew.gov.au/news/2021/03/02/amendments-epbc-act-list-threatened-species>
- DCCEEWb. 2024 Conservation Advice for the Christmas Island Frigatebird - *Fregata andrews*. Accessed June 2024. URL: <http://www.environment.gov.au/biodiversity/threatened/species/pubs/1011-conservation-advice-19102020.pdf>
- DCCEEWc. 2024. Protected Matters Search Tool. Accessed July 2024. URL: <https://www.dcceew.gov.au/environment/epbc/protected-matters-search-tool>
- DCCEEWd. 2024 Conservation Advice for *Ramphotyphlops exocoeti* - Christmas Island Blind Snake. Accessed July 2024. URL: <https://www.environment.gov.au/biodiversity/threatened/species/pubs/1262-conservation-advice-15072016.pdf>
- DCCEEW. 2024 Conservation advice for *Pteropus melanotus natalis* – Christmas Island Flying Fox. Accessed July 2024. URL: <https://www.dcceew.gov.au/sites/default/files/env/pages/2cd790fe-e19b-4c90-8589-d5f7026b5477/files/64801-conservation-advice.pdf>



- DEH (now DCCEEW) 2004. National Recovery Plan for the Abbott's Booby *Papasula abbotti*. Department of the Environment and Heritage, Canberra, Australian Capital Territory.
- Director of National Parks. 2014. DRAFT Christmas Island Biodiversity Conservation Plan. Department of the Environment, Canberra, Australian Capital Territory.
- Director of National Parks. 2014b. Christmas Island National Park Management Plan 2014 – 2024. Director of National Parks, Canberra, Australian Capital Territory.
- Drew, M., Harzsch, S., Stensmyr, M., Erland, S. and Hansson, B.S. (2010). A review of the biology and ecology of the robber crab, *Birgus latro* (Linnaeus, 1767) (Anomura: Coenobitidae). Zoologischer Anzeiger. 2010.03.001.
- EPA. 2016. Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment in Western Australia, 2016. Environmental Protection Authority, Perth, Western Australia.
- EPA. 2020. Statement of Environmental Principles, Factors and Objectives. Environmental Protection Authority, Perth, Western Australia.
- Grimes, K. (2001). Karst features of Christmas Island (Indian Ocean). Helictite, 37(2), 41-58.
- Hill, R. 2004a. National Recovery Plan for the Christmas Island Goshawk *Accipiter fasciatus natalis*. Commonwealth of Australia, Canberra.
- Hill, R. 2004b. National recovery plan for the Christmas Island Hawk-Owl *Ninox natalis*. [Online]. Department of the Environment and Heritage. Canberra, Commonwealth of Australia. Available from: <http://www.environment.gov.au/biodiversity/threatened/publications/recovery/n-natalis/index.html>.
- Hollingsworth, I. (2003). Soils and geomorphology in relation to proposed new mining leases, Christmas Island, Indian Ocean. In Draft Environmental Impact Statement for the Proposed Christmas Island Phosphate Mines, Technical Appendices Vol 1 (2005 ed.). Perth: Phosphate Resources Ltd.
- Holmes, J., & Holmes, G. (2002). Conservation status of the flora of Christmas Island, Indian Ocean. Unpublished report to Parks Australia.
- JBS&G. 2022. Phosphate Resources Limited Christmas Island Mountain Bike Track – Clearing Permit Application and Supporting Documentation. Perth, Australia.
- James, D., Green, P., Humphreys, W., & Woinarski, J. (2019). Endemic species of Christmas Island, Indian Ocean. Records of the Western Australian Museum, 34(2), pp. 55-114.
- Keighery, B.J. 1994. Bushland Plant Survey: A Guide to Plant Community Survey for the Community. Wildflower Society of Western Australia (Inc). Nedlands, Western Australia.
- McKenzie, N. L., May, J. E. and McKenna, S. 2002. Bioregional Summary of the 2002 Biodiversity Audit for Western Australia. Department of Conservation and Land Management. Perth.
- Mitchell, B.A. 1985. A vegetation survey of Christmas Island. Report to Australian Nature Conservation Agency, Canberra.
- Nelson, J.B. and Powell, D. 1986. The breeding ecology of Abbott's booby. *Emu* 86:33-46.
- Ng, P. K. L. and Davie, P. J. F. 2012. The Blue Crab of Christmas Island, *Discoplax celeste*, new species (Crustacea: Decapoda: Brachyura: Gecarcinidae). Raffles Bulletin of Zoology, 60: 89–100.
- NVIS Technical Working Group. (2017). Australian Vegetation Attribute Manual Version 7.0. Canberra: NVIS Technical Working Group, Department of the Environment and Energy.
- Ohlsen, D., Perrie, L., Shepherd, & Bayly, M. (2015). Taxonomic status and distribution of the critically endangered Christmas Island spleenwort (*Asplenium listeri*, Aspleniaceae): it is not as rare as we thought. Australian Systemic Botany(27), 372-377.
- Parks Australia, Christmas Island National Park/ Animals. Accessed June 2024. URL: <https://parksaustralia.gov.au/christmas/discover/nature/animals/>
- PRL, 2024, Christmas Island Social and Economic Impact, Accessed May 2024. URL: <https://prlgroup.com.au/community-sustainability/christmas-island-impact/>
- Puhalovich, A., Jacobsen, N., & Overall, R. (2003). Surface water and groundwater hydrology in relation to proposed new mining leases, Christmas Island, Indian Ocean. In Draft Environmental Impact



Statement for the Proposed Christmas Island Phosphate Mines, Technical Appendices Vol. 1 (2005 ed.). Phosphate Resources Ltd.

- Range to Reef Environmental. 2012. Flora and Fauna Risk Assessment - Report to support Environment Protection (Biodiversity Conservation) Act 1999 referral for Proposed Phosphate Mining in South Point Christmas Island. Perth, Australia.
- Range to Reef Environmental. 2013. Fauna and Flora Survey, Block 2, South Point, Christmas Island. Perth, Western Australia.
- Range to Reef Environmental. 2014. Clearing Permit Application Supporting Documentation. Unpublished report prepared for Christmas Island Phosphates by Range to Reef Environmental.
- Regional Development Organisation Australia. 2022. Indian Ocean Territories Strategic Regional Plan 2030. Indian Ocean Territories Regional Development Organisation, under Regional Development Australia.
- Reddell, P., Zimmerman, A., & Milnes, A.R. (2020). Vegetation surveys to assess possible impacts of phosphate mining, Christmas Island, Indian Ocean. Technical Report published on ResearchGate. Accessed June 2024. URL: https://www.researchgate.net/publication/342673319_Vegetation_surveys_to_assess_possible_impacts_of_phosphate_mining_Christmas_Island_Indian_Ocean
- Shire of Christmas Island. 2015. Christmas Island Local Planning Strategy. Shire of Christmas Island.
- Shire of Christmas Island. 2016 Town Planning Scheme Map No. 2. Shire of Christmas Island.
- Shire of Christmas Island. (2016, February 17). Local Planning Scheme No. 2 District Zoning Scheme. Government Gazette, WA (No. 27 Special), pp. 480-514.
- Swarbrick, J.T. and Hart. R. 2000. Environmental weeds of Christmas Island (Indian Ocean) and their management. Plant Protection Quarterly Vol 16 (2): 54-57



Appendices

Appendix I Ground Truthing Maps

Appendix II Site Photos and Condition Survey Score

Appendix III Protected Matters Search Report