



GMA Mining Australia

CPS 8825/1 Documentation for a Native Vegetation Clearing Permit Amendment Application



M70/204 Supporting Information

GMA Mining Australia

Contents

1.	Introduction	4
1.1	Background	4
1.2	Document Purpose	4
2.	Clearing description details	5
2.1.1	Mining Expansion	5
3.	Environmental Setting	6
3.1	Climate	6
3.2	Land use	6
3.2.1	Reserves	6
3.3	Landforms, geology and soils	7
3.4	Hydrogeology and Hydrology	7
3.4.1	Surface water	7
3.4.2	Groundwater	8
3.4.3	Public Drinking Water Source Areas	8
3.5	Flora and Vegetation	8
3.5.1	Broad Vegetation mapping and Extent	8
3.5.2	Mapped vegetation types and conditions	9
3.5.3	Ecological Communities	9
3.5.4	Flora Diversity	9
3.5.5	Conservation significant flora	9
3.5.6	Environmentally Sensitive Area	11
3.6	Fauna	11
3.6.1	Fauna Diversity	11
4.	Environmental Risk Assessment and management	13
4.1	Identifying Environmental Threats	13
4.2	Risk Assessment	13
4.3	Specific Management Actions to Address Impacts from Clearing	16
4.3.1	Clearing of Native Vegetation	16
4.3.2	Dust	16
5.	Clearing and Rehabilitation Status	16
5.1	Clearing Status	16
5.2	Rehabilitation Status	17
6.	Assessment of the Ten Clearing Principles	17
7.	Reference	21
Appendix A. Environmental Surveys		22
Appendix B. Risk Assessment		23
Appendix C. GMA Dust and Management Plan		27

M70/204 Supporting Information

GMA Mining Australia

Abbreviations

Abbreviation	Definition
BAM Act	Biosecurity and Agricultural Management Act 2007
BoM	Bureau of Meteorology
DAWE	Department of Agriculture, Water and Environment
DBCA	Department of Biodiversity, Conservation and Attractions
DP	Declared Pest
DWER	Department of Water and Environmental Regulation
EP Act	<i>Environmental Protection Act 1986</i>
EPBC Act	Environmental Protection and Biodiversity Conservation Act 1999
ESA	Environmentally Sensitive Area
NVCP	Native Vegetation Clearing Permit
PEC	Priority Ecological Community
RIWI Act	<i>Rights in Water and Irrigation Act 1914</i>
TEC	Threatened Ecological Community
BC Act	<i>Biodiversity and Conservation Act 2016</i>

1. Introduction

1.1 Background

GMA Garnet Pty Ltd (GMA) is a wholly owned subsidiary of Garnet International Resources Pty Ltd. GMA owns and operates the garnet mineral sand mining and processing operations in the Mid-West Region, Port Gregory, Western Australia. GMA operates two open cut alluvial garnet mines, the Hose Mine (tenements G70/171, M70/856, M70/926 and M70/927) and the Lynton Mine (tenements M70/204, M70/259, M70/968, M70/1330 and M70/1331). Mining is currently undertaken within M70/204 (Lynton north and south pit) and M70/926. All ore is processed at the wet separation plant (wet plant) located on M70/856.

The proposal involves seeking approval to amend the clearing permit boundary and expand the area to be cleared under CPS 8825/1. The proposed amendment includes an increase in the approved development envelope by approximately 4.6 hectares and an extension of the clearing area by 2.6 hectares. The proposed changes are outlined in Figure 1.

A clearing permit is required under the *Environmental Protection (Clearing of Native Vegetation) Regulation 2004* and the *Environmental Protection Act 1986* (EP Act), which contains provisions that protect native vegetation while allowing the approved clearing activities. The proposed amendment extends the permit's duration from 15 May 2025 to 15 May 2030.

1.2 Document Purpose

The purpose of this document is to provide supporting information to amend the CPS 8825/1 clearing permit under Section 51k of Part V of the *Environmental Protection Act 1986* to amend the existing clearing permit boundary by 4.6 hectares and increase the clearing extent by 2.6 hectares(ha) within the application area in M70/204.

This document comprises the following:

- A description of the clearing details.
- Environmental Setting.
- Summary of rehabilitation undertaken within M70/204.
- Risk assessment and management.
- Assessment of the Ten Clearing Principles as defined in Schedule 5 of the EP Act.

GMA commissioned GHD Pty Ltd (GHD, 2020a) to undertake a flora, vegetation and fauna survey and a targeted flora survey (GHD, 2020b) of the application area. The information contained within the flora, vegetation and fauna survey informed the environmental assessment component of this report (Appendix A).

Both surveys supporting this NVCP application were previously submitted to the Index of Biodiversity Surveys for Assessments (IBSA). The submission details are summarised in Table 1.

M70/204 Supporting Information

GMA Mining Australia

Table 1 IBSA Submission Details

Report name	IBSA number
GMA Garnet Pty Ltd Lynton Mine Expansion Biological Survey	IBSA-2020-0538
Targeted Flora Survey conducted for GMA Garnet Pty Ltd, for Lynton.	IBSA-2020-0405

2.

Clearing description details

The clearing activities within the application area are outlined in the subsections below and mapped in Figure. 1.

Table 2 provides a summary of the clearing activities within the application area.

Activity	Ha
Expansion of the Lynton 2b pit	2.6
Increase in development envelope	4.6

2.1.1 Mining Expansion

GMA plans to expand the existing mining pit progressively. It is anticipated that 15 hectares of native vegetation will be cleared annually. The proposed clearing method is a dozer with scrub rake.

This mining method allows the mining voids to be progressively backfilled and rehabilitated at the trailing edge of the pit, while mining activities continue at the leading edge, progressing northwards.

The mining area will be progressively rehabilitated and returned to native vegetation as per the Notice of Intent – Mining Lease M70/204 (NOI 3461) and Port Gregory Project – Revised Mine Closure Plan (Reg. ID: 85076).

The Notice of Intent is available in Appendix A

3. Environmental Setting

3.1 Climate

The application area is located within the Mid-West Region of Western Australia. The climate of Mid-West is considered warm semi-arid to Mediterranean climate with 400 to 500 mm of rainfall per annum (Desmond and Chant, 2002). The region experiences short mild, wet winter and the remainder of the year is warm to hot, dry to windy.

Annual Evaporation rate in the area is around 2,500 mm.

The nearest Bureau of Meteorological (BoM) station that provides reliable wind data is the Geraldton Airport (Site No. 8051). The BoM's Geraldton Airport 2007 meteorological file indicates dominant wind blows from the south and south-east direction, with a secondary prevailing wind from the north-east direction (Chart 1). Wind speeds between 2 and 6 m/s are most often observed, with wind speed reaching 8 m/s from the south-east direction.





Chart 1 Wind rose (9 am and 3 pm) BoM 2007

3.2 Land use

3.2.1 Reserves

A search of the *NatureMap* database identified one DBCA listed reserve – Utcha Well Nature Reserve within 11 km of the application area (GHD, 2020a).



M70/204 Supporting Information

GMA Mining Australia

The Hutt Lagoon is listed as a wetland of national importance on the Directory of Important Wetlands in Australia (DIWA) (DEC 2009). Hutt Lagoon is a macroscale elongate sump land aligned northwest to south-east, parallel to the coast. The Lagoon is usually partly filled with hypersaline water during winter for the remainder of the year, the Lagoon is usually dry. The Lagoon contains the world's largest microalgae production plan, a 250 hectares series of artificial ponds used to produce beta- carotene. During summer and in dry seasons, the Lagoon is mostly empty except the artificial ponds used for algal cultivation (URS, 2013). The Hutt Lagoon neighbor's a macroscale elongate floodplain (to the north-west and the south-east) that include more than twenty microscale elongate sump lands such as Utcha Swamp (Jaensch 1992). Water supply for the Hutt Lagoon derives from direct precipitation, surface inform from several minor creeks and seepage of groundwater (DEC, 2009).

3.4.2 Groundwater

The Department of Water and Environmental Regulation (DWER) Perth Groundwater Map indicates the survey area is in within the Gascoyne Groundwater Area.

A superficial aquifer underlies the Application Area with superficial formation present are up to 15 m thick and become progressively thinner to the east. Sub-surface flows are from east to west and discharge into the Hutt Lagoon. The flows discharge over a hypersaline saltwater wedge extending from the eastern portion of the Hutt Lagoon. Groundwater salinity within the application area varies from 800 mg/L to 1,500 mg/L. Groundwater salinities are higher toward the Utcha Swamp (up to 30,000 mg/L) and the Hutt Lagoon perimeter (up to 150,000 mg/L). Groundwater standing levels vary of 15 m below ground levels (m bgl) towards the western boundary of the tenement to 35 m bgl (URS, 2013).

3.4.3 Public Drinking Water Source Areas

There are no public drinking water source areas within 10 km of the application area. The nearest public drinking water source is 60 km north of the application area – Kalbarri Water Reserve (Department of Water and Environmental Regulation, 2020).

3.5 Flora and Vegetation

3.5.1 Broad Vegetation mapping and Extent

Broadscale mapping (1:1,000,000) pre-European vegetation mapping (Beard, 1976) indicates two Beard Vegetation Associations (BVA) was mapped within the application area including:

- BVA 371 - Low forest.

Shephard et al. (2002) adapted and digitised the pre-European mapping. The extent of vegetation associations has been determined by the State-Wide vegetation extents calculations maintained by the DBCA (current as of March 2019—GoWA, 2019).

As shown in Table 3, the current extent of BVA 371 is below the 30% retention target of the pre-clearing size at all levels except Local Government Authority Levels.

Table 3 Pre-European Vegetation Extent Association (GoWA, 2020)

Pre-European Vegetation Extent Association	Pre-European (ha)	Current extent (ha)	Remaining pre-European extent (%)
Greenough_371			
State	32,816.04	3,499.60	10.66
IBRA Bioregion: Geraldton Sandplains	32,807.53	3,499.10	10.67
Sub-IBRA: Geraldton Hills	32,807.53	3,499.10	10.67
LGA: Shire of Northampton	5,749.92	2,142.08	36.94

M70/204 Supporting Information

GMA Mining Australia

3.5.2 Mapped vegetation types and conditions

GHD (2020a) mapped one vegetation type within the application area.

Vegetation type 1: *Acacia rostellifera* open woodland to woodland.

There are areas within the application area that have either been degraded to completely degraded (GHD, 2020a). The vegetation types mapped within the application area are shown in Figure 3.

The vegetation condition within the application area ranged from good to degraded (GHD 2020a). The application area has been subject to historical grazing and clearing. Figure 4 depicts the mapped vegetation conditions within the application area.

Some of the area has not been previously mapped. The extrapolated vegetation type is vegetation type 1 and is consistent with the vegetation type mapped south and west of the proposed of the proposed clearing area (GHD 2020).

GHD (2020a) undertook a comparison of mapped BVA with the vegetation types recorded within the applications area and concluded the following:

☐ The vegetation type mapped within the application area- *Acacia rostellifera* open woodland to woodland with brown to orange sands and Shrublands on seasonally wet brackish drainage flats.

☐ BVA 371 (*Acacia* low forest) located on some flats north of the Hutt River and is a taller version of the *A. rostellifera* thicket exceeding 10m in height, and it is very dense. *Acacia rostellifera* seems to be a pure stand of that species (Beard and Burns 1976).

3.5.3 Ecological Communities

GHD (2020a) desktop searches did not identify Threatened Ecological Communities within 10 km of the application area. Two Priority Ecological Communities PECs were identified within 10 km of the application, and these include:

- *The Kalbarri Ironstone Community* (P1) – 8 km east of the application area.
- *Shrubland of the Northampton Area, dominated by Melaleuca species over exposed Kockatea shale* (Priority 1 – PEC) – 5 km south-east of the application area.

No PECs or TECs were delineated from the application area (GHD, 2020a).

3.5.4 Flora Diversity

Sixty-four flora taxa (including subspecies and varieties) representing 26 families and 50 genera were recorded from the survey area during the field survey (GHD, 2020a).

3.5.5 Conservation significant flora


A review of the *Nature Map*, EPBC PMST and purchase DBCA database indicate the potential presence of 48 conservation significant flora occurring within 10 km of the application area (GHD 2020a).

No Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) or Biodiversity Conservation Act 2016 (BC Act) or Department of Biodiversity Conservation and Attraction listed flora were recorded within the survey area.

The likelihood of occurrence assessment post-field survey concluded three species considered possible to occur, five species unlikely to occur, and 40 species highly unlikely to occur in the survey area. The species considered possible to occur within the mapped vegetation type of the application area included *Anthocercis intricata* (P3) and *Balladonia aervoides* (P3) (GHD 2020a).

3.5.6 Environmentally Sensitive Area

One Environmentally Sensitive Area (ESA) was identified 450 meters west of the application area (GHD 2020a).



Legend

- CPS 8825 Amendment.shp
CPS 8825-1 Amendment
- GMA Mining Tenements
M70/204
M70/259
M70/968
- Environmental Surveys
SUR_BF_2C_VegetationTypes_MGA50_20200115.shp
Cleared
Prev cleared regrowth
Rehabilitated
VT01
VT02



50 m 250 m 450 m


Map Projection: Transverse Mercator
Horizontal Datum and Grid: GDA 1994 MGA Zone 50
Survey: GHD (2020)

N

Figure 3
GMA Garnet
Port Gregory Mine Site
Mapped Vegetation Types

6880000 m

6880000 m



Legend

CPS 8825 Amendment.shp

CPS 8825-1 Amendment

GMA Mining Tenements

M70/204

M70/259

M70/968

Environmental Surveys

GHD (2020) Mapped Vegetation Condition

Cleared

Completely Degraded

Degraded

Good

6880000 m

50 m250 m450 m

Map Projection: Transverse Mercator
Horizontal Datum and Grid: GDA 1994 MGA Zone 50
Survey: GHD (2020)

Figure 4
GMA Garnet
Port Gregory Mine Site
Proposed Clearing Application
Mapped Vegetation Conditions

6880000 m

M70/204 Supporting Information

GMA Mining Australia

Table 6 Risk assessment and management

Environmental Threat	Cause	Potential impact	C	L	Inherent Risk Level	Management	C	L	Treated risk
Clearing of vegetation	Clearing works undertaken to the proposal.	Clearing of native vegetation in unapproved areas and/or outside the tenement boundary. Environmental factor: <ul style="list-style-type: none"> Biodiversity/flora/fauna and biodiversity. 	minor	possible	9 - medium	<ul style="list-style-type: none"> Awareness training highlights the procedure for clearing and Clearing and Ground Disturbance permit systems and procedures are in place. 	minor	Infrequent	5 – low
Dust	Vehicle and machinery movement	Dust resulting from the movement of vehicles and the operation of machinery settles on adjacent vegetation and causes plant death. Environmental Factors: <ul style="list-style-type: none"> Biodiversity/flora/fauna/ecosystem 	minor	possible	9 - medium	<ul style="list-style-type: none"> Survey control of areas to be cleared. Post-clearing checks to ensure clearing has been undertaken following the permit conditions. 	minor	Infrequent	5 - low
	Wind	Dust from wind blowing across cleared areas and stockpiles settles on adjacent vegetation and causes plant death. Environmental Factors: <ul style="list-style-type: none"> Biodiversity/flora/fauna/ecosystem 	moderate	likely	17- high	<p>Dust management will be undertaken following the GMA's Dust Management Procedure provided in Appendix C.</p> <p>The following management measures are proposed:</p> <ul style="list-style-type: none"> Water carts will undertake dust suppression on haul roads Dust suppressant additives (mulches or polymer additives) will be used if water applicates is insufficient to ameliorate dust generation. Any mining activities will cease in the event dust suppression controls fail to mitigate dust emissions. 	minor	infrequent	5 - low

M70/204 Supporting Information

GMA Mining Australia

Environmental Threat	Cause	Potential impact	C	L	Inherent Risk Level	Management	C	L	Treated risk
						Potential impacts associated with wind erosion can be minimised by progressive rehabilitation.			
Native fauna and habitat	Clearing of vegetation.	Impact on native fauna and habitat. Environmental factor: • Biodiversity and habitat loss.	minor	likely	10-medium	Clearing is managed through the clearing and ground disturbance procedure.	minor	infrequent	5 - Low
Native fauna and habitat	Interaction with fauna.	Interaction with native fauna. Environmental Factors: Biodiversity/flora/fauna/ecosystem	minor	likely	10-medium	<ul style="list-style-type: none"> Fauna awareness training. One directional clearing to allow fauna to abscond. Speed limits impose on access and haul roads. 	minor	infrequent	5 - Low
Introduced flora	New weed species introduced to site.	Any machinery brought to site must have completed a weed and seed certificate. Environmental Factors: • Biodiversity/flora/fauna/ecosystem	Minor	Likely	10 – medium	<ul style="list-style-type: none"> Weed and hygiene procedure. Machinery or equipment brought to sit clean before mobilisation. Inspection of machinery on arrival. Weed surveys to be undertaken. 	minor	Infrequent	5- low

M70/204 Supporting Information

GMA Mining Australia

4.3 Specific Management Actions to Address Impacts from Clearing

4.3.1 Clearing of Native Vegetation

Management Action

To ensure clearing is managed appropriately, management actions will include:

- Vegetation clearing procedures and processes to prevent unauthorised clearing, including as part of the Site induction training.
- A clearing and ground disturbance permit system that requires authorisation from the Environmental and Land Access and Compliance departments.
- Survey control to ensure the clearing extent is demarcated with flagging tape.
- All site personnel undertaking clearing activities have been inducted and understand the clearing procedure and permitting processes.

4.3.2 Dust

The main environmental impact of dust emission is vegetation death from high dust settling on plant leaves, preventing photosynthesis and respiration.

The main dust sources are vegetation clearing, vehicle movement, machinery operations and strong winds.

Newly cleared areas, along with existing open areas such as Run-of-Mines, laydown areas, and stockpiles, have the potential to generate dust, particularly in windy conditions.

Management Actions

As per our Environmental Licence, dust generation management actions include:

- Staged clearing of areas to minimise open areas.
- Land clearing and topsoil handling avoid windy conditions, particularly where they may impact sensitive receptors.
- Water cart to suppress dust.
- Progressive rehabilitation of mined areas.

5. Clearing and Rehabilitation Status

5.1 Clearing Status

Table 7 summarises the clearing activities and the extent undertaken under CPS 8825/1, and Figure 4 identifies the areas where clearing activities have been undertaken.

Table 7 Area of Clearing Under CPS 8825/1 since 2020

Reporting Periods	Clearing Extent (ha)	Date cleared	Clearing Purpose
1 July to 30 June			
2019-2020	10.04	15/06/2020	Pit progression
2020-2021	1.16	30/12/2020	Pit Progression
2021-2022	1.49	27/07-31/07/2022	Pit Progression
	2.0	23/02-4/03/2022	Pit Progression
2022-2023	0.8	6/06-18/06/2023	Topsoil stockpile access to progress rehabilitation and pit progression
2023-2024	1.54	13/10-27/11/2023	Pit Progression

M70/204 Supporting Information

GMA Mining Australia

5.2 Rehabilitation Status

Rehabilitation works undertaken are summarized below and shown in Figure 6:

Table 8 summarizes the equivalent area rehabilitated within the cross-hatched red on the attached Plan 8825/1. In total, 1.31 hectares underwent topsoiling, progressive ripping, and application of both organic matter during and weed management this reporting period.

Table 8 Rehabilitation Works Under CPS 8825/1

Reporting Periods 1 July to 30 June	Area of Rehab. (ha)	Date of Rehab.	Rehab Work Undertaken
2020-2021	1.28	1/05/2021-30/06/2021	Topsoil application and progressive ripping.
2021-2022	1.28	01/07-30/07/2021	Organic matter, weed management
2022-2023	1.28	15/05-30/05/2023	Weed management
2022-2023	1.28	07/09/2023	Weed management
	0.03	05/05/2024	Topsoil application and vegetation matter



230000 m

230000 m



Map Projection: Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 50
Survey: GMA (30 June 2024)




Figure 6
GMA Garnet
Port Gregory Mine Site
CPS 8825-1 Amendment
Rehabilitation Status 2020 - 2024

6. **Assessment of the Ten Clearing Principles**

The clearing is required to improve the line of sight at the Lynton entrance and facilitate Lynton bore field upgrade works. An assessment of the proposed clearing action against the ten clearing principles, as outlined in Schedule 5 of the EP Act, is provided in Table 6.

The assessment indicates the clearing is 'not considered to be at variance with the Ten Clearing Principles'.



Table 6 Assessment of the Ten Clearing Principles

Clearing Principle	Assessment	Conclusion
Principle (a) – Native vegetation should not be cleared if it comprises a high level of biological diversity.	<p>The application area is in the Geraldton Hill sub-region of the Geraldton Sandplains IBRA. One Beard Vegetation Association has been mapped in the application area BVA 371.</p> <p>The pre-European extent of remaining for BVA 371 is 10.66 to 10.67% at a State, IBRA, Sub-IRA level. At an LGA level, 36.9% of the native vegetation's extent remains.</p> <p>Mapping results from the vegetation and flora survey conducted by GHD (2020a) described one vegetation types (<i>Acacia rostellifera</i> open woodland to woodland) within the application area, consistent with BVA 17.</p> <p>Sixty-four flora taxa (including subspecies and varieties) representing 26 families and 50 genera were recorded from the survey area during the field survey. This total comprised 49 native taxa and 15 introduced flora taxa. The species diversity ranged from 14 taxa per 100 m². As such, the species diversity is comparatively lower than that known within a 10 km radius, as according to <i>NatureMap</i> 455 flora taxa have been recorded (GHD 2020a).</p> <p>The application area is not within a TEC or PEC.</p> <p>Based on available range and habitat type, two priority flora species are considered to potentially occur in the application area. No threatened or priority flora taxa were recorded from the application area (GHD 2020a and 2020b).</p> <p>The application area is mostly cleared. Where native vegetation is present, vegetation conditions were rated good to completely degraded. Much of the understorey comprises weeds (GHD, 2020a).</p> <p>A total of 31 fauna species were recorded within the broader survey area. Of these, 24 are native, and seven introduced. One Migratory/Marine listed EPBC Act fauna species – <i>Pandion cristatus</i> (Osprey) was recorded nesting outside the application area. A 100 metre buffer has been implemented around the nesting site to ensure clearing will not impact on the nesting site (Figure 5).</p> <p>The proposal includes clearing up to 2.6 hectares of degraded to completely degraded vegetation for purpose of mine pit expansion. The clearing area will be rehabilitated as per Mine Closure requirements.</p>	The proposed clearing not considered to be at variance with this Principle.
Principle (b) – Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant	<p>The EPBC Act PMST, <i>NatureMap</i> and DBCA Threatened and Priority Flora databases identified the presence/potential presence of 48 conservation significant flora taxa within 10 km of the survey area. Of these two were considered as likely to occur including <i>Apus pacificus</i> (Fork-tailed Swift) and <i>Falco peregrinus</i> (Peregrine Falcon).</p>	The proposed clearing is not considered to be at variance with this Principle.

M70/204 Supporting Information

GMA Mining Australia

Clearing Principle	Assessment	Conclusion
habitat for fauna indigenous to Western Australia.	One Migratory/Marine listed fauna species <i>Pandion cristatus</i> (Osprey) nesting site was recorded outside the application area. A 100-metre buffer has been implemented around the nesting site. The nesting site is approximately 1.36 kms north of the amendment area.	
Principle (c) – Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, rare flora.	No Threatened (Declared Rare) flora was recorded from the application area (GHD 2020a and 2020b).	The proposed clearing is not considered to be at variance with this Principle.
Principle (d) – Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a threatened ecological community.	There are no known TECs within the application area. The vegetation types mapped within the application area are not considered to be representative of the TEC or PEC (GHD, 2020a).	The proposed clearing is not considered to be at variance with this Principle.
Principle (e) – Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.	The application area comprises small tracts of remnant vegetation, one vegetation types were described by GHD (2020a) including <i>A. rostellifera</i> open woodland to woodland. A comparison of vegetation types with the Beard Vegetation Associations mapped within the application area, indicates that vegetation type 1 closely aligns with BVA 17 (<i>Acacia rostellifera</i> dense thicket at 6 m in height, principal species comprise of <i>Alyogyne cuneiformis</i> , <i>Pimelea floribunda</i> and <i>Melaleuca cardiophylla</i>). The current extent of vegetation association Greenough 17 is greater than 30% of its pre-European extent at State, IBRA regional and sub-regional, and LGA levels.	The proposed clearing is not considered to be at variance with this Principle.
Principle (f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.	There are no watercourses within the application area. The nearest watercourse is the Hutt River, located 4 km south of the application area (GHD 2020). There are no wetlands within the application area. The nearest wetland is the Hutt Lagoon, which is approximately 450 metres from the area (GHD 2020).	The proposed clearing is not considered to be at variance with this Principle.
Principle (g) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation	The deep sands of the area have a high to very high wind erosion risk. GMA proposes to expand the existing mining pit southwards progressively. Given the proposed activities' disturbance footprint includes a narrow tract of vegetation (regrowth) it is unlikely to cause appreciable land degradation due to the narrow tract of vegetation being cleared.	The proposed clearing is not considered to be at variance with this Principle.

M70/204 Supporting Information

GMA Mining Australia

Clearing Principle	Assessment	Conclusion
Principle (h) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.	There are no conservation areas within the application area. The nearest reserve is Utcha Well, located approximately 12.5 kilometres north of the application area. There is no direct linkage between the reserve and the application area; therefore, it is unlikely that clearing will impact the environmental values of the nearby reserve. Also, the proposed clearing is temporary and will be returned to pre-mining vegetation assemblages as per the Mine Closure Plan and Notice of Intent conditions.	The proposed clearing is not considered to be at variance with this Principle.
Principle (i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water	<p>Due to the porous nature of the soils, any rainfall rapidly infiltrates directly through limestone. It is expected most of the surface water will rapidly infiltrate.</p> <p>The progressive and final rehabilitation of the mining pit area will incorporate re-contouring to blend in with the surrounding landscape and ensure any pre-mining landforms are reinstated. As a result, with this management approach, there will be no effect on surface water flow.</p> <p>The clearing is not considered likely to alter the quality of surface or groundwater within the application area. Mining operations are above the groundwater table as per Mine Closure Plan and Notice of Intent. The water table is too deep (greater than 16 to 35 m bgl) to support root systems of any species (URS 2013).</p>	The proposed clearing is not considered to be at variance with this Principle.
Principle (j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence of flooding.	<p>The climate of the application is semi-arid to Mediterranean climate with 400 to 500 mm of rainfall per annum (Desmond and Chant, 2002). The region experiences short mild wet winter and the remainder of the year being warm to hot, dry to windy. Due to the porous nature of the soils, any rainfall rapidly infiltrates directly through limestone. It is expected most of the surface water will rapidly infiltrate.</p> <p>Clearing of native vegetation is not expected to cause or exacerbate the incidence or intensity of flooding. The application area occurs on sandy soils which are not prevalent to flooding events.</p>	The proposed clearing is not considered to be at variance with this Principle.

7. Reference

- Beard and Burns (1976) the Vegetation of Geraldton Area Western Australia, Map and Explanatory Memoir
- Desmond, A and Chant, A (2001) Geraldton Sandplains (GS2 – Geraldton Hills Subregion). A Biodiversity Audit of Western Australia's 53 Biogeographical Subregions in 2002.
- EPA (2016) Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment
- GHD (2020a) *Lynton Mine Expansion Biological Survey*. Unpublished. Prepared for GMA Garnet
- GHD (2020b) Targeted *Caladenia bryceana* subsp. *cracens* survey and conservation listed flora survey of proposed haul road. Unpublished. Prepared for GMA Garnet.
- URS (2013) *Hose Mine Hydrological Assessment*. Unpublished. Prepared for GMA Garnet.



M70/204 Supporting Information

GMA Mining Australia

Appendix A. Environmental Surveys



GMA Garnet Pty Ltd
Lynton Mine Expansion
Biological Survey

February 2020

Executive summary

GMA Garnet Pty Ltd (GMA) currently own and operate the open pit Hose and Lynton Mines, located near Port Gregory, Western Australia.

GHD Pty Ltd (GHD) was commissioned to undertake a biological assessment across three tenements M70/204, M70/1330 and M70/259. The purpose of the survey was to delineate key flora, vegetation and fauna aspects.

The outcome of the survey and information supplied in the biological survey will be used to inform the environmental assessment and approvals process. This report is subject to, and must be read in conjunction with, the limitations set out in section 1.7 and the assumptions and qualifications contained throughout the report.

Key flora findings

- Three vegetation types were identified in the survey area, not including previously cleared areas (mining areas, tracks, cleared areas with no native species)
- The condition of the vegetation ranged from Good to Completely Degraded. Areas mapped as Good had vegetation that was largely intact with native species present across each structural layer, although had high weed cover and signs of high grazing impacts from pigs and kangaroos. Much of the survey area had undergone historical clearing (exploration) and rehabilitation of some of these areas
- No vegetation communities identified in the survey area were consistent with Threatened or Priority Environmental Communities
- Sixty-four flora taxa (including subspecies and varieties) representing 26 families and 50 genera were recorded from the survey area during the field survey
- No *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) or *Biodiversity Conservation Act 2016* (BC Act) listed flora were recorded within the survey area. No Priority flora, as listed by the Department of Biodiversity Conservation and Attraction, were recorded within the survey area
- The likelihood of occurrence assessment post-field survey concluded three species are considered possible to occur, five species unlikely to occur, and 40 species highly unlikely to occur in the survey area. The species considered possible to occur are; *Caladenia Bryceana* subsp. *cracens*, *Anthocercis intricata* (P3) and *Balladonia aervoides* (P3).

Key fauna findings

- Five broad habitat types (including rehabilitated and cleared areas) were recorded during the survey
- Thirty-one fauna species were recorded within the survey area, including 21 bird, 8 mammal and 2 reptile species. Of these, 24 are native and seven are introduced/feral
- No Threatened fauna listed under the EPBC Act and/or BC Act or Priority fauna species listed by the DBCA was recorded during the survey
- The Eastern Osprey (*Pandion cristatus*) which is listed as Migratory and Marine under the EPBC Act and under International Agreement under the BC Act was recorded during the survey

- Of the 35 conservation significant fauna identified in the desktop searches one species has been identified as present (Osprey), two are considered likely to occur and the remaining species are considered unlikely or highly unlikely to occur within the survey area.

Table of contents

1.	Introduction	1
1.1	Project background	1
1.2	Purpose of this report	1
1.3	Study area.....	1
1.4	Survey area.....	1
1.5	Scope of works.....	1
1.6	Relevant legislation, conservation codes and background information	2
1.7	Report limitations and assumptions	2
2.	Methodology	4
2.1	Desktop assessment	4
2.2	Field survey.....	4
2.3	Limitations.....	7
3.	Desktop assessment	11
3.1	Regional biogeography.....	11
3.2	Climate.....	11
3.3	Hydrology.....	12
3.4	Geology, landforms and soils.....	12
3.5	Land use	12
3.6	Vegetation and flora	12
3.7	Fauna.....	15
3.8	Previous survey results	16
4.	Field results	18
4.1	Flora and vegetation.....	18
4.2	Fauna.....	26
5.	Recommendations	33
5.1	Recommendations	33
6.	References	34

Table index

Table 1	Data collected during the flora and vegetation field survey	5
Table 2	Flora and fauna survey limitations	8
Table 3	Extents of vegetation associations mapped within the survey area (GoWA 2020)	14
Table 4	Vegetation types identified within the survey area	19
Table 5	Extent of vegetation condition ratings mapped in the survey area	25
Table 6	Fauna habitat types identified within the survey area	27

Table 7 Conservation listed fauna Osprey location coordinates31

Table 8 Conservation significant fauna present or likely to occur within the survey area.....32

Appendices

Appendix A – Figures

Appendix B – Relevant legislation, background information and conservation code

Appendix C – Desktop searches

Appendix D – Flora data

Appendix E – Fauna data

1. Introduction

1.1 Project background

GMA Garnet Pty Ltd (GMA) currently own and operate the open pit Hose and Lynton Mines, located near Port Gregory, Western Australia (WA). Mining activities are currently undertaken within M70/926, M70/204 and M70/968. Mining is undertaken using an open-cut sand mining methods. Mobile earthmoving equipment, including front-end loaders, excavator and dump trucks are used for pit excavation and backfilling. Soil and overburden are moved ahead of ore excavation and replaced in their original stratigraphic order over the backfilled tailings.

GMA are currently in the process of planning for the expansion of their operations within the Lynton Mine, located to the east of Hutt Lagoon, near Port Gregory in WA. Biological surveys are required to be undertaken to inform this expansion.

1.2 Purpose of this report

GMA commissioned GHD Pty Ltd (GHD) to undertake a biological assessment across three tenements M70/204, M70/1330 and M70/259. The purpose of the survey was to delineate key flora, vegetation and fauna aspects.

The outcome of the survey and information supplied in the biological survey will be used to inform the environmental assessment and approvals process.

1.3 Study area

The study area of the project is located in Geraldton, and encapsulates an area of 10 km around the survey area.

1.4 Survey area

The survey area for this project is located at Port Gregory, approximately 96 kilometres (km) north of Geraldton in the mid-west of WA. The survey area is 413 hectare (ha) in size and occurs across three tenements (M70/204, M70/1330 and M70/259). The survey area boundary is shown in Figure 1, Appendix A.

1.5 Scope of works

The scope of works was to undertake a desktop assessment and biological survey of the survey area. The following actions were completed to fulfil the scope:

- A desktop assessment of the survey area prior to the field survey to identify biological features and constraints, which may be in, or near the survey area
- A review of relevant publicly available or supplied by GMA environmental reports
- A field survey to verify/ground truth the desktop assessment findings through a detailed (single-season) vegetation and flora survey and level 1 fauna survey
- Identification and mapping of vegetation types to a scale appropriate for the bioregion and described according to the National Vegetation Information System (NVIS) structure and floristics
- Identification and mapping of Threatened or Priority Ecological Communities (TECs or PECs) inferred through the use of quadrats and relevés

- Assessment of the survey area's flora species diversity, density, composition, structure and weed cover, recording the percentage of each in nominated quadrats
- Delineation and mapping of fauna habitat types
- A flora and fauna likelihood of occurrence assessment based on the vegetation units and fauna habitat present within the survey area and known species distribution and habitat requirements
- Mapping using Geographic Information Systems (GIS) mapping software
- A concise report (this document) on the findings of the biological survey and targeted flora assessment.

1.6 Relevant legislation, conservation codes and background information

In WA some ecological communities, flora and fauna are protected under both Federal and State Government legislation. In addition, regulatory authorities also provide a range of guidance and information on expected standards and protocols for environmental surveys.

An overview of key legislation and guidelines, conservation codes and background information relevant to this biological survey is provided in Appendix B.

1.7 Report limitations and assumptions

This report has been prepared by GHD for GMA and may only be used and relied on by GMA for the purpose agreed between GHD and the GMA as set out in section 1.4 of this report.

GHD otherwise disclaims responsibility to any person other than GMA arising in connection with this report. GHD also excludes implied warranties and conditions, to the extent legally permissible.

The services undertaken by GHD in connection with preparing this report were limited to those specifically detailed in the report and are subject to the scope limitations set out in the report.

The opinions, conclusions and any recommendations in this report are based on conditions encountered and information reviewed at the date of preparation of the report. GHD has no responsibility or obligation to update this report to account for events or changes occurring subsequent to the date that the report was prepared.

The opinions, conclusions and any recommendations in this report are based on assumptions made by GHD described in this report (refer section(s) 1.4 of this report). GHD disclaims liability arising from any of the assumptions being incorrect.

GHD has prepared this report on the basis of information provided by GMA and others who provided information to GHD (including Government authorities), which GHD has not independently verified or checked beyond the agreed scope of work. GHD does not accept liability in connection with such unverified information, including errors and omissions in the report which were caused by or omissions in that information.

The opinions, conclusions and any recommendations in this report are based on information obtained from, and testing undertaken at or in connection with, specific sample points. Site conditions at other parts of the site may be different from the site conditions found at the specific sample points.

Site conditions may change after the date of the field survey. GHD does not accept responsibility arising from, or in connection with, any change to the site conditions. GHD is also not responsible for updating this report if the site conditions change.

This report has assessed the flora and fauna within the survey area (Figure 1, Appendix A). Should the survey area change or be refined, further assessment may be required.

2. Methodology

2.1 Desktop assessment

Prior to the commencement of the field survey, a desktop assessment was undertaken to identify relevant environmental information pertaining to the survey area and within 10 km of the survey area (referred to herein as the study area). This included a review of:

- The Department of the Environment and Energy (DotEE) Protected Matters Search Tool (PMST) to identify communities and species listed under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) potentially occurring within the study area (DotEE 2019) (Appendix C)
- The Department of Biodiversity, Conservation and Attractions (DBCA) TEC and PEC database to determine the potential for conservation significant communities to be present within the study area
- The DBCA *NatureMap* database for flora and fauna species previously recorded within the study area (DBCA 2019) (Appendix C)
- The DBCA Threatened (Declared Rare) and Priority Flora (TPFL) database and the WA Herbarium database (WAHERB) for Threatened flora listed under the *Biodiversity Conservation Act 2016* (BC Act) and listed as Priority by the DBCA, previously recorded within the study area
- Existing datasets including previous pre-European vegetation mapping of the survey area (Beard 1976), aerial photography, hydrology information to provide background information on the variability of the environment, likely vegetation units and fauna habitats and to identify areas that potentially contain TECs and PECs
- Existing flora, fauna and vegetation reports and/or data:
 - GMA Port Gregory Mine Tenement M70/968 Vegetation, Flora and Fauna Assessment (GHD 2013)
 - GMA Garnet Port Gregory Mine Targeted Flora Survey (GHD 2014)
 - GMA Garnet Mining Lease M70/926 Biological Survey (GHD 2016)
 - GMA Port Gregory Mine Site M70/1380 Biological Survey (GHD 2019).

The mapped biological constraints within 10 km of the survey area is provided in Figure 2, Appendix A.

2.2 Field survey

2.2.1 Flora and vegetation

Two GHD botanists/ecologists completed a detailed (single-season) flora and vegetation survey from 8 - 12 December 2019. The field survey was undertaken to verify the results of the desktop assessment, identify and describe the dominant vegetation units, assess vegetation condition, and identify and record vascular flora taxa present at the time of survey. The survey seasonal timing did not allow for targeted searches for key conservation significant flora species, however potential habitat for significant flora were identified and mapped where present.

The survey methodology employed by GHD was undertaken with reference to the Environmental Protection Authority (EPA) Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment (EPA 2016a).

Data collection

Field survey methods involved a combination of sampling quadrats and transects located in identified vegetation units and traversing the survey area by vehicle and foot. Twenty non-permanent quadrats and seven relevés were placed within the survey area, which is deemed suitable for the represented vegetation types identified. Transects were spaced at 10 m intervals when traversing a specified vegetation type. The degraded condition of vegetation in much of the survey area caused transects to become spaced further apart (<50 m) as the survey progressed.

Quadrats (measuring 10 m x 10 m – area of 100 m²) were located within each identified vegetation unit. Field data at each quadrat was recorded on a pro-forma data sheet and included the parameters detailed in Table 1.

Table 1 Data collected during the flora and vegetation field survey

Aspect	Measurement
Collection attributes	Site code, personnel/recorder, date, quadrat dimensions, photograph of the quadrat, marking method
Physical features	Landform, aspect, slope, soil attributes, ground surface cover, leaf and wood litter
Location	Coordinates recorded in GDA94 datum (Zone 50) using a hand-held Global Positioning System (GPS) tool to accuracy approximately ±5 m
Vegetation condition	Vegetation condition in accordance with the vegetation condition rating scale for the South-West Interzone Botanical Province (EPA 2016)
Disturbance	Level and nature of disturbances (e.g. weed presence, fire and time since last fire, impacts from grazing, exploration activities).
Flora	List of dominant flora from each structural layer, list of all species within the quadrat including stratum, average height and cover (using National Vegetation Information System (NVIS)).

Quadrat data is provided in Appendix D. A flora inventory was compiled from taxa listed in described quadrats, relevés and opportunistic floristic records throughout the survey area (Appendix D).

Vegetation units

Vegetation units were identified and boundaries delineated using a combination of aerial photography, topographical features, field data/observations and statistical analyses.

Vegetation units were described based on structure, dominant taxa and cover characteristics as defined by quadrat data and field observations. Vegetation unit descriptions follow the NVIS and are consistent with NVIS Level V (Association). At Level V, three (or more) taxa per stratum are used to describe the association (NVIS Technical Working Group 2017).

Statistical analyses

PRIMER version 6 (Clarke and Gorley 2006) was used to examine the similarity between sites using collected data. A presence/absence matrix was created of all taxa (including perennials and annuals) present in GHD quadrats. The dissimilarity between quadrats was determined using the Bray-Curtis measure and the Resemblance function in PRIMER. A Cluster analysis (using Agglomerative Hierarchical Clustering technique) based on group average was undertaken using the Bray-Curtis similarity matrix and results presented as a dendrogram. In addition, a nonmetric multi-dimensional scaling analysis (MDS) was undertaken using the Bray-Curtis similarity matrix and results presented as a two dimensional scatter plot. The analysis was repeated using removing all singleton taxa. The outputs of the PRIMER analysis were used to inform decisions on vegetation units.

Vegetation condition

The vegetation condition of the survey area was assessed and mapped in accordance with the vegetation condition rating scale for the South-West Interzone Botanical Province of WA (devised by Keighery (1994) and adapted by the EPA (2016a)). The scales recognise the intactness of vegetation and consists of six rating levels as outlined in Appendix B.

Flora identification and nomenclature

Species that were well known to the survey botanists were identified in the field; all other species were collected and assigned a unique collection number to facilitate tracking. Flora collections were made under Joel Collin's DBCA Scientific Flora License (#FB620000200). All specimens collected during the field assessment were dried and processed in accordance with the requirements of the WA Herbarium. Species were identified by a qualified taxonomist using taxonomic literature, electronic keys and online electronic databases.

The conservation status of all recorded flora was compared against the current lists available on *FloraBase* (WA Herbarium 2020) and the EPBC Act Threatened species database provided by DotEE (2020). Nomenclature used in this report follows that used by the WA Herbarium as reported on *FloraBase* (WA Herbarium 2020).

2.2.2 Fauna

GHD ecologists undertook a Level 1 fauna survey (reconnaissance survey) in conjunction with the flora and vegetation survey from 8 - 12 December 2019. The survey area was traversed on foot over the course of the survey to identify and describe the dominant fauna habitat types present and their condition, assess habitat connectivity, and identify and record fauna species within the survey area. An assessment of the likelihood of conservation significant fauna occurring within the survey area was also undertaken.

The survey methodology employed by GHD was undertaken in accordance with the EPA *Technical Guidance – Sampling methods for terrestrial vertebrate fauna* (EPA 2016b) and *Technical Guidance – Terrestrial Fauna Surveys* (EPA 2016c).

Opportunistic fauna searches

Opportunistic fauna searches were conducted across the survey area. Opportunistic searches involved:

- Searching the survey area for tracks, scats, bones, diggings and feeding areas for both native and introduced/feral species
- Visual and aural surveys, which accounted for many bird species potentially utilising the survey area
- Recording GPS locations of any conservation significant fauna species observed.

Fauna species identification

Identification of fauna species was made in the field using available field guides and electronic guides (e.g. Morcombe 2011). Where identification was not possible, photographs of specimens were collected to be later identified.

Fauna nomenclature

Nomenclature used in this report follows that used by the Western Australian Museum and the DBCA NatureMap database (DBCA 2019) with the exception of birds, where Christidis & Boles (2008) was used.

2.3 Limitations

2.3.1 Desktop limitations

The EPBC Act PMST is based on bioclimatic modelling for the potential presence of species. As such, this does not represent actual records of the species within the area. The records from the DBCA searches of Threatened fauna provide more accurate information for the general area and local occurrence. However, some collection, sighting or trapping records cannot be dated and often misrepresent the current range of Threatened species

2.3.2 Field survey limitations

The EPA (2016a, b) states that flora and fauna survey reports for environmental impact assessment in WA should contain a section describing the limitations of the survey methods used. The limitations and constraints associated with this field survey are discussed in Table 2.

Table 2 Flora and fauna survey limitations

Aspect	Constraint	Comment
Sources of information and availability of contextual information.	Nil	<p>Adequate information is available for the survey area.</p> <ul style="list-style-type: none"> • Pre-European vegetation mapping (Beard 1976) • GHD (2019) GMA Garnet Port Gregory Mine Mining Tenement M70/1380 Biological Survey • GHD (2016) GMA Garnet Mining Lease M70/926 • GHD (2014) GMA Garnet Port Gregory Mine Targeted Flora Survey • GHD (2013) GMA Garnet Port Gregory Mine M70/968 Vegetation, Flora and Fauna Assessment.
Scope (what life forms were sampled etc.)	Nil	<p>Vascular flora and terrestrial vertebrate fauna were sampled during the survey. Non-vascular flora, invertebrate and aquatic fauna were not surveyed.</p> <p>Adequate time was available to complete the biological survey to the required standard.</p>
Proportion of flora collected and identified (based on sampling, timing and intensity) Proportion of fauna identified, recorded and/or collected	Moderate	<p>The flora and vegetation survey was undertaken from 8 - 12 December 2019. Spring is considered the most optimal time to undertake vegetation surveys in the Geraldton bioregion. This survey is considered an out of season survey as seasonal conditions at the time of the survey are deemed unsatisfactory for some annuals and ephemeral species, such as orchids. The survey sampling and intensity was considered adequate. The vegetation survey was a broad scale and targeted assessment, undertaken to identify and describe the dominant vegetation units and map conservation significant flora. The portion of flora collected and identified was considered appropriate for the level of experience of the Senior Botanist undertaking the survey. All taxonomic groups were considered to be represented. The portion of flora collected and identified was considered moderate; and it is likely the survey under-recorded some grass species (Poaceae), annuals and herbs due to lower than average rainfall and out of season timing. However, based on the likelihood assessment it is unlikely these species would be conservation significant.</p> <p>The reconnaissance fauna survey was undertaken from 8 - 12 December 2019. The fauna assessment sampled those species that can be easily seen, heard or have distinctive signs, such as tracks, scats, diggings, etc. Many cryptic species would not have been identified during a reconnaissance survey and seasonal variation within species often requires targeted surveys at a particular time of the year. Of the fauna species recorded during the survey, all were identified to species level.</p>
Flora determination	Minor	<p>Flora determination was undertaken by GHD botanist/ecologist in the field and at the WA Herbarium by Botanist Frank Obbens.</p> <p>Four taxa could be identified to genus level only, due to lack of flowering and/or fruiting material required for identification. None of these taxa were considered to be conservation significant species.</p> <p>The taxonomy and conservation status of the WA flora is dynamic. This report was prepared with reliance on taxonomy and conservation status current at the time of report development, but it should be noted this may change in response to ongoing research and review of the International Union for Conservation Nature criteria.</p>

Aspect	Constraint	Comment
Completeness and further work which might be needed (e.g. was the relevant area fully surveyed)	Minor	The majority of the survey area was accessible and was accessed by foot and vehicle. There were some areas of large piles of dead Acacia logs and branches which restricted movement by foot, however, these areas could still be ground-truthed.
Mapping reliability	Minor	<p>The vegetation was mapped using high-resolution ESRI aerial imagery obtained from Landgate, topographical features, previous broad scale mapping (Beard 1976) and field data.</p> <p>Data was recorded in the field using hand-held GPS tools (e.g. Samsung tablet and Garmin GPS). Certain atmospheric factors and other sources of error can affect the accuracy of GPS receivers. The Garmin GPS units used for this survey are accurate to within ± 5 metres on average. Therefore the data points consisting of coordinates recorded from the GPS may contain inaccuracies.</p>
Timing/weather/season/cycle	Moderate	<p>The Detailed Flora and Level 1 fauna field survey was conducted in December 2019. In the six months prior to the flora survey (June -November), Lynton weather station (BoM 2020) recorded a total of 270 mm of rainfall. This rainfall total is slightly lower than the long-term average for the same period (June to November; 277 mm) (BoM 2020).</p> <ul style="list-style-type: none"> • The weather conditions during the field survey included: • Daily maximum temperatures ranging from 25 to 35 °C • Daily minimum temperature ranging from 14 to 22 °C • No rainfall occurred during the survey. <p>This survey is considered and an out of season flora survey as seasonal conditions at the time of the survey are deemed unsatisfactory for some annuals and ephemeral species, such as orchids. For majority of the other flora species the timing of detailed flora survey was considered appropriated due to a number of flora flowering or fruiting at the time of the survey.</p>
Disturbances (e.g. fire, flood, accidental human intervention)	Minor	Much of the survey area had undergone historical clearing (exploration) and rehabilitation of some of these areas. Newly cleared or disturbed areas (roads and tracks) were also prevalent, as well as completely cleared areas, where the mine development has progressed. These disturbances did not limit the survey.
Resources	Nil	Adequate resources were employed during the field survey. Two staff over five days were spent undertaking the flora and fauna survey using a dedicated botanist and ecologist.
Access restrictions	Nil	No access problems were encountered during the survey. There were some areas of large piles of dead Acacia logs and branches which restricted movement by foot, however, these areas could still be ground-truthed.
Experience levels	Nil	The botanist/ecologists who executed the survey were practitioners suitably qualified in their respective fields. Joel Collins, is suitably qualified with over 16 years' experience in undertaking flora and fauna surveys and assessments in Western Australia. Joel has extensive experience undertaking flora and

Aspect	Constraint	Comment
		assessments on the Geraldton Sandplains. Sarah Flemington (Ecologist) has three years experience undertaking flora and fauna surveys across the South West and across the arid region and interzone.

3. Desktop assessment

3.1 Regional biogeography

The survey area is located within the Geraldton Sandplains bioregion and Geraldton Hills sub-region as described by the Interim Biogeographic Regionalisation of Australia (IBRA).

This region comprises of sandy earths of an extensive undulating and lateritic sandplain mantling Permian to Cretaceous strata. This region occurs within the southern end of the Carnarvon Basin and the northern end of the Perth Basin, with exposed areas of Permian/Silurian siltstone and Jurassic sandstones mostly overlain by sandplains, alluvial plains and coastal limestone. The vegetation consists primarily of proteaceous heath with *Banksia* - York gum woodlands on alluvial plains and *Acacia* scrub on limestone (Desmond and Chant, 2002).

3.2 Climate

The survey area experiences a Mediterranean type climate, characterised by warm to hot dry summers and mild wet winters. The Bureau of Meteorology (BoM) Kalbarri weather station (Number 8251) is the nearest active weather station to the survey area with continuous long-term temperature data (approximately 54 km from the survey area). Climatic data from this site indicates the mean maximum temperature of the area ranges from 21.8 degrees Celsius (°C) in July to 34.2 °C in February, and the mean minimum temperature ranges from 9.7 °C in July to 20.6 °C in February (Plate 1) (BoM 2020). Rainfall data has been sourced from the Lynton weather station (Number 8075), which was likely to better represent the survey area. The average annual rainfall measured at Lynton is 405.9 mm with the average monthly rainfall ranging 3.5 mm in December to 93.7 mm in June. Rainfall was significantly higher than the average for the area in June 2019, at 171.5 mm (Plate 1) (BoM 2020). Rainfall was not recorded at Lynton in 2019 for the months September to December. Results from 2018 have been supplemented for those months. The majority of rainfall occurs in the winter months and is generally associated with frontal systems from the south west. The summer rains are associated with isolated thunderstorms and tropical lows.




Plate 1 Mean rainfall for Lynton and temperatures for Kalbarri

3.3 Hydrology

3.3.1 Groundwater

The Department of Water and Environmental Regulation (DWER) Perth Groundwater Map indicates the survey area is located in within the Gascoyne Groundwater Area.

3.3.1 Surface water

There are no watercourses or wetlands located within the survey area. The closest watercourse is the Hutt River, which is located approximately 4 km south of the survey area and flows west into the ocean.

The Hutt Lagoon, which is located directly west of the survey area, is listed as a wetland of national importance on the Directory of Important Wetlands in Australia (DIWA) (DEC 2009). Hutt Lagoon is a macroscale elongate sumpland aligned northwest to southeast, parallel to the coast. It neighbours macroscale elongate floodplains (to the northwest and southeast) that include more than twenty microscale elongate sumplands such as Utcha Swamp (Jaensch 1992). Water supply for the Hutt Lagoon derives from direct precipitation, surface inform from several minor creeks and seepage of groundwater (DEC 2009).

3.4 Geology, landforms and soils

The survey area is located within the Tumblagooda Sandstone, which is characterised by sandstone, with minor siltstone and granulate to pebble conglomerate. The majority of the survey area is located on the Tamala North Land System, described as low hills with relict dunes and some limestone outcrop, which forms a coastal band 3 to 7 km wide. Parts of the western boundary of the survey area are located within the Grey Land System, described as river beds, terraces and alluvial flats, includes dissected margins of relic alluvial plains (Rogers et al. 1996).

3.5 Land use

3.5.1 Conservation reserves and estates

There are no conservation reserves or estates located within or immediately adjacent to the survey area. The closest conservation reserve, the Utcha Well Nature Reserve (R 640), is located approximately 3 km north of the survey area (Figure 2, Appendix A).

3.5.2 Environmentally sensitive areas

One Environmentally Sensitive Areas (ESAs) is located approximately 200 m west of the survey area. This ESA is associated with the Hutt Lagoon and does not intersect the survey area (Figure 2, Appendix A).

No PECs or TECs are located within the survey area. The nearest PEC, Kalbarri ironstone community (P1) is located approximately 8 km east of the survey area.

3.6 Vegetation and flora

3.6.1 Broad vegetation mapping and extents

Broad scale (1:250,000) pre-European vegetation mapping of the survey area has been completed by Beard (1975) at an association level. The mapping indicates that the survey area intersects two broad vegetation associations (BVA):

- Shrublands; *Acacia rostellifera* thicket: wattle, casuarina and teatree acacia-allocauarina melaleuca alliance (association 17)
- Low forest; *Acacia rostellifera*: Acacia, Rottnest pine, coastal moort or mixed tropical forest, *Acacia rostellifera*, *Callitris preissii*, *Eucalyptus lehmannii*, *E. cornuta* (association 371).

The extents of these associations within the survey area are shown in Table 3.

The pre-European mapping has been adapted and digitised by Shepherd *et al.* (2002). The extent of the vegetation associations have been determined by the state-wide vegetation remaining extent calculations maintained by the DBCA (latest update March 2019 –Government of Australia (GoWA) 2020). As shown in Table 3 the current extent of vegetation association 17 is above 30 % of its pre-European extent at the State, IBRA bioregion, IBRA subregion and Local Government Authority (LGA) levels. Vegetation association 371 is below the 30% of the pre-European extents at all levels except for the LGA.

Table 3 Extents of vegetation associations mapped within the survey area (GoWA 2020)

Vegetation association	Scale	Pre-European extent (ha)	Current extent (ha)	Remaining (%)	Remaining within DBCA Managed lands (%)	Hectares (ha) within the survey area	% of current extent within the survey area
17	State: WA	76,633.84	67,605.49	88.22	13.06	52.70	0.07
	IBRA Bioregion: Geraldton Sandplains	54,078.08	45,159.85	83.51	13.44		0.11
	Sub-region: Geraldton Hills	49,605.04	42,016.28	84.70	13.26		0.12
	LGA: Shire of Northampton	49,549.89	41,939.33	84.64	13.29		0.12
371	State: WA	32,816.04	3,499.60	10.66	6.92	360.16	10.29
	IBRA Bioregion: Geraldton Sandplains	32,807.53	3,499.10	10.67	6.92		10.29
	Sub-region: Geraldton Hills	32,807.53	3,499.10	10.67	6.92		10.29
	LGA: Shire of Northampton	5,749.92	2,142.08	36.94	10.69		16.81

3.6.2 Conservation significant ecological communities

The PMST (DotEE 2019) did not identify any TECs listed under the EPBC Act within the study area. The DBCA TEC/PEC database identified two Priority 1 PECs within the study area, however none are known to occur within the survey area (Figure 2, Appendix A). The two PECs identified are (DBCA 2019a):

- Kalbarri Ironstone Community (Priority 1 PEC) – winter wet, mallee/Melaleuca over herbs. Dense shrubland when burnt. Surrounded by sandplain. Yerina springs and north Euardy Station. Z-bend loop, Junga Dam. The taxon *Eremophila microtheca* (previously declared rare flora) occurs in this community (located approximately 8 km east of the survey area)
- Shrubland of the Northampton Area, dominated by *Melaleuca* species over exposed Kockatea Shale (Priority 1 PEC) – Heath on breakaways located in Port Gregory, west of Northampton. Community includes priority taxa; *Ptilotus chortophytum* (P1), *Leucopogon* sp. Port Gregory, *Ozothamnus* sp. Northampton, *Gastrolobium propinquum* (P1), outlier of *Ptilotus helichrysoides*. Unusual geology (Kockatea Shale) outcropping at surface (closest record is approximately 5 km south-east of the survey area).

3.6.3 Flora diversity

The *NatureMap* database (DBCA 2019) identified 455 flora taxa, representing 85 families and 235 genera previously recorded within 10 km of the survey area. This total comprised 403 native flora taxa and 52 naturalised (introduced) flora taxa. Dominant families recorded included Myrtaceae (48 taxa), Asteraceae (36 taxa), Fabaceae (36 taxa) and Poaceae (30 taxa).

The *NatureMap* database search is provided in Appendix C.

3.6.4 Conservation significant flora

The EPBC Act PMST, *NatureMap* and DBCA Threatened and Priority Flora databases identified the presence/potential presence of 48 conservation significant flora taxa within 10 km of the survey area (Appendix C). The desktop searches recorded:

- 13 Threatened flora taxa listed under the EPBC Act and/or BC Act
- One taxa listed as Threatened under the EPBC Act and Priority 4 by DBCA
- Six Priority 1 taxa
- Seven Priority 2 taxa
- 14 Priority 3 taxa
- Seven Priority 4 taxa.

The locations of conservation significant flora registered on the DBCA databases are mapped on Figure 2, Appendix A. Previously recorded conservation significant flora in report *GMA Garnet Port Gregory Mine Mining Tenement M70/1380 Biological Survey* (see section 3.8) have also been represented in Figure 2, due to the proximity of these records to the survey area.

3.7 Fauna

3.7.1 Fauna diversity

The *NatureMap* database identified 176 terrestrial vertebrate fauna species previously recorded within 10 km of the survey area. This total comprised of four amphibians, 151 birds, 14 reptiles and seven mammals. Of the 176 fauna species previously recorded, 172 are native species and

four are naturalised (introduced) species (under the *Biosecurity and Agriculture Management Act 2007* (BAM Act)).

The *NatureMap* database search is provided in Appendix C.

3.7.2 Conservation significant fauna

The EPBC Act PMST and DBCA databases identified the potential presence of 33 conservation significant fauna within 10 km of the survey area (Appendix C). This total does not include species identified as exclusively marine (e.g. marine mammals and reptiles) as no marine habitat is present within the survey area.

The species listed include:

- 25 listed as Threatened under the EPBC Act and/or BC Act
- Five bird species listed as Migratory (terrestrial or wetland) only, under the EPBC Act and/or Schedule 5 (Migratory birds protected under an international agreement) of the BC Act
- One species listed as Schedule 7 (Specially Protected) under the BC Act
- Two listed as Priority 4 by the DBCA.

3.8 Previous survey results

GMA Garnet Port Gregory Mine Mining Tenement M70/1380 Biological Survey

The vegetation, flora and fauna assessment was undertaken by GHD between July and August 2019. The survey area was located within mining tenement M70/1380, approximately 1,465 ha in size. The key findings of the survey include:

- Eight vegetation types were delineated from the survey area. None of the vegetation types were considered representative of either a conservation significant ecological community or other significant vegetation community
- Six broad fauna habitat types were recorded from the survey area. Marginal foraging habitat was recorded that included 31.4 ha of Banksia woodland
- One hundred and sixty-five flora taxa (including subspecies and varieties) representing 61 families and 121 genera were recorded from the survey area. This total included 18 introduced flora
- One flora species (*Senna planitiicola*) considered to be an extension of its known range and has not been previously recorded within the Geraldton Sandplains IBRA however the species was recorded within a farming paddock and was considered to have been planted
- One EPBC Act / BC Act listed flora and three DBCA Priority listed flora species were identified within the survey area:
 - *Caladenia bryceana* subsp. *cracens* (Vulnerable – EPBC Act, Endangered – BC Act)
 - *Melaleuca huttensis* (Priority 3)
 - *Anthocercis intricata* (Priority 3)
 - *Diuris recurva* (Priority 4)
- A total of 42 fauna species, including one amphibian, 28 birds, nine mammals and four reptiles were recorded within the survey area of which eight were introduced
- No conservation significant fauna was identified during the survey.

GMA Garnet Mining Lease M70/926 Biological Survey

A biological survey was undertaken by GHD in August 2016 to identify environmental constraints within mining tenement M70/926, located approximately 7 km north of M70/1380. The key findings include:

- A total of four vegetation types were delineated from the study location including:
 - *Acacia rostellifera* Low Forest
 - *Acacia rostellifera* Tall Open Shrubland
 - *Acacia rostellifera* Low Shrubland on Shallow Soils
 - Cleared and Degraded.
- A total of 60 flora taxa from 28 families were recorded at the study location, of which 26 were introduced
- Twenty-one fauna taxa were recorded from the study location, including 17 birds and six mammals
- No conservation significant communities, flora or fauna were recorded.

GMA Garnet Port Gregory Mine Targeted Flora Survey

A targeted flora survey was undertaken by GHD in August 2014 for the Threatened flora species *Caladenia bryceana* subsp. *cracens* at M70/968. The key findings include:

- No Threatened Flora was recorded during the survey
- Habitat for the threatened orchid was considered extremely marginal
- Evidence of disturbances such as wild pigs, which further reduced the likelihood of the species occurrence
- Two State-listed Priority species were recorded including 23 individual plants of *Melaleuca huttensis* (Priority 1) and 54 individual plants of *Anthocercis intricata* (Priority 3).

GMA Garnet Port Gregory Mine Tenement M70/968 Vegetation, Flora and Fauna Assessment

The vegetation, flora and fauna assessments were undertaken by GHD in August 2013 to identify environmental constraints within mining tenement M70/968. The key findings include:

- Six vegetation types were recorded from the study location, none of which were considered conservation significant
- A total of 75 flora taxa from 39 families were recorded from the study location, No conservation significant communities or flora were present
- The study location was considered to be very marginal habitat for the Threatened *Caladenia bryceana* subsp. *cracens*
- There was evidence of current disturbances from feral fauna (rabbits and pigs)
- A total of five birds and two mammals were recorded from the study location, none were conservation significant.

4. Field results

4.1 Flora and vegetation

4.1.1 Vegetation types

Three vegetation types were identified in the survey area, not including previously cleared areas and regrowth/rehabilitated areas, which were mapped separately from the vegetation types (Table 4). Much of the survey area had undergone historical clearing (exploration) and rehabilitation of these areas. Newly cleared or disturbed areas (roads and tracks) were also prevalent, as well as completely cleared areas, where the mine development has progressed. Rehabilitated areas contained fragmented vegetation resembling VT01 (*Acacia rostellifera* open woodland), but with an understory dominated by introduced grasses.


A significant portion of the survey area consisted of *Acacia rostellifera* open woodland to woodland (VT01) (67.9% 280.43 ha). This vegetation type occurred in low-lying and middle to upper slopes. Both BVA 371 (Low forest, *Acacia rostellifera*) and BVA 17 (Shrublands, *Acacia rostellifera*) intersected VT01. The description of BVA 17 by Beard & Burns (1976) aligns closely with VT01 (*Acacia rostellifera* dense thicket at 6 m in height, principal species comprise *Alyogyne cuneiformis*, *Pimelea floribunda* and *Melaleuca cardiophylla*). *Melaleuca cardiophylla* shrubland to open shrubland (VT02) (6.8% 28.30 ha) occurred on the upper slopes, west facing of the survey area, to the north, on limestone. Beard & Burns (1976) describes *M. cardiophylla* as dominant, more or less as a sole species on the rockiest and steepest places in BVA 17, which strongly aligns with VT02. *Myoporum insulare* shrubland (VT03) (0.1% 0.45 ha) was isolated in occurrence, and consisted of chenopod shrubland with *Frankenia pauciflora* and *Threlkeldia diffusa*, due to saline influence. All 0.45 ha of VT03 was mapped within BVA 371.


The vegetation types VT01 and VT02 generally align with BVA 17, where VT01 contains mostly wooded areas (Low forest), and VT02 contains *Melaleuca* shrublands (thicket). VT03 does not align with BVA 371, which is likely, considering the small size of the vegetation type. However, VT03 does align with association 125 (Salt lake, lagoon, clay pan) (Beard 1975) that describes the Hutt Lagoon.


Floristic analysis


The similarity between the quadrats were examined using PRIMER with all species recorded in the quadrats analysed based on presence/absence. Quadrat Lyn12 (VT03) was removed from the analysis as it was very different from the other quadrats it impacted the MDS results, which did not allow for a useful analysis. A stress value of 0.16 was produced indicating a fair representation (Plate 2). The MDS scatter plot for VT01 and VT02 quadrats loosely grouped together, particularly for VT01. The vegetation types were mapped using a combination of statistical analysis, dominant species, landforms and field observations.

Table 4 Vegetation types identified within the survey area

Vegetation Types	Vegetation type description and landform	Representative quadrat/releve and Total area (ha)	Representative photograph
VT01 - <i>Acacia rostellifera</i> open woodland to woodland	<i>Acacia rostellifera</i> open woodland to woodland over <i>Rhagodia preissii</i> subsp. <i>obovata</i> , <i>Pimelea microcephala</i> subsp. <i>microcephala</i> , <i>Olearia</i> sp. Kennedy Range (G. Byrne 66) and <i>Stylobasium spathulatum</i> open shrubland over <i>Austrostipa elegantissima</i> and <i>*Ehrharta longiflora</i> open grassland to grassland. Other common species include <i>Alyogyne hakeifolia</i> , <i>Roepera fruticulosa</i> , <i>Commicarpus australis</i> and <i>Euphorbia boophthona</i> . Occurs over lower and middle slopes on brown to orange sands. Previously disturbed through historic clearing and heavily disturbed by grazing.	Lyn04, Lyn05, Lyn06, Lyn07, Lyn08, Lyn09, Lyn13 (releve), Lyn14 (releve), Lyn17, Lyn19, Lyn20, Lyn22 (releve), Lyn 23 (releve), Lyn25, Lyn26, Lyn27 280.43 ha	

Vegetation Types	Vegetation type description and landform	Representative quadrat/releve and Total area (ha)	Representative photograph
VT02 - <i>Melaleuca cardiophylla</i> shrubland to open shrubland	<p><i>Melaleuca cardiophylla</i> shrubland to open shrubland over <i>Alyogyne hakeifolia</i>, <i>Pimelea microcephala</i> subsp. <i>microcephala</i> and <i>Rhagodia preissii</i> subsp. <i>obovata</i> open shrubland over <i>Ptilotus divaricatus</i> scattered forbland. Other common species include <i>Roepera fruticulosa</i>, <i>Pimelea gilgiana</i> and <i>*Bromus diandrus</i>. Areas that contain deeper soils <i>Acacia rostellifera</i> was also recorded. Occurs on upper mid slopes on white-brown sand with limestone outcropping. Disturbances include high grazing impacts from feral pigs and other native species (kangaroo).</p>	<p>Lyn01, Lyn02, Lyn03, Lyn10, Lyn11, Lyn15 (releve), Lyn16 (releve), Lyn18</p> <p>28.30 ha</p>	

Vegetation Types	Vegetation type description and landform	Representative quadrat/releve and Total area (ha)	Representative photograph
VT03 – <i>Myoporum insulare</i> shrubland	<i>Myoporum insulare</i> shrubland over <i>Frankenia pauciflora</i> and <i>Threlkeldia diffusa</i> open chenopod shrubland over <i>Sporobolus virginicus</i> open grassland. Occurs on light brown clay on seasonally wet brackish drainage flats.	Lyn12 0.45 ha	

Vegetation Types	Vegetation type description and landform	Representative quadrat/releve and Total area (ha)	Representative photograph
Rehabilitation areas	Rehabilitation areas consisting of <i>Acacia rostellifera</i> , <i>Alyogyne hakeifolia</i> , <i>Pimelea microcephala</i> subsp. <i>microcephala</i> , <i>Stylobasium spathulatum</i> and <i>Olearia</i> sp. Kennedy Range (G. Byrne 66) on lower and middle slopes on brown to orange sands. The undertsorey is dominated by introduced grasses including <i>Avena barbata</i> and <i>Ehrharta calycina</i> .	Lyn21, Lyn24 (releve) 34.52 ha	

Vegetation Types	Vegetation type description and landform	Representative quadrat/releve and Total area (ha)	Representative photograph
Cleared areas (including mine areas, tracks, cleared areas containing no native species)	N/A	N/A 69.16 ha	




Plate 2 MDS showing broad clusters of quadrats for VT01 and VT02

4.1.2 Conservation significant ecological communities

Based on the results of the desktop searches, dominant species, landform features, field observations, and coupled with the statistical analyses no vegetation communities identified in the survey area were consistent with any TECs or PECs.

4.1.3 Other significant vegetation

No other vegetation considered significant as per EPA (2016a) was recorded from the survey area.

4.1.4 Vegetation condition

The condition of the vegetation within the survey area ranged from Good to Completely Degraded. The extents of the vegetation condition within the survey area are detailed in Table 5 and mapped in Figure 4, Appendix A.

Areas mapped as Good had vegetation that was largely intact with native species present across each structural layer, however, also had high weed cover and signs of high grazing impacts from pigs and kangaroos. Much of the survey area had undergone historical clearing (exploration) and rehabilitation of some of these areas. Newly cleared or disturbed areas (roads and tracks) were also prevalent, as well as completely cleared areas, where the mine development has progressed. Rehabilitated areas contained fragmented vegetation resembling VT01.

Table 5 Extent of vegetation condition ratings mapped in the survey area

Vegetation Condition	Extent in the survey area (ha) (%)
Cleared	69.83 (16.9)
Completely Degraded	34.34 (8.3)
Degraded	105.15 (25.4)
Good	203.54 (49.3)

4.1.5 Flora diversity

Sixty-four flora taxa (including subspecies and varieties) representing 26 families and 50 genera were recorded from the survey area during the field survey. This total comprised 49 native taxa and 15 introduced flora taxa.

Dominant families recorded from the survey area included:

- Poaceae (10 taxa)
- Chenopodiaceae (6 taxa)

Based on described quadrats, species diversity ranged from 9 to 21 (average 14) taxa per 100 m².

The full list of flora identified within the survey area compiled by quadrat and species inventory by family is provided in Appendix D.

4.1.6 Introduced flora

Fifteen introduced flora species were recorded from the survey area. No introduced flora species recorded are listed as Declared Pests under the BAM Act or WoNS. All introduced flora species recorded are considered environmental weeds and all have been previously recorded on the Geraldton Sandplains bioregion.

4.1.7 Conservation significant flora

No EPBC Act or BC Act listed flora were recorded within the survey area. No Priority flora, as listed by the DBCA, were recorded within the survey area.

Likelihood of occurrence assessment

A likelihood of occurrence assessment was conducted post-field survey for all conservation significant flora taxa identified in the desktop assessment (Appendix D). This assessment took into account previous records, habitat requirements, efficacy of the survey, intensity of the survey, flowering times and any uniquely defining characteristics or interactions of species. Due to the size of the survey area and some seasonal limitation some of the potentially occurring species may be present, but not observed during the survey.

The likelihood of occurrence assessment post-field survey concluded that three species are considered possible to occur, 40 species highly unlikely and five species unlikely to occur in the survey area. The species considered possible to occur are:

- The Threatened *Caladenia bryceana* subsp. *cracens* is considered possible to occur in VT02 *Melaleuca cardiophylla* shrubland to open shrubland
- *Anthocercis intricata* (P3)
- *Balladonia aervoides* (P3).

4.1.8 Other significant flora

No other significant flora, such as significant range extensions, were recorded from the survey area.

4.2 Fauna

4.2.1 Fauna habitat

Five broad habitat types (including rehabilitated and cleared areas) were recorded during the survey (Figure 5, Appendix A). The five habitat types described in Table 6 closely align with vegetation types described in section 4.1.1. The five broad habitat types include:


- Acacia woodlands
- Melaleuca shrublands on limestone
- Shrublands on seasonally wet brackish drainage flats
- Rehabilitation areas
- Cleared areas.



4.2.2 Habitat corridors and linkages


The survey area comprises remnant vegetation much of which has been previously disturbed, cleared land for mine exploration and activity, and cleared agricultural land. Much of the land outside of the survey area is cleared for paddocks, or developed for the GMA processing area and existing roads. There is little contiguous vegetation remaining in the survey area and wider study area.


Within the survey area, regrowth vegetation and rehabilitated patches surround largely cleared areas, or are adjacent drive tracks. There is significant evidence of foraging within the survey area by feral pigs, European rabbits and sheep. Evidence of feral cat, red fox and domestic dog were also recorded. The patchy vegetation and drive tracks and roads may increase the likelihood of use by feral species, due to the access between habitats for these fauna.

Table 6 Fauna habitat types identified within the survey area

Fauna habitat type	Extent within survey area	Representative photo
<p>Acacia woodlands</p> <p>This habitat type was recorded over the majority of the survey area and associated with lower and middle slopes on brown to orange sands. The vegetation type comprises <i>Acacia rostellifera</i> over chenopod shrubs (<i>Rhagodia preissii</i> subsp. <i>obovata</i>) and other mixed low shrubs, native and introduced grasses. The habitat contains a high level of wood and branches through previously cleared Acacia trees providing suitable habitat for reptiles and birds. There is evidence of high grazing impacts, including from feral pigs within this habitat type.</p> <p>Conservation significant fauna</p> <p>A nesting record of the Eastern Osprey (<i>Pandion cristatus</i>) was recorded within this habitat, with the species utilising the nearby coastline and saline system of the Hutt Lagoon for foraging. The habitat type is considered very rarely used by other conservation significant species, such as the Peregrine Falcon and Fork-tailed Swift.</p>	280.43 ha	

Fauna habitat type	Extent within survey area	Representative photo
<p>Melaleuca shrublands on limestone</p> <p>This habitat type was restricted to the shallow limestone upper mid slopes on white-brown sand with limestone outcropping on the eastern side of the survey area. This habitat type is dominated by <i>Melaleuca cardiophylla</i> on shallow limestone and in areas of deeper soils scattered <i>Acacia rostellifera</i> was present. The environment had areas of good ground cover, litter and debris. Some areas of outcropping with exfoliating rock and crevices was present and would provide excellent cover for a range of fauna species. There is evidence of high grazing impacts, including from feral pigs within this habitat type.</p> <p>Conservation significant fauna</p> <p>No conservation significant fauna were recorded within this habitat type. The habitat type is considered very rarely to be used by conservation significant species, such as the Peregrine Falcon and Fork-tailed Swift.</p>	28.30 ha	
<p>Shrubland on seasonally wet brackish drainage flats</p> <p>This habitat type occurred over a small area on the western boundary of the survey area. This habitat type was dominated by <i>Myoporum insulare</i>, <i>Frankenia pauciflora</i> and <i>Threlkeldia diffusa</i> shrubs with native marine couch grass. Occurs on light brown clay on seasonally wet brackish drainage flats. The dense vegetation provides ideal habitat for reptiles and birds.</p> <p>Conservation significant fauna</p> <p>No conservation significant fauna were recorded within this habitat type. The habitat type is considered to be marginal habitat for migratory bird species given the small size of the area and the major inundation occurring outside of the migratory bird species summer patterns. The habitat would also be very rarely to be used by conservation significant species, such as the Peregrine Falcon and Fork-tailed Swift.</p>	0.45 ha	

Fauna habitat type	Extent within survey area	Representative photo
<p>Rehabilitation areas</p> <p>Rehabilitation areas consisting of mixed trees and shrubs of <i>Acacia rostellifera</i>, <i>Alyogyne hakeifolia</i>, <i>Pimelea microcephala</i> subsp. <i>microcephala</i>, <i>Stylobasium spathulatum</i> and <i>Olearia</i> sp. Kennedy Range (G. Byrne 66) on lower and middle slopes on brown to orange sands. The undertsorey is dominated by introduced grasses including <i>Avena barbata</i> and <i>Ehrharta calycina</i>. The habitat contains moderate level of wood and branches with more open areas. There is evidence of high grazing impacts, including from feral pigs within this habitat type.</p> <p>Conservation significant fauna</p> <p>No conservation significant fauna were recorded within this habitat type. The habitat would also be very rarely to be used by conservation significant species, such as the Peregrine Falcon and Fork-tailed Swift.</p>	34.52 ha	

Fauna habitat type	Extent within survey area	Representative photo
<p>Cleared areas</p> <p>The cleared and degraded habitat type was associated with previously cleared areas, access tracks and firebreaks. Much of the cleared degraded areas comprised of introduced grasses .</p> <p>Conservation significant fauna</p> <p>No conservation significant fauna were recorded within this habitat type. The habitat would also be very rarely to be used by conservation significant species, such as the Peregrine Falcon and Fork-tailed Swift.</p>	69.83 ha	

4.2.3 Fauna diversity

During the field survey 31 fauna species were recorded within the survey area, including 21 bird, 8 mammal and 2 reptile species. Of these, 24 are native and seven are introduced/feral (BAM Act) including:

- *Canis lupis* (Domestic Dog)
- *Felis catus* (Feral Cat)
- *Capra hircus* (Feral Goat)
- *Ovis aries* (Sheep)
- *Oryctolagus cuniculus* (European Rabbit)
- *Sus scrofa* (Wild Boar)
- *Streptopelia senegalensis* (Laughing Turtle-dove).

A full list of fauna recorded during the survey is provided in Appendix E.

4.2.4 Conservation significant fauna

No Threatened fauna listed under the EPBC Act and/or BC Act or Priority fauna species listed by the DBCA was recorded during the survey.

The Eastern Osprey (*Pandion cristatus*), listed as Migratory and Marine under the EPBC Act was recorded during the survey. A pair was recorded feeding a chick in a nest on the south-western portion of the survey area. The location of the Osprey nest is shown on Figure 5 with the coordinates presented in Table 7. Ospreys are generally found on or near the coast but also range inland along large rivers, mainly in northern Australia.

Table 7 Conservation listed fauna Osprey location coordinates

Species	EPBC Act	BC Act/ DBCA	Coordinates	
			Eastings	Northings
Eastern Osprey (<i>Pandion cristatus</i>)	MI	IA	230871.59	6882759.92

Likelihood of occurrence assessment

A likelihood of occurrence assessment was conducted for all conservation significant fauna species identified in the desktop assessment. This assessment was based on species biology, habitat requirements, the likely quality and availability of suitable habitat (based on vegetation associations present within the survey area) and records of the species in the vicinity of the survey area. The assessment is provided in Appendix E.

Of the 35 conservation significant fauna identified in the desktop searches one species has been identified as present (Osprey), two are considered likely to occur and the remaining species are considered unlikely or highly unlikely to occur within the survey area. The fauna species identified as present and likely to occur within the survey area are listed in Table 8.

Table 8 Conservation significant fauna present or likely to occur within the survey area

Species	EPBC Act	BC Act/ DBCA	Likelihood of occurrence
Osprey (<i>Pandion cristatus</i>)	MI	IA	Known – The survey area is situated near the coastline. This species was observed nesting within the survey area.
Fork-tailed Swift (<i>Apus pacificus</i>)	MI	IA	Likely – There are a number of records along the coast at Port Gregory and near Hutt Lagoon. This is a widespread species of coastal and sub-coastal areas. Fork-tailed Swifts are almost exclusively aerial and is likely to only utilise the survey area opportunistically.
Peregrine Falcon (<i>Falco peregrinus</i>)		OS	Likely – There are records of this species occurring around the Port Gregory area. This species may occur as an infrequent visitor, foraging within the survey area, however lacks suitable breeding habitat. Therefore likely to occur at least on an occasional basis for foraging.

No species of conservation significance are likely to be solely dependent on the habitats present within the survey area.

5. Recommendations

5.1 Recommendations

The following recommendations are provided to manage and minimise impacts to native vegetation and fauna:

- Minimise native vegetation clearing as much as practical
- Undertake further targeted flora surveys, in particular for *Caladenia bryceana* subsp. *cracens* (Threatened) as it is considered possible to occur in VT02 *Melaleuca cardiophylla* shrubland to open shrubland
- Revegetation should be undertaken by using local “provenance” native seed and / or seedlings
- Implement weed management during project activities to avoid spread of weeds
- Machinery to be maintained and cleaned to reduce the spread of weeds throughout the survey areas
- Restrict movement of machines and other vehicles to the limits of the areas cleared
- If any native fauna is disturbed during clearing it should be allowed to make its own way to adjacent vegetated areas
- Avoid disturbing habitat surrounding Osprey nest site, in particular during the nesting season (autumn – spring, April to February, and later in the season in the southern portion of Australia) (DotEE 2020; Morcombe 2011). An approximate buffer of 100 m around the nest site would be suitable, due to the discreet nature of breeding Osprey. The nest site is currently providing security through vegetation cover.
- Any injured wildlife as a result of vegetation clearing should be taken to a designated veterinary clinic, a DBCA nominated wildlife carer or suitable euthanasia by an appropriately experienced person.

6. References

- Beard JS, 1975, Vegetation Survey of Western Australia: Geraldton Sheet 1:250000, Vegmap Publications, Perth, Western Australia.
- Beard JS, & Burns AC 1976, Vegetation Survey of Western Australia: The Vegetation of the Geraldton Area Map and Explanatory Memoir 1:250,000 Series, Vegmap Publications, Perth, Western Australia
- Bureau of Meteorology (BoM) 2020, *Climate Data Online*, retrieved January 2020, from <http://www.bom.gov.au/climate/data/?ref=fr>
- Christidis, L and Boles, WE 2008, *Systematics and Taxonomy of Australian Birds*, Melbourne, CSIRO Publishing.
- Clarke, KR and Gorley, RN 2006, *PRIMER v6: User Manual/Tutorial*, Plymouth, PRIMER-E
- Department of Biodiversity, Conservation and Attractions (DBCA) 2019, *NatureMap: Mapping Western Australia's biodiversity*, Department of Parks and Wildlife, retrieved November 2019, from <http://NatureMap.dpaw.wa.gov.au/default.aspx>.
- Department of Biodiversity, Conservation and Attractions (DBCA) 2019a, *Priority Ecological Communities for Western Australia Version 28*, Species and Communities Program, DBCA, 17th January 2019.
- Department of Environment and Conservation (DEC) 2009, *Resource Condition Report for a Significant Western Australian Wetland: Hutt Lagoon*, Department of Environment and Conservation, Perth, Western Australia.
- Department of Environment and Conservation (DEC) 2012a, *Chuditch (Dasyurus geoffroii) National Recovery Plan*, Department of Environment and Conservation, Perth, WA.
- Department of Environment and Conservation (DEC) 2012b, *Western Spiny-tailed Skink (Egernia stokesii) National Recover Plan*, Department of Environment and Conservation, Perth, WA.
- Department of the Environment and Energy (DotEE) 2019, *Environmental Protection and Biodiversity Conservation Act 1999 Protected Matters Search Tool Results*, retrieved November 2019, from <http://www.environment.gov.au/epbc/pmst/index.html>.
- Department of the Environment and Energy (DotEE) 2020, *Environment Protection and Biodiversity Act 1999 List of Threatened Species Database*, retrieved January 2020, from <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>.
- Department of Sustainability, Environment, Water, Population and Communities (DSEWPaC) 2012, *Environment Protection and Biodiversity Act 1999 referral guidelines for three threatened black cockatoo species: Carnaby's Black Cockatoo (endangered) Calyptorhynchus latirostris, Baudin's Black Cockatoo (vulnerable) Calyptorhynchus baudinii and Forest Red-tailed Black Cockatoo (vulnerable) Calyptorhynchus banksia naso*, Australian Government Canberra.
- Desmond, A and Chant, A 2002, *Geraldton Sandplains 2 (GS2 – Geraldton Hills subregion)*, retrieved January 2020, from <https://www.dpaw.wa.gov.au/about-us/science-and-research/ecoinformatics-research/117-a-biodiversity-audit-of-wa>.
- EPA 2016a, *Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment*, Perth, Environmental Protection Authority.
- EPA 2016b, *Technical Guidance – Sampling methods for terrestrial vertebrate fauna*, Perth, Environmental Protection Authority.

EPA 2016c, *Technical Guidance, Terrestrial Fauna Surveys*, Perth, Environmental Protection Authority.

GHD 2013, *GMA Port Gregory Mine Vegetation, Flora and Fauna Assessment*. October 2014.

GHD 2014, *GMA Garnet Pty Ltd, Report for Port Gregory Mine Targeted Flora Survey*. October 2014.

GHD 2016, *GMA Garnet Pty Ltd, Mining Lease M70/926 Biological Survey*. October 2016.

GHD 2019, *GMA Garnet Pty Ltd, Port Gregory Mine Mining Tenement M70/1380*. November 2019.

Government of Western Australia (GoWA) 2020, *2019 Statewide Vegetation Statistics incorporating the CAR Reserve Analysis (Full report)*, Current as of March 2019, Perth, Australia, Department of Biodiversity, Conservation and Attractions, retrieved January 2020, from <https://data.wa.gov.au/>.

Higgins P.J. and Davies S.J.J.F. (Eds) 1996, *Handbook of Australian, New Zealand and Antarctic Birds. Volume 3: Snipe to Pigeons*. Oxford University Press, Melbourne

Hoffman, N and Brown, A 1998, *Orchids of South-west Australia*. Revised 2nd edition with supplement. University of Western Australia Press, Nedlands.

Johnstone, R. E., Storr, G. M. 2004, '*Handbook of Western Australian Birds. Volume 1. Nonpasserines (Emu to Dollarbird)*.' (Western Australia Museum: Perth.

Marchant, S. and P.J. Higgins, eds. 1993, *Handbook of Australian, New Zealand and Antarctic Birds. Volume 2 - Raptors to Lapwings*, Melbourne, Victoria: Oxford University Press

Morcombe, M 2011, *Field Guide to Australian Birds*, Queensland, Australia, Steve Parish Publishing Archer Field.

Nevill, S 2013, *Birds of Western Australia*, Perth, Australia, Simon Nevill Publications

NVIS Technical Working Group 2017, *Australian Vegetation Attribute Manual: National Vegetation Information System, Version 7.0*, Department of the Environment and Energy, Canberra.

Pizzey, G. and Knight, F. 2012, *The field guide to the birds of Australia. 9th Edition*, Sydney Australia, Harper Collins Publishing.

Prendergast, H.D.V., A. Brooks & I.M. Taylor (1985). Summer wader counts on Tryon Island, Capricorn Group. *Sunbird*. 15:80-83

Rogers, L G, and National Landcare Program (Australia) 1996, *Geraldton Region Land Resources Survey*, Department of Agriculture and Food, Western Australia, Perth, Report 13.

Shepherd, DP, Beeston, GR, and Hopkins, AJM 2002, *Native Vegetation in Western Australia – Extent, Type and Status*, Resource Management Technical Report 249, Perth, Department of Agriculture, Western Australia.


Van Dyke S and Strahan R 2008, *The Mammals of Australia, Third Edition*, New Holland Publishing, Sydney Australia

Western Australian (WA) Herbarium 2020, *FloraBase—the Western Australian Flora*, Department of Parks and Wildlife, from <http://florabase.dpaw.wa.gov.au/>.


Appendices

Appendix A – Figures

- Figure 1 Survey area**
- Figure 2 Environmental constraints**
- Figure 3 Vegetation types**
- Figure 4 Vegetation condition**
- Figure 5 Fauna habitats**



Map Projection: Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 50





GMA Garnet Pty Ltd
Lynton Mine Expansion Biological Survey


Survey Location and Survey Effort

Project No. 61-37808-05
Revision No. 0
Date 06 Feb 2020

FIGURE 1



Map Projection: Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 50




GMA Garnet Pty Ltd
Lynton Mine Expansion Biological Survey

Environmental Constraints

Project No. 61-38125-05
Revision No. 0
Date 06 Feb 2020

FIGURE 2





Legend

— Major road
— Minor road
— Track

Survey area


Vegetation Condition

Cleared


Good

Degraded

Completely Degraded



Map Projection: Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 50



GMA Garnet Pty Ltd
Lynton Mine Expansion Biological Survey

Vegetation Condition


Project No. 61-38125-05
Revision No. 0
Date 07 Feb 2020

FIGURE 4



Paper Size ISO A3
0 250 500 750 1,000
Metres

Map Projection: Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 50



GMA Garnet Pty Ltd
Lynton Mine Expansion Biological Survey

Fauna Habitats

Project No. 61-38125-05
Revision No. 0
Date 07 Feb 2020

FIGURE 5

Appendix B – Relevant legislation, background information and conservation code

Relevant legislation

Federal *Environment Protection and Biodiversity Conservation Act 1999*

The *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) is the Federal Government's central piece of environmental legislation. It provides a legal framework to protect and manage nationally and internationally important flora, fauna, ecological communities and heritage places, which are defined in the EPBC Act as Matters of National Environmental Significance (MNES).

The biological aspects listed as MNES include:

- Nationally threatened flora and fauna species and ecological communities
- Migratory species

A person must not undertake an action that has, will have, or is likely to have a significant impact (direct or indirect) on MNES, without approval from the Federal Minister for the Environment.

The EPBC Act is administered by the Department of the Environment and Energy (DEE).

State *Environmental Protection Act 1986*

The *Environmental Protection Act 1986* (EP Act) is the primary legislative Act dealing with the protection of the environment in Western Australia. The Act allows the Environmental Protection Authority (EPA), to prevent, control and abate pollution and environmental harm, for the conservation, preservation, protection, enhancement and management of the environment and for matters incidental to or connected with the foregoing. Part IV of the EP Act is administered by the EPA and makes provisions for the EPA to undertake environmental impact assessment of significant proposals, strategic proposals and land use planning schemes.

The Department of Water and Environment Regulation (DWER) is responsible for administering the clearing provisions of the EP Act (Part V). Clearing of native vegetation in Western Australia requires a permit from the DWER, unless exemptions apply. Applications for clearing permits are assessed by the Department and decisions are made to grant or refuse the application in accordance with the Act. When making a decision the assessment considers clearing against the ten clearing principles as specified in Schedule 5 of the EP Act:

- a) Native vegetation should not be cleared if it comprises a high level of biodiversity.
- b) Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of a significance habitat for fauna indigenous to Western Australia.
- c) Native vegetation should not be cleared if it includes, or is necessary, for the continued existence of rare flora.
- d) Native vegetation should not be cleared if it comprises the whole or part of native vegetation in an area that has been extensively cleared.
- e) Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.
- f) Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.
- g) Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.
- h) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.

- i) Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.
- j) Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence of flooding.

Exemptions for clearing include clearing that is a requirement of a written law or authorised under certain statutory processes (listed in Schedule 6 of the EP Act) and exemptions for prescribed low impact day-to-day activities (prescribed in the Environmental Protection (Clearing of Native Vegetation) Regulations 2004); these exemptions do not apply in environmentally sensitive areas (ESAs).

State Biodiversity and Conservation Act 2016

The *Biodiversity Conservation Act 2016* (BC Act) provides for the conservation and protection of biodiversity and biodiversity components, as well as the promotion of the ecologically sustainable use of biodiversity components in Western Australia. The BC Act replaces both the repealed *Wildlife Conservation Act 1950* (WC Act) and the *Sandalwood Act 1929* (Sandalwood Act), as well as their associated regulations. To attain the objectives of the BC Act, principles of ecological sustainable development have been established:

- Decision-making processes should effectively integrate both long-term and short-term economic, environmental, social and equitable considerations
- If there are threats of serious or irreversible environmental damage, lack of full scientific certainty should not be used as a reason for postponing measures to prevent environmental degradation
- The present generation should ensure that the health, diversity and productivity of the environment is maintained or enhanced for the benefit of future generations
- The conservation of biodiversity and ecological integrity should be a fundamental consideration in decision-making
- Improved valuation, pricing and incentive mechanisms should be promoted.

The BC Act is administered by the Department of Biodiversity Conservation and Attractions (DBCA).

State Biosecurity and Agriculture Management Act 2007

The *Biosecurity and Agriculture Management Act 2007* (BAM Act) and associated regulations are administered by the Department of Primary Industries and Regional Development (DPIRD) and replace the repealed *Agriculture and Related Resources Protection Act 1976*. The main purposes of the BAM Act and its regulations are to:

- Prevent new animal and plant pests (vermin and weeds) and diseases from entering WA
- Manage the impact and spread of those pests already present in the state
- Safely manage the use of agricultural and veterinary chemicals
- Increased control over the sale of agricultural products that contain violative chemical residues.

The Western Australian Organism List (WAOL) provides the status of organisms which have been categorised under the BAM Act. A Declared Pest is a prohibited organism or an organism for which a declaration under Section 22(2) of the Act is in force. Declared Pests may be assigned a control category including: C1 (exclusion), C2 (eradication) and C3 (management). The category may apply to the whole of the State, LGAs, districts, individual properties or even paddocks, and all landholders are obliged to comply with the specific category of control. Categories of control are defined below.

DPIRD Categories for Declared Pests under the BAM Act

Control class code	Description
C1 (Exclusion)	Pests will be assigned to this category if they are not established in Western Australia and control measures are to be taken, including border checks, in order to prevent them entering and establishing in the State.
C2 (Eradication)	Pests will be assigned to this category if they are present in Western Australia in low enough numbers or in sufficiently limited areas that their eradication is still a possibility.
C3 (Management)	Pests will be assigned to this category if they are established in Western Australia but it is feasible, or desirable, to manage them in order to limit their damage. Control measures can prevent a C3 pest from increasing in population size or density or moving from an area in which it is established into an area which currently is free of that pest.

Background information

Environmentally Sensitive Areas

Environmentally Sensitive Areas (ESAs) are declared by the Minister for Environment under Section 51B of the EP Act. The Table below outlines the aspects of areas declared as ESA in the Environmental Protection (Environmentally Sensitive Areas) Notice 2005.

Aspects of ESAs

Aspects of Environmentally Sensitive Areas
A declared World Heritage property as defined in Section 13 of the EPBC Act.
An area that is included on the Register of the National Estate (RNE), because of its natural values, under the <i>Australian Heritage Commission Act 1975</i> of the Commonwealth (the RNE was closed in 2007 and is no longer a statutory list – all references to the RNE were removed from the EPBC Act on 19 February 2012).
A defined wetland and the area within 50 m of the wetland. Defined wetlands include Ramsar wetlands, conservation category wetlands and nationally important wetlands.
The area covered by vegetation within 50 m of rare flora, to the extent to which the vegetation is continuous with the vegetation in which the rare flora is located.
The area covered by a Threatened Ecological Community.
A Bush Forever Site listed in “Bush Forever” Volumes 1 and 2 (2000), published by the Western Australia Planning Commission, except to the extent to which the site is approved to be developed by the Western Australia Planning Commission.
The areas covered by the <i>Environmental Protection (Gnangara Mound Crown Land) Policy 1992</i> .
The areas covered by the <i>Environmental Protection (Western Swamp Tortoise Habitat) Policy 2002</i> .
The areas covered by the lakes to which the <i>Environmental Protection (Swan Coastal Plain Lakes) Policy 1992</i> (EPP Lakes) applies.
Protected wetlands as defined in the <i>Environmental Protection (South West Agricultural Zone Wetlands) Policy 1998</i> .

Reserves and conservation areas

Department of Biodiversity, Conservation and Attractions managed lands and waters

DBCA manages lands and waters throughout Western Australia to conserve ecosystems and species, and to provide for recreation and appreciation of the natural environment. DBCA managed lands and waters include national parks, conservation parks and reserves, marine parks and reserves, regional parks, nature reserves, State forest and timber reserves. DBCA managed conservation estate, is vested with the Conservation Commission of Western Australia. Access to, or through, some areas of DBCA managed lands may require a permit or could be restricted due to management activities. Proposed land use changes and development proposals that about DBCA managed lands will generally be referred to DBCA throughout the assessment process.

Wetlands

Wetlands include not only lakes with open water, but areas of seasonally, intermittently or permanently waterlogged soil.

Ramsar Listed Wetlands

The Convention of Wetlands of International Importance was signed in 1971 at the Iranian town of Ramsar. The Convention has since been referred to as the Ramsar Convention. Ramsar Listed wetlands are “sites containing representative, rare or unique wetlands, or wetlands that are important for conserving biological diversity ... because of their ecological, botanical, zoological, limnological or hydrological importance” (DEE 2019b). Once a Ramsar Listed Wetland is designated, the country agrees to manage its conservation and ensure its wise use. Under the Convention, wise use is broadly defined as “maintaining the ecological character of a wetland” (DEE 2019b).

Nationally important wetlands

Wetlands of national significance are listed under the Directory of Important Wetlands in Australia. Nationally important wetlands are wetlands which meet at least one of the following criteria (DEE 2019a):

- It is a good example of a wetland type occurring within a biogeographic region in Australia
- It is a wetland which plays an important ecological or hydrological role in the natural functioning of a major wetland system/complex
- It is a wetland which is important as the habitat for animal taxa at a vulnerable stage in their life cycles, or provides a refuge when adverse conditions such as drought prevail
- The wetland supports one percent or more of the national populations of any native plant or animal taxa
- The wetland supports native plant or animal taxa or communities which are considered endangered or vulnerable at the national level
- The wetland is of outstanding historical or cultural significance

Vegetation extent and status

The National Objectives and Targets for Biodiversity Conservation 2001–2005 (Commonwealth of Australia 2001) recognise that the retention of 30 percent or more of the pre-clearing extent of each ecological community is necessary if Australia’s biological diversity is to be protected. This is the threshold level below which species loss appears to accelerate exponentially and loss below this level should not be permitted. This level of recognition is in keeping with the targets recommended in the review of the National Strategy for the Conservation of Australia’s Biological Diversity (ANZECC 2000).

The extent of remnant native vegetation in WA has been assessed by Shepherd et al. (2002) and the GoWA (2018), based on broadscale vegetation association mapping by Beard (various publications). The GoWA produces Statewide Vegetation Statistics Reports that are used for a number of purposes including conservation planning, land use planning and when assessing development applications. The reports are updated at least every two years.

Vegetation condition

The vegetation condition can be assessed in accordance with the vegetation condition rating scale for the South West and Interzone Botanical Provinces (EPA 2016a). The scale recognises the intactness of vegetation and consists of six rating levels as outlined below.

Vegetation condition rating scale for the South West and Interzone Botanical Provinces

Condition	South West and Interzone Botanical Provinces description
Pristine	Pristine or nearly so, no obvious signs of damage caused by human activities since European settlement.
Excellent	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species. Damage to trees caused by fire, the presence of non-aggressive weeds and occasional vehicle tracks.
Very Good	Vegetation structure altered, obvious signs of disturbance. Disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.
Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.
Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds at high density, partial clearing, dieback and grazing.
Completely Degraded	The structure of vegetation is no longer intact and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees or shrubs.

Conservation codes

Species of significant flora, fauna and communities are protected under both Federal and State Acts. The Federal EPBC Act provides a legal framework to protect and manage nationally important flora and communities. The State BC Act is the primary wildlife conservation legislation in Western Australia. Information on the conservation codes is summarised in the following sections.

Ecological communities

Conservation significant communities

Ecological communities are defined as naturally occurring biological assemblages that occur in a particular type of habitat (English and Blyth 1997). Federally listed Threatened Ecological Communities (TECs) are protected under the EPBC Act. The BC Act provides for the Minister to list an ecological community as a TEC (section 27), or as a collapsed ecological community (section 31) statutory listing of State TECs by the Minister. The legislation also describes statutory processes for preparing recovery plans for TECs, the registration of their critical habitat, and penalties for unauthorised modification of TECs.

Possible TECs that do not meet survey criteria are added to the DBCA Priority Ecological Community (PEC) List under Priorities 1, 2 and 3. These are ecological communities that are adequately known; are rare but not threatened, or meet criteria for Near Threatened. PECs that have been recently removed from the threatened list are placed in Priority 4. These ecological communities require regular monitoring. Conservation dependent ecological communities are placed in Priority 5. PECs are not listed under any formal Federal or State legislation, however, may be listed as TECs under the EPBC Act.

Conservation codes and definitions for TECs listed under the EPBC Act and/ or BC Act

Categories	Definition
Federal Government Conservation Categories (EPBC Act)	
Critically Endangered (CR)	An ecological community if, at that time, is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria (as outlined in Environment Protection and Biodiversity Conservation Regulations 2000)
Endangered (EN)	An ecological community if, at that time: A) is not critically endangered; and B) is facing a very high risk of extinction in the wild in the near future, as determined in accordance with the prescribed criteria (as outlined in Environment Protection and Biodiversity Conservation Regulations 2000)
Vulnerable (VU)	An ecological community if, at that time: A) is not critically endangered or endangered; and B) is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria (as outlined in Environment Protection and Biodiversity Conservation Regulations 2000)
Western Australia Conservation Categories (BC Act)	
<u>Threatened Ecological Communities</u>	

Categories	Definition
Critically Endangered (CR)	An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or that was originally of limited distribution and is facing severe modification or destruction throughout its range in the immediate future, or is already severely degraded throughout its range but capable of being substantially restored or rehabilitated.
Endangered (EN)	An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or was originally of limited distribution and is in danger of significant modification throughout its range or severe modification or destruction over most of its range in the near future.
Vulnerable (VU)	An ecological community that has been adequately surveyed and is found to be declining and/or has declined in distribution and/or condition and whose ultimate security has not yet been assured and/or a community that is still widespread but is believed likely to move into a category of higher threat in the near future if threatening processes continue or begin operating throughout its range.

Collapsed ecological communities

An ecological community is eligible for listing as a collapsed ecological community at a particular time if, at that time –

- (a) there is no reasonable doubt that the last occurrence of the ecological community has collapsed); or
- (b) the ecological community has been so extensively modified throughout its range that no occurrence of it is likely to recover –
 - (i) its species composition or structure; or
 - (ii) its species composition and structure.

Section 33 of the BC Act provides for a collapsed ecological community to be regarded as a threatened ecological community if it is discovered in a state that no longer makes it eligible for listing as a collapsed ecological community.

Conservation categories and definitions for PECS as listed by the DBCA

Category	Description
Priority 1	<p>Poorly known ecological communities.</p> <p>Ecological communities that are known from very few occurrences with a very restricted distribution (generally ≤ 5 occurrences or a total area of ≤ 100 ha). Occurrences are believed to be under threat either due to limited extent, or being on lands under immediate threat (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) or for which current threats exist. May include communities with occurrences on protected lands. Communities may be included if they are comparatively well-known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.</p>
Priority 2	<p>Poorly known ecological communities.</p> <p>Communities that are known from few occurrences with a restricted distribution (generally ≤ 10 occurrences or a total area of ≤ 200 ha). At least some occurrences are not believed to be under immediate threat of destruction or degradation. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.</p>

Category	Description
Priority 3	<p>Poorly known ecological communities.</p> <p>(i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or:</p> <p>(ii) communities known from a few widespread occurrences, which are either large or with significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or;</p> <p>(iii) communities made up of large, and/or widespread occurrences, that may or may not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, and inappropriate fire regimes.</p> <p>Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.</p>
Priority 4	<p>Ecological communities that are adequately known, rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.</p> <p>(i) Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands.</p> <p>(ii) Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable.</p> <p>(iii) Ecological communities that have been removed from the list of threatened communities during the past five years.</p>
Priority 5	<p>Conservation Dependent ecological communities.</p> <p>Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.</p>

Other significant vegetation

Vegetation may be significant for a range of reasons other than a statutory listing. The EPA (2016b) states that significant vegetation may include vegetation that includes the following:

- Restricted distribution
- Degree of historical impact from threatening processes
- Local endemism in restricted habitats
- Novel combinations of taxa
- A role as a refuge
- A role as a key habitat for Threatened species or large population representing a significant proportion of the local to regional total population of a species
- Being representative of a vegetation unit in 'pristine' condition in a highly cleared landscape, recently discovered range extensions, or isolated outliers of the main range)
- Being poorly reserved.

This may apply at a number of levels, so the unit may be significant when considered at the fine-scale (intra-locality), intermediate-scale (locality or inter-locality) or broad-scale (local to region).

Flora and fauna

Conservation significant flora and fauna

Species of significant flora are protected under both Federal and State legislation. Any activities that are deemed to have a significant impact on species that are recognised by the EPBC Act, and/or the BC Act can warrant referral to the DEE and/or the EPA.

The Federal conservation level of flora and fauna species and their significance status is assessed under the EPBC Act. The significance levels for flora and fauna used in the EPBC Act align with the International Union for Conservation of Nature (IUCN) Red List criteria, which are internationally recognised as providing best practice for assigning the conservation status of species. The EPBC Act also protects land and migratory species that are listed under International Agreements. The list of migratory species established under section 209 of the EPBC Act comprises:

- Migratory species which are native to Australia and are included in the appendices to the Bonn Convention (Convention on the Conservation of Migratory Species of Wild Animals Appendices I and II)
- Migratory species included in annexes established under the Japan-Australia Migratory Bird Agreement (JAMBA) and the China–Australia Migratory Bird Agreement (CAMBA)
- Native, migratory species identified in a list established under, or an instrument made under, an international agreement approved by the Minister, such as the republic of Korea–Australia Migratory Bird Agreement (ROKAMBA)

The State conservation level of flora and fauna species and their significance status also follows the IUCN Red List criteria. Under the BC Act flora and fauna can be listed as Threatened, Extinct and as Specially Protected species.

Threatened species are those species which have been adequately searched for and are deemed to be, in the wild, either rare, under identifiable threat of extinction, or otherwise in need of special protection, and have been gazetted as such. The assessment of the conservation status of Threatened species is based on their national extent and ranked according to their level of threat using IUCN Red List categories and criteria. Specially protected species meet one or more of the following categories: species of special conservation interest; migratory species; cetaceans; species subject to international agreement; or species otherwise in need of special protection. Species that are listed as Threatened or Extinct species under the BC Act cannot also be listed as Specially Protected species.

Possibly threatened species that do not meet survey criteria, or are otherwise data deficient, are added to the Priority Fauna or Priority Flora Lists under Priorities 1, 2 or 3. These three categories are ranked in order of priority for survey and evaluation of conservation status so that consideration can be given to their declaration as threatened flora or fauna.

Species that are adequately known, are rare but not threatened, or meet criteria for near threatened, or that have been recently removed from the threatened species or other specially protected fauna lists for other than taxonomic reasons, are placed in Priority 4. These species require regular monitoring.

Assessment of Priority codes is based on the Western Australian distribution of the species, unless the distribution in WA is part of a contiguous population extending into adjacent States, as defined by the known spread of locations.

For the purposes of this assessment, all species listed under the EPBC Act, BC Act and DBCA Priority species are considered conservation significant.

Conservation categories and definitions for EPBC Act and BC Act listed flora and fauna species

Conservation category	Definition
Threatened species	
Critically Endangered (CR)	<p>Threatened species considered to be “facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with criteria set out in the ministerial guidelines”.</p> <p>Listed as critically endangered under section 19(1)(a) of the BC Act in accordance with the criteria set out in section 20 and the ministerial guidelines.</p>
Endangered (EN)	<p>Threatened species considered to be “facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines”.</p> <p>Listed as endangered under section 19(1)(b) of the BC Act in accordance with the criteria set out in section 21 and the ministerial guidelines.</p>
Vulnerable (VU)	<p>Threatened species considered to be “facing a high risk of extinction in the wild in the medium term future, as determined in accordance with criteria set out in the ministerial guidelines”.</p> <p>Listed as vulnerable under section 19(1)(c) of the BC Act in accordance with the criteria set out in section 22 and the ministerial guidelines.</p>
Extinct species	
Extinct (EX)	Species where “there is no reasonable doubt that the last member of the species has died”, and listing is otherwise in accordance with the ministerial guidelines (section 24 of the BC Act).
Extinct in the Wild (EW)	Species that “is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; and it has not been recorded in its known habitat or expected habitat, at appropriate seasons, anywhere in its past range, despite surveys over a time frame appropriate to its life cycle and form”, and listing is otherwise in accordance with the ministerial guidelines (section 25 of the BC Act).
Specially protected species	
Migratory (MI)	<p>Fauna that periodically or occasionally visit Australia or an external Territory or the exclusive economic zone; or the species is subject of an international agreement that relates to the protection of migratory species and that binds the Commonwealth; and listing is otherwise in accordance with the ministerial guidelines (section 15 of the BC Act).</p> <p>Includes birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and fauna subject to the Convention on the Conservation of Migratory Species of Wild Animals (Bonn Convention), an environmental treaty under the United Nations Environment Program. Migratory species listed under the BC Act are a subset of the migratory animals, that are known to visit Western Australia, protected under the international agreements or treaties, excluding species that are listed as Threatened species.</p>

Conservation category	Definition
Species of special conservation interest (conservation dependent fauna) (CD)	Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened.
Other specially protected fauna (OS)	Fauna otherwise in need of special protection to ensure their conservation, and listing is otherwise in accordance with the ministerial guidelines (section 18 of the BC Act).

Conservation codes for DBCA listed Priority flora and fauna

Priority category	Definition
Priority 1	<p>Poorly-known taxa</p> <p>Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.</p>
Priority 2	<p>Poorly-known taxa</p> <p>Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.</p>
Priority 3	<p>Poorly-known taxa</p> <p>Species that are known from several locations, and the species does not appear to be under imminent threat, or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. Such species are in need of further survey.</p>
Priority 4	<p>Rare, Near Threatened and other taxa in need of monitoring</p> <p>A. Rare: Taxa that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These taxa are usually represented on conservation lands.</p> <p>B. Near Threatened. Taxa that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable.</p> <p>C. Taxa that have been removed from the list of threatened taxa during the past five years for reasons other than taxonomy.</p>

Other significant flora

Flora species, subspecies, varieties, hybrids and ecotypes may be significant for a range of reasons, other than a statutory listing. The EPA (2016b) states that significant flora may include taxa that have:

- A keystone role in a particular habitat for threatened or Priority flora or fauna species, or large populations representing a considerable proportion of the local or regional total population of a species
- Relictual status, being representation of taxonomic or physiognomic groups that no longer occur widely in the broader landscape
- Anomalous features that indicate a potential new discovery
- Being representative of the range of a species (particularly, at the extremes of range, recently discovered range extensions, or isolated outliers of the main range)
- The presence of restricted subspecies, varieties, or naturally occurring hybrids
- Local endemism (a restricted distribution) or association with a restricted habitat type (e.g. surface water or groundwater dependent ecosystems)
- Being poorly reserved

Other significant fauna

Fauna species may be significant for a range of reasons other than those protected by international agreement or treaty, Specially Protected or Priority Fauna. Significant fauna may include short-range endemic species, species that have declining populations or declining distributions, species at the extremes of their range, or isolated outlying populations, or species which may be undescribed (EPA 2010).

Introduced plants (weeds)

Declared Pests

Information on species considered to be Declared Pests is provided under *State Biosecurity and Agriculture Management Act 2007*.

Weeds of National Significance

The spread of weeds across a range of land uses or ecosystems is important in the context of socio-economic and environmental values. The assessment of Weeds of National Significance (WoNS) is based on four major criteria:

- Invasiveness
- Impacts
- Potential for spread
- Socio-economic and environmental values

Australian state and territory governments have identified thirty-two Weeds of National Significance (WoNS); a list of 20 WoNS was endorsed in 1999 and a further 12 were added in 2012.

References

- ANZECC 2000, *Core Environmental Indicators for Reporting on the State of Environment*, ANZECC State of the Environment Reporting Task Force.
- Commonwealth of Australia 2001, *National Targets and Objectives for Biodiversity Conservation 2001–2005*, Canberra, AGPS.
- DEE 2019a, *Criteria for determining nationally important wetlands*, retrieved 2019, from <http://www.environment.gov.au/topics/water/water-our-environment/wetlands/australian-wetlands-database/directory-important>.
- DEE 2019b, *The Ramsar Convention on Wetlands*, retrieved 2019, from <http://www.environment.gov.au/topics/water/water-our-environment/wetlands/ramsar-convention-wetlands>.
- English, V and Blyth, J 1997, *Identifying and Conserving Threatened Ecological Communities in the South West Botanical Province*, Perth, Department of Conservation and Land Management.
- EPA 2010, *Technical Guide – Terrestrial Fauna Surveys*, EPA, Perth, WA.
- EPA 2016a, *Technical Guide – Flora and Vegetation Surveys for Environmental Impact Assessment*, EPA, Perth, WA.
- EPA 2016b, *Environmental Factor Guideline - Flora and Vegetation*, EPA, Perth, WA.
- GoWA 2018, *Statewide Vegetation Statistics incorporating the CAR Reserve Analysis (Full report)*, Current as of December 2017, Perth Western Australia, Department of Environment and Conservation, from <https://www2.landgate.wa.gov.au/web/guest/downloader>.
- Shepherd, DP, Beeston, GR & Hopkins, AJM 2002, *Native Vegetation in Western Australia – Extent, Type and Status*, Resource Management Technical Report 249, Perth, Department of Agriculture.

Appendix C – Desktop searches

EPBC Act PMST (10 km)

Naturemap Flora report (10 km)

Naturemap Fauna report (10 km)



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 20/01/20 13:03:54

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)


[Acknowledgements](#)



This map may contain data which are
©Commonwealth of Australia
(Geoscience Australia), ©PSMA 2010

[Coordinates](#)

[Buffer: 10.0Km](#)



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	None
Wetlands of International Importance:	None
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	1
Listed Threatened Ecological Communities:	None
Listed Threatened Species:	48
Listed Migratory Species:	46

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	None
Commonwealth Heritage Places:	None
Listed Marine Species:	76
Whales and Other Cetaceans:	11
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Australian Marine Parks:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	2
Regional Forest Agreements:	None
Invasive Species:	15
Nationally Important Wetlands:	1
Key Ecological Features (Marine)	2

Details

Matters of National Environmental Significance

Commonwealth Marine Area

[Resource Information]

Approval is required for a proposed activity that is located within the Commonwealth Marine Area which has, will have, or is likely to have a significant impact on the environment. Approval may be required for a proposed action taken outside the Commonwealth Marine Area but which has, may have or is likely to have a significant impact on the environment in the Commonwealth Marine Area. Generally the Commonwealth Marine Area stretches from three nautical miles to two hundred nautical miles from the coast.

Name

EEZ and Territorial Sea

Marine Regions

[Resource Information]

If you are planning to undertake action in an area in or close to the Commonwealth Marine Area, and a marine bioregional plan has been prepared for the Commonwealth Marine Area in that area, the marine bioregional plan may inform your decision as to whether to refer your proposed action under the EPBC Act.

Name

[South-west](#)

Listed Threatened Species

[Resource Information]

Name	Status	Type of Presence
Birds		

Anous tenuirostris melanops Australian Lesser Noddy [26000]	Vulnerable	Species or species habitat may occur within area
--	------------	--

Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
--	------------	--

Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
---	-----------------------	---

Calyptorhynchus latirostris Carnaby's Cockatoo, Short-billed Black-Cockatoo [59523]	Endangered	Species or species habitat likely to occur within area
--	------------	--

Charadrius leschenaultii Greater Sand Plover, Large Sand Plover [877]	Vulnerable	Species or species habitat known to occur within area
--	------------	---

Diomedea amsterdamensis Amsterdam Albatross [64405]	Endangered	Species or species habitat may occur within area
--	------------	--

Diomedea epomophora Southern Royal Albatross [89221]	Vulnerable	Species or species habitat may occur within area
---	------------	--

Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
---	------------	--

Leipoa ocellata Malleefowl [934]	Vulnerable	Species or species habitat likely to occur
---	------------	--

Name	Status	Type of Presence
		within area
Limosa lapponica baueri Bar-tailed Godwit (baueri), Western Alaskan Bar-tailed Godwit [86380]	Vulnerable	Species or species habitat may occur within area
Limosa lapponica menzbieri Northern Siberian Bar-tailed Godwit, Bar-tailed Godwit (menzbieri) [86432]	Critically Endangered	Species or species habitat may occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pterodroma mollis Soft-plumaged Petrel [1036]	Vulnerable	Species or species habitat may occur within area
Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat may occur within area
Sternula nereis nereis Australian Fairy Tern [82950]	Vulnerable	Species or species habitat known to occur within area
Thalassarche carteri Indian Yellow-nosed Albatross [64464]	Vulnerable	Foraging, feeding or related behaviour may occur within area
Thalassarche cauta cauta Shy Albatross [82345]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta steadi White-capped Albatross [82344]	Vulnerable	Foraging, feeding or related behaviour likely to occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Mammals		
Balaenoptera musculus Blue Whale [36]	Endangered	Species or species habitat likely to occur within area
Dasyurus geoffroii Chuditch, Western Quoll [330]	Vulnerable	Species or species habitat likely to occur within area
Eubalaena australis Southern Right Whale [40]	Endangered	Species or species habitat likely to occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Species or species habitat known to occur within area
Neophoca cinerea Australian Sea-lion, Australian Sea Lion [22]	Vulnerable	Species or species habitat may occur within

Name	Status	Type of Presence area
Other		
Idiosoma nigrum Shield-backed Trapdoor Spider, Black Rugose Trapdoor Spider [66798]	Vulnerable	Species or species habitat may occur within area
Plants		
Androcalva bivillosa Straggling Androcalva [87807]	Critically Endangered	Species or species habitat likely to occur within area
Caladenia barbarella Small Dragon Orchid, Common Dragon Orchid [68686]	Endangered	Species or species habitat may occur within area
Caladenia bryceana subsp. cracens Northern Dwarf Spider-orchid [64556]	Vulnerable	Species or species habitat may occur within area
Caladenia elegans Elegant Spider-orchid [56775]	Endangered	Species or species habitat known to occur within area
Caladenia hoffmanii Hoffman's Spider-orchid [56719]	Endangered	Species or species habitat known to occur within area
Diuris drummondii Tall Donkey Orchid [4365]	Vulnerable	Species or species habitat known to occur within area
Drakaea concolor Kneeling Hammer-orchid [56777]	Vulnerable	Species or species habitat known to occur within area
Drummondita ericoides Morseby Range Drummondita [9193]	Endangered	Species or species habitat may occur within area
Eucalyptus cuprea Mallee Box [56773]	Endangered	Species or species habitat likely to occur within area
Hypocalymma angustifolium subsp. Hutt River (S.Patrick 2982) [85023]	Endangered	Species or species habitat known to occur within area
Pterostylis sinuata Northampton Midget Greenhood, Western Swan Grrenhood [84991]	Endangered	Species or species habitat known to occur within area
Stachystemon nematophorus Three-flowered Stachystemon [81447]	Vulnerable	Species or species habitat known to occur within area
Reptiles		
Caretta caretta Loggerhead Turtle [1763]	Endangered	Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Foraging, feeding or related behaviour known to occur within area
Egernia stokesii badia Western Spiny-tailed Skink, Baudin Island Spiny-tailed Skink [64483]	Endangered	Species or species habitat may occur within area

Name	Status	Type of Presence
Natator depressus Flatback Turtle [59257]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Sharks		
Carcharias taurus (west coast population) Grey Nurse Shark (west coast population) [68752]	Vulnerable	Species or species habitat likely to occur within area
Carcharodon carcharias White Shark, Great White Shark [64470]	Vulnerable	Species or species habitat known to occur within area
Rhincodon typus Whale Shark [66680]	Vulnerable	Species or species habitat may occur within area
Listed Migratory Species		
[Resource Information]		
* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.		
Name	Threatened	Type of Presence
Migratory Marine Birds		
Anous stolidus Common Noddy [825]		Species or species habitat likely to occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardenna carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [82404]		Foraging, feeding or related behaviour likely to occur within area
Diomedea amsterdamensis Amsterdam Albatross [64405]	Endangered	Species or species habitat may occur within area
Diomedea epomophora Southern Royal Albatross [89221]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
Fregata ariel Lesser Frigatebird, Least Frigatebird [1012]		Species or species habitat likely to occur within area
Hydroprogne caspia Caspian Tern [808]		Foraging, feeding or related behaviour known to occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Onychoprion anaethetus Bridled Tern [82845]		Foraging, feeding or related behaviour likely to occur within area
Thalassarche carteri Indian Yellow-nosed Albatross [64464]	Vulnerable	Foraging, feeding or related behaviour may occur within area
Thalassarche cauta Shy Albatross [89224]	Vulnerable*	Species or species habitat may occur within

Name	Threatened	Type of Presence
area		
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche steadi White-capped Albatross [64462]	Vulnerable*	Foraging, feeding or related behaviour likely to occur within area
Migratory Marine Species		
Balaena glacialis australis Southern Right Whale [75529]	Endangered*	Species or species habitat likely to occur within area
Balaenoptera edeni Bryde's Whale [35]		Species or species habitat may occur within area
Balaenoptera musculus Blue Whale [36]	Endangered	Species or species habitat likely to occur within area
Carcharodon carcharias White Shark, Great White Shark [64470]	Vulnerable	Species or species habitat known to occur within area
Caretta caretta Loggerhead Turtle [1763]	Endangered	Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Foraging, feeding or related behaviour known to occur within area
Lamna nasus Porbeagle, Mackerel Shark [83288]		Species or species habitat may occur within area
Manta alfredi Reef Manta Ray, Coastal Manta Ray, Inshore Manta Ray, Prince Alfred's Ray, Resident Manta Ray [84994]		Species or species habitat known to occur within area
Manta birostris Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray, Oceanic Manta Ray [84995]		Species or species habitat may occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Species or species habitat known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Orcinus orca Killer Whale, Orca [46]		Species or species habitat may occur within area
Rhincodon typus Whale Shark [66680]	Vulnerable	Species or species habitat may occur within area
Migratory Terrestrial Species		
Motacilla cinerea Grey Wagtail [642]		Species or species

Name	Threatened	Type of Presence
Migratory Wetlands Species		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat known to occur within area
Arenaria interpres Ruddy Turnstone [872]		Species or species habitat known to occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris alba Sanderling [875]		Species or species habitat known to occur within area
Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Calidris ruficollis Red-necked Stint [860]		Species or species habitat known to occur within area
Charadrius leschenaultii Greater Sand Plover, Large Sand Plover [877]	Vulnerable	Species or species habitat known to occur within area
Limosa lapponica Bar-tailed Godwit [844]		Species or species habitat known to occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Breeding known to occur within area
Pluvialis fulva Pacific Golden Plover [25545]		Species or species habitat known to occur within area
Tringa brevipes Grey-tailed Tattler [851]		Species or species habitat known to occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat known to occur within area

Other Matters Protected by the EPBC Act

Listed Marine Species		[Resource Information]
* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.		
Name	Threatened	Type of Presence
Birds		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat known to occur within area
Anous stolidus Common Noddy [825]		Species or species habitat likely to occur within area
Anous tenuirostris melanops Australian Lesser Noddy [26000]	Vulnerable	Species or species habitat may occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba Great Egret, White Egret [59541]		Species or species habitat known to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Arenaria interpres Ruddy Turnstone [872]		Species or species habitat known to occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris alba Sanderling [875]		Species or species habitat known to occur within area
Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Calidris ruficollis Red-necked Stint [860]		Species or species habitat known to occur within area
Catharacta skua Great Skua [59472]		Species or species habitat may occur within area
Charadrius leschenaultii Greater Sand Plover, Large Sand Plover [877]	Vulnerable	Species or species habitat known to occur within area
Charadrius ruficapillus Red-capped Plover [881]		Species or species habitat known to occur

Name	Threatened	Type of Presence
Chrysococcyx osculans Black-eared Cuckoo [705]		within area Species or species habitat likely to occur within area
Diomedea amsterdamensis Amsterdam Albatross [64405]	Endangered	Species or species habitat may occur within area
Diomedea epomophora Southern Royal Albatross [89221]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
Fregata ariel Lesser Frigatebird, Least Frigatebird [1012]		Species or species habitat likely to occur within area
Haliaeetus leucogaster White-bellied Sea-Eagle [943]		Species or species habitat known to occur within area
Heteroscelus brevipes Grey-tailed Tattler [59311]		Species or species habitat known to occur within area
Himantopus himantopus Pied Stilt, Black-winged Stilt [870]		Species or species habitat known to occur within area
Larus pacificus Pacific Gull [811]		Foraging, feeding or related behaviour known to occur within area
Limosa lapponica Bar-tailed Godwit [844]		Species or species habitat known to occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Breeding known to occur within area
Pluvialis fulva Pacific Golden Plover [25545]		Species or species habitat known to occur within area
Pterodroma mollis Soft-plumaged Petrel [1036]	Vulnerable	Species or species habitat may occur within area

Name	Threatened	Type of Presence
Puffinus assimilis Little Shearwater [59363]		Foraging, feeding or related behaviour known to occur within area
Puffinus carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [1043]		Foraging, feeding or related behaviour likely to occur within area
Recurvirostra novaehollandiae Red-necked Avocet [871]		Species or species habitat known to occur within area
Rostratula benghalensis (sensu lato) Painted Snipe [889]	Endangered*	Species or species habitat may occur within area
Sterna anaethetus Bridled Tern [814]		Foraging, feeding or related behaviour likely to occur within area
Sterna caspia Caspian Tern [59467]		Foraging, feeding or related behaviour known to occur within area
Thalassarche carteri Indian Yellow-nosed Albatross [64464]	Vulnerable	Foraging, feeding or related behaviour may occur within area
Thalassarche cauta Shy Albatross [89224]	Vulnerable*	Species or species habitat may occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche steadi White-capped Albatross [64462]	Vulnerable*	Foraging, feeding or related behaviour likely to occur within area
Thinornis rubricollis Hooded Plover [59510]		Species or species habitat may occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat known to occur within area
Fish		
Acentronura australe Southern Pygmy Pipehorse [66185]		Species or species habitat may occur within area
Campichthys galei Gale's Pipefish [66191]		Species or species habitat may occur within area
Choeroichthys suillus Pig-snouted Pipefish [66198]		Species or species habitat may occur within area
Halicampus brocki Brock's Pipefish [66219]		Species or species habitat may occur within area
Hippocampus angustus Western Spiny Seahorse, Narrow-bellied Seahorse [66234]		Species or species habitat may occur within area

Name	Threatened	Type of Presence
Hippocampus breviceps Short-head Seahorse, Short-snouted Seahorse [66235]		Species or species habitat may occur within area
Hippocampus subelongatus West Australian Seahorse [66722]		Species or species habitat may occur within area
Lissocampus fatiloquus Prophet's Pipefish [66250]		Species or species habitat may occur within area
Maroubra perserrata Sawtooth Pipefish [66252]		Species or species habitat may occur within area
Mitotichthys meraculus Western Crested Pipefish [66259]		Species or species habitat may occur within area
Nannocampus subosseus Bonyhead Pipefish, Bony-headed Pipefish [66264]		Species or species habitat may occur within area
Phycodurus eques Leafy Seadragon [66267]		Species or species habitat may occur within area
Phyllopteryx taeniolatus Common Seadragon, Weedy Seadragon [66268]		Species or species habitat may occur within area
Pugnaso curtirostris Pugnose Pipefish, Pug-nosed Pipefish [66269]		Species or species habitat may occur within area
Solegnathus lettiensis Gunther's Pipehorse, Indonesian Pipefish [66273]		Species or species habitat may occur within area
Stigmatopora argus Spotted Pipefish, Gulf Pipefish, Peacock Pipefish [66276]		Species or species habitat may occur within area
Stigmatopora nigra Widebody Pipefish, Wide-bodied Pipefish, Black Pipefish [66277]		Species or species habitat may occur within area
Syngnathoides biaculeatus Double-end Pipehorse, Double-ended Pipehorse, Alligator Pipefish [66279]		Species or species habitat may occur within area
Urocampus carinirostris Hairy Pipefish [66282]		Species or species habitat may occur within area
Vanacampus margaritifer Mother-of-pearl Pipefish [66283]		Species or species habitat may occur within area
Mammals		
Arctocephalus forsteri Long-nosed Fur-seal, New Zealand Fur-seal [20]		Species or species habitat may occur within area
Neophoca cinerea Australian Sea-lion, Australian Sea Lion [22]	Vulnerable	Species or species habitat may occur within area
Reptiles		
Aipysurus pooleorum Shark Bay Seasnake [66061]		Species or species habitat may occur within

Name	Threatened	Type of Presence
Caretta caretta Loggerhead Turtle [1763]	Endangered	area Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Foraging, feeding or related behaviour known to occur within area
Disteira kingii Spectacled Seasnake [1123]		Species or species habitat may occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Pelamis platurus Yellow-bellied Seasnake [1091]		Species or species habitat may occur within area

Whales and other Cetaceans		[Resource Information]
Name	Status	Type of Presence
Mammals		
Balaenoptera acutorostrata Minke Whale [33]		Species or species habitat may occur within area
Balaenoptera edeni Bryde's Whale [35]		Species or species habitat may occur within area
Balaenoptera musculus Blue Whale [36]	Endangered	Species or species habitat likely to occur within area
Delphinus delphis Common Dophin, Short-beaked Common Dolphin [60]		Species or species habitat may occur within area
Eubalaena australis Southern Right Whale [40]	Endangered	Species or species habitat likely to occur within area
Grampus griseus Risso's Dolphin, Grampus [64]		Species or species habitat may occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Species or species habitat known to occur within area
Orcinus orca Killer Whale, Orca [46]		Species or species habitat may occur within area
Stenella attenuata Spotted Dolphin, Pantropical Spotted Dolphin [51]		Species or species habitat may occur within area
Tursiops aduncus Indian Ocean Bottlenose Dolphin, Spotted Bottlenose Dolphin [68418]		Species or species habitat likely to occur within area
Tursiops truncatus s. str. Bottlenose Dolphin [68417]		Species or species habitat may occur within area

Extra Information

State and Territory Reserves		[Resource Information]
Name		State
Port Gregory		WA
Utcha Well		WA

Invasive Species[Resource Information]

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resouces Audit, 2001.

Name	Status	Type of Presence
Birds		
Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Streptopelia senegalensis Laughing Turtle-dove, Laughing Dove [781]		Species or species habitat likely to occur within area
Mammals		
Canis lupus familiaris Domestic Dog [82654]		Species or species habitat likely to occur within area
Capra hircus Goat [2]		Species or species habitat likely to occur within area
Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
Mus musculus House Mouse [120]		Species or species habitat likely to occur within area
Oryctolagus cuniculus Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
Rattus rattus Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
Sus scrofa Pig [6]		Species or species habitat likely to occur within area

Name	Status	Type of Presence
Vulpes vulpes Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
Cenchrus ciliaris Buffel-grass, Black Buffel-grass [20213]		Species or species habitat may occur within area
Lantana camara Lantana, Common Lantana, Kamara Lantana, Large-leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White Sage, Wild Sage [10892] Lycium ferocissimum African Boxthorn, Boxthorn [19235]		Species or species habitat likely to occur within area
Prosopis spp. Mesquite, Algaroba [68407]		Species or species habitat likely to occur within area
Tamarix aphylla Athel Pine, Athel Tree, Tamarisk, Athel Tamarisk, Athel Tamarix, Desert Tamarisk, Flowering Cypress, Salt Cedar [16018]		Species or species habitat likely to occur within area

Nationally Important Wetlands		[Resource Information]
Name	State	
Hutt Lagoon System	WA	

Key Ecological Features (Marine)	[Resource Information]
Key Ecological Features are the parts of the marine ecosystem that are considered to be important for the biodiversity or ecosystem functioning and integrity of the Commonwealth Marine Area.	

Name	Region
Commonwealth marine environment within and	South-west
Western rock lobster	South-west

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-28.109489 114.219784,-28.199066 114.294036,-28.199947 114.294436,-28.199947 114.294436

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [-Office of Environment and Heritage, New South Wales](#)
- [-Department of Environment and Primary Industries, Victoria](#)
- [-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [-Department of Environment, Water and Natural Resources, South Australia](#)
- [-Department of Land and Resource Management, Northern Territory](#)
- [-Department of Environmental and Heritage Protection, Queensland](#)
- [-Department of Parks and Wildlife, Western Australia](#)
- [-Environment and Planning Directorate, ACT](#)
- [-Birdlife Australia](#)
- [-Australian Bird and Bat Banding Scheme](#)
- [-Australian National Wildlife Collection](#)
- [-Natural history museums of Australia](#)
- [-Museum Victoria](#)
- [-Australian Museum](#)
- [-South Australian Museum](#)
- [-Queensland Museum](#)
- [-Online Zoological Collections of Australian Museums](#)
- [-Queensland Herbarium](#)
- [-National Herbarium of NSW](#)
- [-Royal Botanic Gardens and National Herbarium of Victoria](#)
- [-Tasmanian Herbarium](#)
- [-State Herbarium of South Australia](#)
- [-Northern Territory Herbarium](#)
- [-Western Australian Herbarium](#)
- [-Australian National Herbarium, Canberra](#)
- [-University of New England](#)
- [-Ocean Biogeographic Information System](#)
- [-Australian Government, Department of Defence](#)
- [Forestry Corporation, NSW](#)
- [-Geoscience Australia](#)
- [-CSIRO](#)
- [-Australian Tropical Herbarium, Cairns](#)
- [-eBird Australia](#)
- [-Australian Government – Australian Antarctic Data Centre](#)
- [-Museum and Art Gallery of the Northern Territory](#)
- [-Australian Government National Environmental Science Program](#)
- [-Australian Institute of Marine Science](#)
- [-Reef Life Survey Australia](#)
- [-American Museum of Natural History](#)
- [-Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [-Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- [-Other groups and individuals](#)

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.

NatureMap Species Report

Created By Guest user on 04/12/2019

Current Names Only Yes
Core Datasets Only Yes
Method 'By Circle'
Centre 114° 15' 04" E, 28° 07' 59" S
Buffer 10km
Group By Kingdom

Kingdom	Species	Records
Animalia	353	3102
Chromista	10	13
Fungi	18	33
Plantae	304	516
TOTAL	685	3664

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
Animalia				
1.	<i>Abudefduf sexfasciatus</i>			
2.	24559 <i>Acanthagenys rufogularis</i> (Spiny-cheeked Honeyeater)			
3.	<i>Acanthistius pardalotus</i>			
4.	24261 <i>Acanthiza chrysorrhoa</i> (Yellow-rumped Thornbill)			
5.	<i>Acariformes</i> sp.			
6.	25536 <i>Accipiter fasciatus</i> (Brown Goshawk)			
7.	<i>Acritoptila globosa</i>			
8.	41323 <i>Actitis hypoleucos</i> (Common Sandpiper)		IA	
9.	<i>Adversaeschna brevistyla</i>			
10.	<i>Aeshnidae</i> sp.			
11.	<i>Agraptocorixa eurynome</i>			
12.	<i>Agraptocorixa</i> sp.			
13.	<i>Alboa worooa</i>			
14.	<i>Allotrochosina karri</i>			
15.	<i>Alona rigidicaudis</i>			
16.	<i>Alotanypus dalyupensis</i>			
17.	24312 <i>Anas gracilis</i> (Grey Teal)			
18.	24315 <i>Anas rhynchotis</i> (Australasian Shoveler)			
19.	24316 <i>Anas superciliosa</i> (Pacific Black Duck)			
20.	<i>Anax papuensis</i>			
21.	47414 <i>Anhinga novaehollandiae</i> (Australasian Darter)			
22.	<i>Anisops elstoni</i>			
23.	<i>Anisops nasutus</i>			
24.	<i>Anisops thienemanni</i>			
25.	<i>Anopheles annulipes</i> s.l.			
26.	25634 <i>Anous stolidus</i> (Common Noddy)		IA	
27.	24562 <i>Anthochaera lunulata</i> (Western Little Wattlebird)			
28.	<i>Apocyclops dengizicus</i>			
29.	<i>Apogon doederleini</i>			
30.	<i>Apogon victoriae</i>			
31.	25554 <i>Apus pacificus</i> (Fork-tailed Swift, Pacific Swift)		IA	
32.	24285 <i>Aquila audax</i> (Wedge-tailed Eagle)			
33.	<i>Arcella</i> sp. a (SAP)			
34.	<i>Arcella</i> sp. c (SAP)			
35.	<i>Archaeosynthemis occidentalis</i>			
36.	25558 <i>Ardea ibis</i> (Cattle Egret)			
37.	41324 <i>Ardea modesta</i> (great egret, white egret)			
38.	24610 <i>Ardeotis australis</i> (Australian Bustard)			
39.	25736 <i>Arenaria interpres</i> (Ruddy Turnstone)		IA	
40.	<i>Armatalona macrocopa</i>			
41.	<i>Arrenurus (Truncaturus)</i> sp. 25 (TST)			
42.	25566 <i>Artamus cinereus</i> (Black-faced Woodswallow)			
43.	24355 <i>Artamus minor</i> (Little Woodswallow)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
44.	24356 <i>Artamus personatus</i> (Masked Woodswallow)			
45.	<i>Artemia</i> sp.			
46.	<i>Artoria cingulipes</i>			
47.	<i>Australocypris insularis</i>			
48.	<i>Australoeucyclops darwini</i> (ex <i>Paracyclops</i> sp 1 nr <i>timmsi</i>)			
49.	<i>Austrochilonia subtenuis</i>			
50.	<i>Austrolestes aleison</i>			
51.	<i>Austrolestes annulosus</i>			
52.	24318 <i>Aythya australis</i> (Hardhead)			
53.	<i>Barnardius zonarius</i>			
54.	<i>Bdelloidea</i> sp.			
55.	<i>Berosus approximans</i>			
56.	<i>Berosus discolor</i>			
57.	<i>Berosus</i> sp.			
58.	<i>Bezzia</i> sp. 2 (SAP)			
59.	24319 <i>Biziura lobata</i> (Musk Duck)			
60.	<i>Brachionus plicatilis</i> s.l.			
61.	<i>Brentidae</i> sp.			
62.	42307 <i>Cacomantis pallidus</i> (Pallid Cuckoo)			
63.	24269 <i>Calamanthus campestris</i> (Rufous Fieldwren)			
64.	<i>Calamoecia clitellata</i>			
65.	24779 <i>Calidris acuminata</i> (Sharp-tailed Sandpiper)		IA	
66.	24780 <i>Calidris alba</i> (Sanderling)		IA	
67.	25738 <i>Calidris canutus</i> (Red Knot, knot)		IA	
68.	24784 <i>Calidris ferruginea</i> (Curlew Sandpiper)		T	
69.	24786 <i>Calidris melanotos</i> (Pectoral Sandpiper)		IA	
70.	24788 <i>Calidris ruficollis</i> (Red-necked Stint)		IA	
71.	24789 <i>Calidris subminuta</i> (Long-toed Stint)		IA	
72.	24790 <i>Calidris tenuirostris</i> (Great Knot)		T	
73.	24734 <i>Calyptrorhynchus latirostris</i> (Carnaby's Cockatoo, White-tailed Short-billed Black Cockatoo)		T	
74.	<i>Candonocypris novaezelandiae</i>			
75.	<i>Ceinidae</i> sp.			
76.	<i>Centropyxis</i> cf <i>kahlui</i>			Y
77.	<i>Cephalodella gibba</i>			
78.	<i>Ceratopogonidae</i> sp.			
79.	25575 <i>Charadrius leschenaultii</i> (Greater Sand Plover)		T	
80.	25576 <i>Charadrius mongolus</i> (Lesser Sand Plover)		T	
81.	24377 <i>Charadrius ruficapillus</i> (Red-capped Plover)			
82.	24321 <i>Chenonetta jubata</i> (Australian Wood Duck, Wood Duck)			
83.	47909 <i>Cheramoeca leucosterna</i> (White-backed Swallow)			
84.	<i>Chironominae</i> sp.			
85.	<i>Chironomus</i> aff. <i>alternans</i> (V24) (CB)			
86.	<i>Choeroichthys suillus</i>			
87.	<i>Chroicocephalus novaehollandiae</i>			
88.	24288 <i>Circus approximans</i> (Swamp Harrier)			
89.	<i>Cladopelma curtivalva</i>			
90.	24774 <i>Cladorhynchus leucocephalus</i> (Banded Stilt)			
91.	<i>Cleidopus gloriamaris</i>			
92.	<i>Cletocamptus dietersi</i>			
93.	<i>Cloeon</i> sp.			
94.	25675 <i>Colluricincla harmonica</i> (Grey Shrike-thrush)			
95.	24399 <i>Columba livia</i> (Domestic Pigeon)	Y		
96.	<i>Colurodontis paxmani</i>			
97.	25568 <i>Coracina novaehollandiae</i> (Black-faced Cuckoo-shrike)			
98.	<i>Corduliidae</i> sp.			
99.	<i>Corixidae</i> sp.			
100.	24416 <i>Corvus bennetti</i> (Little Crow)			
101.	25592 <i>Corvus coronoides</i> (Australian Raven)			
102.	<i>Corynoneura</i> sp.			
103.	<i>Corynoneura</i> sp. (V49) (SAP)			
104.	<i>Coxiella striatula</i>			
105.	24420 <i>Cracticus nigrogularis</i> (Pied Butcherbird)			
106.	25595 <i>Cracticus tibicen</i> (Australian Magpie)			
107.	25596 <i>Cracticus torquatus</i> (Grey Butcherbird)			
108.	25401 <i>Crinia pseudinsignifera</i> (Bleating Froglet)			
109.	<i>Cristiceps australis</i>			
110.	24881 <i>Ctenophorus maculatus</i> subsp. <i>maculatus</i> (Spotted Military Dragon)			
111.	24886 <i>Ctenophorus reticulatus</i> (Western Netted Dragon)			
112.	<i>Culicidae</i> sp.			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
113.	<i>Culicoides</i> sp.			
114.	24322 <i>Cygnus atratus</i> (Black Swan)			
115.	<i>Cypretta baylyi</i>			
116.	<i>Dasyhelea</i> sp.			
117.	<i>Diacypris spinosa</i>			
118.	25607 <i>Dicaeum hirundinaceum</i> (Mistletoebird)			
119.	<i>Dicrotendipes conjunctus</i>			
120.	<i>Diffugia</i> sp.			
121.	<i>Diplacodes bipunctata</i>			
122.	<i>Dipulus caecus</i>			
123.	24470 <i>Dromaius novaehollandiae</i> (Emu)			
124.	<i>Dytiscidae</i> sp.			
125.	<i>Egretta novaehollandiae</i>			
126.	<i>Elanus axillaris</i>			
127.	47937 <i>Elseyornis melanops</i> (Black-fronted Dotterel)			
128.	<i>Enchytraeidae</i> sp.			
129.	<i>Eolophus roseicapillus</i>			
130.	24651 <i>Eopsaltria australis</i> subsp. <i>griseogularis</i> (Western Yellow Robin)			
131.	<i>Ephydriidae</i> sp. 6 (SAP)			
132.	<i>Epinephelus coioides</i>			
133.	24567 <i>Epthianura albiglans</i> (White-fronted Chat)			
134.	24568 <i>Epthianura aurifrons</i> (Orange Chat)			
135.	24570 <i>Epthianura tricolor</i> (Crimson Chat)			
136.	24379 <i>Erythronyx cinctus</i> (Red-kneed Dotterel)			
137.	<i>Euchlanis deflexa</i>			Y
138.	<i>Eucyclops australiensis</i>			
139.	<i>Eviota bimaculata</i>			
140.	25621 <i>Falco berigora</i> (Brown Falcon)			
141.	25622 <i>Falco cenchroides</i> (Australian Kestrel, Nankeen Kestrel)			
142.	25623 <i>Falco longipennis</i> (Australian Hobby)			
143.	25624 <i>Falco peregrinus</i> (Peregrine Falcon)		S	
144.	<i>Forcypomyia</i> sp.			
145.	24761 <i>Fulica atra</i> subsp. <i>australis</i> (Eurasian Coot)			
146.	24959 <i>Gehyra variegata</i>			
147.	47954 <i>Gelochelidon nilotica</i> (Gull-billed Tern)		IA	
148.	24401 <i>Geopelia cuneata</i> (Diamond Dove)			
149.	25585 <i>Geopelia striata</i> (Zebra Dove)			
150.	24443 <i>Grallina cyanoleuca</i> (Magpie-lark)			
151.	<i>Gymnothorax woodwardi</i>			
152.	25627 <i>Haematopus fuliginosus</i> (Sooty Oystercatcher)			
153.	24487 <i>Haematopus longirostris</i> (Pied Oystercatcher)			
154.	24293 <i>Haliaeetus leucogaster</i> (White-bellied Sea-Eagle)			
155.	24295 <i>Haliastur sphenurus</i> (Whistling Kite)			
156.	<i>Halichoeres brownfieldi</i>			
157.	<i>Halicyclops ambiguus</i>			Y
158.	<i>Halplus fuscatus</i>			
159.	<i>Helcogramma decurrens</i>			
160.	<i>Helochares tenuistriatus</i>			
161.	<i>Hemicordulia tau</i>			
162.	24961 <i>Heteronotia binoei</i> (Bynoe's Gecko)			
163.	47965 <i>Hieraaetus morphnoides</i> (Little Eagle)			
164.	25734 <i>Himantopus himantopus</i> (Black-winged Stilt)			
165.	24491 <i>Hirundo neoxena</i> (Welcome Swallow)			
166.	<i>Hydrachna australica</i>			
167.	<i>Hydrachna</i> nr. <i>approximata</i> (SAP)			
168.	<i>Hydroglyphus leai</i>			
169.	<i>Hydrophilidae</i> sp.			
170.	48587 <i>Hydroprogne caspia</i> (Caspian Tern)		IA	
171.	<i>Hydropsychidae</i> sp.			
172.	<i>Hyphydrus elegans</i>			
173.	<i>Hyphydrus</i> sp.			
174.	<i>Ischnura aurora aurora</i>			
175.	<i>Ischnura heterosticta heterosticta</i>			
176.	<i>Istiblennius meleagris</i>			
177.	<i>Kennethia cristata</i>			
178.	<i>Keratella procurva</i>			
179.	<i>Kiefferulus intertinctus</i>			
180.	<i>Labracinus lineatus</i>			
181.	25638 <i>Larus pacificus</i> (Pacific Gull)			
182.	<i>Lecane bulla</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
183.	<i>Lecane luna</i>			
184.	<i>Lecane thalera</i>			
185.	<i>Lepadichthys sandaracatus</i>			
186.	<i>Lepidoptera</i> sp.			
187.	<i>Leptoceridae</i> sp.			
188.	25148 <i>Lerista lineopunctulata</i>			
189.	25165 <i>Lerista praepedita</i>			
190.	<i>Lestidae</i> sp.			
191.	<i>Lethrinus genivittatus</i>			
192.	<i>Lethrinus miniatus</i>			
193.	25005 <i>Lialis burtonis</i>			
194.	<i>Libellulidae</i> sp.			
195.	25661 <i>Lichmera indistincta</i> (Brown Honeyeater)			
196.	<i>Limnichidae</i> sp.			
197.	<i>Limnocythere mowbrayensis</i>			
198.	25415 <i>Limnodynastes dorsalis</i> (Western Banjo Frog)			
199.	<i>Limnophyes vestitus</i> (V41)			
200.	30932 <i>Limosa lapponica</i> (Bar-tailed Godwit)		IA	
201.	25388 <i>Litoria moorei</i> (Motorbike Frog)			
202.	<i>Lophoictinia isura</i>			
203.	<i>Lourinidae</i> sp.			Y
204.	24132 <i>Macropus fuliginosus</i> (Western Grey Kangaroo)			
205.	<i>Macrothrix breviseta</i>			
206.	24326 <i>Malacorhynchus membranaceus</i> (Pink-eared Duck)			
207.	25651 <i>Malurus lamberti</i> (Variegated Fairy-wren)			
208.	24544 <i>Malurus lamberti</i> subsp. <i>assimilis</i> (Variegated Fairy-wren)			
209.	25652 <i>Malurus leucopterus</i> (White-winged Fairy-wren)			
210.	24551 <i>Malurus pulcherrimus</i> (Blue-breasted Fairy-wren)			
211.	25654 <i>Malurus splendens</i> (Splendid Fairy-wren)			
212.	24583 <i>Manorina flavigula</i> (Yellow-throated Miner)			
213.	25758 <i>Megalurus gramineus</i> (Little Grassbird)			
214.	<i>Megaporus</i> sp.			
215.	24598 <i>Merops ornatus</i> (Rainbow Bee-eater)			
216.	<i>Mesochra baylyi</i>			
217.	<i>Mesocyclops brooksi</i>			
218.	<i>Mesocyclops</i> sp.			
219.	<i>Mesostigmata</i> sp.			
220.	<i>Microcarbo melanoleucos</i>			
221.	<i>Micronecta robusta</i>			
222.	<i>Microvelia</i> (<i>Austromicrovelia</i>) <i>peramoena</i>			
223.	<i>Microvelia</i> (<i>Pacificovelia</i>) <i>oceanica</i>			
224.	<i>Microvelia</i> sp.			
225.	25191 <i>Morethia lineocellata</i>			
226.	48008 <i>Morus serrator</i> (Australasian Gannet)			
227.	<i>Muraenichthys</i> sp.			
228.	24223 <i>Mus musculus</i> (House Mouse)	Y		
229.	25420 <i>Myobatrachus gouldii</i> (Turtle Frog)			
230.	<i>Mytilocypris mytiloides</i>			
231.	<i>Naididae</i> (ex <i>Tubificidae</i>)			
232.	<i>Nannophya occidentalis</i>			
233.	<i>Necterosoma penicillatus</i>			
234.	<i>Necterosoma</i> sp.			
235.	<i>Nematoda</i> sp.			
236.	<i>Nemertini</i> sp.			
237.	<i>Neohydrocoptus subfasciatus</i>			
238.	33984 <i>Neopasiphae simplicior</i> (a short-tongued bee)		T	
239.	<i>Nitocra</i> sp. 3 (SAP)			Y
240.	<i>Nitocra</i> sp. 5 (nr <i>reducta</i>) (SAP)			
241.	<i>Notolabrus parilus</i>			
242.	<i>Notonectidae</i> sp.			
243.	24799 <i>Numenius minutus</i> (Little Curlew, Little Whimbrel)		IA	
244.	25742 <i>Numenius phaeopus</i> (Whimbrel)		IA	
245.	24194 <i>Nyctophilus geoffroyi</i> (Lesser Long-eared Bat)			
246.	24742 <i>Nymphicus hollandicus</i> (Cockatiel)			
247.	24407 <i>Ocyphaps lophotes</i> (Crested Pigeon)			
248.	<i>Oecetis</i> sp.			
249.	<i>Oligochaeta</i> sp.			
250.	<i>Onychocamptus bengalensis</i>			
251.	<i>Onychohydus</i> sp.			
252.	<i>Oribatida</i> sp.			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
253.	<i>Orthetrum caledonicum</i>			
254.	<i>Orthocladinae</i> sp.			
255.	<i>Orthocladinae</i> sp. 1 (SAP)			
256.	<i>Oxyethira</i> sp.			
257.	25680 <i>Pachycephala rufiventris</i> (Rufous Whistler)			
258.	48591 <i>Pandion cristatus</i> (Osprey, Eastern Osprey)		IA	
259.	<i>Paracyclops chiltoni</i>			
260.	<i>Paracyclops</i> sp.			
261.	<i>Paradoxostoma</i> sp.			Y
262.	<i>Parakiefferiella variegatus</i>			
263.	<i>Paralimnophyes pullulus</i> (V42)			
264.	<i>Paramerina levidensis</i>			
265.	<i>Parapercis haackei</i>			
266.	<i>Parma occidentalis</i>			
267.	24648 <i>Pelecanus conspicillatus</i> (Australian Pelican)			
268.	<i>Pempheris mangula</i>			
269.	<i>Pescecyclops</i> sp. 462			
270.	48060 <i>Petrochelidon ariel</i> (Fairy Martin)			
271.	48061 <i>Petrochelidon nigricans</i> (Tree Martin)			
272.	25697 <i>Phalacrocorax carbo</i> (Great Cormorant)			
273.	24667 <i>Phalacrocorax sulcirostris</i> (Little Black Cormorant)			
274.	25699 <i>Phalacrocorax varius</i> (Pied Cormorant)			
275.	24801 <i>Phalaropus lobatus</i> (Red-necked Phalarope)		IA	
276.	24409 <i>Phaps chalcoptera</i> (Common Bronzewing)			
277.	24802 <i>Philomachus pugnax</i> (Ruff, reeve)		IA	
278.	<i>Platynectes</i> sp.			
279.	<i>Plectorhinchus flavomaculatus</i>			
280.	<i>Plotiopsis</i> sp.			
281.	<i>Plotosus lineatus</i>			
282.	24382 <i>Pluvialis fulva</i> (Pacific Golden Plover)		IA	
283.	24383 <i>Pluvialis squatarola</i> (Grey Plover)		IA	
284.	25703 <i>Podargus strigoides</i> (Tawny Frogmouth)			
285.	24681 <i>Poliiocephalus poliocephalus</i> (Hoary-headed Grebe)			
286.	<i>Polypedilum nr. convexum</i> (SAP)			
287.	<i>Polypedilum nubifer</i>			
288.	<i>Pomacentrus milleri</i>			
289.	24769 <i>Porzana fluminea</i> (Australian Spotted Crake)			
290.	24771 <i>Porzana tabuensis</i> (Spotless Crake)			
291.	<i>Priolepis nuchifasciata</i>			
292.	<i>Pristina sima</i>			
293.	<i>Procladius paludicola</i>			
294.	<i>Procladius villosimanus</i>			
295.	<i>Pseudochromis wilsoni</i>			
296.	42416 <i>Pseudonaja mengdeni</i> (Western Brown Snake)			
297.	<i>Pyrallidae nr. sp. 39/40 of JHH</i> (SAP)			
298.	24245 <i>Rattus rattus</i> (Black Rat)	Y		
299.	24776 <i>Recurvirostra novaehollandiae</i> (Red-necked Avocet)			
300.	48096 <i>Rhipidura albiscapa</i> (Grey Fantail)			
301.	25614 <i>Rhipidura leucophrys</i> (Willie Wagtail)			
302.	<i>Robertsonia</i> sp.			Y
303.	<i>Sargocentron rubrum</i>			
304.	<i>Sarscyridopsis aculeata</i>			
305.	<i>Schuettea woodwardi</i>			
306.	<i>Scirtidae</i> sp.			
307.	<i>Scorpaena sumptuosa</i>			
308.	<i>Scorpaenodes steenei</i>			
309.	25534 <i>Sericornis frontalis</i> (White-browed Scrubwren)			
310.	<i>Sillago robusta</i>			
311.	<i>Simocephalus elizabethae</i>			
312.	<i>Simocephalus</i> sp.			
313.	<i>Simuliidae</i> sp.			
314.	<i>Simulium ornatipes</i>			
315.	<i>Solegnathus lettiensis</i>			
316.	<i>Staphylinidae</i> sp.			
317.	<i>Stegastes obreptus</i>			
318.	48594 <i>Sternula nereis</i> (Fairy Tern)			
319.	<i>Stratiomyidae</i> sp.			
320.	25590 <i>Streptopelia senegalensis</i> (Laughing Turtle-Dove)	Y		
321.	24942 <i>Strophurus spinigerus</i> subsp. <i>spinigerus</i>			
322.	33992 <i>Synemon gratiosa</i> (Graceful Sunmoth)			

P4

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
323.	<i>Tabanidae</i> sp.			
324.	25705 <i>Tachybaptus novaehollandiae</i> (Australasian Grebe, Black-throated Grebe)			
325.	24682 <i>Tachybaptus novaehollandiae</i> subsp. <i>novaehollandiae</i> (Australasian Grebe, Black-throated Grebe)			
326.	24331 <i>Tadorna tadornoides</i> (Australian Shelduck, Mountain Duck)			
327.	30870 <i>Taeniopygia guttata</i> (Zebra Finch)			
328.	<i>Tanypodinae</i> sp.			
329.	<i>Tanytarsus barbitarsis</i>			
330.	<i>Tanytarsus fuscithorax/semibarbitarsus</i>			
331.	<i>Tanytarsus palmatus</i>			
332.	<i>Tanytarsus</i> sp. D (SAP)			
333.	<i>Tanytarsus</i> sp. G (SAP)			
334.	24167 <i>Tarsipes rostratus</i> (Honey Possum, Noolbenger)			
335.	48597 <i>Thalasseus bergii</i> (Crested Tern)		IA	
336.	<i>Thalassoma septemfasciata</i>			
337.	<i>Thienemanniella</i> sp. (V19) (SAP)			
338.	<i>Tipulidae</i> sp.			
339.	<i>Tipulidae</i> type F (SAP)			
340.	25549 <i>Todiramphus sanctus</i> (Sacred Kingfisher)			
341.	48141 <i>Tribonyx ventralis</i> (Black-tailed Native-hen)			
342.	24803 <i>Tringa brevipes</i> (Grey-tailed Tattler)		P4	
343.	24806 <i>Tringa glareola</i> (Wood Sandpiper)		IA	
344.	24808 <i>Tringa nebularia</i> (Common Greenshank, greenshank)		IA	
345.	24809 <i>Tringa stagnatilis</i> (Marsh Sandpiper, little greenshank)		IA	
346.	<i>Triplectides australis</i>			
347.	<i>Turbellaria</i> sp.			
348.	<i>Veliidae</i> sp.			
349.	<i>Venatrix pullastra</i>			
350.	<i>Venonia micarioides</i>			
351.	<i>Xanthagrion erythroneurum</i>			
352.	41351 <i>Xenus cinereus</i> (Terek Sandpiper)		IA	
353.	25765 <i>Zosterops lateralis</i> (Grey-breasted White-eye, Silvereye)			

Chromista

354.	35910 <i>Canistrocarpus crispatus</i>			
355.	26586 <i>Caulocystis uvifera</i>			
356.	26766 <i>Dictyopteris muelleri</i>			
357.	26778 <i>Dictyota furcellata</i>			
358.	26810 <i>Encyothalia cliftonii</i>			
359.	26949 <i>Hydroclathrus clathratus</i>			
360.	27043 <i>Lobophora variegata</i>			
361.	27246 <i>Sargassum lacerifolium</i>			
362.	35911 <i>Scytosiphon lomentaria</i>			
363.	27373 <i>Zonaria turneriana</i>			

Fungi

364.	27574 <i>Acarospora citrina</i>			
365.	31099 <i>Caloplaca kantvilasii</i>			
366.	48963 <i>Caloplaca lithophila</i>			
367.	48176 <i>Cladia beaugleholei</i>			
368.	48177 <i>Cladia muelleri</i>			
369.	28208 <i>Cladonia cervicornis</i> subsp. <i>verticillata</i>			
370.	27753 <i>Fulgensia bracteata</i>			
371.	27754 <i>Fulgensia subbracteata</i>			
372.	<i>Lecanora</i> sp.			
373.	27815 <i>Lecanora sphaerospora</i>			
374.	<i>Lecidea</i> sp.			
375.	30457 <i>Notocladonia cochleata</i>			
376.	27935 <i>Peltula euploca</i>			
377.	49073 <i>Peziza austrogeaster</i>			
378.	27999 <i>Psora crystallifera</i>			
379.	28000 <i>Psora decipiens</i>			
380.	28060 <i>Siphula coriacea</i>			
381.	28070 <i>Thysanothecium hookeri</i>			

Plantae

382.	16111 <i>Acacia alata</i> var. <i>biglandulosa</i>			
383.	3225 <i>Acacia ashbyae</i>			
384.	3376 <i>Acacia idiomorpha</i>			
385.	11611 <i>Acacia lasiocarpa</i> var. <i>lasiocarpa</i>			
386.	14612 <i>Acacia latipes</i> subsp. <i>licina</i>		P3	
387.	14134 <i>Acacia pelophila</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
			P1	
388.	15481 <i>Acacia pulchella</i> var. <i>glaberrima</i>			
389.	3521 <i>Acacia ridleyana</i>		P3	
390.	3525 <i>Acacia rostellifera</i> (Summer-scented Wattle)			
391.	3532 <i>Acacia scirpifolia</i>			
392.	3549 <i>Acacia spathulifolia</i>			
393.	15484 <i>Acacia sphacelata</i> subsp. <i>sphacelata</i>			
394.	1208 <i>Acanthocarpus preissii</i>			
395.	20797 <i>Acanthocarpus</i> sp. <i>Ajana</i> (C.A. Gardner 8596)			
396.	1775 <i>Adenanthos cygnorum</i> (Common Woollybush)			
397.	11837 <i>Adenanthos cygnorum</i> subsp. <i>cygnorum</i> (Common Woollybush)			
398.	17422 <i>Adriana tomentosa</i> var. <i>tomentosa</i>			
399.	36277 <i>Aloe vera</i> var. <i>officinalis</i>	Y		
400.	4905 <i>Alyogyne hakeifolia</i>			
401.	4906 <i>Alyogyne huegelii</i> (Lilac Hibiscus)			
402.	42940 <i>Alyogyne</i> sp. <i>Geraldton</i> (R. Davis 3487)			
403.	13267 <i>Amyema linophylla</i> subsp. <i>linophylla</i>			
404.	13266 <i>Amyema miraculosa</i> subsp. <i>miraculosa</i>			
405.	40914 <i>Androcalva gaudichaudii</i>			
406.	11434 <i>Anigozanthos humilis</i> subsp. <i>humilis</i>			
407.	1410 <i>Anigozanthos kalbarriensis</i> (Kalbarri Catspaw)			
408.	11565 <i>Anigozanthos manglesii</i> subsp. <i>quadrans</i>			
409.	11725 <i>Anthocercis ilicifolia</i> subsp. <i>ilicifolia</i>			
410.	6949 <i>Anthocercis littorea</i> (Yellow Tailflower)			
411.	3180 <i>Aphanopetalum clematideum</i>			
412.	26486 <i>Asparagopsis taxiformis</i>			
413.	20695 <i>Astroloma</i> sp. <i>Kalbarri</i> (D. & B. Bellairs 1368)		P2	
414.	16369 <i>Atriplex canescens</i>	Y		Y
415.	2452 <i>Atriplex cinerea</i> (Grey Saltbush)			
416.	2463 <i>Atriplex isatidea</i> (Coast Saltbush)			
417.	2470 <i>Atriplex paludosa</i> (Marsh Saltbush)			
418.	11525 <i>Atriplex paludosa</i> subsp. <i>baudinii</i>			
419.	17237 <i>Austrostipa elegantissima</i>			
420.	17240 <i>Austrostipa flavescens</i>			
421.	17244 <i>Austrostipa macalpinei</i>			
422.	17246 <i>Austrostipa nitida</i>			
423.	233 <i>Avena barbata</i> (Bearded Oat)	Y		
424.	48221 <i>Balladonia aevoides</i>		P3	
425.	32524 <i>Banksia fraseri</i> var. <i>ashbyi</i>			
426.	11386 <i>Banksia leptophylla</i> var. <i>melletica</i>			
427.	32079 <i>Banksia sessilis</i> var. <i>flabellifolia</i>			
428.	1852 <i>Banksia telmatiaea</i> (Swamp Fox Banksia)			
429.	743 <i>Baumea juncea</i> (Bare Twigrush)			
430.	747 <i>Baumea rubiginosa</i>			
431.	748 <i>Baumea vaginalis</i> (Sheath Twigrush)			
432.	31606 <i>Blackallia nudiflora</i> (Wedge-leaved Cryptandra)		P3	
433.	11381 <i>Boronia ramosa</i> subsp. <i>anethifolia</i>			
434.	3719 <i>Bossiaea spinescens</i>			
435.	8661 <i>Brachypodium distachyon</i> (False Brome)	Y		
436.	249 <i>Bromus diandrus</i> (Great Brome)	Y		
437.	6213 <i>Bupleurum semicompositum</i>	Y		
438.	29439 <i>Caesia</i> sp. <i>Wongan</i> (K.F. Kenneally 8820)			
439.	13618 <i>Caladenia elegans</i>		T	
440.	15348 <i>Caladenia flava</i> subsp. <i>flava</i>			
441.	15349 <i>Caladenia flava</i> subsp. <i>maculata</i>			
442.	17760 <i>Caladenia nobilis</i>			
443.	2860 <i>Calandrinia polyandra</i> (Parakeelya)			
444.	19304 <i>Calectasia browneana</i>		P2	
445.	5401 <i>Calothamnus blepharospemus</i>			
446.	35758 <i>Calothamnus quadrifidus</i> subsp. <i>homalophyllus</i> (Murchison Clawflower)			
447.	5450 <i>Calytrix depressa</i>			
448.	5460 <i>Calytrix fraseri</i> (Pink Summer Calytrix)			
449.	7911 <i>Carthamus lanatus</i> (Saffron Thistle)	Y		
450.	2948 <i>Cassytha aurea</i>			
451.	12073 <i>Cassytha aurea</i> var. <i>aurea</i>			
452.	2952 <i>Cassytha glabella</i> (Tangled Dodder Laurel)			
453.	2957 <i>Cassytha racemosa</i> (Dodder Laurel)			
454.	11799 <i>Cassytha racemosa</i> forma <i>racemosa</i>			
455.	48455 <i>Caulerpa geminata</i>			
456.	258 <i>Cenchrus ciliaris</i> (Buffel Grass)	Y		

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
457.	1124	<i>Centrolepis cephaloformis</i>			
458.	17685	<i>Chaetanthus aristatus</i>			
459.	2494	<i>Chenopodium murale</i> (Nettle-leaf Goosefoot)	Y		
460.	29619	<i>Chondrophycus brandenii</i>			
461.	4853	<i>Clematicissus angustissima</i>			
462.	26683	<i>Codium spongiosum</i>			
463.	4550	<i>Comesperma calymega</i> (Blue-spike Milkwort)			
464.	4554	<i>Comesperma flavum</i>			
465.	4564	<i>Comesperma virgatum</i> (Milkwort)			
466.	40872	<i>Commersonia borealis</i>			
467.	2776	<i>Commicarpus australis</i> (Perennial Tar Vine)			
468.	15607	<i>Conospermum acerosum</i> subsp. <i>acerosum</i>			
469.	15608	<i>Conospermum acerosum</i> subsp. <i>hirsutum</i>			
470.	15513	<i>Conospermum boreale</i> subsp. <i>boreale</i>			
471.	15611	<i>Conospermum stoechadis</i> subsp. <i>stoechadis</i> (Common Smokebush)			
472.	12028	<i>Conostylis aculeata</i> subsp. <i>septentrionora</i>			
473.	1423	<i>Conostylis aurea</i> (Golden Conostylis)			
474.	1446	<i>Conostylis prolifera</i> (Mat Cottonheads)			
475.	1456	<i>Conostylis stylidioides</i>			
476.	6614	<i>Convolvulus remotus</i>			
477.	3137	<i>Crassula colorata</i> (Dense Stonecrop)			
478.	35839	<i>Cristonia stenophylla</i>			
479.	4802	<i>Cryptandra mutila</i>			
480.	6663	<i>Cuscuta epithymum</i> (Lesser Dodder, Greater Dodder)	Y		
481.	283	<i>Cynodon dactylon</i> (Couch)	Y		
482.	794	<i>Cyperus gymnocaulos</i> (Spiny Flat-sedge)			
483.	801	<i>Cyperus laevigatus</i>	Y		
484.	7421	<i>Dampiera altissima</i> (Tall Dampiera)			
485.	11723	<i>Dampiera incana</i> var. <i>incana</i>			
486.	7459	<i>Dampiera oligophylla</i> (Sparse-leaved Dampiera)			
487.	5522	<i>Darwinia pauciflora</i>			
488.	5534	<i>Darwinia virescens</i> (Murchison Darwinia)			
489.	18561	<i>Daviesia divaricata</i> subsp. <i>lanulosa</i>			
490.	1287	<i>Dichopogon capillipes</i>			
491.	1290	<i>Dichopogon tyleri</i>			
492.	15270	<i>Diplolaena geraldtonensis</i>			
493.	4456	<i>Diplolaena grandiflora</i> (Wild Rose)			
494.	4748	<i>Diplopeltis petiolaris</i>			
495.	7961	<i>Dittrichia graveolens</i> (Stinkwort)	Y		
496.	10796	<i>Diuris drummondii</i> (Tall Donkey Orchid)		T	
497.	12936	<i>Diuris recurva</i>		P4	
498.	13633	<i>Drakaea concolor</i>		T	
499.	3113	<i>Drosera neesii</i> (Jewel Rainbow)			
500.	3116	<i>Drosera omissa</i> (Bright Sundew)			
501.	346	<i>Ehrharta brevifolia</i> (Annual Veldt Grass)	Y		
502.	347	<i>Ehrharta calycina</i> (Perennial Veldt Grass)	Y		
503.	353	<i>Eleusine indica</i> (Crowsfoot Grass)	Y		
504.	378	<i>Eragrostis dielsii</i> (Mallee Lovegrass)			
505.	5538	<i>Eremaea brevifolia</i>			
506.	5539	<i>Eremaea ebracteata</i>			
507.	14102	<i>Eremaea ebracteata</i> var. <i>ebracteata</i>			
508.	17175	<i>Eremophila glabra</i> subsp. <i>albicans</i>			
509.	14193	<i>Eremophila glabra</i> subsp. <i>carcosa</i>			
510.	7241	<i>Eremophila microtheca</i> (Heath-like Eremophila)		P4	
511.	4333	<i>Erodium cicutarium</i> (Common Storksbill)	Y		
512.	12740	<i>Erymophyllum tenellum</i>			
513.	12895	<i>Eucalyptus arachnaea</i> subsp. <i>arachnaea</i>			
514.	5730	<i>Eucalyptus oraria</i> (Ooragmandee)			
515.	4620	<i>Euphorbia boophthona</i> (Gascoyne Spurge)			
516.	4644	<i>Euphorbia sharkoensis</i>			
517.	4648	<i>Euphorbia terracina</i> (Geraldton Carnation Weed)	Y		
518.	5193	<i>Frankenia confusa</i>		P4	
519.	5209	<i>Frankenia pauciflora</i> (Seaheath)			
520.	907	<i>Gahnia trifida</i> (Coast Saw-sedge)			
521.	12780	<i>Gilberta tenuifolia</i>			
522.	3938	<i>Glycine canescens</i> (Silky Glycine)			
523.	7983	<i>Gnaphalium indutum</i> (Tiny Cudweed)			
524.	3957	<i>Gompholobium tomentosum</i> (Hairy Yellow Pea)			
525.	7495	<i>Goodenia berardiana</i>			
526.	18116	<i>Grevillea commutata</i> subsp. <i>commutata</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
527.	2032 <i>Grevillea leucopteris</i> (White Plume Grevillea)			
528.	8838 <i>Grevillea pinaster</i>			
529.	2113 <i>Grevillea triloba</i>		P3	
530.	5011 <i>Guichenotia ledifolia</i>			
531.	6696 <i>Halgania sericiflora</i>			
532.	47213 <i>Halimeda versatilis</i>			
533.	5120 <i>Hibbertia desmophylla</i>			
534.	5171 <i>Hibbertia spicata</i>			
535.	450 <i>Hordeum marinum</i>	Y		
536.	6234 <i>Hydrocotyle medicaginoides</i> (Trefoil Pennywort)			
537.	35071 <i>Hypocalymma angustifolium</i> subsp. Hutt River (S. Patrick 2982)		T	
538.	5821 <i>Hypocalymma longifolium</i>		T	
539.	8086 <i>Hypochaeris glabra</i> (Smooth Catsear)	Y		
540.	1070 <i>Hypolaena exsulca</i>			
541.	7396 <i>Isotoma hypocrateriformis</i> (Woodbridge Poison)			
542.	3992 <i>Isotropis cuneifolia</i> (Granny Bonnets)			
543.	14780 <i>Jacksonia arenicola</i>			
544.	14785 <i>Jacksonia rigida</i>			
545.	1175 <i>Juncus acutus</i> (Spiny Rush)	Y		
546.	1178 <i>Juncus bufonius</i> (Toad Rush)	Y		
547.	11922 <i>Juncus kraussii</i> subsp. <i>australiensis</i>			
548.	4044 <i>Kennedia prostrata</i> (Scarlet Runner)			
549.	5043 <i>Lasiopetalum oldfieldii</i>		P3	
550.	13289 <i>Lawrencella davenportii</i>			
551.	4955 <i>Lawrencia glomerata</i>			
552.	4959 <i>Lawrencia squamata</i>			
553.	4960 <i>Lawrencia viridigrisea</i>			
554.	7572 <i>Lechenaultia expansa</i>			
555.	7580 <i>Lechenaultia linarioides</i> (Yellow Leschenaultia)			
556.	946 <i>Lepidosperma striatum</i>			
557.	6487 <i>Limonium companyonis</i>	Y		
558.	41780 <i>Limonium hyblaenum</i>	Y		
559.	9289 <i>Lobelia anceps</i> (Angled Lobelia)			
560.	7403 <i>Lobelia heterophylla</i> (Wing-seeded Lobelia)			
561.	1227 <i>Lomandra hastilis</i>			
562.	1231 <i>Lomandra maritima</i>			
563.	4060 <i>Lotus australis</i> (Austral Trefoil)			
564.	18049 <i>Lyginia imberbis</i>			
565.	36375 <i>Lysimachia arvensis</i> (Pimpernel)	Y		
566.	2839 <i>Macarthuria australis</i>			
567.	19384 <i>Melaleuca bisulcata</i>			
568.	5887 <i>Melaleuca cardiophylla</i> (Tangling Melaleuca)			
569.	18112 <i>Melaleuca leuropoma</i>			
570.	5959 <i>Melaleuca raphiophylla</i> (Swamp Paperbark)			
571.	5987 <i>Melaleuca viminea</i> (Mohan)			
572.	13280 <i>Melaleuca viminea</i> subsp. <i>viminea</i>			
573.	4085 <i>Melilotus indicus</i>	Y		
574.	2813 <i>Mesembryanthemum crystallinum</i> (Iceplant)	Y		
575.	2814 <i>Mesembryanthemum nodiflorum</i> (Slender Iceplant)	Y		
576.	4100 <i>Mirbelia spinosa</i>			
577.	19177 <i>Moraea setifolia</i>	Y		
578.	7291 <i>Myoporum insulare</i> (Blueberry Tree, boobialla)			
579.	17158 <i>Myoporum montanum</i> (Native Myrtle)			
580.	138 <i>Najas marina</i> (Prickly Water Nymph)			
581.	2401 <i>Nuytsia floribunda</i> (Christmas Tree, Mudja)			
582.	6138 <i>Oenothera drummondii</i> (Beach Evening Primrose)	Y		
583.	18256 <i>Opercularia spermacoea</i>			
584.	516 <i>Parapholis incurva</i> (Coast Barbgrass)	Y		
585.	12670 <i>Parietaria cardiostegia</i>			
586.	1762 <i>Parietaria debilis</i> (Pellitory)			
587.	2290 <i>Petrophile conifera</i>			
588.	2301 <i>Petrophile macrostachya</i>			
589.	551 <i>Phalaris minor</i> (Lesser Canary Grass)	Y		
590.	4675 <i>Phyllanthus calycinus</i> (False Boronia)			
591.	6008 <i>Phymatocarpus porphyrocephalus</i>			
592.	20220 <i>Pileanthus rubronitidus</i>			
593.	18250 <i>Pileanthus vernicosus</i>			
594.	5246 <i>Pimelea gilgiana</i>			
595.	5256 <i>Pimelea microcephala</i> (Shrubby Riceflower, Banjine)			
596.	573 <i>Poa drummondiana</i> (Knotted Poa)			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
597.	8184	<i>Podotheca gnaphalioides</i> (Golden Long-heads)			
598.	8188	<i>Pogonolepis stricta</i>			
599.	582	<i>Polypogon monspeliensis</i> (Annual Beardgrass)	Y		
600.	1671	<i>Prasophyllum elatum</i> (Tall Leek Orchid)			
601.	1672	<i>Prasophyllum fimbria</i> (Fringed Leek Orchid)			
602.	37460	<i>Pterostylis sinuata</i>		T	
603.	2717	<i>Ptilotus divaricatus</i> (Climbing Mulla Mulla)			
604.	2719	<i>Ptilotus eriotrichus</i>			
605.	2766	<i>Ptilotus villosiflorus</i>			
606.	592	<i>Puccinellia stricta</i> (Marsh Grass)			
607.	41041	<i>Quoya atriplicina</i>			
608.	11728	<i>Rhagodia latifolia</i> subsp. <i>latifolia</i>			
609.	2584	<i>Rhagodia preissii</i>			
610.	13241	<i>Rhodanthe chlorocephala</i> subsp. <i>rosea</i>			
611.	48887	<i>Roepera billardiarei</i>			
612.	48891	<i>Roepera fruticulosa</i>			
613.	40426	<i>Rytidosperma occidentale</i>			
614.	2906	<i>Sagina apetala</i> (Annual Pearlwort)	Y		
615.	48433	<i>Salicornia blackiana</i>			
616.	48430	<i>Salicornia quinqueflora</i>			
617.	6484	<i>Samolus repens</i> (Creeping Brookweed)			
618.	14107	<i>Samolus repens</i> var. <i>paucifolius</i>			
619.	2356	<i>Santalum acuminatum</i> (Quandong, Warnga)			
620.	7614	<i>Scaevola globulifera</i>			
621.	17026	<i>Scaevola kallophylla</i>		P4	
622.	7634	<i>Scaevola phlebopetala</i> (Velvet Fanflower)			
623.	12588	<i>Scaevola virgata</i>			
624.	972	<i>Schoenus armeria</i>			
625.	994	<i>Schoenus humilis</i>			
626.	1011	<i>Schoenus rigens</i>			
627.	1018	<i>Schoenus subfascicularis</i>			
628.	6034	<i>Scholtzia laxiflora</i>			
629.	49129	<i>Scholtzia pentamera</i> subsp. <i>pentamera</i>			
630.	15427	<i>Scholtzia spatulata</i>			
631.	6041	<i>Scholtzia umbellifera</i>			
632.	6544	<i>Sebaea ovata</i> (Yellow Sebaea)			
633.	8207	<i>Senecio glossanthus</i> (Slender Groundsel)			
634.	3069	<i>Sisymbrium erysimoides</i> (Smooth Mustard)	Y		
635.	6988	<i>Solanum americanum</i> (Glossy Nightshade)	Y		
636.	7022	<i>Solanum nigrum</i> (Black Berry Nightshade)	Y		
637.	7025	<i>Solanum oldfieldii</i>			
638.	7037	<i>Solanum symonii</i>			
639.	9367	<i>Sonchus hydrophilus</i> (Native Sowthistle)			
640.	8231	<i>Sonchus oleraceus</i> (Common Sowthistle)	Y		
641.	2915	<i>Spergularia rubra</i> (Sand Spurry)	Y		
642.	635	<i>Sporobolus virginicus</i> (Marine Couch)			
643.	19953	<i>Stachystemon nematophorus</i>		P4	
644.	43601	<i>Stackhousia</i> sp. <i>Mid west coastal</i> (D. & B. Bellairs 6561)			
645.	2316	<i>Stirlingia latifolia</i> (Blueboy)			
646.	27318	<i>Struvea plumosa</i>			
647.	7693	<i>Stylidium brunonianum</i> (Pink Fountain Triggerplant)			
648.	17412	<i>Stylidium kalbarriense</i>			
649.	25837	<i>Stylidium purpureum</i> (Purple Fountain Triggerplant)			
650.	19247	<i>Stylidium septentrionale</i>			
651.	3182	<i>Stylobasium spathulatum</i> (Pebble Bush)			
652.	2639	<i>Suaeda australis</i> (Seablite)			
653.	4220	<i>Swainsona canescens</i> (Grey Swainsona)			
654.	25902	<i>Symphyotrichum squamatum</i> (Bushy Starwort)	Y		
655.	33236	<i>Tecticornia halocnemoides</i> (Shrubby Samphire)			
656.	33319	<i>Tecticornia indica</i> subsp. <i>bidens</i>			
657.	31618	<i>Tecticornia pruinosa</i>			
658.	31716	<i>Tecticornia syncarpa</i>			
659.	31717	<i>Tecticornia undulata</i>			
660.	2820	<i>Tetragonia decumbens</i> (Sea Spinach)	Y		
661.	2823	<i>Tetragonia implexicoma</i> (Bower Spinach)			
662.	673	<i>Themeda triandra</i>			
663.	2644	<i>Threlkeldia diffusa</i> (Coast Bonefruit)			
664.	1339	<i>Thysanotus multiflorus</i> (Many-flowered Fringe Lily)			
665.	1356	<i>Thysanotus teretifolius</i>			
666.	1361	<i>Tricoryne elatior</i> (Yellow Autumn Lily)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
667.	4312 <i>Trifolium striatum</i> (Knotted Clover)	Y		
668.	147 <i>Triglochin mucronata</i>			
669.	151 <i>Triglochin striata</i>			
670.	152 <i>Triglochin trichophora</i>			
671.	98 <i>Typha domingensis</i> (Bulrush, Djandjidi)			
672.	15725 <i>Verbesina encelioides</i>	Y		
673.	7666 <i>Verreauxia reinwardtii</i> (Common Verreauxia)			
674.	12402 <i>Verticordia chrysanthella</i>			
675.	48829 <i>Wahlenbergia capillaris</i>			
676.	13331 <i>Waitzia acuminata</i> var. <i>acuminata</i>			
677.	13330 <i>Waitzia acuminata</i> var. <i>albicans</i>			
678.	13328 <i>Waitzia nitida</i>			
679.	8281 <i>Waitzia podolepis</i>			
680.	8282 <i>Waitzia suaveolens</i> (Fragrant Waitzia)			
681.	6658 <i>Wilsonia backhousei</i> (Narrow-leaf Wilsonia)			
682.	6659 <i>Wilsonia humilis</i> (Silky Wilsonia)			
683.	12072 <i>Wurmbea dioica</i> subsp. <i>alba</i>			
684.	1398 <i>Wurmbea monantha</i>			
685.	1256 <i>Xanthorrhoea preissii</i> (Grass tree, Palga)			

Conservation Codes

T - Rare or likely to become extinct
X - Presumed extinct
IA - Protected under international agreement
S - Other specially protected fauna
1 - Priority 1
2 - Priority 2
3 - Priority 3
4 - Priority 4
5 - Priority 5

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.

Appendix D – Flora data

Flora species list

Quadrat data

Flora likelihood of occurrence

Flora Species List

Family	Taxon	Status
Amaranthaceae	<i>Ptilotus divaricatus</i>	
Amaranthaceae	<i>Ptilotus stirlingii</i> sp. <i>stirlingii</i>	
Apocynaceae	<i>Alyxia buxifolia</i>	
Asparagaceae	<i>Acanthocarpus canaliculatus</i>	
Asparagaceae	<i>Acanthocarpus preissii</i>	
Asparagaceae	<i>Lomandra maritima</i>	
Asparagaceae	<i>Thysanotus ?manglesianus</i>	
Asteraceae	<i>Hypochaeris glabra</i>	*
Asteraceae	<i>Helianthus annuus</i>	*
Asteraceae	<i>Reichardia tingitana</i>	*
Asteraceae	<i>Sonchus oleraceus</i>	*
Asteraceae	<i>Asteraceae</i> sp (insufficient material)	
Asteraceae	<i>Austrostipa nitida</i>	
Asteraceae	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	
Azioaceae	<i>Mesembryanthemum crystallinum</i>	*
Azioaceae	<i>Tetragonia implexicoma</i>	
Brassicaceae	<i>Brassica tournefortii</i>	*
Brassicaceae	<i>Sisymbrium orientale</i>	*
Chenopodiaceae	<i>Atriplex cinerea</i>	
Chenopodiaceae	<i>Enchylaena tomentosa</i>	
Chenopodiaceae	<i>Rhagodia latifolia</i> ssp. <i>latifolia</i>	
Chenopodiaceae	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	
Chenopodiaceae	<i>Salsola australis</i>	
Chenopodiaceae	<i>Threlkeldia diffusa</i>	
Convolvulaceae	* <i>Cuscuta epithymum</i>	*
Euphorbiaceae	<i>Euphorbia boophthona</i>	
Euphorbiaceae	<i>Euphorbia terracina</i>	*
Fabaceae	<i>Acacia rostellifera</i>	
Fabaceae	<i>Acacia saligna</i>	
Fabaceae	<i>Acacia sclerosperma</i> subsp. <i>sclerosperma</i>	
Fabaceae	<i>Glycine canescens</i>	
Fabaceae	<i>Templetonia retusa</i>	
Frankeniaceae	<i>Frankenia tingitana</i>	
Goodeniaceae	<i>Scaevola tomentosa</i>	
Lauraceae	<i>Cassytha aurea</i> var. <i>aurea</i>	
Loranthaceae	<i>Amyema preissii</i>	
Malvaceae	<i>Alyogyne hakeifolia</i>	
Malvaceae	<i>Commersonia boerhavia</i>	
Myrtaceae	<i>Eucalyptus baudiniana</i>	
Myrtaceae	<i>Eucalyptus utilis</i> (planted non-local)	
Myrtaceae	<i>Melaleuca cardiophylla</i>	
Nyctaginaceae	<i>Commicarpus australis</i>	

Phyllanthaceae	<i>Phyllanthus calycinus</i>	
Pittosporaceae	<i>Pittosporum ligustrifolium</i>	
Poaceae	<i>Avena barbata</i>	*
Poaceae	<i>Cenchrus ciliatus</i>	*
Poaceae	<i>Brachypodium distachyon</i>	*
Poaceae	<i>Bromus diandrus</i>	*
Poaceae	<i>Ehrharta longiflora</i>	*
Poaceae	<i>Ehrharta brevifolia</i>	*
Poaceae	<i>Aristida</i> sp (insufficient material)	
Poaceae	<i>Austrostipa elegantissima</i>	
Poaceae	<i>Austrostipa nitida</i>	
Poaceae	<i>Poaceae</i> sp. (insufficient material)	
Poaceae	<i>Sporobolus virginicus</i>	
Proteaceae	<i>Grevillea argyrophylla</i>	
Scrophulariaceae	<i>Myoporum insulare</i>	
Solanaceae	<i>Anthocercis littorea</i>	
Surianaceae	<i>Stylobasium spathulatum</i>	
Thymelaeaceae	<i>Pimelea gilgiana</i>	
Thymelaeaceae	<i>Pimelea microcephala</i> subsp <i>microcephala</i>	
Thymelaeaceae	<i>Pimelea gilgiana</i>	
Zygophyllaceae	<i>Roepera apiculata</i>	
Zygophyllaceae	<i>Roepera fruticulosa</i>	

* Denoted an introduced species

Flora species by site matrix (Site:Lyn_X)

Taxon	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
<i>*Avena barbata</i>	1				1		1	1			1		1		1		1	1	1	1	1	1	1	1		1	
<i>*Brachypodium distachyon</i>	1														1	1			1								
<i>*Brassica tournefortii</i>	1				1		1	1	1	1				1					1	1					1	1	
<i>*Bromus diandrus</i>												1															
<i>*Bromus diandrus</i>	1		1	1	1					1	1																
<i>*Cuscuta epithymum</i>					1						1																
<i>*Ehrharta longiflora</i>							1	1	1	1		1		1			1	1	1	1	1	1	1	1	1	1	1
<i>*Hypochaeris glabra</i>																					1						
<i>*Reichardia tingitana</i>					1																						
<i>*Sisymbrium orientale</i>		1	1	1	1																						
<i>*Sonchus oleraceus</i>				1			1	1				1								1						1	
<i>Acacia rostellifera</i>	1	1	1	1	1	1	1	2	1	1			1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
<i>Acacia saligna</i>										1																	
<i>Acacia sclerosperma subsp. sclerosperma</i>	1	1	1		2							1															
<i>Acanthocarpus canaliculatus</i>		1																									

<i>Acanthocarpus preissii</i>		1												1	1										
<i>Alyogyne hakeifolia</i>	1	1		1		1						1	1	1	1	1	1			1	1	1	1	1	
<i>Alyxia buxifolia</i>									1																
<i>Anthocercis littorea</i>																						1			
<i>Aristida sp (insufficient material)</i>	1				1					1															1
<i>Asteraceae sp (insufficient material)</i>	1	1	1	1														1						1	
<i>Austrostipa nitida</i>						1																			
<i>Austrostipa elegantissima</i>	1		1	1	1	1	1	1	1		1		1		1	1		1	1	2	1		1	1	1
<i>Austrostipa nitida</i>		1	1																						
<i>Cassythia aurea var. aurea</i>			1			1	1												1				1		
<i>Commicarpus australis</i>	1	1		1	1		1	1	1				1	1	1	1	1		1	1	1			1	1
<i>Enchylaena tomentosa</i>		1					1			1		1													
<i>Eucalyptus baudiniana</i>															1										
<i>Euphorbia boophthona</i>	1			1	1														1						1
<i>Frankenia tingitana</i>											1														
<i>Glycine canescens</i>	1				1		1																		

<i>Grevillea argyrophylla</i>											1																	
<i>Lomandra maritima</i>		1																										
<i>Melaleuca cardiophylla</i>	1	1	1							2	1		1		1	1												
<i>Myoporum insulare</i>												1																
<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	1	1	1	1	1		1							1	1	1		1	1		1			1	1	1		
<i>Phyllanthus calycinus</i>																									1			
<i>Pimelea gilgiana</i>	1	1	1							1					1			1							1			
<i>Pimelea microcephala</i> subsp <i>microcephala</i>	1	1	1	1	1	1	1	2	2	1	1		1	1	1	1	1	1	1	1	1	1	1		1	1	1	
<i>Pimelea gilgiana</i>																1												
<i>Pittosporum ligustrifolium</i>																									1			
<i>Poaceae</i> sp. (insufficient material)												1																
<i>Ptilotus divaricatus</i>										1	1				1	1	1									1		
<i>Ptilotus stirlingii</i> sp. <i>stirlingii</i>	1																											
<i>Rhagodia latifolia</i> ssp. <i>latifolia</i>																		1	1									

<i>Rhagodia preissii</i> subsp. <i>obovata</i>	1	1	2	1	1	1	1	1	1	3	2		1	1	1		1	1	2	1	1		1	1	1	1	1
<i>Roepera apiculata</i>										1			1					1									
<i>Roepera fruticulosa</i>	1	1	1	1	1	1	1		1	1	1		1	1	1	1	1	1	1	1			1			1	1
<i>Salsola australis</i>																								1	1		
<i>Scaevola tomentosa</i>													1														
<i>Sporobolus virginicus</i>												1															
<i>Stylobasium spathulatum</i>			1	1	1	1								1	1	1		1	1		1				1		
<i>Templetonia retusa</i>	1	1																									
<i>Tetragonia implexicoma</i>			1		1	1	1				1			1	1	1	1			1						1	1
<i>Threlkeldia diffusa</i>											1	1													1		
<i>Thysanotus ?manglesianus</i>		1																1	1								

Flora site raw data

Site number	Taxon	Cover (%)	Height (m)	Form/Stratum	Opportunistic
Lyn_01	<i>*Avena barbata</i>	0.5	0.25	Other grass (G)	
Lyn_01	<i>*Brachypodium distachyon</i>	2	0.1	Other grass (G)	
Lyn_01	<i>*Brassica tournefortii</i>	0.1	0.1	Forb (G)	
Lyn_01	<i>*Bromus diandrus</i>	50	0.1	Other grass (G)	
Lyn_01	<i>Acacia rostellifera</i>	2	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Acacia sclerosperma subsp. sclerosperma</i>	0.5	0.25	Forb (G)	
Lyn_01	<i>Alyogyne hakeifolia</i>	3	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Aristida sp (insufficient material)</i>	0.5	0.25	Other grass (G)	
Lyn_01	<i>Asteraceae sp (insufficient material)</i>	0.1	0.1	Forb (G)	
Lyn_01	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_01	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Euphorbia boophthona</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Glycine canescens</i>	0.5	0.25	Vine (G)	
Lyn_01	<i>Melaleuca cardiophylla</i>	10	1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Olearia sp. Kennedy Range (G Byrne 66)</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_01	<i>Pimelea microcephala subsp microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_01	<i>Ptilotus stirlingii sp. stirlingii</i>	0.1	0.25	Shrub, cycad, grass-tree, tree-fern (M)	

Lyn_01	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	2	1.25	Chenopod shrub (M)	
Lyn_01	<i>Roepera fruticulosa</i>	4	0.5	Vine (G)	
Lyn_01	<i>Templetonia retusa</i>	1	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	* <i>Sisymbrium orientale</i>	0.1	0.1	Forb (G)	
Lyn_02	<i>Acacia rostellifera</i>	5	4	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Acacia sclerosperma</i> <i>subsp. sclerosperma</i>	0.5	0.25	Forb (G)	
Lyn_02	<i>Acanthocarpus</i> <i>canaliculatus</i>	20	0.1	Other grass (G)	
Lyn_02	<i>Acanthocarpus preissii</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Alyogyne hakeifolia</i>	3	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Asteraceae sp</i> <i>(insufficient material)</i>	0.1	0.1	Forb (G)	
Lyn_02	<i>Austrostipa nitida</i>	0.1	0.25	Other grass (G)	
Lyn_02	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Enchylaena tomentosa</i>	0.1	0.25	Chenopod shrub (M)	
Lyn_02	<i>Lomandra maritima</i>	0.1	0.25	Forb (G)	
Lyn_02	<i>Melaleuca cardiophylla</i>	1	3.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Olearia sp. Kennedy</i> <i>Range (G Byrne 66)</i>	4	0.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_02	<i>Pimelea microcephala</i> <i>subsp. microcephala</i>	5	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	2	1.25	Chenopod shrub (M)	
Lyn_02	<i>Roepera fruticulosa</i>	4	0.5	Vine (G)	

Lyn_02	<i>Templetonia retusa</i>	10	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_02	<i>Thysanotus ?manglesianus</i>	0.1	0.25	Forb (G)	
Lyn_03	* <i>Bromus diandrus</i>	1	0.1	Other grass (G)	
Lyn_03	* <i>Sisymbrium orientale</i>	0.1	0.1	Forb (G)	
Lyn_03	<i>Acacia rostellifera</i>	1	1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_03	<i>Acacia sclerosperma subsp. sclerosperma</i>	0.5	0.5	Forb (G)	
Lyn_03	<i>Asteraceae sp (insufficient material)</i>	0.1	0.1	Forb (G)	
Lyn_03	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_03	<i>Austrostipa nitida</i>	0.5	0.25	Other grass (G)	
Lyn_03	<i>Cassytha aurea var. aurea</i>	0.5	0.25	Vine (G)	
Lyn_03	<i>Melaleuca cardiophylla</i>	50	3.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_03	<i>Olearia sp. Kennedy Range (G Byrne 66)</i>	4	0.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_03	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_03	<i>Pimelea microcephala subsp microcephala</i>	2	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_03	<i>Rhagodia preissii subsp. obovata</i>	2	0.25	Chenopod shrub (M)	
Lyn_03	<i>Rhagodia preissii subsp. obovata</i>	0.1	0.75	Chenopod shrub (M)	
Lyn_03	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_03	<i>Stylobasium spathulatum</i>	2	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_03	<i>Tetragonia implexicoma</i>	0.5	0.25	Forb (G)	
Lyn_04	* <i>Bromus diandrus</i>	25	0.1	Other grass (G)	

Lyn_04	<i>*Sisymbrium orientale</i>	5	0.1	Forb (G)	
Lyn_04	<i>*Sonchus oleraceus</i>	0.1	0.1	Forb (G)	
Lyn_04	<i>Acacia rostellifera</i>	1	4	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Alyogyne hakeifolia</i>	2	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Asteraceae sp (insufficient material)</i>	0.1	0.1	Forb (G)	
Lyn_04	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_04	<i>Commicarpus australis</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Euphorbia boophthona</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Olearia sp. Kennedy Range (G Byrne 66)</i>	4	0.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Pimelea microcephala subsp microcephala</i>	5	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_04	<i>Rhagodia preissii subsp. obovata</i>	5	1	Chenopod shrub (M)	
Lyn_04	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_04	<i>Stylobasium spathulatum</i>	5	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>*Avena barbata</i>	2	0.25	Other grass (G)	
Lyn_05	<i>*Brassica tournefortii</i>	1	0.25	Forb (G)	
Lyn_05	<i>*Bromus diandrus</i>	20	0.1	Other grass (G)	
Lyn_05	<i>*Cuscuta epithymum</i>	0.1	0.1	Vine (G)	
Lyn_05	<i>*Reichardia tingitana</i>	0.1	0.1	Forb (G)	
Lyn_05	<i>*Sisymbrium orientale</i>	5	0.1	Forb (G)	
Lyn_05	<i>Acacia rostellifera</i>	8	4	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Acacia sclerosperma subsp. sclerosperma</i>	2	0.75	Forb (G)	

Lyn_05	<i>Acacia sclerosperma</i> <i>subsp. sclerosperma</i>		1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Aristida</i> sp (insufficient material)	0.1	0.5	Other grass (G)	
Lyn_05	<i>Austrostipa elegantissima</i>		0.25	Other grass (G)	
Lyn_05	<i>Commicarpus australis</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Euphorbia boophthona</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Glycine canescens</i>	0.1	0.25	Vine (G)	
Lyn_05	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	1	0.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	5	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	5	1	Chenopod shrub (M)	
Lyn_05	<i>Roepera fruticulosa</i>	10	0.5	Vine (G)	
Lyn_05	<i>Stylobasium spathulatum</i>	10	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_05	<i>Tetragonia implexicoma</i>	0.5	0.25	Chenopod shrub (M)	
Lyn_06	<i>Acacia rostellifera</i>	60	6	Tree, palm (U)	
Lyn_06	<i>Alyogyne hakeifolia</i>	5	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_06	<i>Austrostipa nitida</i>	0.5	0.75	Other grass (G)	
Lyn_06	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_06	<i>Cassytha aurea</i> var. <i>aurea</i>	3	0.25	Vine (G)	
Lyn_06	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_06	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	5	1	Chenopod shrub (M)	

Lyn_06	<i>Roepera fruticulosa</i>	40	0.5	Vine (G)	
Lyn_06	<i>Stylobasium spathulatum</i>	3	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_06	<i>Tetragonia implexicoma</i>	2	0.25	Chenopod shrub (M)	
Lyn_07	* <i>Avena barbata</i>	5	0.25	Other grass (G)	
Lyn_07	* <i>Brassica tournefortii</i>	0.5	0.25	Forb (G)	
Lyn_07	* <i>Ehrharta longiflora</i>	55	0.25	Other grass (G)	
Lyn_07	* <i>Sonchus oleraceus</i>	0.1	0.1	Forb (G)	
Lyn_07	<i>Acacia rostellifera</i>	30	8	Tree, palm (U)	
Lyn_07	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_07	<i>Cassutha aurea</i> var. <i>aurea</i>	3	0.25	Vine (G)	
Lyn_07	<i>Commicarpus australis</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_07	<i>Enchylaena tomentosa</i>	0.5	0.25	Chenopod shrub (M)	
Lyn_07	<i>Glycine canescens</i>	0.1	0.25	Vine (G)	
Lyn_07	<i>Olearia</i> sp. <i>Kennedy Range</i> (G Byrne 66)	1	0.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_07	<i>Pimelea microcephala</i> subsp. <i>microcephala</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_07	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	35	1	Chenopod shrub (M)	
Lyn_07	<i>Roepera fruticulosa</i>	40	0.5	Vine (G)	
Lyn_07	<i>Tetragonia implexicoma</i>	2	0.25	Chenopod shrub (M)	
Lyn_08	* <i>Avena barbata</i>	1	0.25	Other grass (G)	
Lyn_08	* <i>Brassica tournefortii</i>	2	0.25	Forb (G)	
Lyn_08	* <i>Ehrharta longiflora</i>	20	0.25	Other grass (G)	
Lyn_08	* <i>Sonchus oleraceus</i>	0.1	0.1	Forb (G)	
Lyn_08	<i>Acacia rostellifera</i>	20	8	Tree, palm (U)	

Lyn_08	<i>Acacia rostellifera</i>	5	3	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_08	<i>Austrostipa elegantissima</i>	1	0.25	Other grass (G)	
Lyn_08	<i>Commicarpus australis</i>	5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_08	<i>Pimelea microcephala subsp microcephala</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_08	<i>Pimelea microcephala subsp microcephala</i>	0.5	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_08	<i>Rhagodia preissii subsp. obovata</i>	60	1	Chenopod shrub (M)	
Lyn_09	* <i>Brassica tournefortii</i>	0.1	0.25	Forb (G)	
Lyn_09	* <i>Ehrharta longiflora</i>	30	0.25	Other grass (G)	
Lyn_09	<i>Acacia rostellifera</i>	30	8	Tree, palm (U)	
Lyn_09	<i>Austrostipa elegantissima</i>	1	0.25	Other grass (G)	
Lyn_09	<i>Commicarpus australis</i>	5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_09	<i>Pimelea microcephala subsp microcephala</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_09	<i>Pimelea microcephala subsp microcephala</i>	0.5	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_09	<i>Rhagodia preissii subsp. obovata</i>	60	1	Chenopod shrub (M)	
Lyn_09	<i>Roepera fruticulosa</i>	5	0.5	Vine (G)	
Lyn_10	* <i>Brassica tournefortii</i>	2	0.25	Forb (G)	
Lyn_10	* <i>Bromus diandrus</i>	1	0.1	Other grass (G)	
Lyn_10	* <i>Ehrharta longiflora</i>	30	0.25	Other grass (G)	
Lyn_10	<i>Acacia rostellifera</i>	1	1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_10	<i>Acacia saligna</i>	0.5	0.75	Shrub, cycad, grass-tree, tree-fern (M)	

Lyn_10	<i>Alyxia buxifolia</i>	1	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_10	<i>Aristida sp (insufficient material)</i>	0.1	0.75	Other grass (G)	
Lyn_10	<i>Enchylaena tomentosa</i>	0.5	0.25	Chenopod shrub (M)	
Lyn_10	<i>Melaleuca cardiophylla</i>	10	3.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_10	<i>Melaleuca cardiophylla</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_10	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_10	<i>Pimelea microcephala subsp microcephala</i>	1	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_10	<i>Ptilotus divaricatus</i>	0.5	0.5	Forb (G)	
Lyn_10	<i>Rhagodia preissii subsp. obovata</i>	25	0.25	Chenopod shrub (M)	
Lyn_10	<i>Rhagodia preissii subsp. obovata</i>	8	0.25	Other grass (G)	
Lyn_10	<i>Rhagodia preissii subsp. obovata</i>	5	1	Chenopod shrub (M)	
Lyn_10	<i>Roepera apiculata</i>	5	0.25	Chenopod shrub (M)	
Lyn_10	<i>Roepera fruticulosa</i>	3	0.5	Vine (G)	
Lyn_11	* <i>Avena barbata</i>	60	0.25	Other grass (G)	
Lyn_11	* <i>Bromus diandrus</i>	5	0.1	Other grass (G)	
Lyn_11	* <i>Cuscuta epithymum</i>	0.1	0.1	Vine (G)	
Lyn_11	<i>Austrostipa elegantissima</i>	0.1	0.25	Other grass (G)	
Lyn_11	<i>Grevillea argyrophylla</i>	0.5	1.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_11	<i>Melaleuca cardiophylla</i>	40	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_11	<i>Pimelea microcephala subsp microcephala</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_11	<i>Ptilotus divaricatus</i>	1	0.5	Forb (G)	

Lyn_11	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	5	0.25	Chenopod shrub (M)	
Lyn_11	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	1	1	Chenopod shrub (M)	
Lyn_11	<i>Roepera fruticulosa</i>	20	0.5	Vine (G)	
Lyn_11	<i>Tetragonia</i> <i>implexica</i>	1	0.5	Chenopod shrub (M)	
Lyn_11	<i>Threlkeldia diffusa</i>	0.5	0.5	Chenopod shrub (M)	
Lyn_12	* <i>Bromus diandrus</i>	2	0.25	Other grass (G)	
Lyn_12	* <i>Ehrharta longiflora</i>	40	0.5	Other grass (G)	
Lyn_12	* <i>Sonchus oleraceus</i>	0.5	0.25	Forb (G)	
Lyn_12	<i>Acacia sclerosperma</i> <i>subsp. Sclerosperma</i>	20	2	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_12	<i>Alyogyne hakeifolia</i>	2	2	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_12	<i>Enchylaena tomentosa</i>	5	0.25	Chenopod shrub (M)	
Lyn_12	<i>Frankenia tingitana</i>	5	0.25	Chenopod shrub (M)	
Lyn_12	<i>Myoporum insulare</i>	45	2	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_12	<i>Poaceae sp.</i> (insufficient material)	5	0.25	Other grass (G)	
Lyn_12	<i>Sporobolus virginicus</i>	10	0.1	Other grass (G)	
Lyn_12	<i>Threlkeldia diffusa</i>	10	0.5	Chenopod shrub (M)	
Lyn_13	* <i>Avena barbata</i>	80	0.25	Other grass (G)	
Lyn_13	<i>Acacia rostellifera</i>	20	2	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_13	<i>Alyogyne hakeifolia</i>	15	2	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_13	<i>Austrostipa</i> <i>elegantissima</i>	0.1	0.25	Other grass (G)	
Lyn_13	<i>Commicarpus australis</i>	2	1	Shrub, cycad, grass- tree, tree-fern (M)	

Lyn_13	<i>Melaleuca cardiophylla</i>	5	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_13	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	5	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_13	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	15	1	Chenopod shrub (M)	
Lyn_13	<i>Roepera apiculata</i>	1	0.25	Chenopod shrub (M)	
Lyn_13	<i>Roepera fruticulosa</i>	20	0.5	Vine (G)	
Lyn_13	<i>Scaevola tomentosa</i>	0.5	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	* <i>Brassica tournefortii</i>	0.1	0.25	Forb (G)	
Lyn_14	* <i>Ehrharta longiflora</i>	0.5	0.25	Other grass (G)	
Lyn_14	<i>Acacia rostellifera</i>	20	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Alyogyne hakeifolia</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Commicarpus australis</i>	1	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Olearia sp. Kennedy Range</i> (G Byrne 66)	1	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	2	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	40	1	Chenopod shrub (M)	
Lyn_14	<i>Roepera fruticulosa</i>	5	0.5	Vine (G)	
Lyn_14	<i>Stylobasium spathulatum</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_14	<i>Tetragonia implexicoma</i>	30	0.5	Chenopod shrub (M)	
Lyn_15	* <i>Avena barbata</i>	0.5	0.25	Other grass (G)	
Lyn_15	* <i>Brachypodium distachyon</i>	20	0.1	Other grass (G)	
Lyn_15	<i>Acacia rostellifera</i>	10	3	Tree, palm (U)	

Lyn_15	<i>Acanthocarpus preissii</i>	0.5	0.25	Forb (G)	
Lyn_15	<i>Alyogyne hakeifolia</i>	3	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_15	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_15	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_15	<i>Melaleuca cardiophylla</i>	10	3	Tree, palm (U)	
Lyn_15	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_15	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_15	<i>Pimelea microcephala</i> subsp. <i>microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_15	<i>Ptilotus divaricatus</i>	0.5	0.25	Forb (G)	
Lyn_15	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	2	1.25	Chenopod shrub (M)	
Lyn_15	<i>Roepera fruticulosa</i>	4	0.5	Vine (G)	
Lyn_15	<i>Stylobasium spathulatum</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_15	<i>Tetragonia implexicoma</i>	10	0.5	Chenopod shrub (M)	
Lyn_16	* <i>Brachypodium distachyon</i>	20	0.1	Other grass (G)	
Lyn_16	<i>Acacia rostellifera</i>	0.5	3	Tree, palm (U)	
Lyn_16	<i>Acanthocarpus preissii</i>	0.5	0.25	Forb (G)	
Lyn_16	<i>Alyogyne hakeifolia</i>	3	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_16	<i>Austrostipa elegantissima</i>	0.5	0.25	Other grass (G)	
Lyn_16	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_16	<i>Eucalyptus baudiniana</i>	0.5	3	Mallee shrub (M)	
Lyn_16	<i>Melaleuca cardiophylla</i>	40	3	Tree, palm (U)	

Lyn_16	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_16	<i>Pimelea microcephala</i> subsp <i>microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_16	<i>Pimelea gilgiana</i>	0.5	0.25	Forb (G)	
Lyn_16	<i>Ptilotus divaricatus</i>	0.5	0.25	Forb (G)	
Lyn_16	<i>Roepera fruticulosa</i>	4	0.5	Vine (G)	
Lyn_16	<i>Stylobasium spathulatum</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_16	<i>Tetragonia implexicoma</i>	10	0.5	Chenopod shrub (M)	
Lyn_17	* <i>Avena barbata</i>	0.5	0.25	Other grass (G)	
Lyn_17	* <i>Ehrharta longiflora</i>	25	0.25	Other grass (G)	
Lyn_17	<i>Acacia rostellifera</i>	50	8	Tree, palm (U)	
Lyn_17	<i>Alyogyne hakeifolia</i>	2	1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_17	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_17	<i>Pimelea microcephala</i> subsp <i>microcephala</i>	0.5	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_17	<i>Ptilotus divaricatus</i>	0.5	0.25	Forb (G)	
Lyn_17	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	45	1	Chenopod shrub (M)	
Lyn_17	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_17	<i>Tetragonia implexicoma</i>	30	0.5	Chenopod shrub (M)	
Lyn_18	* <i>Avena barbata</i>	25	0.25	Other grass (G)	
Lyn_18	* <i>Ehrharta longiflora</i>	65	0.25	Other grass (G)	
Lyn_18	<i>Acacia rostellifera</i>	20	8	Tree, palm (U)	
Lyn_18	<i>Alyogyne hakeifolia</i>	4	1.75	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_18	<i>Austrostipa elegantissima</i>	5	0.25	Other grass (G)	

Lyn_18	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	1	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_18	<i>Pimelea gilgiana</i>	1	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_18	<i>Pimelea microcephala</i> subsp <i>microcephala</i>	5	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_18	<i>Rhagodia latifolia</i> ssp. <i>latifolia</i>	5	1.25	Chenopod shrub (M)	
Lyn_18	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	2	1	Chenopod shrub (M)	
Lyn_18	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_18	<i>Stylobasium spathulatum</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_18	<i>Thysanotus ?manglesianus</i>	0.5	1	Forb (G)	
Lyn_19	* <i>Avena barbata</i>	2	0.25	Other grass (G)	
Lyn_19	* <i>Brachypodium distachyon</i>	1	0.25	Other grass (G)	
Lyn_19	* <i>Brassica tournefortii</i>	0.1	0.25	Forb (G)	
Lyn_19	* <i>Ehrharta longiflora</i>	75	0.25	Other grass (G)	
Lyn_19	<i>Acacia rostellifera</i>	30	8	Tree, palm (U)	
Lyn_19	<i>Asteraceae</i> sp (insufficient material)	0.1	0.1	Forb (G)	
Lyn_19	<i>Austrostipa elegantissima</i>	6	0.25	Other grass (G)	
Lyn_19	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_19	<i>Olearia</i> sp. Kennedy Range (G Byrne 66)	1	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_19	<i>Pimelea microcephala</i> subsp <i>microcephala</i>	5	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_19	<i>Rhagodia latifolia</i> ssp. <i>latifolia</i>	1	1.25	Chenopod shrub (M)	

Lyn_19	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	2	1	Chenopod shrub (M)	
Lyn_19	<i>Rhagodia preissii</i> <i>subsp. obovata</i>				
Lyn_19	<i>Roepera apiculata</i>	0.5	0.1	Chenopod shrub (M)	
Lyn_19	<i>Roepera fruticulosa</i>	5	0.5	Vine (G)	
Lyn_19	<i>Stylobasium</i> <i>spathulatum</i>	5	1.25	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_19	<i>Thysanotus</i> <i>?manglesianus</i>	0.1	1	Forb (G)	
Lyn_20	* <i>Avena barbata</i>	1	0.25	Other grass (G)	
Lyn_20	* <i>Brassica tournefortii</i>	1	0.25	Forb (G)	
Lyn_20	* <i>Ehrharta longiflora</i>	75	0.25	Other grass (G)	
Lyn_20	* <i>Sonchus oleraceus</i>	0.1	0.1	Forb (G)	
Lyn_20	<i>Acacia rostellifera</i>	60	6	Tree, palm (U)	
Lyn_20	<i>Austrostipa</i> <i>elegantissima</i>	6	0.25	Other grass (G)	
Lyn_20	<i>Austrostipa</i> <i>elegantissima</i>	5	0.25	Other grass (G)	
Lyn_20	<i>Cassytha aurea</i> var. <i>aurea</i>	1	1	Vine (G)	
Lyn_20	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_20	<i>Euphorbia boophthona</i>	0.1	0.25	Forb (G)	
Lyn_20	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	5	1.25	Shrub, cycad, grass- tree, tree-fern (M)	
Lyn_20	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	2	1	Chenopod shrub (M)	
Lyn_20	<i>Roepera fruticulosa</i>	5	0.5	Vine (G)	
Lyn_20	<i>Tetragonia</i> <i>implexicoma</i>	10	0.5	Chenopod shrub (M)	
Lyn_21	* <i>Avena barbata</i>	70	0.25	Other grass (G)	
Lyn_21	* <i>Ehrharta longiflora</i>	10	0.25	Other grass (G)	


Lyn_21	<i>*Hypochaeris glabra</i>	2	0.1	Forb (G)	
Lyn_21	<i>Acacia rostellifera</i>	30	3	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_21	<i>Alyogyne hakeifolia</i>	30	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_21	<i>Austrostipa elegantissima</i>	2	0.25	Other grass (G)	
Lyn_21	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_21	<i>Olearia sp. Kennedy Range (G Byrne 66)</i>	1	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_21	<i>Pimelea microcephala subsp microcephala</i>	1	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_21	<i>Rhagodia preissii subsp. obovata</i>	2	1	Chenopod shrub (M)	
Lyn_21	<i>Stylobasium spathulatum</i>	2	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_22	<i>*Avena barbata</i>	40	0.25	Other grass (G)	
Lyn_22	<i>*Ehrharta longiflora</i>	30	0.25	Other grass (G)	
Lyn_22	<i>Acacia rostellifera</i>	20	7	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_22	<i>Alyogyne hakeifolia</i>	5	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_22	<i>Pimelea microcephala subsp microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_23	<i>*Avena barbata</i>	5	0.25	Other grass (G)	
Lyn_23	<i>*Ehrharta longiflora</i>	5	0.25	Other grass (G)	
Lyn_23	<i>Acacia rostellifera</i>	40	7	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_23	<i>Alyogyne hakeifolia</i>	2	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_23	<i>Austrostipa elegantissima</i>	2	0.25	Other grass (G)	


Lyn_23	<i>Pimelea microcephala</i> <i>subsp. microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_23	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	10	1	Chenopod shrub (M)	
Lyn_23	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_24	* <i>Avena barbata</i>	85	0.25	Other grass (G)	
Lyn_24	* <i>Ehrharta longiflora</i>	5	0.25	Other grass (G)	
Lyn_24	<i>Acacia rostellifera</i>	30	4	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_24	<i>Alyogyne hakeifolia</i>	2	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_24	<i>Olearia sp. Kennedy Range</i> (G Byrne 66)	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_24	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	2	1	Chenopod shrub (M)	
Lyn_24	<i>Salsola australis</i>	0.5	0.5	Chenopod shrub (M)	
Lyn_25	* <i>Brassica tournefortii</i>	0.1	0.1	Forb (G)	
Lyn_25	* <i>Ehrharta longiflora</i>	1	0.25	Other grass (G)	
Lyn_25	<i>Acacia rostellifera</i>	30	4	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Alyogyne hakeifolia</i>	2	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Anthocercis littorea</i>	0.5	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Asteraceae sp</i> (insufficient material)	0.1	0.1	Forb (G)	
Lyn_25	<i>Austrostipa elegantissima</i>	2	0.25	Other grass (G)	
Lyn_25	<i>Cassytha aurea</i> var. <i>aurea</i>	4	1.5	Vine (G)	
Lyn_25	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Olearia sp. Kennedy Range</i> (G Byrne 66)	5	1	Shrub, cycad, grass-tree, tree-fern (M)	


Lyn_25	<i>Phyllanthus calycinus</i>	0.5	0.5	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Pimelea gilgiana</i>	0.5	0.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Pittosporum ligustrifolium</i>	1	1.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	5	1.25	Chenopod shrub (M)	
Lyn_25	<i>Salsola australis</i>	0.5	0.5	Chenopod shrub (M)	
Lyn_25	<i>Stylobasium spathulatum</i>	1	2.25	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_25	<i>Threlkeldia diffusa</i>	2	0.25	Chenopod shrub (M)	
Lyn_26	* <i>Avena barbata</i>	30	0.25	Other grass (G)	
Lyn_26	* <i>Brassica tournefortii</i>	0.1	0.1	Forb (G)	
Lyn_26	* <i>Ehrharta longiflora</i>	35	0.25	Other grass (G)	
Lyn_26	* <i>Sonchus oleraceus</i>	0.1	0.1	Forb (G)	
Lyn_26	<i>Acacia rostellifera</i>	10	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_26	<i>Austrostipa elegantissima</i>	2	0.25	Other grass (G)	
Lyn_26	<i>Commicarpus australis</i>	0.5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_26	<i>Olearia sp. Kennedy Range</i> (G Byrne 66)	10	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_26	<i>Pimelea microcephala</i> <i>subsp microcephala</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_26	<i>Ptilotus divaricatus</i>	0.1	0.5	Forb (G)	
Lyn_26	<i>Rhagodia preissii</i> <i>subsp. obovata</i>	20	1	Chenopod shrub (M)	
Lyn_26	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	


Lyn_26	<i>Tetragonia implexicoma</i>	20	1.25	Chenopod shrub (M)	
Lyn_27	* <i>Ehrharta longiflora</i>	75	0.25	Other grass (G)	
Lyn_27	<i>Acacia rostellifera</i>	60	2	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_27	<i>Aristida sp (insufficient material)</i>	0.1	0.25	Other grass (G)	
Lyn_27	<i>Austrostipa elegantissima</i>	8	0.25	Other grass (G)	
Lyn_27	<i>Commicarpus australis</i>	2	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_27	<i>Euphorbia boophthona</i>	0.5	0.25	Forb (G)	
Lyn_27	<i>Pimelea microcephala subsp microcephala</i>	5	1	Shrub, cycad, grass-tree, tree-fern (M)	
Lyn_27	<i>Rhagodia preissii subsp. obovata</i>	10	1.25	Chenopod shrub (M)	
Lyn_27	<i>Roepera fruticulosa</i>	1	0.5	Vine (G)	
Lyn_27	<i>Tetragonia implexicoma</i>	20	1.25	Chenopod shrub (M)	
	* <i>Helianthus annuus</i>				x
	* <i>Cenchrus ciliatus</i>				x
	* <i>Ehrharta brevifolia</i>				x
	* <i>Euphorbia terracina</i>				x
	* <i>Mesembryanthemum crystallinum</i>				x
	<i>Amyema preissii</i>				x
	<i>Atriplex cinerea</i>				x
	<i>Commersonia boerhavia</i>				x
	<i>Eucalyptus utilis (planted non-local)</i>				x

Quadrat and point data

Site		Lyn_01	
Type: Quadrat		Size: 10 x 10	
Location: E 229786.45, N 6884508.45			
Vegetation Type: VT02			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Brown/white sand			
Vegetation Condition: n/a			
Disturbances :weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 11-30%		

Site		Lyn_02	
Type: Quadrat		Size: 10 x 10	
Location: E 229700.52 N 6884617.32			
Vegetation Type: VT02			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Brown/white sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 31-70%		

Site		Lyn_03	
Type: Quadrat		Size: 10 x 10	
Location: E 229640.3 N 6884668.98			
Vegetation Type: VT02			
Landform: Hillslope/moderate	Drainage: Good		
Soil Colour & Type: Pale/red brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 31-70%	Litter: 11-30%		
			


Site		Lyn_04	
Type: Quadrat		Size: 10 x 10	
Location: E 229410.34 N 6884757.73			
Vegetation Type: VT01			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 31-70%		


Site		Lyn_05	
Type: Quadrat		Size: 10 x 10	
Location: E 229204.34 N 6885011.92			
Vegetation Type: VT01			
Landform: Hillslope/moderate	Drainage: Good		
Soil Colour & Type: Pale brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 31-70%	Litter: 31-70%		



Site		Lyn_06	
Type: Quadrat		Size: 10 x 10	
Location: E 229015.06 N 6884757.82			
Vegetation Type: VT01			
Landform: Sandplain /gentle	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: <2%	Litter: 11—30%		



Site		Lyn_07	
Type: Quadrat		Size: 10 x 10	
Location: E 228795.5 N 6885075.98			
Vegetation Type: VT01			
Landform: Sandplain /gentle	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: <2%	Litter: 11-30%		


Site		Lyn_08	
Type: Quadrat		Size: 10 x 10	
Location: E 228600.3 N 6885391.62			
Vegetation Type: VT01			
Landform: Sandplain /gentle	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: <2%	Litter: 11-30%		


Site		Lyn_09	
Type: Quadrat		Size: 10 x 10	
Location: E 228096.99 N 6885852.3			
Vegetation Type: VT01			
Landform: Sandplain /negligible		Drainage: Good	
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: <2%		Litter: 31-70%	




Site		Lyn_10	
Type: Quadrat		Size: 10 x 10	
Location: E 227994.53 N 6886401.6			
Vegetation Type: VT02			
Landform: Foothlope /moderate		Drainage: Good	
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%		Litter: 2-10%	





Site		Lyn_11	
Type: Quadrat		Size: 10 x 10	
Location: E 229040.02 N 6885353.38			
Vegetation Type: VT02			
Landform: Boulders/rockpile /moderate	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 11-30%		


Site		Lyn_12	
Type: Quadrat		Size: 10 x 10	
Location: E 227639.13 N 6886366.05			
Vegetation Type: VT03			
Landform: Drainage area/floodplain/ negligible	Drainage: Seasonally wet		
Soil Colour & Type: Light brown clay			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 2-10%		

Site		Lyn_13	
Type: Releve		Size: 10 x 10	
Location: E 227241.68 N 6886969.47			
Vegetation Type: VT01			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 11-30%		


Site		Lyn_14	
Type: Releve		Size: 10 x 10	
Location: E 229622.99 N 6884218.17			
Vegetation Type: VT01			
Landform: Sandplain /gentle	Drainage: Good		
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 31-70%		


Site		Lyn_15	
Type: Releve		Size: 10 x 10	
Location: E 229870.34 N 6884378.17			
Vegetation Type: VT02			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Brown/white sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 31-70%	Litter: 11-30%		

Site		Lyn_16	
Type: Releve		Size: 10 x 10	
Location: E 230131.9 N 6884144.79			
Vegetation Type: VT02			
Landform: Hillslope /moderate	Drainage: Good		
Soil Colour & Type: Brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 31-70%		

Site		Lyn_17	
Type: Quadrat		Size: 10 x 10	
Location: E 231034.5 N 6882513.78			
Vegetation Type: VT01			
Landform: Sandplain /negligible	Drainage: Good		
Soil Colour & Type: Brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 31-70%		

Site		Lyn_18	
Type: Quadrat		Size: 10 x 10	
Location: E 231264.75 N 6882595.25			
Vegetation Type: VT01			
Landform: Foothlope /gentle	Drainage: Good		
Soil Colour & Type: Brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 11-30%		

Site		Lyn_19	
Type: Quadrat		Size: 10 x 10	
Location: E 231529.36 N 6882483.33			
Vegetation Type: VT01			
Landform: Foothill /gentle	Drainage: Good		
Soil Colour & Type: Brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 11-30%		


Site		Lyn_20	
Type: Quadrat		Size: 10 x 10	
Location: E 231844.61 N 6881671.02			
Vegetation Type: VT01			
Landform: Hillcrest/Upper Hillslope	Drainage: Good		
Soil Colour & Type: Brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 2-10%		


Site		Lyn_21	
Type: Quadrat		Size: 10 x 10	
Location: E 232238.85 N 6881343.07			
Vegetation Type: Rehabilitation area			
Landform: Hillslope/moderate		Drainage: Good	
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%		Litter: 11-30%	




Site		Lyn_22	
Type: Releve		Size: 10 x 10	
Location: E 232528.44 N 6880732.71			
Vegetation Type: VT01			
Landform: Hillcrest/upper Hillslope		Drainage: Good	
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%		Litter: 2-10%	




Site		Lyn_23	
Type: Releve		Size: 10 x 10	
Location: E 232917.63 N 6880203.95			
Vegetation Type: VT01			
Landform: Hillcrest/upper Hillslope /negligible	Drainage: Good		
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances: weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 31-70%	Litter: 11-30%		

Site		Lyn_24	
Type: Releve		Size: 10 x 10	
Location: E 232806.62 N 6880438.81			
Vegetation Type: Rehabilitation areas			
Landform: Flat/ Negligible	Drainage: Good		
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 2-10%	Litter: 2-10%		

Site		Lyn_25	
Type: Quadrat		Size: 10 x 10	
Location: E 232681.45 N 6880449.53			
Vegetation Type: Rehabilitation areas			
Landform: Flat /Negligible	Drainage: Good		
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 11-30%		

Site		Lyn_26	
Type: Quadrat		Size: 10 x 10	
Location: E 232038.67 N 6881377.85			
Vegetation Type: VT01			
Landform: Flat /Negligible	Drainage: Good		
Soil Colour & Type: Light brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing/clearing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: 11-30%	Litter: 11-30%		

Site		Lyn_27	
Type: Quadrat		Size: 10 x 10	
Location: E 230656.58 N 6882650.17			
Vegetation Type: VT01			
Landform: Flat /Gentle		Drainage: Good	
Soil Colour & Type: Dark brown sand			
Vegetation Condition: n/a			
Disturbances:weeds/grazing			
Fire Age and Intensity: Old 6+yr			
Bare Ground: <2%		Litter: 31-70%	

Flora likelihood of occurrence assessment guidelines

Likelihood of occurrence	Guideline
Known	Species recorded within study area from field project results (none as this is a desktop search only).
Likely	Species previously recorded within 2 km and large areas of suitable habitat occur in the survey area.
Possible	Species previously recorded within 10 km and areas of suitable habitat occur/may occur in the survey area.
Unlikely	Species previously recorded within 20 km, or suitable habitat does not occur in the survey area.
Highly unlikely	Species not previously recorded within 20 km, suitable habitat does not occur in the survey area and/or the survey area is outside the natural distribution of the species.
Other considerations	Date of known records, cryptic nature of species, anecdotal evidence from previous studies/surveys

Definitions

Term	Description
Study area	A 10 km buffer around the survey area
Survey area	The potential project footprint
Cr	Critically endangered
En	Endangered
T	Threatened
Vu	Vulnerable
P1 – P4	Priority 1 – Priority 4
EPBC Act	Environmental Protection and Biodiversity Conservation Act 1999
DBCA	Department of Biodiversity and Conservation Attractions 2018. WA Government, Department of Parks and Wildlife Threatened (Declared Rare) and Priority Flora List
BC Act	Biodiversity Conservation Act 2016

Flora likelihood of occurrence assessment of conservation significant flora identified in the desktop assessment as potentially occurring within 10 km of the survey area

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Amaranthaceae	<i>Ptilotus chortophytus</i>		P1	Erect perennial herb to 0.15 m high. Flowers yellow. Hillside. Kockatea, breakaway, quartz and shale. The nearest record is approximately 7 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Asteraceae	<i>Balladonia aevroides</i>		P3	Annual herb. Calcareous sand or sandy loam. Flowers August to October. The nearest record is approximately 3 km west of the survey area.	Possible – there is suitable habitat within the survey area.
Asteraceae	<i>Ozothamnus vespertinus</i>		P1	Perennial small shrub, 0.6 m high x 0.5 m wide. White flowers in globular head. Edge of breakaway. Mudstone/shale gravel. Fine white loam/clay. Closest known record is approximately 8 km south-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Colchicaceae	<i>Wurmbea tubulosa</i>	En	T	Cormous, perennial, herb, 0.01-0.03 m high, dioecious or sometimes andromonoecious. Fl. white-pink, Jun to Aug. Clay, loam. River banks, seasonally-wet places. The nearest record is approximately 8 km south.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Dasypogonaceae	<i>Calectasia browneana</i>		P2	Spreading, caespitose perennial, herb, 0.2-0.5 m high, to 0.4 m wide. Flowers blue-purple, Jun to August. White-grey sand, laterite. Adjacent to wet areas of creekline. The nearest recorded in 9 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Ericaceae	<i>Leucopogon</i> sp. Port Gregory (C. Page 33)		P1	Erect shrub to 0.3 m and 0.5 m wide. Flowers white. Shale breakaway. Wet red brown soil on	Highly Unlikely – there is no suitable habitat for this species

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
				shale. The nearest know record approximately 8 km east and south-east of the survey area.	within the survey area.
Ericaceae	<i>Styphelia cernua</i> (previously known as <i>Astroloma</i> sp. Kalbarri (D. & B. Bellairs 1368)		P2	Shrub up to 1.7 m with white flowers, green young fruit. Yellow sandplain, undulating plain, white/grey sand, dunes. Closest record approximately 10 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Euphorbiaceae	<i>Beyeria cinerea</i> subsp. <i>cinerea</i>		P3	Shrub to 1 m high. Flowers yellow. Limestone ridge. Dry, rocky brown sand over limestone, grey sands. The nearest record is approximately 10 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Euphorbiaceae	<i>Stachystemon nematophorus</i>	Vu	P4	Woody, dense shrub, to 1.2 m high. Dry sand, sandy gravel over laterite, sandstone. Exposed rocky sites, disturbed ground. The nearest recorded is approximately 8 km east and south-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Fabaceae	<i>Acacia latipes</i> subsp. <i>licina</i>		P3	Pungent shrub, 0.4-1.2 m high. White sand, granitic soils. Limestone hills, sandplains. Flowers yellow, June to September. The nearest record is approximately 200 m east of the survey area.	Unlikely – suitable survey effort did not record this species
Fabaceae	<i>Acacia pelophila</i>		P1	Dense, spreading shrub, 0.9-2 m high. Clay. Saline creeklines. Flowers yellow, July to August. The nearest recorded is approximately 8 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Fabaceae	<i>Acacia ridleyana</i>		P3	Spreading, sprawling shrub, 0.2-0.9 m high, 0.5-2 m wide. Grey or yellow/brown sand, gravelly clay, granitic loam. Flowers yellow, August to December. The closest record is approximately 10 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Fabaceae	<i>Gastrolobium propinquum</i>		P3	Low, bushy shrub, to 1(-1.8) m high. Flowers orange & yellow & red, June to September. Clay, clay-loam or sandy clay soils, granite, shale. Hills, flats, drainage lines, winter-wet areas. The nearest record is 8 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Frankeniaceae	<i>Frankenia confusa</i>		P4	Low, diffuse shrub, to 0.75 m high, to 0.75 wide. Wet pale brown sand, brown clay, grey soil. Banks of rivers & waterholes, river floodplains. Flowers pink, September. The nearest record is approximately 3 km west of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Goodeniaceae	<i>Scaevola kallophylla</i>		P4	Erect, compact shrub, to 1 m high. Sandy soils over limestone. Coastal plain. Flowers white, May or August to December. The nearest record is located approximately 3 km west of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Goodeniaceae	<i>Scaevola oldfieldii</i>		P3	Erect shrub to 1.5 m and 1.2 m wide. Flowers white. Sandplains, grey sand, brown gravelly loam. The nearest record is 8 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Hydatellaceae	<i>Trithuria australis</i>		P4	Small annual aquatic herb. Seasonally wet poorly drained flat, edge of wetlands, along drying margins, grey and black clayey soils.	Highly Unlikely – there is no suitable habitat for this species

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
				The nearest record is approximately 7.5 km south of the survey area.	within the survey area.
Lamiaceae	<i>Hemigenia pimelifolia</i>		P2	Shrub, 0.2-1 m high. Flowers blue-purple/violet, July to October. Gravelly soils. The nearest record is approximately 7 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Lamiaceae	<i>Prostanthera scutata</i>		P2	Erect, compact shrub, 0.2 – 0.3 m high. Flowers blue-violet, October or December to January. Gravelly sand. The nearest record is approximately 7 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Lamiaceae	<i>Teucrium</i> sp. Hutt River (W.H. Butler 54)		P1	No available information. Only one known record of this species (from 1964), located approximately 7 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Malvaceae	<i>Androcalva bivillosa</i>	Cr	T	Low spreading shrub. Occur on flats and slopes, reddish-brown or yellow sand with lateritic gravel. Road verge lateritic gravel and orange brown clayey sand. Recent soil disturbance. White flowers July to October. Fruit present during late October to December. The closest known record is approximately 38 km south-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Malvaceae	<i>Androcalva microphylla</i>		P2	Spreading recumbent dwarf shrub 30 cm x 100 cm. White flowers. White grey sand over sandstone. The nearest record is 10 km north of the survey area.	Highly Unlikely – there is no suitable habitat for this species

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
					within the survey area.
Malvaceae	<i>Guichenotia quasicalva</i>		P2	Erect, compact shrub, to 0.5 m high. Flowers blue-purple, September to October. Sandy clay over laterite. Drainage line. The nearest recorded in 7 km south-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Malvaceae	<i>Lasiopetalum oldfieldii</i>		P3	Shrub, 0.2-0.8 m high. Sandy soils. Flowers pink, August to October. The nearest record is less than 2 km west of the survey area.	Unlikely – there is suitable habitat within the survey area, however, suitable survey effort did not record this species
Menyanthaceae	<i>Liparophyllum congestiflorum</i>		P4	Small annual herb to 20 cm, yellow petals, green sepals. Occurs in winter wet low lying area, low plain, grey sand over sandstone. The nearest record is approximately 7 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Myrtaceae	<i>Calytrix harvestiana</i>		P2	Shrub, 0.3-0.7 m high. White or yellow sand. Flats. Flowers purple-pink/violet, September to December. Sandplain, yellow sand. Sandstone, brown sand. The nearest record is 4 km south of the survey area.	Unlikely – there is suitable habitat within the survey area, however, suitable survey effort did not record this species
Myrtaceae	<i>Calytrix pimeleoides</i>		P3	Erect, perennial shrub 1 m high x .5 m wide. Flowers yellow. Ridge. Dry, gravelly yellow-brown	Highly Unlikely – there is no suitable habitat

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
				sand. The nearest record is approximately 8 km east of the survey area.	for this species within the survey area.
Myrtaceae	<i>Chamelaucium</i> sp. Coolcalalaya (A.H. Burbidge 4233)		P1	Dense and compact shrub to 1.5 m, red-purple/white flowers. Undulating dunes, white sandplains. Yellow sandy loam. Closest known record is approximately 9 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Myrtaceae	<i>Eucalyptus blaxellii</i>		P4	(Mallee), 1-4 m high, bark smooth. Flowers white-cream, August to November. Grey sand, clay. Rocky hillsides, creek flats. The nearest record is approximately 8 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Myrtaceae	<i>Eucalyptus cuprea</i>	En	T	(Mallee), 2.5-5 m high, bark rough to 1.5 m, box-type. Flowers white, August to November. Shallow soils over granite. The nearest record is 16 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Myrtaceae	<i>Hypocalymma angustifolium</i> subsp. Hutt River (S. Patrick 2982)	Vu	T	Shrub. Moist, brown black peat clay with sand. The nearest record is approximately 9 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Myrtaceae	<i>Hypocalymma longifolium</i>	Vu	T	Open shrub, to 1 m high. Grey sand or clay, sandstone. Rocky breakaways, swampland. Flowers white/cream, August to September. The nearest record is approximately 9 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Myrtaceae	<i>Melaleuca huttensis</i>		P3	Erect shrub to 1.5 m high, flowers cream. Generally occurs on brown sand over laterite, but has been previously recorded in the survey area in orange, white and yellow sands on lower slopes of undulating plains and sandplains (GHD 2019). Flat plain. There area records located in M70/968, located less than 500 m east of the survey area (GHD 2014).	Unlikely – there is suitable habitat within the survey area, however, suitable survey effort did not record this species
Myrtaceae	<i>Verticordia densiflora</i> var. <i>roseostella</i>		P3	Open shrub, 0.4-1.3 m high. Flowers pink-white, September to December. Sandy gravelly soils. The nearest record is approximately 9 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Orchidaceae	<i>Caladenia barbarella</i>	En	T	Tuberous, perennial, herb, 0.08-0.25 m high. Flowers green, August to September. Occur on shallow, grey, dark brown, sandy clayey loam. Rocky ledges, alongside seasonal creeklines, winter-wet depressions. Closest known record is approximately 43 km north-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Orchidaceae	<i>Caladenia bryceana</i> subsp. <i>cracens</i>	Vu	T	Tuberous, perennial, herb, 0.03-0.08 m high. Flowers green-yellow, August to September. Sand over limestone, and shallow beige sands under moist areas beneath <i>M. cardiophylla</i> and <i>Grevillea argyropylla</i> , in the study area (GHD 2019). South of Kalbarri in low heath on limestone hills; north in winter-moist flats. The nearest records are approximately 23 km north and south of the survey area. GHD have recorded this species in adjacent tenements, with the closest record approximately 600 m east (GHD 2019).	Possible – suitable habitat (VT02) is present in the survey area and targeted surveys during the species flowering period may possibly record the species

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Orchidaceae	<i>Caladenia elegans</i>	En	T	Tuberous, perennial, herb, 0.2-0.3 m high. Flowers yellow, July to August. Clayey loam. Winter-wet clay flats. The nearest record is approximately 35 metres from the northern end of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area. The record of this species was recorded in 2009, and the land has since been dramatically altered.
Orchidaceae	<i>Caladenia hoffmanii</i>	En	T	Tuberous, perennial, herb, 0.13-0.3 m high. Flowers green and yellow and red, August to October. Clay, loam, laterite, granite. Rocky outcrops and hillsides, ridges, swamps and gullies. The nearest record is 8 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Orchidaceae	<i>Diuris drummondii</i>	Vu	T	Tuberous, perennial, herb, 0.5-1.05 m high. Flowers yellow, November to December or January. Low-lying depressions, swamps. The nearest record is approximately 9 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Orchidaceae	<i>Diuris recurva</i>		P4	Tuberous, perennial, herb, 0.2-0.3 m high. Flowers yellow & brown, July to August. Loam. Winter-wet areas. In the study area, has been recorded on pale brown/yellow sand in sparsely vegetated areas with <i>M. huttensis</i> and <i>Grevillea argyrophylla</i> heath, and <i>M. huttensis</i> and <i>Acacia rostellifera</i> shrubland (GHD 2019). The nearest DBCA record is approximately 8 km east of the survey area. The species has been recorded recently approximately 600 m east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Orchidaceae	<i>Drakaea concolor</i>	Vu	T	Tuberous, perennial, herb, 0.25-0.3 m high. Flowers red and yellow, August to September. Grows in moist sandy sites in the Northampton region along the Murchison and Hutt River. The nearest record is approximately 3.5 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Orchidaceae	<i>Pterostylis sinuata</i>	En	T	Small tuberous herb 5-10 cm tall. The flower spike emerges from a basal rosette of leaves and bears between two and twenty pale green 'greenhood' flowers, each of which are approximately 5 x 5 mm (Hoffman and Brown 1998). Fl. August to early September. Prefers open <i>Melaleuca uncinata</i> and <i>Hakea recurva</i> low scrub over low heath in winter-wet clay soils over laterite. The nearest record is 9 km south-east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Polygalaceae	<i>Comesperma rhadinocarpum</i>		P3	Perennial, herb. Flowers blue, October to November. Sandy soils. The nearest record is approximately 8 km north of the survey area.	Unlikely – suitable survey effort did not record this species
Proteaceae	<i>Grevillea triloba</i>		P3	Diffuse or spreading shrub, (0.4-) 0.9-1.5 (-2.5) m high. Flowers white/pink-white, June to October. Sandy loam on sandstone or limestone, lateritic soils. The nearest record is from the Port Gregory area near the Murchison (GPS record incorrect on Naturemap).	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Rhamnaceae	<i>Blackallia nudiflora</i>		P3	Shrub, 0.3-1 m high, often with spinescent branchlets. Clay or sandy clay with granite. On hills or breakaways, plains. The nearest record is approximately 4 km east of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.

Family	Taxa	Status		Description and closest record information (if available) (WA Herbarium 2020, DBCA 2019)	Likelihood of occurrence
		Federal	State		
Rutaceae	<i>Drummondita ericoides</i>	En	T	Divaricately branched shrub, 0.3-1 m high. Flowers yellow and white and violet/purple, September to October. Occur on rocky places. Closest known record is approximately 44 km south of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Scrophulariaceae	<i>Eremophila microtheca</i>		P4	Erect shrub, 0.7-1.6 m high. Sandy clay. Winter wet flats, saline flats, sandplains. Flowers blue-purple, August to September. The nearest record is approximately 3 km west of the survey area.	Highly Unlikely – there is no suitable habitat for this species within the survey area.
Solanaceae	<i>Anthocercis intricata</i>		P3	Dense, spinescent shrub, 0.9-3 m high. Flowers white-cream, June to September. Sand or loam over limestone. In the study area, orange/yellow sand in <i>M. huttensis</i> and <i>Acacia rostellifera</i> shrubland (GHD 2019). Consolidated sand dunes. The nearest DBCA record is 5 km south however GHD has recorded this species in adjacent tenements, with records less than 500 m from the survey area (GHD 2014, GHD 2019).	Possible – suitable habitat occurs within the survey area, however, the habitat is degraded.

Appendix E – Fauna data

Fauna species list

Fauna likelihood of occurrence

Fauna species recorded during the survey

Family	Genus	Species	Common Name	Status
Birds				
Acanthizidae	<i>Acanthiza</i>	<i>chrysorrhoa</i>	Yellow-rumped Thornbill	
Accipitridae	<i>Aquila</i>	<i>audax</i>	Wedge-tailed Eagle	
Accipitridae	<i>Pandion</i>	<i>cristatus</i>	Eastern Osprey	Mi, IA
Accipitrinae	<i>Elanus</i>	<i>axillaris</i>	Black-Shouldered Kite	
Artimidae	<i>Artamus</i>	<i>minor</i>	Little Woodswallow	
Artimidae	<i>Cracticus</i>	<i>nigrogularis</i>	Pied Butcherbird	
Campephagidae	<i>Coracina</i>	<i>novaeollandiae</i>	Black-faced Cuckoo-shrike	
Corvidae	<i>Corvus</i>	<i>coronoides</i>	Australian Raven	
Columbidae	<i>Streptopelia</i>	<i>senegalensis</i>	Laughing Turtle Dove	*
Falconidae	<i>Falco</i>	<i>cenchroides</i>	Australian Kestrel	
Halcyonidae	<i>Todiramphus</i>	<i>sanctus</i>	Sacred Kingfisher	
Hirundinidae	<i>Hirundo</i>	<i>neoxena</i>	Welcome Swallow	
Hirundinidae	<i>Petrochelidon</i>	<i>nigricans</i>	Tree Martin	
Maluridae	<i>Malurus</i>	<i>lamberti</i>	Variegated Fairy-wren	
Maluridae	<i>Malurus</i>	<i>splendid</i>	Splendid Fairy-wren	
Meliphagidae	<i>Gavicalis</i>	<i>virscens</i>	Singing Honeyeater	
Meliphagidae	<i>Lichmera</i>	<i>indistincta</i>	Brown Honeyeater	
Meropidae	<i>Merops</i>	<i>ornatus</i>	Rainbow Bee-eater	
Pachycephalidae	<i>Pachycephala</i>	<i>pectoralis</i>	Golden Whistler	
Petroicidae	<i>Eopsaltria</i>	<i>georgiana</i>	White-breasted Robin	
Pomatostomidae	<i>Pomatostomus</i>	<i>superciliosus</i>	White-browed Babbler	
Timaliidae	<i>Zosterops</i>	<i>lateralis</i>	Silvereye	
Tytonidae	<i>Tyto</i>	<i>alba</i>	Barn Owl	
Mammals				
Bovidae	<i>Ovis</i>	<i>aries</i>	Sheep	*
Canidae	<i>Canis</i>	<i>lupis</i>	Domestic Dog	*
Canidae	<i>Vulpes</i>	<i>vulpes</i>	Red Fox	*
Felidae	<i>Felis</i>	<i>catus</i>	Feral Cat	*
Leporidae	<i>Oryctolagus</i>	<i>cuniculus</i>	European Rabbit	*
Macropodidae	<i>Macropus</i>	<i>fuliginosus</i>	Western Grey Kangaroo	
Macropodidae	<i>Osphranter</i>	<i>rufus</i>	Red Kangaroo	
Suidae	<i>Sus</i>	<i>scrofa</i>	Wild Boar	*
Reptiles				
Agamidae	<i>Pogona</i>	<i>minor minor</i>	Bearded Dragon	
Scincidae	<i>Tiliqua</i>	<i>rugosa</i>	Bobtail Skink	

* Introduced (BAM Act)

Parameters of fauna likelihood of occurrence assessment

Assessment outcome	Description
Present	Species recorded during the field survey or from recent, reliable records from within or close proximity to the survey area.
Likely	Species are likely to occur in the survey area where there is suitable habitat within the survey area and there are recent records of occurrence of the species in close proximity to the survey area. OR Species known distribution overlaps with the survey area and there is suitable habitat within the survey area.
Unlikely	Species assessed as unlikely include those species previously recorded within 10 km of the survey area however: <ul style="list-style-type: none"> • There is limited (i.e. the type, quality and quantity of the habitat is generally poor or restricted) habitat in the survey area. • The suitable habitat within the survey area is isolated from other areas of suitable habitat and the species has no capacity to migrate into the survey area. OR Those species that have a known distribution overlapping with the survey area however: <ul style="list-style-type: none"> • There is limited habitat in the survey area (i.e. the type, quality and quantity of the habitat is generally poor or restricted). • The suitable habitat within the survey area is isolated from other areas of suitable habitat and the species has no capacity to migrate into the survey area.
Highly unlikely	Species that are considered highly unlikely to occur in the survey area include: <ul style="list-style-type: none"> • Those species that have no suitable habitat within the survey area. • Those species that have become locally extinct, or are not known to have ever been present in the region of the survey area.

Definitions

Term	Description
Study area	A 20 km buffer around the survey area
Survey area	The potential project footprint
Cr	Critically endangered
En	Endangered
Vu	Vulnerable
IA	International agreement
Mi, Ma	Migratory, Marine
CD	Conservation dependent
OS	Other specially protected fauna
P1 – P4	Priority 1 – Priority 4
EPBC Act	Environmental Protection and Biodiversity Conservation Act 1999
DBCA	Department of Biodiversity and Conservation Attractions 2019 WA Government, Department of Parks and Wildlife Threatened and Priority fauna rankings
BC Act	<i>Biodiversity Conservation Act 2016</i>

Fauna likelihood of occurrence assessment of conservation significant fauna identified in the desktop assessment as potentially occurring within the study area

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Anous tenuirostris melanops</i> Australian Lesser Noddy	Vu	En			X	The Australian Lesser Noddy is usually found only around its breeding islands in the Houtman Abrolhos Islands in Western Australia. There are also some records north of the breeding islands, for example at the Wallabi Group of islands, in the northern Houtman Abrolhos Islands, on Barrow Island, and at Webb Island. The species usually occupies coral-limestone islands that are densely fringed with White Mangrove <i>Avicennia marina</i> . It occasionally occurs on shingle or sandy beaches (Higgins & Davies 1996). The Australian Lesser Noddy roosts mainly in mangroves, especially at night but may sometimes rest on beaches.	Highly unlikely There is no suitable habitat within the survey area. The closest known records are from the Abrolhos Islands, over 60 km off the coast.
<i>Calidris canutus</i> Red knot, Knot	En	En, IA			X	In Australasia the Red Knot mainly inhabits intertidal mudflats, sandflats and sandy beaches of sheltered coasts, in estuaries, bays, inlets, lagoons and harbours; sometimes on sandy ocean beaches or shallow pools on exposed wave-cut rock platforms or coral reefs. They are occasionally seen on terrestrial saline wetlands near the coast, such as lakes, lagoons, pools and pans, and recorded on sewage ponds and saltworks, but rarely use freshwater swamps. They rarely use inland lakes or swamps (DEE 2017). They are found near mudflats and estuaries from Murchison to Bunbury but are then uncommon from Wilson Inlet to Esperance. In the Perth region they are mainly found in Alfred Cove and Peel Inlet (Nevill 2013).	Unlikely The species has been recorded from the nearby Hutt Lagoon, however there is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Calidris ferruginea</i> Curlew Sandpiper	Cr	Cr	X		X	Curlew Sandpipers mainly occur on intertidal mudflats in sheltered coastal areas, such as estuaries, bays, inlets and lagoons, and also around non-tidal swamps, lakes and lagoons near the coast, and ponds in saltworks and sewage farms. They are also recorded inland, though less often, including around ephemeral and permanent lakes, dams, waterholes and bore drains, usually with bare edges of mud or sand. They occur in both fresh and brackish waters. Occasionally they are recorded around floodwaters (Higgins & Davies 1996). Curlew Sandpipers forage on mudflats and nearby shallow water. They forage at the edges of shallow pools and drains of intertidal mudflats and sandy shores. At high tide, they forage among low sparse emergent vegetation, such as saltmarsh, and sometimes forage in flooded paddocks or inundated saltflats. Curlew Sandpipers generally roost on bare dry shingle, shell or sand beaches, sandspits and islets in or around coastal or near-coastal lagoons and other wetlands, occasionally roosting in dunes during very high tides and sometimes in saltmarsh (Higgins & Davies 1996).	Unlikely The species has been recorded multiple times from Hutt Lagoon and nearby coastal areas, however there is no suitable habitat within the survey area.
<i>Calidris tenuirostris</i> Great Knot	Cr	Cr	X			In Australasia, the species typically prefers sheltered coastal habitats, with large intertidal mudflats or sandflats. This includes inlets, bays, harbours, estuaries and lagoons. They are occasionally found on exposed reefs or rock platforms, shorelines with mangrove vegetation, ponds in saltworks, at swamps near the coast, saltlakes and non-tidal lagoons. The Great Knot rarely occurs on inland lakes and swamps. Typically, the Great Knot roosts in large groups in	Unlikely The species has been recorded from Hutt Lagoon, however there is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
						open areas, often at the waters edge or in shallow water close to feeding grounds (DotEE 2020).	
<i>Calyptrorhynchus latirostris</i> Carnaby's Cockatoo	En	En	X	X	X	Carnaby's Black-cockatoo occurs in uncleared or remnant native eucalypt woodlands, especially those that contain salmon gum, wandoo, marri, jarrah and karri, and in shrubland or kwongan heathland dominated by Hakea, Dryandra, Banksia and Grevillea species. Breeding activity is restricted to eucalypt woodlands mainly in the semiarid and subhumid interior, from Kalbarri in the north, Three Springs District south to the Stirling Range, west to Cockleshell Gully and east to Manmanning. The species has expanded its breeding range westward and south into the jarrah-marri forests of the Darling Scarp and into the tuart forests of the Swan Coastal Plain, including the Yanchep area, Lake Clifton and near Bunbury. It nests in trees older than 120-150 years (DotEE 2020).	Unlikely The survey area is located within the non-breeding range of the modelled distribution of the Carnaby's Cockatoo (DSEWPaC 2012). However, there is no suitable habitat (foraging or roosting habitat) present within the survey area.
<i>Charadrius leschenaultii</i> Greater Sand Plover	Vu	Vu	X		X	In the non-breeding grounds in Australasia, the species is almost entirely coastal, inhabiting littoral and estuarine habitats. They mainly occur on sheltered sandy, shelly or muddy beaches with large intertidal mudflats or sandbanks, as well as sandy estuarine lagoons and inshore reefs, rock platforms, small rocky islands or sand cays on coral reefs. They are occasionally recorded on near-coastal saltworks and saltlakes, including marginal saltmarsh, and on brackish swamps (DotEE 2020).	Unlikely The species has been recorded multiple times from Hutt Lagoon and nearby coastal areas, however there is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Charadrius leschenaultii</i> Lesser Sand Plover	En, Mi	En	X			In non-breeding grounds in Australia, the Lesser Sand Plover usually occurs in coastal littoral and estuarine environments. It inhabits large intertidal sandflats or mudflats in sheltered bays, harbours and estuaries, and occasionally sandy ocean beaches, coral reefs, wave-cut rock platforms and rocky outcrops. It also sometimes occurs in short saltmarsh or among mangroves, in saltworks and near-coastal salt pans, brackish swamps and sandy or silt islands in river beds. The species is seldom recorded away from the coast, at margins of lakes, soaks and swamps associated with artesian bores (DotE 2016). The Lesser Sand Plover mainly occurs in northern regions, and becomes more scarce in the south west (Nevill 2013).	Unlikely The species has been recorded multiple times from Hutt Lagoon and nearby coastal areas, however there is no suitable habitat within the survey area.
<i>Diomedea amsterdamensis</i> (Amsterdam Albatross)	En, Mi	Cr			X	All the Albatross species have been grouped together as they are all primarily marine, pelagic, aerial birds.	Highly Unlikely No suitable habitat is present within the survey area.
<i>Diomedea epomophora</i> (Southern Royal Albatross)	Vu, Mi	Vu					
<i>Diomedea exulans</i> (Wandering Albatross)	Vu, Mi	Vu					
<i>Thalassarche carteri</i> (Indian Yellow-nosed Albatross)	Vu, Mi	En					
<i>Thalassarche cauta cauta</i> (Shy Albatross)	Vu, Mi	Vu					
<i>Thalassarche cauta steadi</i> (White-capped Albatross)	Vu, Mi	Vu					

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Thalassarche impavida</i> Campbell Albatross	Vu, Mi	Vu					
<i>Thalassarche melanophris</i> Black-browed Albatross	Vu, Mi	En					
<i>Leipoa ocellata</i> Malleefowl	Vu	Vu		X	X	The Malleefowl generally occurs in semi-arid areas of WA, in shrublands and low woodlands that are dominated by mallee vegetation, as well as native pine <i>Callitris</i> woodlands, <i>Acacia</i> shrublands, paperbark, skheoak, Broombush <i>Melaleuca uncinata</i> vegetation, eucalypt woodlands, or coastal heathlands. Mostly they are found where there are sandy or gravel soils. The nest is a large mound of sand or soil and organic matter (Jones & Goth 2008; Morcombe 2011; Nevill 2013). In WA they are found from the southwest Nullarbor to Albany, north, and then west from Moore River up to Shark Bay, past Cue, across to Wiluna and east to the northern Victoria Desert south of the Blackstone Ranges (Nevill 2013; Pizzey & Knight 2012).	Unlikely The distribution of this species within this region is restricted to the Kalbarri National Park, which is located approximately 40 km north of the survey area. No evidence of this species was observed during the survey.
<i>Limosa lapponica</i> Bar-tailed Godwit / Northern Siberian Bar-tailed Godwit	Vu or Cr, Mi	Vu or Cr, IA			X	The Bar-tailed Godwit is found mainly in coastal habitats such as large intertidal sandflats, banks, mudflats, estuaries, inlets, harbours, coastal lagoons and bays. It is found often around beds of seagrass and, sometimes, in nearby saltmarsh (Morcombe 2011). They usually forage near the edge of water or in shallow water, mainly in tidal estuaries and harbours and roost on sandy beaches, sandbars, spits and also in near-coastal saltmarshes (Marchant & Higgins 1993).	Unlikely The species has been recorded from Hutt Lagoon, however there is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Macronectes giganteus</i> Southern Giant-Petrel	En, Mi	IA			X	The Southern Giant-Petrel is marine bird that occurs in Antarctic to subtropical waters. In summer, it mainly occurs over Antarctic waters, and it is widespread south as far as the pack-ice and onto the Antarctic continent (Marchant & Higgins 1990). The species is not known to breed in Australia.	Highly unlikely There is no suitable habitat within the survey area.
<i>Macronectes halli</i> Northern Giant Petrel	Vu, Mi	Mi				The Northern Giant Petrel breeds in the sub-Antarctic and visits areas off the Australian mainland during the winter months (May-Oct). They are usually seen in waters off the south of Australia (DotEE 2020). The species is primarily Marine.	Highly unlikely There is no suitable habitat within the survey area.
<i>Numenius madagascariensis</i> Eastern Curlew	Cr, Mi	Cr			X	The Eastern Curlew is most commonly associated with sheltered coasts, especially estuaries, bays, harbours, inlets and coastal lagoons, with large intertidal mudflats or sandflats, often with beds of seagrass (Marchant & Higgins 1993).	Highly unlikely There is no suitable habitat within the survey area.
<i>Pterodroma mollis</i> Soft-plumaged Petrel	Vu				X	The Soft-plumaged Petrel is a marine, oceanic species. Soft-plumaged Petrels are mainly subantarctic, but occur over a wide range of sea surface-temperatures. Soft-plumaged Petrels breed on Maatsuyker Island off southern Tasmania. Beachcast birds have been found from Maryborough, Queensland, south to NSW, Tasmania, Victoria, South Australia and south-west Western Australia (DotEE 2020).	Highly unlikely There is no suitable habitat within the survey area.
<i>Rostratula australis</i> Australian Painted Snipe	En	En			X	The Australian Painted Snipe generally inhabits shallow terrestrial freshwater (occasionally brackish) wetlands, including temporary and permanent lakes, swamps and claypans. Australian Painted Snipe breeding habitat requirements may be quite specific: shallow wetlands with areas of bare wet mud and both	Highly unlikely There is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
						upper and canopy cover nearby. The species rarely occurs in south-western Australia, where it was once more common (Marchant & Higgins 1993; Garnett and Crowley 2000).	
<i>Sternula nereis nereis</i> Australian Fairy Tern	Vu	Vu			X	The Fairy Tern occurs along the coast of WA as far north as the Dampier Archipelago near Karratha, but mostly in the southern part of Australia including most of the coastline in the south west. It nests on sheltered sandy beaches, coastal inlets, spits and banks above the high tide line and below vegetation. It has been found in embayments of a variety of habitats including offshore, estuarine or lacustrine (lake) islands, wetlands, and mainland coastline (DotEE 2020; Nevill 2013). They can also be seen in saltfields, saline or brackish lakes, and sewage ponds near the coast (Pizzey and Knight 2012).	Highly unlikely There is no suitable habitat within the survey area.
<i>Pandion cristatus</i> Osprey	Mi	IA	X	X	X	Ospreys occur in littoral and coastal habitats and terrestrial wetlands of tropical and temperate Australia and offshore islands. They are mostly found in coastal areas but occasionally travel inland along major rivers, particularly in northern Australia. They require extensive areas of open fresh, brackish or saline water for foraging. They frequent a variety of wetland habitats including inshore waters, reefs, bays, coastal cliffs, beaches, estuaries, mangrove swamps, broad rivers, reservoirs and large lakes and waterholes. They exhibit a preference for coastal cliffs and elevated islands in some parts of their range but may also occur on low sandy, muddy or rocky shores and over coral cays. The distribution of the species around the northern coast (south-western WA to	Present An individual was observed nesting in a dead <i>Acacia</i> tree within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
						south-eastern NSW) appears continuous except for a possible gap at Eighty Mile Beach (DotEE 2020).	
<i>Apus pacificus</i> Fork-tailed Swift	Mi	IA	X	X	X	The Fork-tailed Swift are widespread in coastal and sub-coastal areas between Augusta and Carnarvon, including some on nearshore and offshore islands. This species is almost exclusively aerial, flying less than 1 m to at least 300 m above ground. Occupies low to very high airspace over varied habitat, rainforest to semi-desert; most active just ahead of summer storm fronts. They do not breed in Australia (DotEE 2020)	Likely There are a number of records along the coast at Port Gregory and near Hutt Lagoon.
<i>Pluvialis fulva</i> Pacific Golden Plover	Mi	IA	X	X	X	In Australia the Pacific Golden Plover usually inhabits coastal habitats, on beaches, mudflats and sandflats (sometimes in vegetation such as mangroves, low saltmarsh such as <i>Sarcocornia</i> , or beds of seagrass) in sheltered areas including harbours, estuaries and lagoons, and also in saltworks. It is sometimes recorded on islands, sand and coral cays and exposed reefs and rocks. They are less often recorded in terrestrial habitats, but can be seen in habitats with short grass in paddocks, crops or airstrips, or ploughed or	Unlikely This species is largely restricted to coastal areas. There are records of this species around Port Gregory and Hutt Lagoon however it is considered unlikely to utilise habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
						recently burnt areas. This species does not breed in Australia (DotEE 2020).	
<i>Falco peregrinus</i> Peregrine Falcon		OS	X			The Peregrine Falcon is found on and near cliffs, gorges, timbered watercourses, riverine environments, wetlands, plains, open woodlands, and pylons and spires of buildings, though less frequently in desert regions (Morcombe 2011; Pizzey & Knight 2012). They are not common but can be found almost anywhere throughout WA and in the southwest, including particularly at Fitzgerald River, Stirling Range, Porongurup National Parks, Kondinin, and Peak Charles, with many more locations north of Perth (Nevill 2013).	Likely There are records of this species occurring around the Port Gregory area. This species may occur as an infrequent visitor, foraging within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Calidris acuminata</i> Sharp-Tailed Sandpiper	Mi	IA	X	X	X	In Australasia, the Sharp-tailed Sandpiper prefers muddy edges of shallow fresh or brackish wetlands, with inundated or emergent sedges, grass, saltmarsh or other low vegetation. This includes lagoons, swamps, lakes and pools near the coast, and dams, waterholes, soaks, bore drains and bore swamps, saltpans and hypersaline saltlakes inland. They also occur in saltworks and sewage farms. They use flooded paddocks, sedgelands and other ephemeral wetlands, but leave when they dry. They use intertidal mudflats in sheltered bays, inlets, estuaries or seashores, and also swamps and creeks lined with mangroves. Sometimes they occur on rocky shores. They are widespread from Cape Arid to Carnarvon, around coastal and subcoastal plains of Pilbara Region to south-west and east Kimberley Division (DotEE 2020).	Unlikely There are multiple records of this species occurring within Hutt Lagoon and Port Gregory, however there is no suitable habitat within the survey area.

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Calidris melanotos</i> Pectoral Sandpiper	Mi	IA	X	X	X	In Australia, the Pectoral Sandpiper prefers shallow fresh to saline wetlands. The species is found at coastal lagoons, estuaries, bays, swamps, lakes, inundated grasslands, saltmarshes, river pools, creeks, floodplains and artificial wetlands. The species is usually found in coastal or near coastal habitat but occasionally found further inland. It prefers wetlands that have open fringing mudflats and low, emergent or fringing vegetation, such as grass or samphire. The species has also been recorded in swamp overgrown with lignum. In Western Australia the species is rarely recorded (DotEE 2020).	Unlikely There are records of this species occurring from Port Gregory, however this species is unlikely to utilise the habitat within the survey area.
<i>Tringa brevipes</i> Grey-tailed Tattler		P4	X			Habitat coastal; forages in inter-tidal pools, shallows, soft surfaces of mudflats and sand beaches as well as rock ledges, reefs. Often perches on branches, posts or jetties. Common summer migrant to northern Australia and uncommon in the south (Morcombe 2011).	Unlikely There is one record from the Gregory/Hutt Lagoon area. The survey area does not contain suitable habitat for this species.

Mammals

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Dasyurus geoffroii</i> Chuditch, Western Quoll	Vu	Vu		X	X	The Chuditch inhabits eucalypt forest (especially Jarrah, <i>E. marginata</i>), dry woodland, mallee shrublands, heaths, and desert, particularly in the south coast of WA. They also occur at lower densities in drier woodland and mallee shrubland in the goldfields and wheatbelt, as well as in Kalbarri National Park (translocated). Chuditch require adequate numbers of suitable den and refuge sites (horizontal hollow logs or earth burrows) to survive (DEC 2012a). In Jarrah forest, Chuditch populations occur in both moist, densely vegetated, steeply sloping forest and drier, open, gently sloping forest (Van Dyck and Strahan 2008). The species can travel large distances, and for this reason requires habitats that are of a suitable size and not excessively fragmented (DEC 2012a).	Highly unlikely The closest known record is approximately 20 km north-east of the survey area (dated 2008). Chuditch have been successfully translocated to Kalbarri National Park, however they are considered a geographically isolated population. The area surrounding the survey area has been largely cleared/fragmented due to agriculture. Given the lack of key habitat for this species, it is considered unlikely to occur.
<i>Notamacropus eugenii</i> subsp. <i>derbianus</i> Tammar Wallaby		P4	X	X		The Tammar Wallaby inhabits dense, low vegetation for daytime shelter and open grassy areas for feeding. It inhabits coastal scrub, heath, dry sclerophyll (leafy) forest and thickets in mallee and woodland. The tammar wallaby is currently known to inhabit three islands in the Houtman Abrolhos group, Garden Island near Perth, Middle and North Twin Peak Islands in the Archipelago of the Recherche, and at least nine sites on the mainland including Dryandra, Boyagin, Tutanning Batalling (reintroduced) Perup, private property near Pingelly, Jaloran Road timber reserve near Wagin, Hopetown, Stirling Range National Park, and Fitzgerald River National Park (DEC 2012; Van Dyck and Strahan 2008).	Unlikely There are two historic records approximately 7 and 18 km south east of the survey area. The species was considered locally extinct however they have been successfully re-introduced into the Kalbarri National Park.

Reptiles

Species	Status		Desktop Search			Ecology and habitat	Likelihood of occurrence
	Federal	State	NM	DBCA Data	PMST		
<i>Egernia stokesii badia</i> Western Spiny-tailed Skink	En	Vu			X	<i>Egernia stokesii badia</i> occurs in open eucalypt woodlands and Acacia-dominated shrublands in semi-arid to arid areas of south-western WA (Geraldton Sandplains and Yalgoo IBRA) and, depending on taxonomic clarification, around Shark Bay including Peron Peninsula, Edel Land and Dirk Hartog Island (Geraldton Sandplain and Carnarvon IBRA). It tends to shelter in logs, in cavities in the trunks and branches of shrubs, as well as in houses and ruins, especially in accumulations of old corrugated iron (DEC 2012b).	Highly Unlikely The closest known record is more than 90 km south east of the survey area.

GHD

Level 10

999 Hay Street

T: 61 8 6222 8222 F: 61 8 9463 6012 E: permail@ghd.com



© GHD 2020

This document is and shall remain the property of GHD. The document may only be used for the purpose for which it was commissioned and in accordance with the Terms of Engagement for the commission. Unauthorised use of this document in any form whatsoever is prohibited.

6138125-

89649/[https://projects.ghd.com/oc/WesternAustralia2/gmagarnetmcpandmp/Delivery/Documents/6138125_REP-A_GMA Garnet Biological Survey.docx](https://projects.ghd.com/oc/WesternAustralia2/gmagarnetmcpandmp/Delivery/Documents/6138125_REP-A_GMA%20Garnet%20Biological%20Survey.docx)

Document Status

Revision	Author	Reviewer		Approved for Issue		
		Name	Signature	Name	Signature	Date
0	S Flemington	D. Farrar		D. Farrar		5/02/2020

www.ghd.com





Memorandum

09 September 2020

To GMA Garnet Pty Ltd

Copy to

From	Sarah Flemington	Tel	+61 8 62228638
Subject	Targeted <i>Caladenia bryceana</i> subsp. <i>cracens</i> survey and conservation listed flora survey of proposed haul road	Job no.	12528268

1 Introduction

1.1 Background

GMA Garnet Pty Ltd (GMA) are in the planning phase for expanding their operations within the Lynton Mine, located east of Hutt Lagoon, in Port Gregory, WA (the project). The expansion includes excavation for expansion of an open-cut mine and associated haul road. GMA has identified it will be necessary to clear vegetation to facilitate the expansion works.

GHD (2020) undertook a survey in December 2019 for the Project, with the purpose of assessing the vegetation and potential environmental values. Based on the results of this survey, GMA and GHD have determined that a targeted Threatened orchid survey is required to support the environmental approval, anticipated to be a Native Vegetation Clearing Permit (NVCP) under Part V of the *Environmental Protection Act 1986* (EP Act).

Caladenia bryceana subsp. *cracens* is known from 15 populations between Northampton and Kalbarri. The habitat and distribution is distinctly different to that of the subspecies *bryceana*, where it is geographically isolated from subsp. *cracens*. *C. bryceana* subsp. *cracens* is known to grow scattered in low heath in shallow soil on coastal limestone. The northern section of its distribution forms populations on winter-wet flats or in swales beneath thickets of *Melaleuca uncinata* in pale red-brown sandy loam or brown sandy clay (Commonwealth of Australia, 2013).

1.2 Purpose

GMA commissioned GHD to undertake a targeted survey for the Threatened orchid, *Caladenia bryceana* subsp. *cracens* (Northern Dwarf Spider-orchid), listed as Vulnerable under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) and Endangered under the *Biodiversity Conservation Act 2016* (BC Act 2016). The purpose of the survey was to support a NVCP application. The outcomes of the survey will be provided to the Department of Mines Industry Regulation and Safety (DMIRS) to inform their environmental assessment. This memorandum should be read in conjunction with GHD (2020) *GMA Garnet Pty Ltd Lynton Mine Expansion Biological*

12528268-98180-11/12528268_Rev A GMA Targeted Orchid Survey Memo.docx



Memorandum

Survey report which contains an assessment of the flora and vegetation values of the entire Project area.

1.3 Scope of works

The scope of works involved a targeted orchid survey of potential habitat for *Caladenia bryceana* subsp. *cracens* within the survey area (Figure 1). The survey area in total is approximately 28.3 hectares (ha). GHD implemented the following scope of works to achieve the purpose of the commission:

- Assessment of the desktop results as summarised in GHD (2020) prior to undertaking the field survey
- Undertake a targeted survey for *Caladenia bryceana* subsp. *cracens* (Threatened) in relevant habitat
- Undertake a targeted survey for conservation listed flora for the proposed haul road
- Draft a memorandum (this report) that documents the methods and results of the field survey
- Provide spatial data suitable to support the submission of a NVCP application to DMIRS.





Memorandum

1.4 Limitations and assumptions

This memorandum has been prepared by GHD for GMA Garnet Pty Ltd and may only be used and relied on by GMA Garnet Pty Ltd for the purpose agreed between GHD and the GMA Garnet Pty Ltd as set out in section 1.2 of this memorandum.

GHD otherwise disclaims responsibility to any person other than GMA Garnet Pty Ltd arising in connection with this memorandum. GHD also excludes implied warranties and conditions, to the extent legally permissible.

The services undertaken by GHD in connection with preparing this memorandum were limited to those specifically detailed in the memorandum and are subject to the scope limitations set out in the memorandum.

The opinions, conclusions and any recommendations in this memorandum are based on conditions encountered and information reviewed at the date of preparation of the memorandum. GHD has no responsibility or obligation to update this memorandum to account for events or changes occurring subsequent to the date that the memorandum was prepared.

The opinions, conclusions and any recommendations in this memorandum are based on information obtained from specific sample points. Site conditions at other areas of the site may be different from the site conditions found at the specific sample points. GHD disclaims liability arising from any of the assumptions being incorrect.

GHD has prepared this memorandum on the basis of information provided by GMA Garnet Pty Ltd and others who provided information to GHD (including Government authorities)], which GHD has not independently verified or checked beyond the agreed scope of work. GHD does not accept liability in connection with such unverified information, including errors and omissions in the memorandum which were caused by errors or omissions in that information.

2 Methodology

The targeted orchid survey was carried out by GHD Senior Botanist Joel Collins and Ecologist Sarah Flemington over four days from 11 – 14 August 2020. The targeted survey methodology has been conducted with reference to the Environmental Protection Authority (EPA) 2016 *Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment* and the Commonwealth of Australia 2013 *Survey Guidelines for Australia's Threatened Orchids*.

Prior to the survey a known population of *Caladenia bryceana* subsp. *cracens* west of the survey area, recorded during a previous survey (GHD 2019), was visited to confirm the species was in flower and assess habitat type. The population outside of the survey area was confirmed as being in full flower (Plate 1).

Areas of potentially suitable habitat occurring within the survey site (VT02 *Melaleuca cardiophylla* shrubland to open shrubland) that was previously mapped (GHD 2020) were surveyed by undertaking systematic transects spaced 10 meters apart (Figure 2) across all mapped VT02 areas within the survey area.



Memorandum

A targeted survey for conservation listed flora was also undertaken across and adjacent the proposed haul road in the north-eastern section of the survey area (Figure 2). The haul road and nearby areas were traversed using the same methodology described above.

2.1 Field survey limitations

The EPA (2016) states flora survey reports for environmental impact assessment in WA should contain a section describing the limitations of the survey methods used. The limitations and constraints associated with this field survey are discussed in Table 1.


Table 1 Field survey limitations

Aspect	Constraint	Comment
Sources of information and availability of contextual information.	Nil	Adequate information is available for the survey area.
Scope (what life forms were sampled etc.)	Nil	Vascular flora were sampled during the survey. Non-vascular flora were not surveyed. The survey focused on <i>Caladenia bryceana</i> subsp. <i>cracens</i> (Threatened) and other conservation listed flora species for the proposed haul road.
Proportion of flora collected and identified (based on sampling, timing and intensity)	Nil	The survey focused on <i>Caladenia bryceana</i> subsp. <i>cracens</i> (Threatened) and other conservation listed flora species for the proposed haul road.
Flora determination	Nil	Flora determination was undertaken by GHD Botanist/Ecologist's in the field. All taxa could be identified to species level. The taxonomy and conservation status of the WA flora is dynamic. This report was prepared with reliance on taxonomy and conservation status current at the time report development, but it should be noted this may change in response to ongoing research and review of International Union for Conservation Nature criteria.
Completeness and further work which might be needed (e.g. was the relevant area fully surveyed)	Nil	The complete survey area was traversed on foot (Figure 2). All areas of the survey area were adequately surveyed for the purpose of the assessment.



Memorandum

Aspect	Constraint	Comment
Mapping reliability	Nil	Data was recorded in the field using hand-held GPS tools (e.g. Samsung tablet and Garmin GPS). Certain atmospheric factors and other sources of error can affect the accuracy of GPS receivers. The GPS units used for this survey are accurate to within ± 4 metres on average. Therefore the data points consisting of coordinates recorded from the GPS may contain inaccuracies.
Timing/weather/season/cycle	Nil	<p>The field survey was conducted on 11-14 August 2020. The timing of the survey is not considered to be a limitation as all flora present was actively growing with flowering on many species observed. In addition <i>Caladenia bryceana</i> subsp. <i>cracens</i> at the time of the survey was flowering. The rainfall in the three months prior to timing of the survey (August), were drier than average for the area (Lynton -station 8075) (BoM, 2020), however, significantly higher than average rainfall was experienced in August (103mm compared to 56.6mm). This is sufficient for active growth and flowering of flora species.</p> <p>The survey timing was considered appropriate for the field survey.</p> <p>The weather conditions recorded during the survey were considered unlikely to have impacted upon the vegetation and flora survey.</p>
Disturbances (e.g. fire, flood, accidental human intervention)	Nil	Much of the survey area has been subjected to historical disturbance events (e.g. clearing, tracks, feral grazing, weeds); however, these disturbances did not impact the survey.
Intensity (in retrospect, was the intensity adequate)	Nil	The survey area was sufficiently covered by the GHD Botanist/Ecologist's during the survey.
Resources	Nil	Adequate resources were employed during the field surveys. Four person day was spent surveying the survey area.
Access restrictions	Nil	All areas of the survey area was accessed on foot during the survey with no restrictions.
Experience levels	Nil	The GHD Botanist/Ecologist's who executed the field survey is suitably qualified and experienced in his field. Joel Collins (Senior Botanist) has over 17 years' experience in undertaking flora and vegetation surveys and assessments in Western Australia. Joel has extensive experience undertaking targeted flora assessments on the Geraldton Sandplains and within the local area. Sarah Flemington (Ecologist) has previously undertaken flora surveys on the Geraldton Sandplains.





Memorandum

3 Results

The field survey did not record any individuals of *Caladenia bryceana* subsp. *cracens* within VT02. This was attributed to the habitat type, which did not align with the habitats containing known records of the species (GHD 2019). In addition there was the presence of invasive weeds throughout the majority of the survey area and significant wild pig grazing (Plate 3).

No conservation listed flora species were recorded within the proposed haul road. The haul road followed an existing track, and contained adjacent areas of previously cleared vegetation, mostly inundated by weeds. For these reasons, the habitats traversed within and adjacent the proposed haul road were not suitable for *C. bryceana* subsp. *cracens*.

4 Conclusion

The location of the orchids visited northeast of the survey area, were noted to be occurring on shallow beige sand on moist rocky limestone areas, higher in the landscape, (Plate 4). The vegetation type at this population is a dense shrubland of *Allocasuarina campestris*, *Melaleuca cardiophylla* and *Grevillea argyrophylla* over *Ecdeiocolea monostachya* sedgeland. These species were identified as the dominant and/or indicator species for the orchid habitat. Other individuals were growing on yellow sandy-loam soils also under *Allocasuarina campestris* (Plate 4).

This vegetation type did not occur in the survey area. *C. bryceana* subsp. *cracens* does not occur in the survey area. No conservation listed flora species occur within or adjacent the proposed haul road.



Memorandum



Plate 1 *Caladenia bryceana* subsp. *cracens*



Plate 2 Limestone habitat at known population of *Caladenia bryceana* subsp. *cracens*



Memorandum



Plate 3 Condition of vegetation within traversed habitat



Plate 4 *Caladenia bryceana* subsp. *cracens* occurring on sandy-loam soil



Memorandum

5 References

Bureau of Meteorology 2020, Climate Data Online. Commonwealth of Australia.

Commonwealth of Australia 2013, Survey Guidelines for Australia's Threatened Orchids Guidelines for detecting orchids listed as 'Threatened' under the Environment Protection and Biodiversity Conservation Act 1999.

Department of Agriculture, Water and the Environment 2020, Approved Conservation Advice for *Caladenia bryceana* subsp. *cracens* (Northern Dwarf Spider-orchid).

GHD 2019, GMA Garnet Pty Ltd, Port Gregory Mine Mining Tenement M70/1380. November 2019.

GHD 2020, GMA Garnet Pty Ltd Lynton Mine Expansion Biological Survey. February 2020.

M70/204 Supporting Information

GMA Mining Australia

Appendix B. Risk Assessment

M70/204 Supporting Information

GMA Mining Australia

The Australian and New Zealand Standard on Risk Management (AS/NZS 4360) defines risk as the product of the likelihood of an event occurring and its consequences. The risk matrix has been developed based on the AS/NZS 4360:2004 to assess the level of risk from activities undertaken in this project. To maximise the benefit of environmental management, manpower and other resources must be allocated to issues on a priority basis. It is generally accepted that the highest-risk issues receive the highest priority.

GMA implements the following mitigation strategy to help manage the risks including:

- Avoid – avoid impacts where possible
- Minimise – if impacts cannot be avoided, minimise, and manage appropriately.
- Rectify – repair, rehabilitate and restore affected areas as soon as possible.
- Reduce – reduce affected areas through preservation and maintenance throughout the life of the mine.
- Offset – where negative impacts still occur, develop an offset package to achieve a net benefit.

Each risk identified was assessed against the DEMIRS environmental factors, as shown in Table B1. Each factor was considered relevant throughout all phases of the project.

Table B1 Objectives for Environmental Factors

Factor	Objective
Biodiversity	To maintain representation, diversity, viability and ecological function at the species, population and community level.
Water Resources	To maintain the hydrological regimes, quality and quantity of groundwater and surface water to the extent that existing and potential uses, including ecosystem maintenance, are protected.
Land and Soils	To maintain the quality of land and soils so that environmental values are protected.
Rehabilitation and Mine Closure	Mining activities are rehabilitated and closed in a manner to make them physically safe to humans and animals, geo-technically stable, geochemically non-polluting/noncontaminating, and capable of sustaining an agreed post-mining land use, and without unacceptable liability to the State.

Table B2 Likelihood Descriptor

Descriptor	Frequency	Probability
Almost Certain	Twice or more per year	Event will occur during the Project / period under review.
		High number of known incidents.
Likely	Once per year	Event likely to occur during the Project / period under review.
		Regular incidents known
Possible	Once in 5 years	Event may occur in some instances during the Project / period under review
		Occasional incidents known.
Unlikely	Once in 10 years	Event is not likely to occur during the Project / period under review
		Some occurrences known.
Rare	Once in 20 years	Event will occur in exceptional circumstances during the Project / period under review.
		Very few or no known occurrences.

M70/204 Supporting Information

GMA Mining Australia

Table B3 Consequence Descriptor

Factor	Insignificant	Minor	Moderate	Major	Severe
Biodiversity	Alteration or disturbance to an isolated area with no effect on habitat or ecosystem. Loss of an individual plant / animal of conservation significance.	Alteration or disturbance to <10% of a habitat or ecosystem resulting in a recoverable impact within 2 years. Loss of multiple plants / animals of conservation significance.	Alteration or disturbance to 10-40% of a habitat or ecosystem resulting in a recoverable impact within 2-5 years. Loss of <50% known local population of plant / animal of conservation significance.	Alteration or disturbance to 40- 70% of a habitat or ecosystem resulting in a recoverable impact within 5-15 years. Loss of >50% known local population of plant / animal species with possible loss of entire local population.	Alteration or disturbance to >70% of a habitat or ecosystem resulting in a recoverable impact >15 years. Local loss of conservation significant or listed species. Extinction of a species.
Water Resources	Negligible change to hydrological processes, water availability or water quality	Short-term modification of hydrological processes, water availability and quality within project tenure, but no change in beneficial use.	Medium-term modification of hydrological processes, water availability and water quality within project tenure, but no change in beneficial use. Short-term modification of hydrological processes, water availability and water quality outside project tenure, but no change in beneficial use.	Long-term modification of hydrological processes, water availability and water quality within project tenure, but no change in beneficial use. Medium-term modification of hydrological processes, water availability and water quality outside project tenure, with change in beneficial use.	Long-term or permanent modification of hydrological processes, water availability or water quality outside project tenure, with impacts to a water-dependent environmental value and/or change in beneficial use.
Land and Soils	Clean-up by site personnel, rectified immediately. Confined to immediate area around source.	Clean-up by site personnel, remediation within 1 year. Confined to operational area.	Clean-up by site personnel, remediation within 1-3 years. Minor impact outside disturbance envelope or minor impact to soil stockpiles.	Clean-up requiring external specialist, remediation within 3-10 years. Impact has migrated outside the disturbance envelope or contamination of soil stockpiles.	Clean-up requiring external specialist. Remediation >10 years, or permanent residual impact. Impact outside the tenement boundary.

M70/204 Supporting Information

GMA Mining Australia

Factor	Insignificant	Minor	Moderate	Major	Severe
Rehabilitation and Mine Closure	Site is safe, stable a non-polluting. Post mining land use is not adversely affected.	Site is safe, all major landforms are stable, and any stability or pollution issues are contained and require no residual management. Post mining land use is not adversely affected	Site is safe, and any stability or pollution issues require minor, ongoing maintenance by end land-user. Post mining land use cannot proceed without some management.	Site cannot be considered safe, stable or non-polluting without long-term management or intervention. Post mining land use cannot proceed without ongoing management.	Site is unsafe, unstable and/or causing pollution or contamination that will cause an ongoing residual affect. Post mining land use cannot be achieved.

Table B4 Risk Matrix

	Risk Matrix	Insignificant	Minor	Moderate	Major	Severe
Likelihood	5 Almost Certain	M	H	H	E	E
	4 Likely	M	M	H	H	E
	3 Possible	L	M	M	H	H
	2 Unlikely	L	L	M	H	H
	1 Rare	L	L	L	M	M

Table B5 Level of Consequence

Descriptor	Explanation
Low	Risk rating is based on subjective opinion or relevant past experience. Baseline data/information has limitations, with only general conclusions possible and further work is required.
Medium	Risk rating is based on similar conditions being observed previously. Baseline data/information has some gaps or minor further work required
High	Risk rating is based on testing, modelling or experiments. Baseline data/information is complete and analysis appropriate for level of data.

Table B6 Acceptability of Risk Level (Inherit)

Risk Level	Acceptability	Treatment
Extreme	Unacceptable	Risk will not be tolerated. Modification of activity required, and Mining Proposal amended.
High	May be acceptable, with specific risk treatments.	Risk may be tolerated with application of high reliability risk treatments. Environmental outcome / Closure objective required
Moderate	Acceptable, with relevant risk treatments.	Risk is tolerable with application of appropriate risk treatments. Environmental outcome / Closure objective required.
Low	Acceptable	Risk is acceptable, but still requires industry best practice environmental management.

M70/204 Supporting Information

GMA Mining Australia

Appendix C. GMA Dust and Management Plan

Dust Management

Mining Australia Procedure

Contents

1. Purpose and Scope	2
2. Roles and Responsibilities.....	2
3. Definitions	2
4. Legal requirements	3
5. Process	3
5.1. Document and Communicate Dust Requirements.....	3
5.2. Plan Activities	3
5.3. Minimise Dust during Operations.....	3
5.4. Undertake Monitoring.....	5
5.5. Report Incident or Complaint	5
6. Training and Competency	6
7. Supporting Documents	6
8. Related Documents.....	6
9. References	6
10. Revision	6
Appendix A. Additional Dust Control Guidance	7
A.1. Factors Influencing the Levels of Dust and other Air Pollutants	7
A.2. Dust Control Measures.....	7
A.2.1. Limit Cleared Areas and Maximise Vegetation	7
A.2.2. Timing of Development and Development Staging	8
A.2.3. Wind Barriers	8
A.2.4. Earthmoving Management	8
A.2.5. Management of Material Stockpiles	9
A.2.6. Watering Road	9
A.2.7. Reducing the Traffic and Speed.....	9
A.2.8. Improving Road Design	10
A.2.9. Hydromulch.....	10
A.2.10. Chemical Stabilisation	10
A.2.11. Covering or Sealing Unpaved Surfaces.....	10
A.2.12. Wind monitoring	11
Appendix B. Procedure Acknowledgement Form.....	12

Dust Management

Mining Australia Procedure



1. Purpose and Scope

Dust can be generated through activities undertaken by GMA Garnet Pty Ltd (GMA). Dust Management provides guidance to successfully manage dust to ensure the impact on the environment and communities in which we operate is minimised.

This procedure details management measures to:

- Minimise the emission of dust associated with the operations
- Prevent negative impacts on sensitive receptors (the surrounding environment and local communities)
- Comply with relevant environmental legal and other requirements.

This procedure applies to all personnel employed by GMA and Sites.

2. Roles and Responsibilities

Role	Responsibilities
General Manager	<ul style="list-style-type: none">• Accountable for ensuring adequate resources are available for the implementation and management of this procedure
Operations Manager	<ul style="list-style-type: none">• Managing the implementation of this procedure for their Site
Supervisors/Superintendent	<ul style="list-style-type: none">• Manage the implementation of the requirement of this procedure with their teams and areas of responsibility
Environmental Team	<ul style="list-style-type: none">• Providing advice and assistance to the Division with the implementation of this Procedure.• Undertake monitoring of the requirements within this Procedure.• Periodic reporting of results internally and externally as defined under legislative requirements.

3. Definitions

Term	Definition
Aspect	Element of GMA's activities, products or services that can interact with the environment.
BoM	Bureau of Meteorology
Dust	Fine soil/material particles emitted into the atmosphere from mining and other activities.
Dust Exceedance	In the absence of environmental monitoring data, this could be dust above standard operating, that could impact sensitive receptors and that is more than just a once-off occurrence.
DWER	Department of Water and Environmental Regulations
Environment	Living things, their physical, biological and social surroundings, and interactions between these.
Environmental Impact	Any change to the environment, whether adverse or beneficial, wholly or partially resulting from an organisation's aspects.
EPA Licence	<i>Environment Protection Act 1986</i> Environmental Licence to Operate
Licensed premise	A place that is prescribed under the under the <i>Environment Protection Act 1986</i> .

Term	Definition
Risk	The probability (likelihood) of harm or damage occurring from exposure to a hazard, and the likely consequences of that harm or damage.
Sensitive Receptor	Locations, such as residential buildings or other premises, communities, flora, fauna or habitats, where health or property or environmental values may be affected by emissions above background levels.
Shall	The term "Shall" means mandatory.

4. Legal requirements

Port Gregory Site is an *Environmental Protection Act 1986* (EPA) –Licenced premise (L8561/2011/1). The Licence Premises includes Lynton (M70/204, M70/259, M70/968, M70/1330 and M70/1331), Hose (M70/856 and G70/171) and Utcha (M70/926 and M70/927).

Geraldton Site is also an EPA Licenced premise (L6145/1983/11). As a Licenced premise GMA is required to implement dust management measures to minimise dust and impacts to sensitive receptors. The management measures are outline in Section 5.

5. Process

5.1. Document and Communicate Dust Requirements

The requirements of this procedure and any project specific requirements shall be communicated to personnel involved in dust causing activities during the site inductions, pre-start meetings and during toolbox meetings. The Toolbox Topic: Dust can be used to communicate these requirements.

5.2. Plan Activities

Activities involving the generation of dust shall be planned to minimise emissions and impacts to sensitive receptors:

- Areas to be disturbed shall be identified, minimised and disturbance shall be a staged process as per the requirements outlined in Clearing and Ground Disturbance (HSE-172).
- Use hardstand areas to minimise dust emissions where feasible.
- Discuss activities that have the potential to generate high levels of dust at pre-start meetings and agree on reduction methods before undertaking works.
- Monitor daily weather forecasts for temperature and wind speed and communicate the forecast information to persons involved in dust generating and dust suppression activities, where there is a risk of impacting sensitive receptors. At the Port Gregory mine site monitor the wind station.

5.3. Minimise Dust during Operations

During operations dust shall be minimised by:

- Operating proactively subject to weather forecasting over a 24 hour period (refer to Appendix A.2.12).
- Monitoring Port Gregory wind station (refer to Appendix A.2.12).
- Maintaining roads throughout the Site, including watering the roads to maintain moisture on the surface of roads/haul routes
- Use of water trucks and/or water cannons to dampen areas identified as being potentially dust generating (sandy soils, soil stockpiles, unsealed access roads etc.). The dampening frequency shall be determined based on weather conditions and dust emissions (refer to Appendix A.2.12).
- Restricting all vehicles to dedicated roads and tracks

- Depending on the situation reduce speed limits to minimise dust generation.
- Introducing dust suppression additives where required and practicable
- Maintaining dust suppression systems on conveyor belts.
- Operating dust suppression sprinkler system at Geraldton as per SOP-40.
- Covering vehicles transporting soils off-site to minimise dust generation during transport.
- Implementing regular inspections and preventative maintenance strategies for dust control equipment.
- Maintaining adequate spares at the Site for critical items of control equipment, such as water pumps for dust suppression sprays, spray heads etc.
- Undertaking staged vegetation clearing to minimise open areas
- Undertaking vegetation rehabilitation as soon as practicable to reduce open areas
- Scheduling topsoil stripping to avoid periods of high winds from unfavourable directions relative to sensitive receptors (including George Grey Drive and Utcha Well Nature Reserve).
- Cease/suspending topsoil stripping operations during high wind conditions where there is a risk of dust affecting sensitive receptors.
- Dust suppressant applied proactively to overburden/topsoil stockpiles.
- Dust suppressant reapply proactively subject to visual inspection and weather forecasting.
- Cease activity causing visible dust lift-off where dust management measures have not prevented dust lift-off, and there is a risk of dust affecting sensitive receptors.

Alternative dust control measures, e.g. hydro-mulching, wind fencing, hard standing or chemical dust suppressants may be used and shall be investigated on a case by case basis to determine suitability before implementation.

Additional dust management measures for consideration are documented within Appendix A. Additional Dust Control Guidance.

5.3.1. Product Stockpile Management – Narngulu Operations

Release of fugitive from stockpiled material shall be minimised by:

- Operating dust suppression sprinkler system at Geraldton, as per SOP-40
- Keeping stockpile heights to a minimum. A stockpile shall not exceed the height of the top of the cab of the loader (generally 3 m).
- Scheduling of material cartage so that that stockpiling of material can be kept to a minimum
- Shaping stockpiles with a gentle slope to reduce erosion and sedimentation in the surrounding area
- Maintaining surrounding areas so they are kept free of material build up
- Maintaining an even surface around stockpiles, to reduce material spillage from the loader bucket when in operation
- Reducing loader bucket load volume, so that spillage does not occur.

5.3.2. Mid-West Ports GMA Sheds

The following dust management strategies shall be implemented:

- All trucks loads shall be covered, carting material to the Port.
- In the event the product shed is full, the Contractor shall seek authorisation from Mid West Ports Authority to load from outside the shed on commencement of ship loading.
- Sheds that are at capacity shall have the roller doors lowered until ship loading commences.
- Street sweeping contractor shall be engaged by the Contractor to mitigate the garnet outside the shed areas.

Dust Management

Mining Australia Procedure



5.4. Undertake Monitoring

Monitoring activities and frequencies are summarised in the table below:

Monitoring Activity	Description of Monitoring Activity	Frequency	Responsibility
Report Exceedances	Any evidence of dust exceedances shall be reported to the Supervisor / Superintendent to enable it to be rectified.	Throughout operations	All Personnel
Inspection	Dust produced by work areas shall be inspected, and if dust levels could impact sensitive receptors, mitigation measures shall be put in place to reduce impact.	Daily	Supervisor/ Superintendents
Monitoring	Port Gregory (only) - Superintendent/Supervisors shall monitor the weather station located on the monitor in the lunch room.	Daily	Supervisor/ Superintendents
Monitoring	Monitoring of sensitive receptors	Mining in M70/926 between October and May.	Environmental Team

5.5. Report Incident or Complaint

If an incident occurs, or a complaint is received report, this needs to be reported in skytrust.

6. Training and Competency

Role	Type	Requirement
All Personnel	Awareness	Induction covering the requirements of this procedure.
Supervisors/Superintendents	Awareness	Completed Appendix A "Procedure Acknowledgement Form".

7. Supporting Documents

Document No.	Document Title or Information Source
	Environmental Toolbox Topic: Dust
SOP-40	Dust Suppression Sprinkler System

8. Related Documents

Document No.	Document Title or Information Source
HSE-172	Clearing and Ground Disturbance Procedure

9. References

Document No.	Document Title or Information Source
	<i>Environmental Protection Act 1986</i>
	Environmental Protection National Environmental Protection (Ambient Air Quality) Measure
	Environmental Protection (Unauthorised Discharges) Regulations 2004
	A Guideline for the Development and Implementation of a Dust Management Program (2008) Department of Environment and Conservation
	The dust suppression choice (23 May 2012) Mining Australia
	GHD (2020) GMA Garnet Dust and Noise Modelling. Air Quality Assessment.

10. Revision

Date	Revision	Created/ Amended By	Amendment	Approved By (Document Owner)
1/12/2020	A	Steven Petts	Draft Preliminary – Issued for Review	Ross Avard
15/02/2022	B	Steven Petts	Update plan	Ross Avard

Appendix A. Additional Dust Control Guidance

A.1. Factors Influencing the Levels of Dust and other Air Pollutants

The following factors influence the risk associated with dust and other air pollutants and should be considered when planning and undertaking works.

The soil type and properties of a site will have a considerable impact on the amount of dust generated. In general soils with a dominant particle size corresponding to gravel size or larger have less potential of becoming airborne than finer particles such as fine sand, silt and clay. However, soil may comprise a mixture of different soil particles, for example, fine contaminated dust, such as heavy metals, mixed with coarse particles.

An assessment of soil particle size distribution can help to determine the potential for particles to become airborne. As a general guide, particle sizes of 50µm or more tend not to become airborne.

Soil moisture content is also important. Dry or non-wetting soils are more likely to become air borne. A soil profile will also provide information on the different soil layers and their potential for particle lift off.

Sites with a larger exposed area are identified as having a greater dust generating potential.

The longer the project, the greater the dust risk as the potential for exposure increases.

The proximity of a site to sensitive receptors has a significant influence on the dust risk potential of a site. A site that is located close to sensitive receptors, such as, residential housing, children's day care, schools, hospitals, sports fields etc., will generally require more preventative measures compared to a site in an isolated remote location.

The direction of the prevailing winds can also influence the risk potential of a site for dust and other air pollutants. Suppose the prevailing winds (predominant wind direction) are blowing towards sensitive receptors. In that case, the risk potential increases because the sensitive receptors are more likely to be impacted then if the winds are blowing away from the sensitive receptors. The higher the wind speed, the greater the potential for dust lift. Daily and seasonal variation of wind speed and direction should be considered, refer to Appendix A.2.12.

The nature of works to be conducted will affect the dust levels, for example, land clearing and stockpiling may generate more dust than site levelling.

The topography of the Site may influence wind behaviour at the Site, which could impact the dispersion of dust and other air pollutants from the Site.

A.2. Dust Control Measures

A.2.1. Limit Cleared Areas and Maximise Vegetation

Before the commencement of any works and during the operation, as much vegetation as possible should be retained, including patches and strips to minimise dust. This can be done by implementing the following:

- Before any works commence, identify areas of vegetation cover that need to be retained.
- Protect this vegetation by fencing or blocking off from the rest of operations
- In other areas, maintain the original vegetation cover for as long as possible.
- Avoid clearing the entire area at once, instead clear areas as required in stages of the operation.

Retaining original trees, shrubs and grasses is one of the most efficient and effective ways of minimising dust emissions. Even low or sparse scrub can be very effective at dissipating wind velocity at the ground surface, where dust lift off occurs.

The following should be considered:

- Retain as much existing vegetation as possible

- If an area needs to be cleared, transplant established plants that must be disturbed to areas that need vegetation
- If trees and plants must be removed and it is not possible for them to be replanted, consider chipping and using the material as mulch – the advantage is that reseedling of original vegetation can occur. Where possible, restore vegetation that is native to the area to maximise plant success and improve environmental conditions.

A.2.2. Timing of Development and Development Staging

Activities with high dust-causing potential, such as topsoil stripping, should not be carried out near sensitive receptors during adverse wind conditions. When necessary, topsoil should be stripped in discrete sections, allowing buffer strips (windbreaks) between clearings.

Dust generated by bulk earthworks being performed during the summer months, particularly with sensitive receptors in proximity, can adversely impact the community/environment.

When planning the staging of developments, the impact on personnel including but not limited to the camp, offices, crib rooms and work areas should be taken into account in relation to the cleared areas and the prevailing winds.

A.2.3. Wind Barriers

Having appropriate wind barriers can be an effective measure for the control of dust over short distances. Wind barriers provide a positive visual impact and offer a protection against the movement and impact of dust on nearby land users. Wind barriers should be considered before commencement of works and when it is apparent that one is required during the next phase of the operation. Consider the following options when placing barriers to prevent dust emissions:

- Wind barriers are most effective when placed perpendicular to the direction of the prevailing wind but will have little or no effect when the wind direction is parallel.
- When choosing wind barriers, it has been observed that solid barriers provide significant reductions in wind velocity for relatively short leeward distances, whereas porous barriers provide smaller reductions in velocity for more extended distances.
- Wind barriers should be at least two metres high.

Windbreaks are barriers designed to slow the speed and redirect the flow of wind. These are not widely used but may be useful in some locations. Effective windbreaks do not stop the wind but break its forward movement, to slow it down. Good windbreaks will not create excessive turbulence or wind eddies.

Windbreak materials may include fences, berms and plants. Windbreaks are most useful when designed for specific wind directions. The effective zone of protection created by a windbreak is approximately 25 times its height, although maximum-protection wind reduction occurs in a range of five to eight times the height of the screen.

A.2.4. Earthmoving Management

Earth-moving activities have the potential to generate large amounts of dust. Planning earth-moving activities particularly at the start of an operation can reduce dust emissions by limiting the time the area is exposed. Options for dust control can include the following:

- Plan earth-moving so they are completed just prior to the time they are needed to limit the length of time ground is exposed
- Observe weather conditions and do not commence or continue earth moving if conditions are unsuitable e.g. under conditions of strong winds.
- Reduce off-site hauling via balanced cut and fill operations
- Pre-water areas to be disturbed.

A.2.5. Management of Material Stockpiles

Material stockpiles can generate large amounts of dust. Fine materials stored in stockpiles can be subject to dust pick-up. Materials being loaded onto conveyor belts or into trucks, rail cars or marine vessels are also potential sources of dust emissions. Dust emissions from material stockpiles can be minimised using the following processes:

- Locate stockpiles in sheltered areas where possible. Alternatively, stockpiles may be covered.
- Where stockpiles are located in open areas, limit the height and slope of the stockpiles to reduce wind pick up, orient stockpiles lengthwise into the wind so they offer the minimum cross-sectional area to prevailing winds, install wind barriers on three sides of the stockpile.
- Limit activity to the downwind side of the stockpile
- Limit drop heights from loading facilities and use closed conveyors where possible. Transfer points should also be minimised. Sprinkler systems could also be used on conveyor systems. Alternatively, dust collection systems, such as, cartridge or baghouse systems could be used instead of sprinklers, where moisture is of concern, for example, with mineral concentrates.

A.2.6. Watering Road

Moisture in the surface of dirt roads causes particles to stick together. The moisture content of dirt roads can be increased by watering the road surface. Depending on weather conditions, a single watering may be effective for hours. When water is applied alone, it provides a short-term reduction in dust. Regular, light watering is better than less frequent, heavy watering.

Watering assists with reducing dust lift off from roads and other traffic areas and during earthworks, to controlling dust during movement of materials such as loading/offloading and transportation of materials.

Watering is a very effective short-term measure; however, its efficiency decreases as wind velocity and evaporation rate increase. Dust emissions can be minimised using the following watering processes:

- The surface should be dampened to prevent dust from becoming airborne but should not be wet to the extent of producing run-off. Alternatively, wetting agents could be used, particularly for non-wetting soils.
- Watering is more effective when undertaken prior to strong breezes
- Use watering sprays on materials to be loaded and during loading.

The use of scheme water should be discouraged, and alternative supplies used whenever possible. However, care must be taken to ensure that the quality of water will not have adverse environmental impacts.

Real time automated response systems to turn on water cannon systems in response to dust levels or high wind speeds can be used. These can help save water by only turning on water cannons during the required conditions and help to reduce the possibility of operator error.

A.2.7. Reducing the Traffic and Speed

Vehicles travelling on unpaved roads stir up dust, reducing the number of vehicles or number of vehicle movements can reduce dust. Traffic can be reduced by restricting vehicle weight or type, ensuring vehicles are utilised with maximum passengers (as opposed to one car per person), or by limiting motor vehicle access to dirt roads.

Fast moving vehicles stir up dust. Studies show that particulate matter 10 micrometres or less in diameter (PM10) goes up with vehicle speed. Reducing speed from 65 kilometres per hour (km/h) to 30 km/h reduces dust emissions by 65%. Speed limit signs and enforcement can reduce speeds. Drainage channels across roads and speed bumps can reduce speeds. Speed bumps and drainage will only reduce dust on roadways, not the surrounds.

A.2.8. Improving Road Design

Good road drainage can reduce dust. If a road surface has poor drainage, puddles will form. Water floats the fine particles. With traffic, water and wind spreads the fines as mud or dust. Standing water next to a road may saturate the roadbed, resulting in potholes. When the fines are washed away, or blown away, the larger particles are left unanchored. These larger particles are pushed to the side of the road, resulting in a need for expensive road resurfacing. If a road is treated with a dust suppressant, the performance of the suppressant depends on the road shedding water and having a smooth driving surface.

A.2.9. Hydromulch

Hydromulch is a very effective measure for preventing dust lift-off from areas where bulk earthworks have been completed and little or no further vehicular or traffic is likely. It is a versatile tool, as the constituents of spray mulch can be varied to suit the requirements of the user and the project. The following processes for hydromulch can be utilised to reduce dust emissions:

- Vehicular and pedestrian access to treated areas should be restricted to prevent disturbance to the hydromulch layer
- Wind barriers placed in isolated locations or where long-term effectiveness is required to control access and achieve maximum benefit
- For short-term stabilisation, hydromulch without grass seed should be sufficient stabilisation.
- For longer-term stabilisation, hydromulch with grass seed and fertiliser should be included in the spray. Organic stabiliser can also be added to the mix to provide a more stable base for the germination of seeds.

Recommended application rates for hydromulch should be sought from suppliers to ensure that application rates and the constituents of the mulch are appropriate to the task.

A.2.10. Chemical Stabilisation

Chemical stabilisers provide immediate coverage and protection; they are effective in areas that receive little traffic or disturbance. They provide a longer-term solution compared to watering, although it may be necessary for the chemical ingredients to be evaluated about their environmental effects.

Chemical stabilisers work by binding the soil particles together to create an artificial crust on the soil surface that is less prone to disturbance by wind. The following options should be considered when using chemical stabilisers to reduce dust emissions:

- Physical barriers or other methods of preventing traffic access should be used to protect stabilised areas
- The manufacturer's instructions should be followed to optimise performance.

These chemicals fall into several groups, such as petroleum-based, organic nonpetroleum, electrochemical stabilisers, and synthetic polymers.

A.2.11. Covering or Sealing Unpaved Surfaces

Applying gravel to a dirt road surface can reduce dust. Gravel provides a hard surface protecting soils from vehicle wheels. Gravel does not reduce the strength of air currents caused by vehicles themselves, so traffic can still blow loose soil particles into the air. Without a good road base of crushed aggregate, traffic will push surface gravel down into the ground, especially when the road is wet. If the road surface does not have enough fine material to cement the surface gravel in place, traffic will push the gravel away from the driving lanes.

To be effective over a long period of time, new gravel must be anchored to the road surface. This is done through incorporating gravel with aggregate mixes or soil adhesives. If gravel is lost by being pressed into soils beneath the road, then the use of geotextile fabrics may be necessary. These fabrics are constructed of

polymer threads with very high tensile strength and are available in designs that either form water barriers or allow water, but not fine soil, to migrate through.

Paving or bituminising is the most effective (and most expensive) method to control dust from unpaved roads. Asphalt and Portland concrete provide durable and effective surfaces that prevent the breakdown of soil surfaces. Paved roads may still accumulate dust as vehicles enter from unpaved roads.

A.2.12. Wind monitoring

Port Gregory Mine Site

Dust modelling undertaken by GHD (2020) shows the Port Gregory Mine Site is prone to dust lift-off when wind speeds exceed 5.5 m/s (30 to 39 km/hr). Under these conditions, wind erosion will be a high risk for dust emissions (GHD 2020). Wind directions that arcs between 45 and 180 degrees are likely to impact sensitive receptors. The weatherstation is displayed on the SCADA under the FPP and the borefield page. This shall be reviewed before undertaken activities such as clearing or planning for mining activities.




Figure 1 Geraldton Windrose

Geraldton Site

The dust lift-off threshold applicable to the Geraldton Site, are wind speeds greater than 5 to 6 m/s (30 to 39 km/hr). Under these conditions dust is likely to lead to dust breaching the licenced premises boundary.



Appendix B. Procedure Acknowledgement Form

This form shall be completed by personnel who have a responsibility identified in Section 2 Roles and Responsibilities, of this procedure.

I confirm that I have read and am aware of the requirements within this procedure:

PROJECT / FUNCTIONAL AREA	
PROJECT No	

Name	
Signature	
Date	

Return completed form to the Training Department for record keeping.