




Lombadina Campground Biological Assessment



Prepared for Tourism Western Australia

March 2021

Biota
Environmental
Sciences



This document has been prepared to the requirements of the client identified on the cover page and no representation is made to any third party. It may be cited for the purposes of scientific research or other fair use, but it may not be reproduced or distributed to any third party by any physical or electronic means without the express permission of the client for whom it was prepared or Biota Environmental Sciences Pty Ltd.

This report has been designed for double-sided printing. Hard copies supplied by Biota are printed on recycled paper.

Lombadina Campground Biological Assessment

Contents

1.0	Executive Summary	9
2.0	Introduction	11
2.1	Background	11
2.2	Scope of the Study	11
3.0	Regional Context of the Study Area	13
3.1	IBRA Bioregion and Subregion	13
3.2	Land Systems	13
3.3	Surface Geology, Soils and Hydrology	13
3.4	Beard's Vegetation Units	15
3.5	Conservation Reserves and Protected Areas in the Locality	15
4.0	Methodology	17
4.1	Desktop Study	17
4.2	Field Survey	19
4.3	Specimen Identification, Nomenclature and Data	21
4.4	Limitations	22
5.0	Desktop Study Results	25
5.1	Vegetation and Flora	25
5.2	Terrestrial Vertebrate Fauna	28
6.0	Vegetation Results	35
6.1	Description of the Vegetation	35
6.2	Vegetation Condition	38
6.3	Vegetation of Significance	38
7.0	Flora Results	39
7.1	Overview	39
7.2	Flora of Significance	39
7.3	Introduced Flora	39
8.0	Fauna Results	45
8.1	Fauna Habitats	45
8.2	Field Survey Records	45
8.3	Likelihood of Significant Fauna	45
9.0	Conclusions	47
9.1	Vegetation of Significance	47
9.2	Flora of Significance	47
9.3	Introduced Flora	47
9.4	Fauna of Significance	47
9.5	Assessment Against the Ten Clearing Principles	47
10.0	References	51

Appendix 1

Framework for Conservation Significance Ranking of Species and Communities in WA

Appendix 2

NatureMap Database Search Results

Appendix 3

Atlas of Living Australia Database Search Results

Appendix 4

EPBC Act Protected Matters Database Search Results

Appendix 5

Significant Flora Species Recorded within 40 km and their Likelihood of Occurrence

Appendix 6

Significant Fauna Species Recorded within 40 km and their Likelihood of Occurrence

Appendix 7

Vegetation Structural Classes and Condition Scale

Appendix 8

Survey Effort Map

Appendix 9

Vascular Flora Species List

Appendix 10

Potential Fauna Species List

Appendix 11

Weed Locations

Tables

Table 3.1:	Geological and soil units occurring within the study area.	13
Table 3.2:	Beard's (1979) vegetation mapping unit occurring within the study area and pre-European extent in the subregion (statistics from Government of Western Australia 2019).	15
Table 4.1:	Biota's likelihood ranking system for species that may occur in the study area.	18
Table 4.2:	Summary of survey team qualifications.	19
Table 4.3:	Potential constraints and limitations of the Lombadina biological assessment.	22
Table 5.1:	Overview of vertebrate fauna species with the potential to occur in the study area.	28
Table 5.2:	Invertebrate fauna records from database searches.	32
Table 7.1:	Dominant native families and genera recorded from the study area.	39
Table 7.2:	Weed species recorded from the study area, including WPP rankings.	41
Table 7.3:	Planted (introduced species) recorded in the study area	43
Table 8.1:	Bird species opportunistically observed in the Lombadina study area.	45

Table 10.1: Assessment of remnant native vegetation within the study area against the ten clearing principles.	48
Figures	
Figure 2.1: Location of the proposed Lombadina campground.	12
Figure 3.1: Geological and soil units occurring within the study area and surrounds.	14
Figure 4.1: Total monthly rainfall at Cygnet Bay weather station (#3057) for the 12 months prior to the field survey, compared to the long-term monthly median rainfall.	20
Figure 5.1: Records of TECs and PECs located within 40 km of the study area.	26
Figure 5.2: Historical Priority flora records within 40 km of the study area.	27
Figure 5.3: Historical records of significant mammals and reptiles within 40 km of the study area.	30
Figure 5.4: Historical records of significant birds within 40 km of the study area.	31
Figure 5.5: Historical locations of potential SRE invertebrate fauna recorded from the locality of the study area.	34
Figure 6.1: Vegetation of the Lombadina study area.	36
Figure 6.2: Vegetation condition ranking of the Lombadina study area.	37

This page intentionally blank.

1.0 Executive Summary

Tourism Western Australia is investigating the development of a campground at Lombadina, 175 km northeast of Broome on the Dampier Peninsula, in the Kimberley region of Western Australia. The proposed site is located south of the Lombadina Aboriginal Community and covers an area of 12.35 ha. Biota Environmental Sciences (Biota) was commissioned to undertake a biological assessment of the proposed campground site. The objective of the assessment was to act as a preliminary risk assessment of any potential key issues identified for the study area, and to gather biological information to support a Native Vegetation Clearing Permit (NVCP) application. This was achieved through a desktop study of existing information and data, and a field survey by a botanist on 10 February 2021.

The desktop study was undertaken to identify features of significance known from the study area or the broader locality (within 40 km). The review considered regional information, previous biological surveys in the locality, and the results of various database searches. The results of the desktop study were used to assist in the compilation of a list of flora and fauna species of significance potentially occurring in the study area.

A total of 66 native vascular flora species from 57 genera and 33 families were recorded from the study area during the field survey, all of which are typical of the locality. This number of species is in the range expected for a survey area of this size in this location. Fifteen weed species were recorded including **Jatropha gossypifolia* (Bellyache Bush), which is a Weed of National Significance and listed as a declared pest under the *WA Biosecurity and Agriculture Management Act 2007* and **Azadirachta indica* (Neem Tree) which is also listed as a declared pest. No Threatened flora species were recorded or would be expected to occur. No Priority flora were recorded, and none would be likely to occur within the study area.

One Threatened Ecological Community (TEC), "Monsoon (vine) thickets on coastal sand dunes of Dampier Peninsula", is known to occur in the vicinity of the study area. Two Priority Ecological Communities (PECs) also occur within 40 km of the study area. However, the survey work confirmed that none of these communities were present in the study area.

One fauna habitat was recorded for the study area, *Eucalyptus/Corymbia* woodlands over tussock grassland on orange Pindan sands. This habitat does not represent core habitat for any listed species of significance. Eight bird species were recorded in the study area during the survey, none of which are significant.

Database and literature searches yielded a total of 330 vertebrate fauna species with the potential to occur in the study area, comprising five amphibian species including one introduced amphibian, 53 reptile species including one introduced reptile species, eight native ground-dwelling mammal species, five introduced mammal species, 16 bat species and 243 avifauna species. Of these, 107 are significant species, of which 26 have been considered directly in this report following the exclusion of marine-listed and coastal bird species. The database searches returned 104 invertebrate specimens belonging to three groups known to exhibit short-range endemism (land snails, mygalomorph spiders and scorpions). None were of elevated significance.

Following the field survey, 10 species of fauna of significance were ranked as either 'likely to' or 'may occur' in the study area. However, the small extent of the study area and the wider distributions of the species suggest it would be unlikely that significant impacts on these species would arise as a result of the camp site development, should they be present.

Based on the information available, the proposed clearing is unlikely to be at variance with any of the 10 clearing principles under Schedule 5 of the *Environmental Protection Act 1986*.

This page is intentionally left blank.

2.0 Introduction

2.1 Background

Tourism Western Australia is investigating the development of a campground at Lombadina, 175 km northeast of Broome on the Dampier Peninsula, in the Kimberley region of WA (Figure 2.1). The proposed site (hereafter referred to as the study area) is located south of the Lombadina Aboriginal Community and covers an area of 12.35 ha (Figure 2.1).

Biota was commissioned to undertake a desktop study followed by a biological assessment to identify vegetation, flora and fauna values of the Lombadina study area.

2.2 Scope of the Study

The main objective of the study was to determine if there are any significant flora or fauna species, or vegetation types, that could represent constraints to the potential development of the area, or require specific management. This was achieved through an initial desktop study, a field survey, and preparation of this report to support a Native Vegetation Clearing Permit (NVCP) application.

Specifically, the scope of the study was to:

- review and discuss existing information from the vicinity of the study area, including literature describing previous surveys completed in the locality, to establish the biogeographical context;
- review and discuss the results of relevant database searches to determine fauna, flora and vegetation communities of significance that may occur in the study area; specifically Threatened or Priority listed flora or fauna species, and Threatened Ecological Communities (TECs), Priority Ecological Communities (PECs) or other Environmentally Sensitive Areas (ESAs);
- describe, photograph and map the dominant vegetation units occurring within the study area (including a description of dominant species, structure and vegetation condition, and discussion of their representation in a regional context);
- identify any vegetation units of particular significance within the study area;
- compile a list of vascular flora species recorded in the study area;
- record and photograph flora of significance, including Threatened and Priority species and any other species of interest;
- record any introduced flora species (weeds) occurring in the study area;
- record any opportunistic sightings of fauna whilst conducting flora and vegetation assessments; and
- use the above information to prepare an assessment of the Proposal against the 10 clearing principles, as required for an NVCP application.

The approach and methodology used for the biological field assessment was conducted in accordance with the following:

1. Technical Guidance: Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment (EPA 2020)¹;
2. Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment (EPA 2016c);
3. A guide to the assessment of applications to clear native vegetation under Part V Division 2 of the Environmental Protection Act 1986 (Department of Environment Regulation 2014); and
4. Technical Guidance: Sampling of Short Range Endemic Invertebrate Fauna (Department of Environment Regulation 2014).

The field survey essentially comprised a “reconnaissance” flora survey and a “basic” fauna survey as per EPA (EPA 2016c, 2020).

¹ EPA (2020) Technical Guidance has recently superseded Technical Guidance: Terrestrial Fauna Surveys (EPA 2016a) and Technical Guidance: Sampling Methods for Terrestrial Vertebrate Fauna (EPA 2016b).




Figure 2.1: Location of the proposed Lombadina campground.

3.0 Regional Context of the Study Area

3.1 IBRA Bioregion and Subregion

The Interim Biogeographic Regionalisation for Australia (IBRA) recognises 89 bioregions and 419 subregions within Australia (DSEWPaC 2012). The study area lies within the Dampierland bioregion and the Pindanland subregion, which covers 5,198,904 ha and is described as follows:

*"The Pindanland subregion (DL2) is the coastal, semi-arid, northwestern margin of the Canning Basin. The region has a semi-arid, hot, tropical climate with summer rainfall. Quaternary sandplains overlay Jurassic and Mesozoic sandstones and support Pindan vegetation on the plains and hummock grasslands on hills. Quaternary marine deposits on coastal plains support mangal, samphire, *Sporobolus* grasslands, *Melaleuca alsophila* low forests, and *Spinifex-Crotalaria* strand communities" (Graham 2003).*

3.2 Land Systems

Land systems mapping covering the study area has been prepared by the WA Department of Agriculture (Payne and Schoknecht 2011). The study area falls entirely within the Yeeda land system, which is widespread and well represented through the Pindanland subregion of the Dampierland bioregion. The Yeeda land system is described as "Sandplains with red and yellow sands supporting pindan Acacia shrublands with emergent Eucalypt trees" (Payne and Schoknecht 2011).

3.3 Surface Geology, Soils and Hydrology

The study area occurs within the Canning Basin, the largest sedimentary basin in WA, which consists predominantly of Palaeozoic sedimentary rocks with a thin Mesozoic and Tertiary cover (Paul et al. 2013). The study area encompasses one geological surface unit, Czs, which is the substrate underlying the habitats present within the study area. This unit was mapped at a scale of 1:250,000 by the Geological Survey of Western Australia (2011), (Table 3.1 and Figure 3.1).

The soils of the study area have been mapped, and are available from the Australian Soil Resource Information System (ASRIS) (CSIRO 2014). Two broad soil units are represented in the study area (Table 3.1 and Figure 3.1).

The surface water catchment of the study area is the Cape Leveque Coast Basin of the Western Plateau Division. Drainage within this catchment area is towards the coast, while recharge to the aquifer is via rainfall infiltrating the unconfined Canning Basin (Paul et al. 2013). Sheet flooding is the most widespread pattern of drainage on the Dampier Peninsula, as the gently sloping plains are interrupted by few abrupt rises (McKenzie and Kenneally 1983). No major watercourses or drainage lines occur within the study area.

Table 3.1: Geological and soil units occurring within the study area.

	Unit	Description	Extent within Study Area (ha)
Geology	Czs	Sand or gravel plains; quartz sand sheets commonly with ferruginous pisoliths or pebbles, minor clay; local calcrete, laterite, silcrete, silt, clay, alluvium, colluvium and aeolian sand.	12.35
Soils	AB21	Pindan country: gently undulating sand plain with a few small rocky sandstone residuals; no external drainage: chief soils are red earthy sands and hummocks of siliceous sands.	4.82
	Jw1	Low-lying coastal plains with some sand dunes: chief soils are saline clays (Uf1.41) on the flat to very gently sloping plains. Associated are (Ug5) and (Uf) soils along the inland margin of the plains; areas of saline muds (Um1) on slopes and flats submerged at high tide; and very small areas of calcareous sands (Uc1.1) and/or siliceous sands (Uc1.2) on coastal dunes.	7.53



Figure 3.1: Geological and soil units occurring within the study area and surrounds.

3.4 Beard's Vegetation Units

The vegetation of the Kimberley region was mapped by John Beard (1979) at a scale of 1:1,000,000. The study area lies within the Dampier Botanical District, which broadly corresponds with the Dampierland IBRA bioregion. The district is characterised by pindan vegetation on sandplains, more or less densely wooded according to rainfall; tall grass savanna with or without scattered trees on clay plains; spinifex steppe on sandstone, and limestone outcrops (Beard 1979). The study area occurs in one of Beard's (1979) vegetation units, Dampierland 750 (Table 3.2), described as "*Pindan woodland; Acacia thicket with eucalypt woodland over spinifex; Acacia tumida, Eucalyptus tectifera, Corymbia grandifolia, Triodia pungens, T. bitextura*".

Table 3.2: Beard's (1979) vegetation mapping unit occurring within the study area and pre-European extent in the subregion (statistics from Government of Western Australia 2019).

Beard's Vegetation Mapping Unit	Pre-European Extent within Pindanland Subregion (ha)	Current Extent within Pindanland Subregion (ha)	Extent within Study Area (ha)	Percentage (%) of Pindanland Extent Occurring in the Study Area
Dampierland 750	1,221,727	1,219,556	12.35	< 0.0001

3.5 Conservation Reserves and Protected Areas in the Locality

One conservation reserve occurs within 40 km of the study area; the Swan Island Nature Reserve is located 23.7 km northeast. The study area is also located within the Bardi Jawi Indigenous Protected Area.

This page intentionally blank.

4.0 Methodology

To meet the objectives of this biological assessment an initial desktop study was undertaken (Section 4.1) followed by a field survey (Section 4.2).

4.1 Desktop Study

The desktop study was undertaken to identify features of significance known from the study area or the broader locality (within 40 km). Appendix 1 contains more information regarding the framework for conservation significance ranking of communities and species in WA. The desktop study was also used to assess the level of biological survey work that had been completed previously in the study area.

The review considered regional information, previous biological surveys in the locality, and the results of various database searches (see Appendices 2 to 4), as discussed in the following sections. The results of the desktop study were used to assist in the compilation of a list of flora species of significance potentially occurring in the study area (Appendix 5), as well as lists of fauna, including species of significance, potentially occurring in the study area (Appendix 6).

4.1.1 Database Searches

The following databases were searched for records of fauna, flora and vegetation of significance previously recorded from the study area, or known to occur in the locality²:

1. NatureMap database (<http://NatureMap.dbca.wa.gov.au>): a joint project of the Department of Biodiversity, Conservation and Attractions (DBCA) and the Western Australian Museum (WAM). This database represents the most comprehensive source of information on the distribution of Western Australia's flora and fauna, comprising records from the Fauna Survey Returns database, the WA Threatened Flora and Fauna Databases, the WA Herbarium and WAM Specimen databases, and the BirdLife Australia Atlas. The database search was centered on the point -16.4680, 122.9755 and requested the return of records from a 40 km radius (Appendix 2). The database was queried on 29/10/2020.
2. The DBCA databases of Threatened Ecological Communities (TECs) and Priority Ecological Communities (PECs), Declared Rare and Priority Flora, and Threatened Fauna. These data searches requested the return of records from a 40 km radius around the point -16.4680, 122.9755. The databases were queried on 23/10/2020 but results are not appended here, consistent with normal DBCA data management expectations.
3. The Atlas of Living Australia (ALA) (<http://www.ala.org.au>): a joint project between academic collecting institutions, private individual collectors and community groups. The atlas contains occurrence records, environmental data, images and the conservation status of species throughout Australia. The database search requested the return of fauna records from a 40 km buffer around the study area boundary and was conducted on 29/10/2020 (Appendix 3).
4. The Commonwealth EPBC Act Protected Matters database. The database search requested the return of records within a 40 km buffer from a central point (-16.4680, 122.9755) (Appendix 4). The database was searched on 29/10/2020.
5. eBird (<https://ebird.org/>) - A citizen science database for bird records run by the Cornell Lab of Ornithology. This database was searched for supplementary records of conservation significant bird species within 40 km of the survey area. The database was searched on 02/03/2021.
6. Biota's internal flora and fauna databases. The search was within 40 km of the study area and was conducted on 23/10/2020.

² The search areas for each database included coastal and marine areas. Mammal and reptile species inhabiting these areas were excluded from consideration. Avifauna species inhabiting marine/coastal areas have been included in the potential species list for completeness, but are excluded from the likelihood of occurrence assessment. Erroneous records of species returned from the database searches (i.e. those that were outside of known ranges or would not occur in Australia) were also excluded.

4.1.2 Review of Previous Biological Surveys in the Locality

Publicly available literature was searched to identify relevant flora and vegetation surveys and fauna surveys conducted in the locality of the study area. This was limited to surveys conducted within 40 km of the study area.

No publicly available vegetation or flora surveys have been conducted within 40 km of the study area, with the exception of a broad survey of vine thickets on the Dampier Peninsula (Black et al. 2010). This survey assessed vine thicket patches from Broome to the tip of the Dampier Peninsula, but did not include vine thickets at One Arm Point and Lombadina-Djarindjin. The desktop study therefore focused on existing mapping of significant communities, and historical records of significant flora held in public databases.

No publicly available fauna surveys have been conducted within 40 km of the study area; results were therefore limited to database search returns only. Two fauna surveys were located within 80 km of the study area: the Irvine Island Seasonal Fauna Survey, located 85 km northeast (Biota 2011); and Beagle Bay Big Tree Country Tropical Timber Plantation Project Fauna Assessment, located 60 km southwest (Ecologia 2004). The next closest area in which a detailed biological survey has been conducted is at James Price Point, located 120 km southwest.

4.1.3 Assessment of Likelihood of Occurrence

To determine which species of significance have the potential to occur in the study area, the results of the database and literature searches were examined, while considering the known habitat preferences and distributions of the species identified. Habitats were defined primarily according to the landforms and vegetation that appear to be present in the study area, based on inspection of aerial imagery and knowledge of the locality.

For each species of significance identified for the locality, a set of rankings and criteria that has been developed by Biota was applied to guide the assessment of the likelihood of occurrence within the study area (Table 4.1).

Table 4.1: Biota's likelihood ranking system for species that may occur in the study area.

Rank	Criteria
Recorded	1. The species has been recorded in the study area.
Likely to occur	1. There are existing records of the species in close proximity to the study area (within 20 km); and <ul style="list-style-type: none"> • the species is strongly linked to a specific habitat, which is present in the study area; or • the species has more general habitat preferences, and suitable habitat is present.
May occur	1. There are existing records of the species from the locality (within 40 km), however: <ul style="list-style-type: none"> • the species is strongly linked to a specific habitat, of which only a small amount is present in the study area; or • the species has more general habitat preferences, but only some suitable habitat is present. 2. There is suitable habitat in the study area, but the species is recorded infrequently in the locality.
Unlikely to occur	1. The species is linked to a specific habitat, which is absent from the study area; or 2. Suitable habitat is present, however there are no existing records of the species from the locality despite reasonable previous search effort in suitable habitat; or 3. There is some suitable habitat in the study area, however the species is very infrequently recorded in the locality or the only records are historic (>40 years ago).
Would not occur	1. The species is strongly linked to a specific habitat, which is absent from the study area; or 2. The species' range is very restricted and does not include the study area; or 3. The species is not considered extant in the locality.

Two rankings have been provided:

1. An initial assessment was made during the desktop study (see Appendices 5 and 6). This was based on consideration of the overall distribution of the species, the proximity of the study area

to known populations, the reliability and age of any historic records, and, if the species was known to be linked to particular habitats, whether suitable habitat appeared to be present in the study area based on inspection of aerial photography and/or existing information.

- The likelihood rankings were subsequently revised as necessary based on the findings of the field survey (see Appendices 5 and 6). Where the initial and final likelihood rankings were different, the reason was provided.

4.2 Field Survey

4.2.1 Survey Team

The field survey was undertaken on 10 February 2021 by Tim Willing, a sub-contracted botanist to Biota. Tim has extensive vegetation and flora survey experience in the Kimberley, and is also experienced with identifying fauna habitats and significant vertebrate fauna species. His qualifications are summarised in Table 4.2.

Table 4.2: Summary of survey team qualifications.

Name	Position at Biota	Qualification	Years of Survey Experience	Survey Role
Tim Willing	Principal Botanist (Kimberley specialist)	BA	35	- Vegetation mapping - Rare flora searches - Fauna assessments

The assistance of Darrell Sibosado and family at Lombadina Community Office in completing the field survey is also gratefully acknowledged.

4.2.2 Survey Timing and Conditions

The weather conditions (particularly rainfall) leading up to a field survey are important factors influencing the number and type of species that are recorded from an area, particularly for flora. As an example, more annual species are likely to be present following high rainfall, particularly when this occurs over the summer period, however annual daisies (family Asteraceae) germinate mainly following winter rainfall. Plants are also more likely to bear reproductive material (flowers and/or fruit) following periods of higher rainfall, which means that specimens can be more easily identified. To characterise rainfall leading up to the current survey, total monthly rainfall data for 2020/2021 were sourced from the Bureau of Meteorology's Cygnet Bay weather station³, located approximately 7 km southwest of the study area. Data from the 12 months preceding the survey were compared to the monthly median rainfall for the years 1964-2021 from the Cygnet Bay weather station (Figure 4.1).

The data shows that the rainfall received in the three months prior to the field survey (November 2020 - January 2021) was greater than three-times the long-term median for the same period (928.0 mm compared to 247.7 mm). This was due to well above average rainfall in December 2020, due to several tropical lows.

Weather conditions at the time of field survey were therefore optimal for a post-wet season biological survey. Taking the recent rainfall data into consideration, the conditions at the time of the survey were also considered optimal for the collection of annual and cryptic perennial flora species.

³ Station 3057; sourced from <http://bom.gov.au>




Figure 4.1: Total monthly rainfall at Cygnet Bay weather station (#3057) for the 12 months prior to the field survey, compared to the long-term monthly median rainfall.

4.2.3 Floristic Data Collection

No floristic sampling sites (quadrats or relevés) were assessed, however the vegetation present was described through several mapping notes, where the following parameters were recorded:

1. Location: MGA coordinates were recorded in WGS84 datum using a hand-held Global Positioning System (GPS) unit;
2. Habitat: A description of the landform and habitat;
3. Vegetation Description: A broad description based on the height and estimated cover of dominant species after Aplin's (1979) modification of the vegetation classification system of Specht (1970) (see Appendix 7);
4. Associated Species: A list of non-dominant species within the area;
5. Fire History: An estimate of time since last fire;
6. Soil: A broad description of the soil surface and stony surface mantle;
7. Disturbance Details: Vegetation condition was ranked according to the scale developed by Trudgen (1988), considering evidence of grazing, physical disturbance, weed invasion etc. (see Appendix 7); and
8. Photograph: A representative digital photograph of the vegetation was taken.

A total of 55 mapping notes were taken in the study area during the survey. The location of these, and the GPS track log showing foot traverses, is presented in Appendix 8.

4.2.4 Vegetation Description and Mapping

The scale of vegetation mapping can be influenced by a range of factors including spatial characteristics of the survey area (e.g. the size and variety of habitats present), and other factors such as the scope of the survey and the availability of current, high quality aerial photography. The vegetation units for this study were described at the sub-association level (level VI as per the

National Vegetation Information System; NVIS)⁴, which is the most detailed level under the NVIS. This level of detail would be considered fine-scale (intra-locality) delineation of vegetation units as per EPA (2016c).

Vegetation maps were created and consolidated using Geographical Information System (GIS) software (QGIS and MapInfo Professional), and point locations of mapping note sites, significant flora and weeds were added. All maps in this report were produced by Melissa Robinson (GIS Cartographer at Biota) using MapInfo Professional (version 11).

4.2.5 Searches for Threatened Flora, Priority Flora and Weeds

Targeted, non-systematic searches for significant species were conducted on foot in the entire study area, given its small size. The routes of the foot traverses intersected all major vegetation/habitat units in the study area.

Locations of species of significance, introduced flora species and/or unknown taxa were recorded using a handheld GPS unit (WGS84 datum). The number of individuals and extent of the population were also recorded for each location, along with the habitat and associated species.

4.2.6 Fauna Habitat Assessment

Habitat assessments made in the field were based primarily on the descriptions of landforms and habitat features and were aided by the vegetation descriptions, photographs and aerial imagery.

It is important to note that each broad habitat area defined here cannot be used to map the distribution of any one species or group of taxa, as many species utilise a range of ecological niches for specific activities such as foraging, commuting, breeding and nesting. The resultant habitat map may therefore be viewed as a guide to delineate areas that may be of differing ecological importance to the fauna species utilising the study area.

4.2.7 Fauna Sampling

Non-systematic methods were employed to search for fauna, or secondary evidence thereof, during foot traverses of the study area. The aim was to locate, describe and search habitats that may support fauna of significance, habitat-specific species and other species of interest with the potential to occur based on the results of the desktop study. A range of non-systematic techniques were used, including:

- foot traverses to record evidence of terrestrial vertebrate species of significance;
- identification and recording of secondary signs including tracks, scats and diggings; and
- recording of opportunistic sightings and calls.

4.3 Specimen Identification, Nomenclature and Data

4.3.1 Flora

Common species that were well known to the survey botanist were confirmed in the field. A voucher specimen was collected if the specimen was either difficult to determine without closer examination, belonged to a recognised species complex, was poorly collected or otherwise unusual, or was in very good condition (healthy specimens with flowers and/or fruits are often useful to submit to the WA Herbarium).

Voucher specimens were identified using flora keys, consulting appropriate publications and checking reference collections. The survey botanist identified all specimens.

⁴ <http://www.environment.gov.au/erin/nvis/publications/avam/section-2-1.htm>

A flora species list is provided in Appendix 9. Nomenclature and conservation significance rankings used in this report are consistent with the current listing of WA flora recognised by the WA Herbarium on FloraBase⁵ at the time of preparation of this report.

4.3.2 Fauna

Biota Zoologist, John Graff, reviewed all data relating to fauna observations.

As per the Technical Guidance (EPA 2020), species nomenclature for reptiles, amphibians and mammals follows that of the *Western Australian Museum Checklist of the Vertebrates of Western Australia*, last published in September 2017. Species nomenclature for birds follows that of International Ornithological Congress (Gill et al. 2020), as this is regularly updated and is one of the three major global taxonomies currently in use. It also follows a similar approach to species concepts as the WA Museum reptiles, amphibians and mammals.

4.4 Limitations

As per the EPA's *Technical Guidance for Flora and Vegetation Surveys for Environmental Impact Assessment* (EPA 2016c) and *Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment* (EPA 2020), potential constraints or limitations for this study are addressed in Table 4.3.

Table 4.3: Potential constraints and limitations of the Lombadina biological assessment.

Potential Constraint	Statement of Limitations
1. Sources of information	<ul style="list-style-type: none"> The broader Lombadina locality has not been well surveyed, with no past reports publicly available to be considered as part of the desktop study. However, publicly available databases of rare species and community information were searched, as was the Biota internal database. Regional and local level information is not considered to be a limiting factor for this study.
2. Survey scope	<ul style="list-style-type: none"> The survey objective was to provide the necessary information required to support an NVCP application. Given the size of the study area and the scale of the proposed clearing, a desktop study followed by a reconnaissance flora and basic fauna survey as per EPA (2016c, 2020) was considered appropriate, with some targeted sampling also conducted. No systematic trapping for fauna was performed. This is consistent with a basic fauna survey (EPA 2020).
3. Proportion of flora / fauna collected and identified	<ul style="list-style-type: none"> All vascular flora encountered in the study area were recorded. A total of 66 vascular flora species from 33 families were recorded during the current survey. The basic fauna survey focused on recording significant species from secondary evidence, and did not attempt to record all species (this would require systematic trapping as part of a detailed survey).
4. Completeness of survey	<ul style="list-style-type: none"> The study area was surveyed to a satisfactory level to support an NVCP application. As for any study, additional survey effort would lead to additional species being recorded.
5. Mapping reliability	<ul style="list-style-type: none"> Vegetation units and fauna habitats were described and mapped based on data collected during systematic and targeted foot traverses throughout the study area. The mapping is considered to provide a reliable indication of the vegetation units and fauna habitats in the study area.
6. Timing, weather, season, cycle	<ul style="list-style-type: none"> The survey intensity was adequate for recording the annual, ephemeral and cryptic perennial flora present at the time, and the list of vascular flora documented from the study area is comprehensive for an area of this size (12.35 ha). The survey was conducted at the end of the wet season following above average rainfall and timing was considered optimal for the collection of most flora species.

⁵ <http://florabase.dpaw.wa.gov.au>

Potential Constraint	Statement of Limitations
7. Disturbances	<ul style="list-style-type: none"> There were no disruptions during the survey and it was undertaken as planned.
8. Intensity of survey	<ul style="list-style-type: none"> A reconnaissance/basic survey with targeted sampling was considered adequate to address the requirements for an NVCP application.
9. Resources and experience levels	<ul style="list-style-type: none"> Sufficient time was allocated to the field survey component (a total of 1 person days). The botanist undertaking the survey was suitably qualified and experienced to identify flora and fauna. Additional assistance was sought as required from other personnel at Biota. There were therefore no limitations due to resourcing.
10. Access issues	<ul style="list-style-type: none"> The entirety of the study area could be accessed and surveyed at an appropriate level. There were therefore no limitations due to access issues.

It should also be noted that in relation to vertebrate fauna, only terrestrial fauna species were included in the vertebrate potential species list; species of marine mammals and reptiles were omitted. Birds that rely upon marine, wetland or coastal habitats, including seabirds, wading birds, shorebirds and waterbirds, were included in the potential species list for completeness due to the proximity of marine and coastal habitats to the study area, but were excluded from the likelihood of occurrence assessment (Appendix 6).

This page intentionally blank.

5.0 Desktop Study Results

5.1 Vegetation and Flora

5.1.1 Threatened and Priority Ecological Communities

TECs are defined as biological assemblages that naturally occur in a particular type of habitat, that fit one of the following categories: "Presumed totally destroyed", "Critically Endangered", "Endangered" or "Vulnerable" (DBCAs 2013).

One TEC, the "*Monsoon (vine) thickets on coastal sand dunes of Dampier Peninsula*", is known to occur in the vicinity of the study area. This TEC is listed as Vulnerable at State level under the BC Act and Endangered at Commonwealth level under the EPBC Act. The Vine Thicket TEC is confined to the landward side of coastal dunes, and comprises discontinuous but discrete pockets of dense evergreen and deciduous vine thicket vegetation (DBCAs 2018). The thickets contain many fleshy-fruited plants, which provide an important food resource for native wildlife as well as for Indigenous people (DBCAs 2018).

PECs are defined as possible TECs that do not meet survey criteria or that are not adequately defined. These communities are added to the PEC List under priorities 1, 2 and 3 (DBCAs 2013). Two Priority 3 PECs occur within 40 km of the study area, comprising:

1. *Vegetation Association 37* as defined by John Beard's vegetation mapping for the Kimberley (Beard 1979): Shrublands, teatree thicket. This PEC occurs along the west coast of the Dampier Peninsula, with the closest stand being approximately 3 km southwest of Lombadina (Figure 5.1).
2. *Vegetation Association 67* as defined by John Beard's vegetation mapping for the Kimberley (Beard 1979): Grasslands, tall bunch grass savanna, sparse low tree; ribbon grass and paperbarks. This PEC occurs further south on the Dampier Peninsula, with the closest stand approximately 31 km south of Lombadina (Figure 5.1).

5.1.2 Threatened and Priority Flora

Based on the desktop study, no Threatened species have been recorded from the locality, and none would be expected to occur in the study area.

A total of 10 Priority listed species have been recorded within 40 km (see Figure 5.2 and Figure 5.2). None of these species are considered likely to occur in the study area, however one species may occur: the Priority 3 spinifex *Triodia acutispicula*.

5.1.3 Significant Weed Species

Although databases were not searched for weed species as part of the desktop study, given the location of the study area near a settlement, it is possible that introduced flora listed as Declared Pests or Weeds of National Significance (WONS) may be present.




Figure 5.1: Records of TECs and PECs located within 40 km of the study area.




Figure 5.2: Historical Priority flora records within 40 km of the study area.

5.2 Terrestrial Vertebrate Fauna

5.2.1 Potential Faunal Assemblage

Database and literature searches yielded a total of 330 vertebrate fauna species with the potential to occur in the study area (Table 5.1). This total comprised five amphibian species including one introduced amphibian, 52 reptile species including one introduced reptile species, eight native ground-dwelling mammal species, five introduced mammal species, 16 bat species and 243 avifauna species (Table 5.1). Seventeen of these species are significant species (Table 5.1). A full list of potential fauna species is presented in Appendix 10.

Table 5.1: Overview of vertebrate fauna species with the potential to occur in the study area.

Fauna Group	Number of Species	Number of Conservation Significant Species
Amphibians	4	0
Introduced Amphibians	1	0
Reptiles	52	2
Introduced Reptiles	1	0
Ground-dwelling mammals	8	2
Introduced Mammals	5	0
Bats	16	2
Avifauna	243	101 (20 considered in likelihood of occurrence assessment)
Total	330	107

The four native amphibian species returned in the database searches of the study area comprised three species of tree frog (Pelodyadidae) and one species of burrowing frog (Limnodynastidae). The introduced amphibian species, the Cane Toad, belongs to the family Bufonidae.

The 52 reptile species from the search area comprised 17 species of skink (Scincidae), nine species of gecko (Gekkonidae and Diplodactylidae), six species of front-fanged snake (Elapidae), five species of monitor lizard (Varanidae), five species of dragon (Agamidae), three species of flap-footed lizard (Pygopodidae), three species of python (Pythonidae), two species of blind snake (Typhlopidae), one species of mangrove snake (Homolapsidae), one species of tree snake (Colubridae), and six species of elapid snake (Elapidae). The introduced reptile species was the Flowerpot Blind Snake, in the family Typhlopidae.

The eight species of native ground-dwelling mammals and five introduced mammal species combined comprised four species of rodent (Muridae) (including two introduced rodents), four species of carnivorous marsupial (Dasyuridae), two species of introduced hoofed mammal (Bovidae), one macropod species (Macropodidae), the Bilby (Thylacomyidae), and the introduced Pig (Suidae).

The 16 bat species comprised eight species of evening bat (Vespertilionidae), two species of sheath-tail-bat (Emballonuridae), three species of flying fox (Pteropodidae), one species of Ghost bat (Macrodermatidae), one species of bentwing bat (Minopteridae) and one species of freetail bat (Molossidae).

The 243 bird species returned from the desktop review comprised 66 families, belonging to 21 orders. Twenty-four families represented by 88 species were passerine birds and 42 families, represented by 159 species were non-passerine birds.

5.2.2 Significant Vertebrate Fauna

Native fauna species that are rare, threatened with extinction, or have high conservation value, are specially protected by law under the WA Biodiversity Conservation Act 2016 and/or the Commonwealth EPBC Act. Migratory and Marine species are also protected under the EPBC Act as species of National Environmental Significance. In addition, the DBCA maintains a list of fauna that are deemed a priority for conservation, which have not been assigned statutory protection under the Biodiversity Conservation Act 2016, but are still considered to be of conservation priority, or are considered to be rare but not threatened and are in need of monitoring (DBCA 2020). Appendix 1 details categories of conservation significance recognised under the above frameworks.

A summary of the conservation significant species that were returned for the search area and have potential to occur in the study area is included in Appendix 6, along with an assessment of their likelihood of occurrence. The likelihood of occurrence of each conservation significant species was considered by assessing species habitat preference, potential habitats available within the study area, current known fauna distributions and last known records. Within each fauna group, species are presented in descending order of likelihood to occur. Previous records of significant fauna species are shown in Figure 5.3 and Figure 5.4. Conservation significant species determined not to occur in the study area are not presented.

A total of two reptiles, two ground-dwelling mammals, two bats, and 101 birds are listed as significant species under State or Federal legislation. Of the 101 listed bird species, likelihood assessments were only made for 20 species that had preferred terrestrial habitat potentially available within the study area. The majority of significant bird species were either listed as Marine under the EPBC Act, and were therefore excluded from the likelihood assessment based on the absence of marine habitats in the study area (further details in Appendix 10). Similarly, shorebirds favouring coastal and intertidal environs were not considered in the likelihood assessment. However, these species have been included within the broader potential species list due to the proximity of marine and coastal environs to the study area.




Figure 5.3: Historical records of significant mammals and reptiles within 40 km of the study area.




Figure 5.4: Historical records of significant birds within 40 km of the study area.

5.2.3 Terrestrial Invertebrate Fauna – Short-Range Endemic Invertebrates

Certain terrestrial invertebrate groups that display naturally small distributions (less than 10,000 km²) and are characterised by poor dispersal capabilities, confinement to disjunct habitats and low fecundity, are referred to as Short-range Endemic (SRE) invertebrates (Harvey 2002, Ponder and Colgan 2002). Given this, SRE fauna are more likely to experience population extinctions than more widely distributed taxa, and as such are important in terms of conservation of biodiversity, particularly when considering impact assessment.

The invertebrate taxa that are most likely to contain SRE fauna include (but may not be limited to) land snails, millipedes, scorpions, pseudoscorpions and mygalomorph spiders. The distribution of these taxa are likely to be threatened by processes such as clearing of native vegetation, changes to fire regimes, introduction/spread of weeds and pathogens, habitat fragmentation and changes to surface hydrology (EPA 2009).

The database searches returned 104 invertebrate specimens belonging to three groups known to exhibit short-range endemism (Table 5.2; Appendix 5). The Kimberley region is a relatively understudied area in relation to the majority of the invertebrate groups likely to exhibit short-range endemism. This is not the case for land snails, however, with the Kimberley being recognised as a significant hotspot of land snail diversity since the late 1980s to early 1990s (Gibson and Köhler 2012). This is reflected in the results from the database searches, with 81% of invertebrate records from the database search being land snails (Table 5.2).

Table 5.2: Invertebrate fauna records from database searches.

Invertebrate Group	Number of Records	Number of Species/Taxa
Terrestrial gastropods (land snails)	84	8
Mygalomorph spiders (trapdoor spiders)	8	2
Scorpions	12	2
Total	104	12

5.2.3.1 Land Snails

For the purpose of this assessment, the land snail fauna of the Kimberley region can be categorised as belonging to two groups: camaenid snails (of the family Camaenidae) and the non-camaenids (or micro-snails). The non-camaenids are considered to have relatively broad distributions which encompass northern Australia, extending into Indonesia and Polynesia (Solem 1991, 1997), with only a few that are confined to the Kimberley – Northern Territory region. The camaenids however are considered most likely to support SREs, as some taxa exhibit highly restricted distributions (Solem 1981, 1984, 1991, 1997, Köhler 2010).

Of the 84 land snails recorded from the database searches, all belonged to the family Camaenidae and comprised eight species, of which four are considered to have the potential to occur in the study area (Figure 5.5). None of the species are listed as Threatened, however some may be regarded as potential SREs. The four camaenid species are:

- *Quistrachia leptogramma*: this species is known to occur in woodland, pindan and vine thicket habitat, from Broome in the south to Cape Leveque and One Arm Point in the north (Solem 1997). On the basis of current knowledge, this species is not considered to represent an SRE (R. Teale pers. comm.).
- *Rhagada bulgana* and *Rhagada cygna*: these two species are both distributed in the northern half of the Dampier Peninsula (Burghardt and Köhler 2014), with *R. cygna* replacing *R. bulgana* in the northern part of the peninsula and on several islands to the east (Sunday, Gibbings and Hidden Islands). Uncertainty exists over the species' boundaries; these species may be synonymised in the future (Burghardt and Köhler 2014). The SRE status of these species is uncertain.
- *Rhagada reinga*: limited to the vicinity of Broome (Burghardt and Köhler 2014), this species has previously been collected from vine thicket habitat (Biota 2009). This species is considered likely to represent an SRE species, however it is unlikely to occur in the study area based on its current known distribution.

5.2.3.2 Scorpions

Twelve records of scorpions were returned from the ALA database search comprising five belonging to the genus *Lychas* within the family Buthidae, and seven of the species *Urodacus hoplurus* within the family Urodacidae (Figure 5.5).

The *Lychas* specimens were not identified to species level and as a result could not have their SRE status assessed.

Urodacus hoplurus is distributed throughout Western Australia, South Australia and the Northern Territory and is not considered to be an SRE (Koch 1977).

5.2.3.3 Mygalomorph Spiders

Eight records of mygalomorph spiders were returned from the database search, comprising seven belonging to the genus *Aname* within the family Anamidae (previously Nemesiidae), and one from the genus *Conothele* within the family Halonoproctidae (previously Ctenizidae) (Figure 5.5). None of the eight specimens were identified to species level and so SRE statuses could not be assigned.




Figure 5.5: Historical locations of potential SRE invertebrate fauna recorded from the locality of the study area.

6.0 Vegetation Results

6.1 Description of the Vegetation

Cleared and/or heavily disturbed areas (roads, drains, tracks as well as houses and gardens associated with the Lombadina community) comprised 5.37 ha (44%) of the study area. There were two main native vegetation types recorded in the study area; both were recorded from orange sandy pindan plains (see Figure 6.1) and are described below.

P1: CgCbAtCHpSe *Corymbia greeniana*, *C. bella* very open woodland over *Acacia tumida* tall shrubland over *Chrysopogon pallidus*, *Sorghum ecarinatum* mixed tussock grassland.

This was the dominant vegetation type of the study area, covering 6.23 ha (50%) of the study area (see Plate 6.1). *Corymbia greeniana* (Bilal) and *C. bella* (Maroolal) were co-dominant overstorey trees. *Acacia tumida* (Wanggay) was the only dominant shrub and these were often still recovering from fire. The understorey consisted of *Chrysopogon pallidus* (Irooloo), *Sorghum ecarinatum* (Oonbi) tussock grasses. Typical non-dominant herb species occurring in this vegetation included *Bonamia linearis*, *Calandrinia strophiolata*, *Corchorus pumilio*, *Evolvulus alsinoides*, *Goodenia sepalosa*, *Gyrostemon tepperi*, *Sida rohlenae*, *Spermacoce occidentalis* and *Trianthema pilosum*. Fire age was less than one year, and the condition for this vegetation type was rated as Good (discussed in Section 6.2).

P2: EmCHpSe *Eucalyptus miniata* open woodland over *Chrysopogon pallidus*, *Sorghum ecarinatum* mixed tussock grassland.

This vegetation type was recorded in the southeast corner of the study area over 0.77 ha (6%) of the study area. *Eucalyptus miniata* (Manowan) was the dominant overstorey tree species, and the same dominant tussock grasses of *Chrysopogon pallidus* (Irooloo), *Sorghum ecarinatum* (Oonbi) were present. The condition of this vegetation type was rated as Very Good (discussed in Section 6.2).



Plate 6.1: Vegetation type P1: CgCbAtCHpSe



Plate 6.2: Vegetation type P2: EmCHpSe




Figure 6.1: Vegetation of the Lombadina study area.




Figure 6.2: Vegetation condition ranking of the Lombadina study area.

6.2 Vegetation Condition

The vegetation condition assessment was based on the ranking scale developed by Trudgen (1988) (see Appendix 2). The rankings include degree of invasion by introduced flora (weeds), impact from humans, feral animals and livestock activities, and the structural integrity of the vegetation.

The areas of intact remnant vegetation were mapped and ranged from Poor to Very Good condition (Figure 6.2). Only a small area in the center of the study area was rated as Poor condition, due to a dense Birdwood Grass (*Cenchrus setiger*) infestation. The main disturbance factors recorded for the study area were weeds (15 weed species were recorded; discussed below in Section 9.3) as well as historical human disturbance including dumped items (Plate 6.3) and areas of ground disturbance (Plate 6.4). No pastoral grazing impacts were noted during the survey.



Plate 6.3: Rusty steel hull of a boat dumped in vegetation type P1.



Plate 6.4: Earthy mound recorded in vegetation type P1.

6.3 Vegetation of Significance

6.3.1 TECs and PECs

None of the vegetation described in the study area comprise listed TECs or PECs.

7.0 Flora Results

7.1 Overview

7.1.1 Flora Species Recorded in the Study Area

A total of 66 native vascular flora species from 57 genera and 33 families were recorded from the study area during the current survey (see Appendix 9).

The dominant native plant families and genera recorded from the study area are presented in Table 7.1. These families and genera are typically represented in species lists from this region.

Table 7.1: Dominant native families and genera recorded from the study area.

Family	No. of Native Species	Genus	No. of Native Species
Fabaceae (peas, cassias and wattles)	11	<i>Corymbia</i> (bloodwoods)	3
Malvaceae (hibiscus, sida etc.)	8	<i>Arivela</i>	2
Poaceae (grasses)	5	<i>Corchorus</i>	2
		<i>Crotalaria</i>	2
		<i>Ficus</i> (figs)	2
		<i>Indigofera</i>	2
		<i>Sida</i>	2
		<i>Zornia</i>	2

A total of 15 introduced flora species (weeds) from 13 genera and nine families were recorded in the study area during the current survey, and are discussed separately in Section 7.3.1. In addition, there were a further 11 introduced species that were not considered to be weeds as they were recorded in plantings in established gardens in the study area (Section 7.3.2). All introduced species (weeds and plantings) are listed in the species list in Appendix 9.

7.2 Flora of Significance

No Threatened or Priority species were recorded in the study area during the survey.

No significant flora species, including annual species, were ranked 'likely to occur' or 'may occur' in the flora likelihood assessment following the field survey (see Appendix 5).

7.3 Introduced Flora

7.3.1 Weeds

A total of 15 introduced (weed) species were recorded from the Lombadina study area (Table 7.2, Appendix 11).

One of these species, *Jatropha gossypifolia*, is considered a Weed of National Significance (WoNS; see DoTEE 2018). Two of the weed species, *Jatropha gossypifolia* and *Azadirachta indica* are listed as declared pests under the WA Biosecurity and Agriculture Management Act 2007 (DAFWA 2018).

The then Department of Parks and Wildlife derived a Weed Species Ranking for species in each major region of WA through the Weed Prioritisation Process (WPP) (Department of Parks and Wildlife 2013). This took into account the potential distribution, current distribution, ecological

impact, invasiveness and feasibility of control to derive a broad qualitative weed species ranking corresponding to specific management actions (Department of Parks and Wildlife 2014). According to this methodology, three of the species recorded (**Mesosphaerum suaveolens*, **Azadirachta indica* and **Passiflora foetida* var. *hispida*) have both a 'High' ranking for ecological impact and a 'Rapid' ranking for invasiveness (Table 7.2).

The most significant weed species are discussed in more detail below.

- ****Jatropha gossypifolia* (Bellyache Bush)**

This species is listed as a WoNS, and as a Declared Pest under category "s22(2)" for the whole of WA. Section 22(2) of the BAM Act requires that any applicable import requirements must be satisfied when the species is imported, and the species may also be subject to control and keeping requirements. Its classification as a WoNS means it is considered one of the worst weeds in Australia due to its invasiveness, potential for spread, and economic and environmental impacts, particularly in the northern parts of Western Australia, the Northern Territory and Queensland. This species can form dense thickets and is particularly damaging in riparian habitats, where it replaces or prevents the regeneration of native vegetation (ALA 2021a). It is described as widespread and increasing its range within both the Ord and Fitzroy River catchments in the Kimberley (KRBA 2021).

- ****Azadirachta indica* (Neem Tree)**

Listed as a Declared Pest in the category of "s22(2) Exempt" for Broome, Derby-West Kimberley, Halls Creek, and Wyndham-East Kimberley (see the explanation of Section 22(2) of the BAM Act above for **Jatropha gossypifolia*; the "Exempt" status for **Azadirachta indica* means that no permit or conditions are required for keeping), **Azadirachta indica* is a highly invasive species, and is a serious environmental weed due to its highly competitive nature. This species has a particularly detrimental impact on terrestrial and riparian ecosystems, where it has developed impenetrable forests along waterways in parts of the Kimberley as well as killing boab trees, mature eucalypts and displacing conservation-significant monsoon vine thicket (DPIRD 2021). It is currently known to be rapidly spreading throughout the Ord and Fitzroy catchments (KRBA 2021).



Plate 7.1: **Jatropha gossypifolia* (Bellyache Bush) in the study area.



Plate 7.2: **Azadirachta indica* (Neem Tree) in the study area.

Table 7.2: Weed species recorded from the study area, including WPP rankings.

Descriptions from WA Herbarium (2021), unless otherwise cited.

Family	Species	Description	WPP – Weed Species Ranking		Distribution in the Study Area
			Ecol. Imp.	Inv.	
Amaranthaceae	* <i>Aerva javanica</i> (Kapok Bush)	Erect perennial herb, often occurs on sandy soils. Originally introduced to assist with the revegetation of disturbed bushland; now widespread from the Kimberley to Carnarvon (Hussey et al. 2007).	L	M	Single record from the center of the study area.
	* <i>Gomphrena celosioides</i> (Gomphrena Weed)	A sprawling herb, with greenish flowering spikes. Originally from South America, it is spreading rapidly in irrigated areas of most major settlements in the Kimberley (Hussey et al. 2007). It's been recorded as far south as the Avon Wheatbelt.	L	S	Single record; recorded around the brown house.
Asteraceae	* <i>Cyanthillium cinereum</i> (Little Ironweed)	This annual daisy has pink-purple flowers for most of the year and aromatic foliage. It occurs in a variety of habitats, including near permanent water and in disturbed areas.	Not assessed.		Two records from the center of the study area.
	* <i>Tridax procumbens</i> (Tridax)	Low perennial species with ovate, lobed, stiffly hairy leaves and solitary, erect flower heads with cream ray florets and yellow disc florets (Hussey et al. 2007). Occurs in wet or disturbed ground near settlements in the Pilbara and the Kimberley.	U	S	Single record; recorded around the brown house.
Convolvulaceae	* <i>Distimake dissectus</i> var. <i>dissectus</i>	A twining, hairy plant with large dissected leaves with indented margins and white flowers with a reddish purple centre (Hussey et al. 2007). It has been cultivated as an ornamental and has escaped from cultivation in scattered locations near settlements throughout the Pilbara and the Kimberley.	H	M	Four records in the study area.
Euphorbiaceae	* <i>Jatropha gossypifolia</i> (Bellyache Bush) DP, WoNS	A shrub to small tree growing up to 4 m tall, with purple to red flowers throughout the year but mostly in late summer to autumn. Widespread but scattered throughout northern Australia, from WA to Queensland (ALA 2021b).	H	M	Two records in the western end of the study area, totaling 33 individuals.
Fabaceae	* <i>Macroptilium atropurpureum</i> (Purple Bean)	A trailing or twining, stoloniferous perennial herb climbing to 5 m high, with red-purple flowers most of the year. Recorded from sand, clay and alluvium in creeklines, on dunes, the edges of seasonal wetlands and mangroves, road verges, and other disturbed areas from the Kimberley to Perth.	H	M	Three records in the western end of the study area.
	* <i>Stylosanthes hamata</i> (Verano Stylo)	Erect or decumbent herb or shrub growing to 0.7 m tall with yellow flowers from April to December. Recorded throughout the Kimberley from loam and clay soils in seepage areas, creek banks, pool edges, lawns and in areas of disturbed vegetation; also recorded south as far as Exmouth and Newman, mostly associated with road verges (WA Herbarium 2021).	M	M	Widespread pastoral weed; recorded from four locations in the study area.
	* <i>Stylosanthes scabra</i>	An erect shrub growing to 2 m tall with yellow flowers from February to June. It occurs on alluvial soil, red (pindan) soil and sandstone in	M	M	Widespread pastoral weed; recorded from

Family	Species	Description	WPP – Weed Species Ranking		Distribution in the Study Area
			Ecol. Imp.	Inv.	
		areas adjacent to major rivers, flood prone areas, cultivated ground and road verges.			two locations in the study area.
Lamiaceae	* <i>Mesosphaerum suaveolens</i>	A perennial herb or shrub growing to 2 m tall with a characteristic minty smell when the leaves are crushed (Hussey et al. 2007), with purple-blue flowers from March to August. In WA, only recorded from the Kimberley region.	H	R	Recorded from four locations in the study area where it formed dense patches under trees.
Meliaceae	* <i>Azadirachta indica</i> (Neem Tree) DP	Tree to 20 m tall. Widespread throughout northern Australia, from WA to Queensland (ALA 2021b); typically occurs on cultivated black soil, disturbed pindan, and along creeks. In WA, recorded mainly in the Kimberley, with occasional locations in the northern Pilbara.	H	R	Two individuals recorded from the center of the study area.
Passifloraceae	* <i>Passiflora foetida</i> var. <i>hispida</i>	A woody vine with an unpleasant smell growing up to 9 m high, flowering for most of the year. Typically occurs along river and creek banks, in vine thickets and in coastal areas, from the Kimberley to Shark Bay (WA Herbarium 2021).	H	R	Three records from the western end of the study area.
Poaceae	* <i>Cenchrus biflorus</i> (Gallon's Curse)	Annual grass growing to 60 cm tall, flowering between January and August. This species occurs on red or grey-brown sand in the Central Kimberley, Dampierland, Ord Victoria Plain and Victoria Bonaparte bioregions.	U	R	Single record from around the brown house.
	* <i>Cenchrus setiger</i> (Birdwood Grass)	Perennial tussock grasses growing to 1 m tall and flowering for most of the year. Introduced by pastoralists as a fodder species and now widespread through most of Australia, except for the southernmost regions. Has demonstrated allelopathic capacities, whereby they release chemicals that inhibit the growth of other plants (Cheam 1984a, 1984b, Hussain et al. 2010), and compete aggressively and effectively with native flora. Commonly found along drainage lines, floodplains, in sandy coastal areas and disturbed sites, where they can form dense tussock grasslands. Reproduce by seed and short rhizomes; dispersed primarily by wind and water, but can also be spread through the movement of mammals, birds and vehicles.	H	M	Widespread pastoral weed, recorded from five locations, including a large patch in the center of the study area.
	* <i>Chloris barbata</i> (Purpletop Chloris)	An annual or short-lived perennial grass growing to 1.1 m tall and flowering most of the year. It typically occurs on white or red sand, loam or black clay in sand dunes and river levees.	U	U	Recorded from two locations around the brown house.

WPP = Weed Prioritisation Process (Department of Parks and Wildlife 2013); note that only species with rankings in both categories are listed in Department of Parks and Wildlife (2014).

Ecological Impact Ranking: H = High, L = Low, M = Medium, U = Unknown.

Invasiveness Ranking: M = Moderate, R = Rapid, S = Slow, U = Unknown.

7.3.2 Plantings

A total of 11 planted species were recorded in the study area, around the lawn and gardens of the brown house (see Table 7.3 and species list in Appendix 9). While these species are introduced, they are not considered to be weeds as they are deliberate plantings.

Table 7.3: Planted (introduced species) recorded in the study area

Species Name	Common Name
* <i>Ananas comosus</i>	Pineapple
* <i>Bothriochloa pertusa</i>	-
* <i>Catharanthus roseus</i>	Vinca
* <i>Cocos nucifera</i>	Coconut
* <i>Hibiscus sabdariffa</i>	Rosella
* <i>Ipomoea batatas</i>	Sweet Potato
* <i>Khaya senegalensis</i>	-
* <i>Mangifera indica</i>	Mango
* <i>Ocimum basilicum</i>	Sweet Basil
* <i>Phyla nodiflora</i>	Lippia
* <i>Plumeria rubra</i>	Red Frangipani

This page intentionally blank.

8.0 Fauna Results

8.1 Fauna Habitats

One fauna habitat was identified in the study area, *Eucalyptus/Corymbia* woodlands over tussock grassland on orange Pindan sands. This habitat type encompassed both vegetation types (P1 and P2) described in Section 6.1.

This habitat type provides potentially suitable habitat for the 10 species of significance considered as potentially occurring within the study area. However, the habitat type is widespread in the local area and so the extent within the study area is unlikely to represent significant core habitat for any of these species. The proposed campground development is not expected to have a significant impact on this habitat type at either a local or regional scale, particularly given the small clearing footprint (12.35 ha).

8.2 Field Survey Records

8.2.1 Vertebrate Fauna

During the field survey of the study area, eight species of bird were opportunistically observed. These are listed in Table 8.1. Of the eight species, none are significant.

Table 8.1: Bird species opportunistically observed in the Lombadina study area.

Species	Family	Conservation Status
<i>Geopelia striata</i> (Peaceful Dove)	Columbidae	-
<i>Dacelo leachii</i> (Blue-winged Kookaburra)	Alcedinidae	-
<i>Calyptorhynchus banksii</i> (Red-tailed Black Cockatoo)	Cacatuidae	-
<i>Aprosmictus erythropterus</i> (Red-winged Parrot)	Psittaculidae	-
<i>Philemon argenticeps</i> (Silver-crowned Friarbird)	Meliphagidae	-
<i>Pomatostomus temporalis</i> (Grey-crowned Babbler)	Pomatostomidae	-
<i>Coracina novaehollandiae</i> (Black-faced Cuckoo-shrike)	Campephagidae	-
<i>Dicaeum hirundinaceum</i> (Mistletoebird)	Dicaeidae	-

8.3 Likelihood of Significant Fauna

Following the field survey a final likelihood assessment ranking was made for the 17 significant vertebrate fauna species with the potential to occur in the study area, as identified in the desktop study (Section 5.2).

Based on the final likelihood assessment, 10 of the 26 significant species considered were ranked as 'likely to occur' or 'may occur'. The remaining 16 significant species were ranked as 'unlikely to occur' or "would not occur" (see Appendix 6).

This page intentionally blank.

9.0 Conclusions

9.1 Vegetation of Significance

Two vegetation types were described and mapped for the study area. This is considered to be within the range expected for a study area of this size in this locality, taking into account the habitats present and the scale of vegetation mapping.

The vegetation types in the study area were in Good to Very Good condition. While both have inherent value as relatively intact and good quality examples of native vegetation, neither is representative of any TEC or PEC, or otherwise considered to have any elevated conservation.

9.2 Flora of Significance

A total of 66 native vascular flora species were recorded during the survey. This is within the range expected for a study area of this size in this locality. No Threatened or Priority flora species were recorded during the current survey, and none are considered as likely to occur.

9.3 Introduced Flora

A total of 15 introduced (weed) species were recorded from the study area. **Jatropha gossypifolia* (Bellyache Bush), which was recorded from two locations, is considered to be a WoNS (DotEE 2018) and is listed as a declared pest under the *WA Biosecurity and Agriculture Management Act 2007* (DAFWA 2018). **Azadirachta indica* (Neem Tree), is also listed as a declared pest and was recorded from two locations within the study area.

9.4 Fauna of Significance

No evidence of fauna of significance were recorded in the study area during the survey. On the basis of the desktop study, 10 species of significant fauna were assessed as either likely to or may occur in the study area. The small extent of the study area and the wider distributions of the species suggest it would be unlikely that major impacts on any significant species would arise because of the camp site development, should they be present.

9.5 Assessment Against the Ten Clearing Principles

A general assessment of the proposal to clear land within the survey area against each of the 10 clearing principles, as outlined in Schedule 5 of the *Environmental Protection Act 1986* (EP Act), is provided in Table 9.1.

Table 9.1: Assessment of remnant native vegetation within the study area against the ten clearing principles.

Principle	Assessment	Conclusion
A	Native vegetation should not be cleared if it comprises a high level of biological diversity.	
	<p>Flora</p> <p>The study area is not located within a known biodiversity hotspot for WA (Department of the Agriculture, Water and the Environment 2020).</p> <p>Vascular flora species diversity recorded within the study area was considered to be moderate; a total of 66 native flora species from 33 families were recorded within the 12.35 ha study area. This number is not considered to represent an unusually high level of species diversity given the location and the vegetation types present. No significant flora species were recorded during the survey and none are deemed likely to occur following the field survey.</p> <p>Fauna</p> <p>The scope of the assessment did not include a comprehensive census of vertebrate fauna in the study area however on the basis of the desktop study, 330 vertebrate fauna species have the potential to occur in the study area, of which 107 are significant species. Ten of the 26 species considered as potentially occurring were assessed as either 'likely to occur' or 'may occur' based on the habitats present.</p> <p>The clearing footprint for the proposed campground is a maximum of 12.35 ha; neither the vascular flora species diversity nor the fauna species diversity would be affected by clearing at this small scale.</p>	Unlikely to be at variance
B	Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a significant habitat for fauna indigenous to Western Australia.	
	<p>One fauna habitat was recorded from the study area.</p> <p>The proposal's clearing footprint is minor (12.35 ha) and the remnant vegetation is not considered likely to be significant core habitat for any conservation significant fauna species.</p>	Unlikely to be at variance
C	Native vegetation should not be cleared if it includes, or is necessary for the continued existence of, rare flora.	
	<p>The results of the desktop study show that no records of any Threatened flora species occurring within 40 km of the study area.</p> <p>No Threatened flora, nor any Priority flora, were recorded in the study area during the field survey or are considered to have the potential to occur, given the habitats present.</p>	Unlikely to be at variance
D	Native vegetation should not be cleared if it comprises the whole or a part of, or is necessary for the maintenance of, a threatened ecological community.	
	<p>None of the mapped remnant vegetation within the study area was considered to be analogous with any Threatened Ecological Communities (TECs) under the Commonwealth <i>EPBC Act</i>.</p>	Unlikely to be at variance.
E	Native vegetation should not be cleared if it is significant as a remnant of native vegetation in an area that has been extensively cleared.	
	<p>The study area comprises one broad-scale regional vegetation unit: Dampierland 750, described as "<i>Pindan woodland; Acacia thicket with eucalypt woodland over spinifex; Acacia tumida, Eucalyptus tectifera, Corymbia grandifolia, Triodia pungens, T. bitextura</i>" (Beard 1979).</p> <p>The current extent of this vegetation unit in the Pindanland IBRA subregion is 1,219,556 ha (99.8% of the pre-European extent) (Government of Western Australia 2019).</p> <p>The proposed clearing area is not within a significantly cleared landscape and does not represent a significant ecological linkage. Clearing of 12.35 ha of remnant vegetation would represent a very minor increment on historical clearing.</p>	Unlikely to be at variance.

Principle	Assessment	Conclusion
F	<p>Native vegetation should not be cleared if it is growing in, or in association with, an environment associated with a watercourse or wetland.</p> <p>There are no watercourses or wetlands within the study area.</p> <p>The nearest Ramsar wetland is Roebuck Bay, located approximately 170 km to the southwest (Department of Agriculture, Water and the Environment 2019).</p> <p>The nearest wetland listed in the Directory of Important Wetlands in Australia is Willie Creek, approximately 147 km southwest of the study area (Environment Australia 2001).</p> <p>Given the small scale of proposed clearing it is considered that there would be no impact on these systems.</p>	Unlikely to be at variance.
G	<p>Native vegetation should not be cleared if the clearing of the vegetation is likely to cause appreciable land degradation.</p> <p>The remnant vegetation of the study area is predominantly intact (44% of the study area has been cleared/disturbed for tracks, houses, roads etc.) and encompasses a mostly flat landscape.</p> <p>The soil units mapped for the study area include Pindan sands and saline clays (CSIRO 2014); however field observations show that the Pindan sands are the dominant soil type in the study area.</p> <p>The study area intersects the Yeeda Land System, which is noted to be generally not susceptible to degradation or erosion (Payne and Schoknecht 2011). However, with significant rainfall events, which often occur in the wet season, it is possible that the removal of stabilising vegetation could allow for increased erosion.</p> <p>Fifteen weed species were recorded in the study area including one recognised as a WoNS and two as declared pests. Given the close proximity of the study area to the Lombadina community, the area would be susceptible to weed invasion following disturbance, although this may not be evident until after significant rainfall. Strict weed hygiene measures should be implemented to ensure weeds are not further spread into the study area and surrounds.</p> <p>It is considered unlikely that the proposal would contribute significantly to land degradation in the study area.</p>	Unlikely to be at variance.
H	<p>Native vegetation should not be cleared if the clearing of the vegetation is likely to have an impact on the environmental values of any adjacent or nearby conservation area.</p> <p>No conservation reserves occur within the study area. The closest conservation estate is the Swan Island Nature Reserve, approximately 24 km northeast of the study area (DBCA 2017).</p> <p>Given the small amount of clearing required, it is unlikely to affect any conservation estate.</p>	Unlikely to be at variance.
I	<p>Native vegetation should not be cleared if the clearing of the vegetation is likely to cause deterioration in the quality of surface or underground water.</p> <p>The study area does not intersect any surface water bodies.</p> <p>Given clearing associated with the proposals is small (12.35 ha), and the surrounding vegetation is mostly intact, there is no reason to expect that the clearing would affect the quality of surface or underground water.</p>	Unlikely to be at variance.
J	<p>Native vegetation should not be cleared if clearing the vegetation is likely to cause, or exacerbate, the incidence of flooding.</p> <p>The surface water catchment of the study area is the Cape Leveque Coast Basin of the Western Plateau Division. Drainage within this catchment area is towards the coast.</p> <p>No permanent surface water sources or wetlands occur in the study area. The sandy soil would provide rapid infiltration of rainfall events while any areas of saline clay would be more susceptible to flooding. However, due to the small size of the study area and proposed clearing (12.35 ha), any clearing of vegetation is unlikely to affect the intensity or incidence of flooding.</p>	Unlikely to be at variance.

This page intentionally blank.

10.0 References

- ALA (2021a). WeedsAustralia - Profiles *Jatropha gossypifolia* [WWW Document]. Atlas of Living Australia, . Retrieved from <https://profiles.ala.org.au/opus/weeds-australia>.
- ALA (2021b). Atlas of Living Australia [WWW Document]. Database, . Retrieved from bie.ala.org.au.
- Aplin, T. E. H. (1979). Chapter 3: The Flora. Page in B. J. O'Brien, editor. *Environment and Science*. The University of Western Australia Press.
- Beard, J. S. (1979). *Vegetation Survey of Western Australia: Kimberley. 1:1,000,000 Vegetation Series: Explanatory Notes to Sheet 1*. University of Western Australia.
- Biota (2009). James Price Point Terrestrial Fauna Survey: Wet Season 2009. Unpublished report prepared for the Department of State Development, December 2009, Biota Environmental Sciences, Western Australia.
- Biota (2011). Irvine Island Seasonal Fauna Survey. Unpublished final draft report prepared for Pluton Resources Ltd, December 2011, Biota Environmental Sciences, Western Australia.
- Black, S. J., T. Willing, and D. M. Dureau (2010). A comprehensive survey of the flora, extent and condition of vine thickets on coastal sand dunes of Dampier Peninsula, West Kimberley 2000-2002. Unpublished report prepared for the Broome Botanical Society (Inc.), September 2010, Broome Botanical Society, Broome.
- Burghardt, I., and F. Köhler (2014). *Rhagada* revisited: on the taxonomy of species from the Kimberley and Dampierland, Western Australia (Pulmonata, Camaenidae). *Molluscan Research* 35(1):1-14.
- Cheam, A. H. (1984a). Allelopathy in buffel grass (*Cenchrus ciliaris* L.) Part I. Influence of buffel grass association on calotrope (*Calotropis procera* (Ait) W.T.Ait.). *Australian Weeds* 3:133-136.
- Cheam, A. H. (1984b). Allelopathy in buffel grass (*Cenchrus ciliaris* L.) Part II. Site of release and distribution of allelochemical in the soil profile. *Australian Weeds* 3:137-139.
- CSIRO (2014). ASRIS: Australian Soil Resource Information System [WWW Document]. Retrieved from <http://www.asris.csiro.au/index.html>.
- DAFWA (2018). Western Australian Organism List (WAOL) [WWW Document]. Department of Agriculture and Food, . Retrieved from <https://www.agric.wa.gov.au/organisms>.
- DBCA (2013). Definitions, Categories and Criteria for Threatened and Priority Ecological Communities. Species and Communities Branch, Department of Environment and Conservation, Western Australia, January 2013.
- DBCA (2017). DBCA - Legislated Lands and Waters.
- DBCA (2018). Interim Recovery Plan 2018-2023 for the Monsoon vine thickets on the coastal sand dunes of Dampier Peninsula. Interim Recovery Plan No. 383, February 2018, Department of Biodiversity, Conservation and Attractions, Science and Conservation Service, Perth.
- DBCA (2020). Threatened and Priority Fauna List 2020. Department of Biodiversity, Conservation and Attractions.
- Department of Agriculture, Water and the Environment (2019). Australian Ramsar Wetlands: Internationally Important Wetlands [WWW Document]. Retrieved February 10, 2021, from <https://www.environment.gov.au/water/wetlands/australian-wetlands-database/australian-ramsar-wetlands>.

- Department of Environment Regulation (2014, December). A guide to the assessment of applications to clear native vegetation Under Part V Division 2 of the Environmental Protection Act 1986. Department of Environment Regulation.
- Department of Parks and Wildlife (2013). Weed Prioritisation Process for Department of Parks and Wildlife (formerly DEC) - "An integrated approach to Environmental Weed Management in WA". (As at November 2013). Department of Parks and Wildlife, Western Australia.
- Department of Parks and Wildlife (2014). Kimberley Region - Impact and Invasiveness Ratings.xls. List of weed rankings for the Kimberley derived from the DPaW Invasive Plant Prioritisation Process, downloaded from DBCA website in 2021, Department of Parks and Wildlife, Western Australia.
- Department of the Agriculture, Water and the Environment (2020). Australia's 15 National Biodiversity Hotspots [WWW Document]. Retrieved from <http://www.environment.gov.au/biodiversity/conservation/hotspots/national-biodiversity-hotspots>.
- DotEE (2018). Weeds of National Significance [WWW Document]. Department of the Environment and Energy, . Retrieved from <http://www.environment.gov.au/biodiversity/invasive/weeds/weeds/lists/wons.html>.
- DPIRD (2021). Neem: declared pest [WWW Document]. Department of Primary Industries and Regional Development, . Retrieved from <https://www.agric.wa.gov.au/declared-plants/neem-declared-pest>.
- DSEWPaC (2012). Interim Biogeographic Regionalisation for Australia (IBRA), Version 7 (Subregions) - States and Territories. Department of Sustainability, Environment, Water, Population and Communities, Canberra. Retrieved from <http://www.environment.gov.au/topics/land/national-reserve-system/science-maps-and-data/australias-bioregions-ibra>.
- Ecologia (2004). Beagle Bay Big Tree Country Timber Plantation - Flora Assessment Survey. Ecologia Environmental Consultants, Western Australia.
- Environment Australia (2001). A Directory of Important Wetlands in Australia, 3rd edition. Environment Australia, Canberra. Retrieved from <http://www.environment.gov.au/water/wetlands>.
- EPA (2009). *EPA Guidance Statement No. 20: Sampling of Short Range Endemic Invertebrate Fauna for Environmental Impact Assessment in Western Australia*. Environmental Protection Authority, Western Australia.
- EPA (2016a). *Technical Guidance: Terrestrial Fauna Surveys*. Environmental Protection Authority, Western Australia.
- EPA (2016b). *Technical Guidance: Sampling Methods for Terrestrial Vertebrate Fauna*. Environmental Protection Authority, Western Australia.
- EPA (2016c). *Technical Guidance: Flora and Vegetation Surveys for Environmental Impact Assessment*. Environmental Protection Authority, Western Australia.
- EPA (2016d). *Technical Guidance: Sampling of Short Range Endemic Invertebrate Fauna*. Environmental Protection Authority, Western Australia.
- EPA (2020). *Technical Guidance: Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment*. Environmental Protection Authority, Western Australia.
- Geological Survey of Western Australia (2011, January 13). Surface geology of Australia - 1:1,000,000 scale, Western Australia. Department of Mines and Petroleum, Western Australia.

- Gibson, L. A., and F. Köhler (2012). Determinants of species richness and similarity of species composition of land snail communities on Kimberley islands. *Records of the Western Australian Museum* 81:41–66.
- Gill, F., D. Donsker, and P. Rasmussen (2020). IOC World Bird List (v10.2). doi: 10.14344/IOC.ML.10.1.
- Government of Western Australia (2019). *2018 Statewide Vegetation Statistics incorporating the CAR Reserve Analysis (Full Report)*. Current as of March 2019. Department of Biodiversity, Conservation and Attractions, Perth, Western Australia.
- Graham, G. (2003). Dampierland 2 (DL2 - Pindanland subregion). Pages 179–187 in J. E. May and N. L. McKenzie, editors. *A Biodiversity Audit of Western Australia's 53 Biogeographical Subregions*. Department of Conservation and Land Management, Western Australia.
- Harvey, M. S. (2002). Short-Range Endemism Among the Australian Fauna: Some Examples from Non-Marine Environments. *Invertebrate Systematics* 16:555–570.
- Hussain, F., B. Ahmad, and I. Ilahi (2010). Allelopathic effects of *Cenchrus ciliaris* L. and *Bothriochloa pertusa* (L.) A. Camus. *Pakistan Journal of Botany* 42(5):3587–3604.
- Hussey, B. M. J., G. J. Keighery, R. D. Cousens, J. Dodd, and S. G. Lloyd (2007). *Western Weeds A guide to the weeds of Western Australia*, 2nd edition. The Plant Protection Society of Western Australia (Inc.), Perth.
- Koch, L. E. (1977). The taxonomy, geographic distribution and evolutionary radiation of Australo-Papuan scorpions. *Records of the Western Australian Museum*:83–367.
- Köhler, F. (2010). Uncovering local endemism in the Kimberley, Western Australia: description of new species of the genus *Amplirhagada* Iredale, 1933 (Pulmonata: Camaenidae). *Records of the Australian Museum* 62(2):217–284.
- KRBA (2021). Known Declared Weeds in the Kimberley [WWW Document]. Kimberley Rangelands Biosecurity Association, . Retrieved from <https://krbg.com.au/biosecurity-weeds/>.
- McKenzie, N. L., and K. F. Kenneally (1983). Part I: Background and Environment. Pages 5–23 *Wildlife of the Dampier Peninsula, South-west Kimberley, Western Australia*. Western Australian Wildlife Research Centre, Department of Fisheries and Wildlife.
- Paul, R. J., R. J. George, and P. Gardiner (2013). A review of the Broome Sandstone aquifer in the La Grange area. Department of Agriculture and Food, Western Australia.
- Payne, A. L., and S. Schoknecht (2011). *Technical Bulletin No. 98: Land Systems of the Kimberley Region, Western Australia*. Department of Agriculture and Food, Western Australia.
- Ponder, W. F., and D. J. Colgan (2002). What makes a narrow-range taxon? Insights from Australian freshwater snails. *Invertebrate Systematics* 16:571–582.
- Solem, A. (1981). Camaenid land snails from Western and central Australia (Mollusca: Pulmonata: Camaenidae). II. Taxa from the Kimberley, *Amplirhagada* Iredale 1933. *Records of the Western Australian Museum* 11:321–425.
- Solem, A. (1984). Camaenid land snails from Western and central Australia (Mollusca: Pulmonata: Camaenidae). IV. Taxa from the Kimberley, *Westraichia* Iredale, 1933 and related genera. *Records of the Western Australian Museum* 17:424–705.
- Solem, A. (1991). Land snails of Kimberley rainforest patches and biogeography of all Kimberley landsnails. Pages 145–246 *Kimberley Rainforests of Australia*. Surrey Beatty & Sons, Chipping Norton, Sydney.
- Solem, A. (1997). Camaenid land snails from Western and Central Australia (Mollusca: Pulmonata: Camaenidae). VII Taxa from Dampierland through the Nullarbor. *Records of the Western Australian Museum Supplement* 50:1461–1906.

- Specht, R. L. (1970). Vegetation. Pages 44–67 in G. W. Leeper, editor. *The Australian Environment*, 4th edition. CSIRO in association with Melbourne University Press, Melbourne.
- Trudgen, M. E. (1988). A Report on the Flora and Vegetation of the Port Kennedy Area. Unpublished report prepared for Bowman Bishaw and Associates, West Perth, M.E. Trudgen and Associates, Western Australia.
- WA Herbarium (2021). FloraBase - the Western Australian Flora [WWW Document]. Department of Biodiversity, Conservation and Attractions, . Retrieved from <http://florabase.dpaw.wa.gov.au/>.

Appendix 1

Framework for Conservation Significance Ranking of Species and Communities in WA



A. Definitions, Categories and Criteria for Threatened and Priority Ecological Communities

Species and Communities Branch, Department of Environment and Conservation, December 2010.

1. General Definitions

Ecological Community

A naturally occurring biological assemblage that occurs in a particular type of habitat.

Note: The scale at which biological communities are defined will often depend on the level of detail in the information source, therefore no particular scale is specified.

A **threatened ecological community** (TEC) is one which is found to fit into one of the following categories; "presumed totally destroyed", "critically endangered", "endangered" or "vulnerable".

Possible threatened ecological communities that do not meet survey criteria are added to the Department of Parks and Wildlife's Priority Ecological Community Lists under Priorities 1, 2 and 3. Ecological Communities that are adequately known, are rare but not threatened, or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

An **assemblage** is a defined group of biological entities.

Habitat is defined as the areas in which an organism and/or assemblage of organisms lives. It includes the abiotic factors (e.g. substrate and topography), and the biotic factors.

Occurrence: a discrete example of an ecological community, separated from other examples of the same community by more than 20 metres of a different ecological community, an artificial surface or a totally destroyed community.

By ensuring that every discrete occurrence is recognised and recorded future changes in status can be readily monitored.

Adequately Surveyed is defined as follows:

"An ecological community that has been searched for thoroughly in most likely habitats, by relevant experts."

Community structure is defined as follows:

"The spatial organisation, construction and arrangement of the biological elements comprising a biological assemblage" (e.g. *Eucalyptus salmonophloia* woodland over scattered small shrubs over dense herbs; structure in a faunal assemblage could refer to trophic structure, e.g. dominance by feeders on detritus as distinct from feeders on live plants).

Definitions of **Modification** and **Destruction** of an ecological community:

Modification: "changes to some or all of ecological processes (including abiotic processes such as hydrology), species composition and community structure as a direct or indirect result of human activities. The level of damage involved could be ameliorated naturally or by human intervention."

Destruction: "modification such that reestablishment of ecological processes, species composition and community structure within the range of variability exhibited by the original community is unlikely within the foreseeable future even with positive human intervention."

Note: Modification and destruction are difficult concepts to quantify, and their application will be determined by scientific judgement. Examples of modification and total destruction are cited below:

Modification of ecological processes: The hydrology of Toolibin Lake has been altered by clearing of the catchment such that death of some of the original flora has occurred due to dependence on fresh water. The system may be bought back to a semblance of the original state by redirecting saline runoff and pumping waters of the rising underground watertable away to restore the hydrological balance. Total destruction of downstream lakes has occurred due to hydrology being altered to the point that few of the original flora or fauna species are able to tolerate the level of salinity and/or water logging.

Modification of structure: The understorey of a plant community may be altered by weed invasion due to nutrient enrichment by addition of fertiliser. Should the additional nutrients be removed from the system the balance may be restored, and the original plant species better able to compete. Total destruction may occur if additional nutrients continue to be added to the system causing the understorey to be completely replaced by weed species, and death of overstorey species due to inability to tolerate high nutrient levels.

Modification of species composition: Pollution may cause alteration of the invertebrate species present in a freshwater lake. Removal of pollutants may allow the return of the original inhabitant species. Addition of residual highly toxic substances may cause permanent changes to water quality, and total destruction of the community.

Threatening processes are defined as follows:

"Any process or activity that threatens to destroy or significantly modify the ecological community and/or affect the continuing evolutionary processes within any ecological community."

Examples of some of the continuing threatening processes in Western Australia include: general pollution; competition, predation and change induced in ecological communities as a result of introduced animals; competition and displacement of native plants by introduced species; hydrological changes; inappropriate fire regimes; diseases resulting from introduced micro-organisms; direct human exploitation and disturbance of ecological communities.

Restoration is defined as returning an ecological community to its pre-disturbance or natural state in terms of abiotic conditions, community structure and species composition.

Rehabilitation is defined as the re-establishment of ecological attributes in a damaged ecological community although the community will remain modified.

2. Definitions and Criteria for Presumed Totally Destroyed, Critically Endangered, Endangered and Vulnerable Ecological Communities

ECOLOGICAL COMMUNITIES

Presumed Totally Destroyed (PD)

An ecological community that has been adequately searched for but for which no representative occurrences have been located. The community has been found to be totally destroyed or so extensively modified throughout its range that no occurrence of it is likely to recover its species composition and/or structure in the foreseeable future.

An ecological community will be listed as presumed totally destroyed if there are no recent records of the community being extant and either of the following applies (A or B):

- A) Records within the last 50 years have not been confirmed despite thorough searches of known or likely habitats or
- B) All occurrences recorded within the last 50 years have since been destroyed

Critically Endangered (CR)

An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or that was originally of limited distribution and is facing severe modification or destruction throughout its range in the immediate future, or is already severely degraded throughout its range but capable of being substantially restored or rehabilitated.

An ecological community will be listed as Critically Endangered when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future. This will be determined on the basis of the best available information, by it meeting any one or more of the following criteria (A, B or C):

- A) The estimated geographic range, and/or total area occupied, and/or number of discrete occurrences since European settlement have been reduced by at least 90% and either or both of the following apply (i or ii):
 - i) geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is imminent (within approximately 10 years);
 - ii) modification throughout its range is continuing such that in the immediate future (within approximately 10 years) the community is unlikely to be capable of being substantially rehabilitated.
- B) Current distribution is limited, and one or more of the following apply (i, ii or iii):
 - i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the immediate future (within approximately 10 years);
 - ii) there are very few occurrences, each of which is small and/or isolated and extremely vulnerable to known threatening processes;

iii) there may be many occurrences but total area is very small and each occurrence is small and/or isolated and extremely vulnerable to known threatening processes.

C) The ecological community exists only as highly modified occurrences that may be capable of being rehabilitated if such work begins in the immediate future (within approximately 10 years).

Endangered (EN)

An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or was originally of limited distribution and is in danger of significant modification throughout its range or severe modification or destruction over most of its range in the near future.

An ecological community will be listed as Endangered when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future. This will be determined on the basis of the best available information by it meeting any one or more of the following criteria (A, B, or C):

- A) The geographic range, and/or total area occupied, and/or number of discrete occurrences have been reduced by at least 70% since European settlement and either or both of the following apply (i or ii):
 - i) the estimated geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is likely in the short term future (within approximately 20 years);
 - ii) modification throughout its range is continuing such that in the short term future (within approximately 20 years) the community is unlikely to be capable of being substantially restored or rehabilitated.
- B) Current distribution is limited, and one or more of the following apply (i, ii or iii):
 - i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the short term future (within approximately 20 years);
 - ii) there are few occurrences, each of which is small and/or isolated and all or most occurrences are very vulnerable to known threatening processes;
 - iii) there may be many occurrences but total area is small and all or most occurrences are small and/or isolated and very vulnerable to known threatening processes.
- C) The ecological community exists only as very modified occurrences that may be capable of being substantially restored or rehabilitated if such work begins in the short-term future (within approximately 20 years).

Vulnerable (VU)

An ecological community that has been adequately surveyed and is found to be declining and/or has declined in distribution and/or condition and whose ultimate security has not yet been assured and/or a community that is still widespread but is believed likely to move into a category of higher threat in the near future if threatening processes continue or begin operating throughout its range.

An ecological community will be listed as Vulnerable when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction or significant modification in the medium to long-term future. This will be determined on the basis of the best available information by it meeting any one or more of the following criteria (A, B or C):

- A) The ecological community exists largely as modified occurrences that are likely to be capable of being substantially restored or rehabilitated.
- B) The ecological community may already be modified and would be vulnerable to threatening processes, is restricted in area and/or range and/or is only found at a few locations.
- C) The ecological community may be still widespread but is believed likely to move into a category of higher threat in the medium to long term future because of existing or impending threatening processes.

3. Definitions and Criteria for Priority Ecological Communities

PRIORITY ECOLOGICAL COMMUNITY LIST

Possible threatened ecological communities that do not meet survey criteria or that are not adequately defined are added to the Priority Ecological Community Lists under Priorities 1, 2 and 3. These three categories are ranked in order of priority for survey and/or definition of the community, and evaluation of conservation status, so that consideration can be given to their declaration as threatened ecological communities. Ecological Communities that are adequately known, and are rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

Priority One: Poorly-known ecological communities

Ecological communities with apparently few, small occurrences, all or most not actively managed for conservation (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) and for which current threats exist. Communities may be included if they are comparatively well-known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.

Priority Two: Poorly-known ecological communities

Communities that are known from few small occurrences, all or most of which are actively managed for conservation (e.g. within national parks, conservation parks, nature reserves, State forest, unallocated Crown land, water reserves, etc.) and not under imminent threat of destruction or degradation. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.

Priority Three: Poorly known ecological communities

- (i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or:
- (ii) communities known from a few widespread occurrences, which are either large or within significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or;
- (iii) communities made up of large, and/or widespread occurrences, that may or not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, and inappropriate fire regimes.

Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.

Priority Four: Ecological communities that are adequately known, rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.

- (a) Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands.
- (b) Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable.
- (c) Ecological communities that have been removed from the list of threatened communities during the past five years.

Priority Five: Conservation Dependent ecological communities

Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

B. Categories for Flora and Fauna Species

1. Western Australian Biodiversity Conservation Act 2016, and Priority Species Classification

In Western Australia, 'Threatened', 'Extinct' and 'Specially Protected' fauna and flora species are protected under the *Biodiversity Conservation Act 2016* (the BC Act), making it an offence to take or disturb these species without Ministerial approval. The definition of 'take' is broad, and includes killing, injuring, harvesting or capturing fauna, and gathering, cutting, destroying, harvesting or damaging flora.

Such species are classified within a framework of several categories.

Species of the highest conservation significance are designated as Threatened species and are protected under sections 19(1)(a), 19(1)(b) and 19(1)(c) of the BC Act. Species are listed within one of three categories:

- Critically endangered (CR), Endangered (EN), or Vulnerable (V), representing those species listed in Schedules 1 to 3 respectively of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* or the *Wildlife Conservation (Rare Flora) Notice 2018*.

Presumed extinct species are protected under sections 24 and 25 of the BC Act and are listed in one of two categories:

- Extinct (EX), representing those species listed in Schedule 4 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* or the *Wildlife Conservation (Rare Flora) Notice 2018*; or
- Extinct in the wild (EW); there are currently no listed species under this category.

Specially protected species are protected under section 13(1) of the BC Act, and include species of special conservation interest, migratory species, cetaceans, species subject to international agreement, or species otherwise in need of special protection. Of these:

- Migratory species (MI) are those listed under schedule 5 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*;
- Species of special conservation interest (conservation dependent fauna) (CD) are those listed under schedule 6 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*; and
- Other specially protected fauna (OS) are those listed under schedule 7 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*;

In addition to the species formally designated as protected under the BC Act, the WA Department of Biodiversity, Conservation and Attractions (DBCA) also maintains a list of 'Priority species'.

Species that appear to be rare or threatened, but for which there is insufficient information to properly evaluate their conservation significance, are assigned to one of three Priority categories (Priority 1 to Priority 3), while species that are adequately known but require regular monitoring are assigned to Priority 4.

Note that of the above classifications, only 'Threatened', 'Extinct' and 'Specially Protected' species have statutory standing. The Priority flora and fauna classifications are employed by the WA DBCA to manage and classify their database of species considered potentially rare or at risk, but these categories have no legislative status.

Further explanations of the categories is provided in more detail in the following pages.



CONSERVATION CODES

For Western Australian Flora and Fauna

Threatened, Extinct and Specially Protected fauna or flora¹ are species² which have been adequately searched for and are deemed to be, in the wild, threatened, extinct or in need of special protection, and have been gazetted as such.

The *Wildlife Conservation (Specially Protected Fauna) Notice 2018* and the *Wildlife Conservation (Rare Flora) Notice 2018* have been transitioned under regulations 170, 171 and 172 of the *Biodiversity Conservation Regulations 2018* to be the lists of Threatened, Extinct and Specially Protected species under Part 2 of the *Biodiversity Conservation Act 2016*.

Categories of Threatened, Extinct and Specially Protected fauna and flora are:

T **Threatened species**

Listed by order of the Minister as Threatened in the category of critically endangered, endangered or vulnerable under section 19(1), or is a rediscovered species to be regarded as threatened species under section 26(2) of the *Biodiversity Conservation Act 2016* (BC Act).

Threatened fauna is that subset of 'Specially Protected Fauna' listed under schedules 1 to 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for Threatened Fauna.

Threatened flora is that subset of 'Rare Flora' listed under schedules 1 to 3 of the *Wildlife Conservation (Rare Flora) Notice 2018* for Threatened Flora.

The assessment of the conservation status of these species is based on their national extent and ranked according to their level of threat using IUCN Red List categories and criteria as detailed below.

CR **Critically endangered species**

Threatened species considered to be "*facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with criteria set out in the ministerial guidelines*".

Listed as critically endangered under section 19(1)(a) of the BC Act in accordance with the criteria set out in section 20 and the ministerial guidelines. Published under schedule 1 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for critically endangered fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for critically endangered flora.

EN **Endangered species**

Threatened species considered to be "*facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines*".

Listed as endangered under section 19(1)(b) of the BC Act in accordance with the criteria set out in section 21 and the ministerial guidelines. Published under schedule 2 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for endangered fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for endangered flora.

VU **Vulnerable species**

Threatened species considered to be "*facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with criteria set out in the ministerial guidelines*".

Listed as vulnerable under section 19(1)(c) of the BC Act in accordance with the criteria set out in section 22 and the ministerial guidelines. Published under schedule 3 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for vulnerable fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for vulnerable flora.

Extinct species

Listed by order of the Minister as extinct under section 23(1) of the BC Act as extinct or extinct in the wild.

EX Extinct species

Species where “*there is no reasonable doubt that the last member of the species has died*”, and listing is otherwise in accordance with the ministerial guidelines (section 24 of the BC Act).

Published as presumed extinct under schedule 4 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018* for extinct fauna or the *Wildlife Conservation (Rare Flora) Notice 2018* for extinct flora.

EW Extinct in the wild species

Species that “*is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; and it has not been recorded in its known habitat or expected habitat, at appropriate seasons, anywhere in its past range, despite surveys over a time frame appropriate to its life cycle and form*”, and listing is otherwise in accordance with the ministerial guidelines (section 25 of the BC Act).

Currently there are no threatened fauna or threatened flora species listed as extinct in the wild. If listing of a species as extinct in the wild occurs, then a schedule will be added to the applicable notice.

Specially protected species

Listed by order of the Minister as specially protected under section 13(1) of the BC Act. Meeting one or more of the following categories: species of special conservation interest; migratory species; cetaceans; species subject to international agreement; or species otherwise in need of special protection.

Species that are listed as threatened species (critically endangered, endangered or vulnerable) or extinct species under the BC Act cannot also be listed as Specially Protected species.

MI Migratory species

Fauna that periodically or occasionally visit Australia or an external Territory or the exclusive economic zone; or the species is subject of an international agreement that relates to the protection of migratory species and that binds the Commonwealth; and listing is otherwise in accordance with the ministerial guidelines (section 15 of the BC Act).

Includes birds that are subject to an agreement between the government of Australia and the governments of Japan (JAMBA), China (CAMBA) and The Republic of Korea (ROKAMBA), and fauna subject to the *Convention on the Conservation of Migratory Species of Wild Animals* (Bonn Convention), an environmental treaty under the United Nations Environment Program. Migratory species listed under the BC Act are a subset of the migratory animals, that are known to visit Western Australia, protected under the international agreements or treaties, excluding species that are listed as Threatened species.

Published as migratory birds protected under an international agreement under schedule 5 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

CD Species of special conservation interest (conservation dependent fauna)

Fauna of special conservation need being species dependent on ongoing conservation intervention to prevent it becoming eligible for listing as threatened, and listing is otherwise in accordance with the ministerial guidelines (section 14 of the BC Act).

Published as conservation dependent fauna under schedule 6 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

OS Other specially protected species

Fauna otherwise in need of special protection to ensure their conservation, and listing is otherwise in accordance with the ministerial guidelines (section 18 of the BC Act).

Published as other specially protected fauna under schedule 7 of the *Wildlife Conservation (Specially Protected Fauna) Notice 2018*.

P **Priority species**

Possibly threatened species that do not meet survey criteria, or are otherwise data deficient, are added to the Priority Fauna or Priority Flora Lists under Priorities 1, 2 or 3. These three categories are ranked in order of priority for survey and evaluation of conservation status so that consideration can be given to their declaration as threatened fauna or flora.

Species that are adequately known, are rare but not threatened, or meet criteria for near threatened, or that have been recently removed from the threatened species or other specially protected fauna lists for other than taxonomic reasons, are placed in Priority 4. These species require regular monitoring.

Assessment of Priority codes is based on the Western Australian distribution of the species, unless the distribution in WA is part of a contiguous population extending into adjacent States, as defined by the known spread of locations.

1 **Priority 1: Poorly-known species**

Species that are known from one or a few locations (generally five or less) which are potentially at risk. All occurrences are either: very small; or on lands not managed for conservation, e.g. agricultural or pastoral lands, urban areas, road and rail reserves, gravel reserves and active mineral leases; or otherwise under threat of habitat destruction or degradation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under immediate threat from known threatening processes. Such species are in urgent need of further survey.

2 **Priority 2: Poorly-known species**

Species that are known from one or a few locations (generally five or less), some of which are on lands managed primarily for nature conservation, e.g. national parks, conservation parks, nature reserves and other lands with secure tenure being managed for conservation. Species may be included if they are comparatively well known from one or more locations but do not meet adequacy of survey requirements and appear to be under threat from known threatening processes. Such species are in urgent need of further survey.

3 **Priority 3: Poorly-known species**

Species that are known from several locations, and the species does not appear to be under imminent threat, or from few but widespread locations with either large population size or significant remaining areas of apparently suitable habitat, much of it not under imminent threat. Species may be included if they are comparatively well known from several locations but do not meet adequacy of survey requirements and known threatening processes exist that could affect them. Such species are in need of further survey.

4 **Priority 4: Rare, Near Threatened and other species in need of monitoring**

(a) Rare. Species that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection but could be if present circumstances change. These species are usually represented on conservation lands.

(b) Near Threatened. Species that are considered to have been adequately surveyed and that are close to qualifying for vulnerable but are not listed as Conservation Dependent.

(c) Species that have been removed from the list of threatened species during the past five years for reasons other than taxonomy.

¹ The definition of flora includes algae, fungi and lichens

² Species includes all taxa (plural of taxon - a classificatory group of any taxonomic rank, e.g. a family, genus, species or any infraspecific category i.e. subspecies or variety, or a distinct population).

DEFINITIONS, CATEGORIES AND CRITERIA FOR THREATENED AND PRIORITY ECOLOGICAL COMMUNITIES

1. GENERAL DEFINITIONS

Ecological Community

A naturally occurring biological assemblage that occurs in a particular type of habitat.

Note: The scale at which ecological communities are defined will often depend on the level of detail in the information source, therefore no particular scale is specified.

A **threatened ecological community** (TEC) is one which is found to fit into one of the following categories; “presumed totally destroyed”, “critically endangered”, “endangered” or “vulnerable”.

Possible threatened ecological communities that do not meet survey criteria are added to DEC’s Priority Ecological Community Lists under Priorities 1, 2 and 3. Ecological Communities that are adequately known, are rare but not threatened, or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

An **assemblage** is a defined group of biological entities.

Habitat is defined as the areas in which an organism and/or assemblage of organisms lives. It includes the abiotic factors (eg. substrate and topography), and the biotic factors.

Occurrence: a discrete example of an ecological community, separated from other examples of the same community by more than 20 metres of a different ecological community, an artificial surface or a totally destroyed community.

By ensuring that every discrete occurrence is recognised and recorded future changes in status can be readily monitored.

Adequately Surveyed is defined as follows:

“An ecological community that has been searched for thoroughly in most likely habitats, by relevant experts.”

Community structure is defined as follows:

“The spatial organisation, construction and arrangement of the biological elements comprising a biological assemblage” (eg. *Eucalyptus salmonophloia* woodland over scattered small shrubs over dense herbs; structure in a faunal assemblage could refer to trophic structure, eg. dominance by feeders on detritus as distinct from feeders on live plants).

Definitions of Modification and Destruction of an ecological community:

Modification: “changes to some or all of ecological processes (including abiotic processes such as hydrology), species composition and community structure as a

direct or indirect result of human activities. The level of damage involved could be ameliorated naturally or by human intervention.”

Destruction: “modification such that reestablishment of ecological processes, species composition and community structure within the range of variability exhibited by the original community is unlikely within the foreseeable future even with positive human intervention.”

Note: Modification and destruction are difficult concepts to quantify, and their application will be determined by scientific judgement. Examples of modification and total destruction are cited below:

Modification of ecological processes: The hydrology of Toolibin Lake has been altered by clearing of the catchment such that death of some of the original flora has occurred due to dependence on fresh water. The system may be bought back to a semblance of the original state by redirecting saline runoff and pumping waters of the rising watertable away to restore the hydrological balance. Total destruction of downstream lakes has occurred due to hydrology being altered to the point that few of the original flora or fauna species are able to tolerate the level of salinity and/or water logging.

Modification of structure: The understorey of a plant community may be altered by weed invasion due to nutrient enrichment by addition of fertiliser. Should the additional nutrients be removed from the system the balance may be restored, and the original plant species better able to compete. Total destruction may occur if additional nutrients continue to be added to the system causing the understorey to be completely replaced by weed species, and death of overstorey species due to inability to tolerate high nutrient levels.

Modification of species composition: Pollution may cause alteration of the invertebrate species present in a freshwater lake. Removal of pollutants may allow the return of the original inhabitant species. Addition of residual highly toxic substances may cause permanent changes to water quality, and total destruction of the community.

Threatening processes are defined as follows:

“Any process or activity that threatens to destroy or significantly modify the ecological community and/or affect the continuing evolutionary processes within any ecological community.”

Examples of some of the continuing threatening processes in Western Australia include: general pollution; competition, predation and change induced in ecological communities as a result of introduced animals; competition and displacement of native plants by introduced species; hydrological changes; inappropriate fire regimes; diseases resulting from introduced microorganisms; direct human exploitation and disturbance of ecological communities.

Restoration is defined as returning an ecological community to its pre-disturbance or natural state in terms of abiotic conditions, community structure and species composition.

Rehabilitation is defined as the re-establishment of ecological attributes in a damaged ecological community although the community will remain modified.

2. DEFINITIONS AND CRITERIA FOR PRESUMED TOTALLY DESTROYED, CRITICALLY ENDANGERED, ENDANGERED AND VULNERABLE ECOLOGICAL COMMUNITIES

Presumed Totally Destroyed (PD)

An ecological community that has been adequately searched for but for which no representative occurrences have been located. The community has been found to be totally destroyed or so extensively modified throughout its range that no occurrence of it is likely to recover its species composition and/or structure in the foreseeable future.

An ecological community will be listed as presumed totally destroyed if there are no recent records of the community being extant **and either** of the following applies (A or B):

- A) Records within the last 50 years have not been confirmed despite thorough searches of known or likely habitats **or**
- B) All occurrences recorded within the last 50 years have since been destroyed

Critically Endangered (CR)

An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or that was originally of limited distribution and is facing severe modification or destruction throughout its range in the immediate future, or is already severely degraded throughout its range but capable of being substantially restored or rehabilitated.

An ecological community will be listed as **Critically Endangered** when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future. This will be determined on the basis of the best available information, by it meeting **any one or more of** the following criteria (A, B or C):

- A) The estimated geographic range, and/or total area occupied, and/or number of discrete occurrences since European settlement have been reduced by at least 90% **and either or both** of the following apply (i or ii):
 - i) geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is imminent (within approximately 10 years);
 - ii) modification throughout its range is continuing such that in the immediate future (within approximately 10 years) the community is unlikely to be capable of being substantially rehabilitated.
- B) Current distribution is limited, **and one or more** of the following apply (i, ii or iii):
 - i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the immediate future (within approximately 10 years);

ii) there are very few occurrences, each of which is small and/or isolated and extremely vulnerable to known threatening processes;

iii) there may be many occurrences but total area is very small and each occurrence is small and/or isolated and extremely vulnerable to known threatening processes.

C) The ecological community exists only as highly modified occurrences that may be capable of being rehabilitated if such work begins in the immediate future (within approximately 10 years).

Endangered (EN)

An ecological community that has been adequately surveyed and found to have been subject to a major contraction in area and/or was originally of limited distribution and is in danger of significant modification throughout its range or severe modification or destruction over most of its range in the near future.

An ecological community will be listed as **Endangered** when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future. This will be determined on the basis of the best available information by it meeting **any one or more of** the following criteria (A, B, or C):

A) The geographic range, and/or total area occupied, and/or number of discrete occurrences have been reduced by at least 70% since European settlement **and either or both** of the following apply (i or ii):

i) the estimated geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is likely in the short term future (within approximately 20 years);

ii) modification throughout its range is continuing such that in the short term future (within approximately 20 years) the community is unlikely to be capable of being substantially restored or rehabilitated.

B) Current distribution is limited, **and one or more** of the following apply (i, ii or iii):

i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the short term future (within approximately 20 years);

ii) there are few occurrences, each of which is small and/or isolated and all or most occurrences are very vulnerable to known threatening processes;

iii) there may be many occurrences but total area is small and all or most occurrences are small and/or isolated and very vulnerable to known threatening processes.

C) The ecological community exists only as very modified occurrences that may be capable of being substantially restored or rehabilitated if such work begins in the short-term future (within approximately 20 years).

Vulnerable (VU)

An ecological community that has been adequately surveyed and is found to be declining and/or has declined in distribution and/or condition and whose ultimate security has not yet been assured and/or a community that is still widespread but is believed likely to move into a category of higher threat in the near future if threatening processes continue or begin operating throughout its range.

An ecological community will be listed as **Vulnerable** when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction or significant modification in the medium (within approximately 50 years) to long-term future. This will be determined on the basis of the best available information by it meeting **any one or more of** the following criteria (A, B or C):

A) The ecological community exists largely as modified occurrences that are likely to be capable of being substantially restored or rehabilitated.

B) The ecological community may already be modified and would be vulnerable to threatening processes, is restricted in area and/or range and/or is only found at a few locations.

C) The ecological community may be still widespread but is believed likely to move into a category of higher threat in the medium to long-term future because of existing or impending threatening processes.

3. DEFINITIONS AND CRITERIA FOR PRIORITY ECOLOGICAL COMMUNITIES

Possible threatened ecological communities that do not meet survey criteria or that are not adequately defined are added to the Priority Ecological Community List under priorities 1, 2 and 3. These three categories are ranked in order of priority for survey and/or definition of the community. Ecological communities that are adequately known, and are rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

Priority One: Poorly-known ecological communities

Ecological communities that are known from very few occurrences with a very restricted distribution (generally ≤ 5 occurrences or a total area of $\leq 100\text{ha}$). Occurrences are believed to be under threat either due to limited extent, or being on lands under immediate threat (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) or for which current threats exist. May include communities with occurrences on protected lands. Communities may be included if they are comparatively well-known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.

Priority Two: Poorly-known ecological communities

Communities that are known from few occurrences with a restricted distribution (generally ≤ 10 occurrences or a total area of $\leq 200\text{ha}$). At least some occurrences are not believed to be under immediate threat (within approximately 10 years) of destruction or degradation. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.

Priority Three: Poorly known ecological communities

- (i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or:
- (ii) communities known from a few widespread occurrences, which are either large or with significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat (within approximately 10 years), or;
- (iii) communities made up of large, and/or widespread occurrences, that may or may not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, inappropriate fire regimes, clearing, hydrological change etc.

Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.

Priority Four: Ecological communities that are adequately known, rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.

- (i) Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands.
- (ii) Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for a higher threat category.
- (iii) Ecological communities that have been removed from the list of threatened communities during the past five years.

Priority Five: Conservation Dependent ecological communities

Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

2. Commonwealth *Environment Protection and Biodiversity Conservation Act 1999*

Many of the species that are specially protected at State level are also listed as Threatened species at the Federal level, as one of the Matters of National Environmental Significance (MNES) identified under the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (the EPBC Act). These may be classified as 'critically endangered', 'endangered', 'vulnerable' or 'lower risk', consistent with IUCN categories:

1. **Critically Endangered (CR):** a taxon is Critically Endangered when it is facing an extremely high risk of extinction in the wild in the immediate future.
2. **Endangered (EN):** a taxon is Endangered when it is not Critically Endangered but is facing a very high risk of extinction in the wild in the near future.
3. **Vulnerable (VU):** a taxon is Vulnerable when it is not Critically Endangered or Endangered but is facing a high risk of extinction in the wild in the medium-term future.
4. **Lower Risk (LR):** a taxon is Lower Risk when it has been evaluated, does not satisfy the criteria for any of the categories Critically Endangered, Endangered or Vulnerable. Taxa included in the Lower Risk category can be separated into three subcategories:
 - **Conservation Dependent (CD).** Taxa which are the focus of a continuing taxon-specific or habitat-specific conservation program targeted towards the taxon in question, the cessation of which would result in the taxon qualifying for one of the threatened categories above within a period of five years.
 - **Near Threatened (NT).** Taxa which do not qualify for Conservation Dependent, but which are close to qualifying for Vulnerable.
 - **Least Concern (LC).** Taxa which do not qualify for Conservation Dependent or Near Threatened.

In addition, numerous Migratory species are listed as MNES under the EPBC Act (some of which are also listed as Threatened). Migratory species are those animals that migrate to Australia and its external territories, or pass through or over Australian waters during their annual migrations. The list of migratory species consists of those species listed under the following international conventions:

1. Convention on the Conservation of Migratory Species of Wild Animals (Bonn Convention);
2. China-Australia Migratory Bird Agreement (CAMBA);
3. Japan-Australia Migratory Bird Agreement (JAMBA); and,
4. Republic of Korea-Australia Migratory Bird Agreement (ROKAMBA).

Marine species are also protected under the EPBC Act, and are listed to ensure the long-term conservation of the species. Marine species include all Australian sea snakes, seals, crocodiles, dugongs, marine turtles, seahorses and seabirds that naturally occur in the Commonwealth marine area.

Under the terms of the EPBC Act, an action (e.g. a project or development) is required to be referred to the Australian Government Environment Minister for approval if it has, will have, or is likely to have, a significant impact on an MNES. The term 'action' includes projects and developments subsequent to commencement of the Act, however there are a number of exemptions (e.g. projects in Commonwealth areas). According to Department of the Environment (2013), a 'significant impact' is an impact which is important, notable, or of consequence, having regard to its context or intensity. Whether or not an action is likely to have a significant impact depends upon the sensitivity, value, and quality of the environment which is impacted, and upon the intensity, duration, magnitude and geographic extent of the impacts.

References:

Department of the Environment (2013). Matters of National Environmental Significance - Significant Impact Guidelines 1.1 *Environment Protection and Biodiversity Conservation Act 1999*. Department of the Environment, Canberra, Australia.

Appendix 2

NatureMap Database Search Results



NatureMap Species Report

Created By Nicola Watson on 29/10/2020

Current Names Only Yes
Core Datasets Only Yes
Method 'By Circle'
Centre 122° 58' 32" E, 16° 28' 05" S
Buffer 2. km
Group By Kingdom

Kingdom	Species	Records
Animalia	397	2802
Chromista	26	100
Fungi	22	40
Plantae	628	1897
TOTAL	1073	4839

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
Animalia				
1.	??			
2.	<i>Abudefduf bengalensis</i>			
3.	<i>Abudefduf</i> sp.			
4.	<i>Acanthopagrus latus</i>			
5.	<i>Acanthurus grammoptilus</i>			
6.	25535 <i>Accipiter cirrocephalus</i> (Collared Sparrowhawk)			
7.	24281 <i>Accipiter cirrocephalus</i> subsp. <i>cirrocephalus</i> (Collared Sparrowhawk)			
8.	25536 <i>Accipiter fasciatus</i> (Brown Goshawk)			
9.	<i>Acentrogobius gracilis</i>			
10.	41323 <i>Actitis hypoleucos</i> (Common Sandpiper)		IA	
11.	25544 <i>Aegotheles cristatus</i> (Australian Owlet-nightjar)			
12.	42372 <i>Amalosia rhombifer</i> (Zigzag velvet gecko)			
13.	<i>Amblygobius bynoensis</i>			
14.	30831 <i>Amphibolurus gilberti</i> (Ta-ta, Gilbert's Dragon)			
15.	<i>Amphiprion rubrocinctus</i>			
16.	24316 <i>Anas superciliosa</i> (Pacific Black Duck)			
17.	47414 <i>Anhinga novaehollandiae</i> (Australasian Darter)			
18.	25634 <i>Anous stolidus</i> (Common Noddy)		IA	
19.	24505 <i>Anous stolidus</i> subsp. <i>pileatus</i> (Common Noddy)		IA	
20.	25317 <i>Antaresia childreni</i> (Children's Python)			
21.	25241 <i>Antaresia stimsoni</i> subsp. <i>stimsoni</i> (Stimson's Python)			
22.	<i>Apogon cookii</i>			
23.	<i>Apogon rueppellii</i>			
24.	<i>Apogon</i> sp.			
25.	<i>Apogon timorensis</i>			
26.	24719 <i>Aprosmictus erythropterus</i> (Red-winged Parrot)			
27.	25554 <i>Apus pacificus</i> (Fork-tailed Swift, Pacific Swift)		IA	
28.	24285 <i>Aquila audax</i> (Wedge-tailed Eagle)			
29.	25559 <i>Ardea intermedia</i> (Intermediate Egret)			
30.	41324 <i>Ardea modesta</i> (great egret, white egret)			
31.	24340 <i>Ardea novaehollandiae</i> (White-faced Heron)			
32.	25560 <i>Ardea sacra</i> (Eastern Reef Egret, Eastern Reef Heron)			
33.	24343 <i>Ardea sacra</i> subsp. <i>sacra</i> (Eastern Reef Egret, Eastern Reef Heron)			
34.	24344 <i>Ardea sumatrana</i> (Great-billed Heron)			
35.	24610 <i>Ardeotis australis</i> (Australian Bustard)			
36.	25736 <i>Arenaria interpres</i> (Ruddy Turnstone)		IA	
37.	<i>Arothron hispidus</i>			
38.	25566 <i>Artamus cinereus</i> (Black-faced Woodswallow)			
39.	25567 <i>Artamus leucorhynchus</i> (White-breasted Woodswallow)			
40.	24354 <i>Artamus leucorhynchus</i> subsp. <i>leucopygialis</i> (White-breasted Woodswallow)			
41.	24355 <i>Artamus minor</i> (Little Woodswallow)			
42.	24356 <i>Artamus personatus</i> (Masked Woodswallow)			
43.	<i>Assiculoides desmonotus</i>			

Name ID	Species Name	Naturalised	Conservation Code	Endemic To Query Area
44.	<i>Atelomycterus macleayi</i>			
45.	<i>Atherinid</i> sp.			
46.	<i>Austracantha minax</i>			
47.	24318 <i>Aythya australis</i> (Hardhead)			
48.	<i>Bathygobius fuscus</i>			
49.	<i>Batrachomoeus dahli</i>			
50.	<i>Blennodesmus scapularis</i>			
51.	24251 <i>Bos taurus</i> (European Cattle)	Y		
52.	24359 <i>Burhinus grallarius</i> (Bush Stone-curlew)			
53.	47897 <i>Butorides striata</i> (Striated Heron, Mangrove Heron)			
54.	25716 <i>Cacatua sanguinea</i> (Little Corella)			
55.	24728 <i>Cacatua sanguinea</i> subsp. <i>sanguinea</i> (Little Corella)			
56.	42307 <i>Cacomantis pallidus</i> (Pallid Cuckoo)			
57.	25599 <i>Cacomantis variolosus</i> (Brush Cuckoo)			
58.	24428 <i>Cacomantis variolosus</i> subsp. <i>variolosus</i> (Brush Cuckoo)			
59.	24779 <i>Calidris acuminata</i> (Sharp-tailed Sandpiper)		IA	
60.	24780 <i>Calidris alba</i> (Sanderling)		IA	
61.	25738 <i>Calidris canutus</i> (Red Knot, knot)		IA	
62.	24784 <i>Calidris ferruginea</i> (Curlew Sandpiper)		T	
63.	24788 <i>Calidris ruficollis</i> (Red-necked Stint)		IA	
64.	24790 <i>Calidris tenuirostris</i> (Great Knot)		T	
65.	<i>Callionymus</i> sp.			
66.	25717 <i>Calyptorhynchus banksii</i> (Red-tailed Black-Cockatoo)			
67.	24730 <i>Calyptorhynchus banksii</i> subsp. <i>macrorhynchus</i> (Northern Red-tailed Black-Cockatoo)			
68.	24253 <i>Capra hircus</i> (Goat)	Y		
69.	25015 <i>Carlia munda</i> (Shaded-litter Rainbow Skink)			
70.	25017 <i>Carlia triacantha</i> (Desert Rainbow Skink)			
71.	<i>Centrogenys vaigiensis</i>			
72.	25600 <i>Centropus phasianinus</i> (Pheasant Coucal)			
73.	30884 <i>Centropus phasianinus</i> subsp. <i>phasianinus</i> (Pheasant Coucal)			
74.	47905 <i>Ceyx azureus</i> (Azure Kingfisher)			
75.	<i>Chaetodontoplus mesoleucus</i>			
76.	25583 <i>Chalcophaps indica</i> (Emerald Dove)			
77.	24186 <i>Chalinolobus gouldii</i> (Gould's Wattled Bat)			
78.	24188 <i>Chalinolobus nigrogriseus</i> (Hoary Wattled Bat)			
79.	25575 <i>Charadrius leschenaultii</i> (Greater Sand Plover)		T	
80.	25576 <i>Charadrius mongolus</i> (Lesser Sand Plover)		T	
81.	24377 <i>Charadrius ruficapillus</i> (Red-capped Plover)			
82.	<i>Chelmon marginalis</i>			
83.	25336 <i>Chelonia mydas</i> (Green Turtle)		T	
84.	24862 <i>Chelosania brunnea</i> (Chameleon Dragon)			
85.	24863 <i>Chlamydosaurus kingii</i> (Frill-necked Lizard)			
86.	<i>Choerodon cyanodus</i>			
87.	<i>Choerichthys brachysoma</i>			
88.	<i>Chroicocephalus novaehollandiae</i>			
89.	25602 <i>Chrysococcyx minutillus</i> (Little Bronze Cuckoo)			
90.	24433 <i>Chrysococcyx minutillus</i> subsp. <i>minutillus</i> (Little Bronze Cuckoo)			
91.	24289 <i>Circus assimilis</i> (Spotted Harrier)			
92.	24565 <i>Cissomela pectoralis</i> (Banded Honeyeater)			
93.	<i>Clupeid</i> sp.			
94.	25675 <i>Colluricincla harmonica</i> (Grey Shrike-thrush)			
95.	24611 <i>Colluricincla harmonica</i> subsp. <i>brunnea</i> (Grey Shrike-thrush)			
96.	<i>Congrogadus subducens</i>			
97.	24566 <i>Conopophila rufogularis</i> (Rufous-throated Honeyeater)			
98.	25568 <i>Coracina novaehollandiae</i> (Black-faced Cuckoo-shrike)			
99.	25593 <i>Corvus orru</i> (Torresian Crow)			
100.	25701 <i>Coturnix ypsilophora</i> (Brown Quail)			
101.	24672 <i>Coturnix ypsilophora</i> subsp. <i>cervina</i> (Brown Quail)			
102.	24420 <i>Cracticus nigrogularis</i> (Pied Butcherbird)			
103.	25595 <i>Cracticus tibicen</i> (Australian Magpie)			
104.	<i>Craterocephalus pauciradiatus</i>			
105.	30891 <i>Cryptoblepharus tyttos</i>			
106.	<i>Cryptocentroides insignis</i>			
107.	25033 <i>Ctenotus colletti</i>			
108.	25048 <i>Ctenotus inornatus</i>			
109.	25371 <i>Cyclorana australis</i> (Giant Frog)			
110.	<i>Cymbacephalus bosschei</i>			
111.	<i>Cymbacephalus nematophthalmus</i>			
112.	25547 <i>Dacelo leachii</i> (Blue-winged Kookaburra)			
113.	25673 <i>Daphoenositta chrysoptera</i> (Varied Sittella)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
114.	24605 <i>Daphoenositta chrysoptera</i> subsp. <i>leucoptera</i> (Varied Sittella, White-winged Sittella)			
115.	24996 <i>Delma borea</i>			
116.	25004 <i>Delma tincta</i>			
117.	42390 <i>Demansia angusticeps</i>			
118.	25294 <i>Demansia papuensis</i> (Great Black Whipsnake)			
119.	25325 <i>Dendrelaphis punctulata</i> (Green Tree Snake)			
120.	25607 <i>Dicaeum hirundinaceum</i> (Mistletoebird)			
121.	24441 <i>Dicaeum hirundinaceum</i> subsp. <i>hirundinaceum</i> (Mistletoebird)			
122.	24442 <i>Dicrurus bracteatus</i> subsp. <i>bracteatus</i> (Spangled Drongo)			
123.	<i>Didymothallus mizolepis</i>			
124.	<i>Dinematichthys trilobatus</i>			
125.	24896 <i>Diporiphora pindan</i> (Pindan Dragon)			
126.	<i>Dischistodus darwiniensis</i>			
127.	<i>Dischistodus fasciatus</i>			
128.	24084 <i>Dugong dugon</i> (Dugong)		S	
129.	<i>Egretta garzetta</i>			
130.	<i>Egretta novaehollandiae</i>			
131.	<i>Elanus axillaris</i>			
132.	25540 <i>Elanus caeruleus</i> (Black-shouldered Kite)			
133.	47937 <i>Euseyornis melanops</i> (Black-fronted Dotterel)			
134.	<i>Enneapterygius</i> sp.			
135.	<i>Eolophus roseicapillus</i>			
136.	24653 <i>Eopsaltria pulverulenta</i> (Mangrove Robin)			
137.	25578 <i>Ephippiorhynchus asiaticus</i> (Black-necked Stork)			
138.	24387 <i>Ephippiorhynchus asiaticus</i> subsp. <i>australis</i> (Black-necked Stork)			
139.	<i>Epinephelus coioides</i>			
140.	<i>Epinephelus corallicola</i>			
141.	<i>Epinephelus ongus</i>			
142.	<i>Epinephelus quoyanus</i>			
143.	24569 <i>Epthianura crocea</i> (Yellow Chat)			
144.	42404 <i>Eremiascincus isolepis</i>			
145.	<i>Eremiascincus isolepis</i>			
146.	24632 <i>Erythrura gouldiae</i> (Gouldian Finch)		P4	
147.	47938 <i>Esacus magnirostris</i> (Beach Stone-curlew, Beach Thick-knee)			
148.	24760 <i>Eulabeornis castaneiventris</i> subsp. <i>castaneiventris</i> (Chestnut Rail)			
149.	24368 <i>Eurostopodus argus</i> (Spotted Nightjar)			
150.	25591 <i>Eurystomus orientalis</i> (Dollarbird)			
151.	25621 <i>Falco berigora</i> (Brown Falcon)			
152.	25622 <i>Falco cenchroides</i> (Australian Kestrel, Nankeen Kestrel)			
153.	24472 <i>Falco cenchroides</i> subsp. <i>cenchrus</i> (Australian Kestrel, Nankeen Kestrel)			
154.	25623 <i>Falco longipennis</i> (Australian Hobby)			
155.	25624 <i>Falco peregrinus</i> (Peregrine Falcon)		S	
156.	<i>Favonigobius</i> n. sp.			Y
157.	<i>Fistularia commersonii</i>			
158.	25327 <i>Fordonia leucobalia</i> (White-bellied Mangrove Snake)			
159.	<i>Fowleria aurita</i>			
160.	24478 <i>Fregata ariel</i> (Lesser Frigatebird)		IA	
161.	25301 <i>Furina ornata</i> (Moon Snake)			
162.	24765 <i>Gallirallus philippensis</i> subsp. <i>mellori</i> (Buff-banded Rail)			
163.	42314 <i>Gavicalis virescens</i> (Singing Honeyeater)			
164.	24952 <i>Gehyra australis</i>			
165.	24954 <i>Gehyra nana</i>			
166.	24955 <i>Gehyra occidentalis</i>			
167.	24956 <i>Gehyra pilbara</i>			
168.	24959 <i>Gehyra variegata</i>			
169.	47954 <i>Gelochelidon nilotica</i> (Gull-billed Tern)		IA	
170.	24401 <i>Geopelia cuneata</i> (Diamond Dove)			
171.	24402 <i>Geopelia humeralis</i> (Bar-shouldered Dove)			
172.	25585 <i>Geopelia striata</i> (Zebra Dove)			
173.	24403 <i>Geopelia striata</i> subsp. <i>placida</i> (Peaceful Dove)			
174.	<i>Gerres oyena</i>			
175.	<i>Gerres</i> sp.			
176.	25531 <i>Gerygone levigaster</i> (Mangrove Gerygone)			
177.	24273 <i>Gerygone levigaster</i> subsp. <i>levigaster</i> (Mangrove Gerygone)			
178.	24274 <i>Gerygone magnirostris</i> subsp. <i>magnirostris</i> (Large-billed Gerygone)			
179.	24276 <i>Gerygone tenebrosa</i> (Dusky Gerygone)			
180.	<i>Gnatholepis argus</i>			
181.	<i>Gobiid</i> n. sp.			Y
182.	<i>Gobiid</i> sp.			
183.	<i>Gobiodon quinquestrigatus</i>			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
184.	24443	<i>Grallina cyanoleuca</i> (Magpie-lark)			
185.	24484	<i>Grus rubicunda</i> (Brolga)			
186.	25627	<i>Haematopus fuliginosus</i> (Sooty Oystercatcher)			
187.	24486	<i>Haematopus fuliginosus</i> subsp. <i>ophthalmicus</i> (Sooty Oystercatcher)			
188.	24487	<i>Haematopus longirostris</i> (Pied Oystercatcher)			
189.	24293	<i>Haliaeetus leucogaster</i> (White-bellied Sea-Eagle)			
190.	25541	<i>Haliastur indus</i> (Brahminy Kite)			
191.	24295	<i>Haliastur spheonurus</i> (Whistling Kite)			
192.		<i>Halichoeres melanurus</i>			
193.		<i>Halichoeres nigrescens</i>			
194.		<i>Halophryne diemensis</i>			
195.	24961	<i>Heteronotia binoei</i> (Bynoe's Gecko)			
196.	48587	<i>Hydroprogne caspia</i> (Caspian Tern)		IA	
197.		<i>Hypoatherina temminckii</i>			
198.		<i>Hyporhamphus quoyi</i>			
199.		<i>Istiblennius meleagris</i>			
200.		<i>Istiblennius meleagris?</i>			Y
201.		<i>Istigobius ornatus</i>			
202.	25571	<i>Lalage leucomela</i> (Varied Triller)			
203.	24511	<i>Larus novaehollandiae</i> subsp. <i>novaehollandiae</i> (Silver Gull)			
204.	25343	<i>Lepidochelys olivacea</i> (Olive Ridley Turtle, Pacific Ridley Turtle)		T	
205.	25121	<i>Lerista apoda</i>			
206.	25125	<i>Lerista bipes</i>			
207.	25138	<i>Lerista griffini</i>			
208.	25170	<i>Lerista separanda</i> (Dampierland plain slider, skink)		P2	
209.		<i>Lethrinus laticaudis</i>			
210.	25005	<i>Lialis burtonis</i>			
211.	25239	<i>Liasis olivaceus</i> subsp. <i>olivaceus</i> (Olive Python)			
212.	25661	<i>Lichmera indistincta</i> (Brown Honeyeater)			
213.	24582	<i>Lichmera indistincta</i> subsp. <i>indistincta</i> (Brown Honeyeater)			
214.	30932	<i>Limosa lapponica</i> (Bar-tailed Godwit)		IA	
215.	25741	<i>Limosa limosa</i> (Black-tailed Godwit)		IA	
216.	25380	<i>Litoria caerulea</i> (Green Tree Frog)			
217.	25391	<i>Litoria rothii</i> (Northern Laughing Tree Frog)			
218.		<i>Liza vaigiensis</i>			
219.	25683	<i>Lonchura castaneothorax</i> (Chestnut-breasted Mannikin)			
220.		<i>Lophiochoron trisignatus</i>			
221.		<i>Lutjanus carponotatus</i>			
222.		<i>Lutjanus russellii</i>			
223.	24171	<i>Macroglossus minimus</i> (Northern Blossom-bat)			
224.	24168	<i>Macrotis lagotis</i> (Bilby, Dalgyte, Ninu)		T	
225.	25651	<i>Malurus lamberti</i> (Variegated Fairy-wren)			
226.	25653	<i>Malurus melanocephalus</i> (Red-backed Fairy-wren)			
227.	24051	<i>Megaptera novaeangliae</i> (Humpback Whale)		S	
228.	47997	<i>Melanodryas cucullata</i> (Hooded Robin)			
229.	24585	<i>Melithreptus albogularis</i> (White-throated Honeyeater)			
230.	25665	<i>Melithreptus gularis</i> (Black-chinned Honeyeater)			
231.	24589	<i>Melithreptus gularis</i> subsp. <i>laetior</i> (Black-chinned Honeyeater)			
232.	24220	<i>Melomys burtoni</i> (Grassland Melomys)			
233.	24736	<i>Melopsittacus undulatus</i> (Budgerigar)			
234.	25185	<i>Menetia maini</i>			
235.	24598	<i>Merops ornatus</i> (Rainbow Bee-eater)			
236.		<i>Microcarbo melanoleucos</i>			
237.		<i>Microdesmid</i> sp.			Y
238.	25693	<i>Microeca fascinans</i> (Jacky Winter)			
239.	24654	<i>Microeca fascinans</i> subsp. <i>assimilis</i> (Jacky Winter)			
240.	25694	<i>Microeca flavigaster</i> (Lemon-breasted Flycatcher)			
241.	24657	<i>Microeca flavigaster</i> subsp. <i>tormenti</i> (Kimberley Flycatcher)			
242.	25542	<i>Milvus migrans</i> (Black Kite)			
243.	24298	<i>Milvus migrans</i> subsp. <i>affinis</i> (Black Kite)			
244.	25493	<i>Miniopterus schreibersii</i> (Common Bentwing-bat)			
245.	24190	<i>Miniopterus schreibersii</i> subsp. <i>oriana</i> (Northern Bentwing-bat)			
246.	25195	<i>Morethia storri</i>			
247.		<i>Mormopterus</i> (<i>Ozimops</i>) <i>cobourgianus</i>			
248.	24183	<i>Mormopterus loriae</i> (Little Northern Freetail-bat)			
249.		<i>Muraenichthys</i> sp.			
250.	25609	<i>Myiagra alecto</i> (Shining Flycatcher)			
251.	24445	<i>Myiagra alecto</i> subsp. <i>melvillensis</i> (Shining Flycatcher)			
252.	25610	<i>Myiagra inquieta</i> (Restless Flycatcher)			
253.	25611	<i>Myiagra rubecula</i> (Leaden Flycatcher)			

Name ID	Species Name	Naturalised	Conservation Code	Endemic To Query Area
254.	24449 <i>Myiagra rubecula</i> subsp. <i>concinna</i> (Leaden Flycatcher)			
255.	25612 <i>Myiagra ruficollis</i> (Broad-billed Flycatcher)			
256.	24450 <i>Myiagra ruficollis</i> subsp. <i>mimikae</i> (Broad-billed Flycatcher)			
257.	25666 <i>Myzomela erythrocephala</i> (Red-headed Honeyeater)			
258.	24590 <i>Myzomela erythrocephala</i> subsp. <i>erythrocephala</i> (Red-headed Honeyeater)			
259.	25747 <i>Ninox connivens</i> (Barking Owl)			
260.	<i>Notograpus</i> sp.			
261.	25198 <i>Notoscincus ornatus</i> subsp. <i>wotjulum</i>			
262.	24798 <i>Numenius madagascariensis</i> (Eastern Curlew)		T	
263.	24799 <i>Numenius minutus</i> (Little Curlew, Little Whimbrel)		IA	
264.	25742 <i>Numenius phaeopus</i> (Whimbrel)		IA	
265.	25564 <i>Nycticorax caledonicus</i> (Rufous Night Heron)			
266.	24192 <i>Nyctophilus arnhemensis</i> (Arnhem Land Long-eared Bat)			
267.	42365 <i>Nyctophilus daedalus</i> (Northwestern Long-eared Bat, Pallid Long-eared Bat)			
268.	24975 <i>Oedura gracilis</i>			
269.	<i>Omobranchus germaini</i>			
270.	<i>Omobranchus lineolatus</i>			
271.	41347 <i>Onychoprion anaethetus</i> (Bridled Tern)		IA	
272.	<i>Onychoprion fuscata</i>			
273.	<i>Opistognathus darwiniensis</i>			
274.	24060 <i>Orcaella heinsohni</i> (Australian Snubfin Dolphin)		P4	
275.	24607 <i>Oriolus flavocinctus</i> (Yellow Oriole)			
276.	24608 <i>Oriolus sagittatus</i> (Olive-backed Oriole)			
277.	24620 <i>Pachycephala lanioides</i> (White-breasted Whistler)			
278.	25678 <i>Pachycephala melanura</i> (Mangrove Golden Whistler)			
279.	24621 <i>Pachycephala melanura</i> subsp. <i>melanura</i> (Mangrove Golden Whistler)			
280.	25680 <i>Pachycephala rufiventris</i> (Rufous Whistler)			
281.	24624 <i>Pachycephala rufiventris</i> subsp. <i>rufiventris</i> (Rufous Whistler)			
282.	48591 <i>Pandion cristatus</i> (Osprey, Eastern Osprey)		IA	
283.	<i>Paracentropogon vespa</i>			
284.	<i>Parachaetodon ocellatus</i>			
285.	<i>Paradiplogrammus enneactis</i>			
286.	<i>Paraplotosus albilabris</i>			
287.	24627 <i>Pardalotus rubricatus</i> (Red-browed Pardalote)			
288.	25682 <i>Pardalotus striatus</i> (Striated Pardalote)			
289.	<i>Pelates quadrilineatus</i>			
290.	24648 <i>Pelecanus conspicillatus</i> (Australian Pelican)			
291.	<i>Periophthalmus argentilineatus</i>			
292.	48060 <i>Petrochelidon ariel</i> (Fairy Martin)			
293.	48061 <i>Petrochelidon nigricans</i> (Tree Martin)			
294.	24663 <i>Phaethon rubricauda</i> (Red-tailed Tropicbird)		P4	
295.	25699 <i>Phalacrocorax varius</i> (Pied Cormorant)			
296.	24409 <i>Phaps chalcoptera</i> (Common Bronzewing)			
297.	25667 <i>Philemon argenticeps</i> (Silver-crowned Friarbird)			
298.	24591 <i>Philemon argenticeps</i> subsp. <i>argenticeps</i> (Silver-crowned Friarbird)			
299.	25668 <i>Philemon citreogularis</i> (Little Friarbird)			
300.	24198 <i>Pipistrellus westralis</i> (Northern Pipistrelle)			
301.	24102 <i>Planigale maculata</i> (Common Planigale)			
302.	24842 <i>Platalea regia</i> (Royal Spoonbill)			
303.	<i>Platycephalid</i> sp.			
304.	42305 <i>Platyleptum ornatum</i> (Omate Burrowing Frog)			
305.	<i>Plectropomus maculatus</i>			
306.	24382 <i>Pluvialis fulva</i> (Pacific Golden Plover)		IA	
307.	24383 <i>Pluvialis squatarola</i> (Grey Plover)		IA	
308.	25703 <i>Podargus strigoides</i> (Tawny Frogmouth)			
309.	24643 <i>Poephila acuticauda</i> (Long-tailed Finch)			
310.	24908 <i>Pogona minor</i> subsp. <i>mitchelli</i> (Dwarf Bearded Dragon)			
311.	<i>Pomacentrus littoralis</i>			
312.	<i>Pomacentrus milleri</i>			
313.	25706 <i>Pomatostomus temporalis</i> (Grey-crowned Babbler)			
314.	24771 <i>Porzana tabuensis</i> (Spotless Crane)			
315.	<i>Psammoperca waigiensis</i>			
316.	24104 <i>Pseudantechinus ningbing</i> (Ningbing Pseudantechinus)			
317.	25261 <i>Pseudechis australis</i> (Mulga Snake)			
318.	24234 <i>Pseudomys delicatulus</i> (Delicate Mouse)			
319.	42416 <i>Pseudonaja mengdeni</i> (Western Brown Snake)			
320.	<i>Pterapogon mirifica</i>			
321.	<i>Ptereleotris microlepis</i>			
322.	24172 <i>Pteropus alecto</i> (Black Flying-fox)			
323.	24173 <i>Pteropus scapulatus</i> (Little Red Flying-fox)			

Name ID	Species Name	Naturalised	Conservation Code	Endemic To Query Area
324.	25588 <i>Ptilinopus regina</i> (Rose-crowned Fruit-dove)			
325.	30946 <i>Ptilinopus regina</i> subsp. <i>ewingii</i> (Rose-crowned Fruit-dove)			
326.	25725 <i>Ptilonorhynchus nuchalis</i> (Great Bowerbird)			
327.	24758 <i>Ptilonorhynchus nuchalis</i> subsp. <i>nuchalis</i> (Great Bowerbird)			
328.	42322 <i>Ptilotula flavescens</i> subsp. <i>flavescens</i> (Yellow-tinted Honeyeater)			
329.	24597 <i>Ramsayornis fasciatus</i> (Bar-breasted Honeyeater)			
330.	24245 <i>Rattus rattus</i> (Black Rat)	Y		
331.	48096 <i>Rhipidura albiscapa</i> (Grey Fantail)			
332.	25614 <i>Rhipidura leucophrys</i> (Willie Wagtail)			
333.	24457 <i>Rhipidura phasiana</i> (Mangrove Grey Fantail)			
334.	25616 <i>Rhipidura rufiventris</i> (Northern Fantail)			
335.	24456 <i>Rhipidura rufiventris</i> subsp. <i>isura</i> (Northern Fantail)			
336.	24174 <i>Saccolaimus flaviventris</i> (Yellow-bellied Sheath-tailed Bat)			
337.	<i>Salarias sexfilum</i>			
338.	<i>Scaevius milii</i>			
339.	<i>Scarus ghobban</i>			
340.	<i>Scolecenchelys macroptera</i>			
341.	24200 <i>Scotorepens greyii</i> (Little Broad-nosed Bat)			
342.	24201 <i>Scotorepens sanborni</i> (Northern Broad-nosed Bat)			
343.	25605 <i>Scythrops novaehollandiae</i> (Channel-billed Cuckoo)			
344.	<i>Siganus doliaus</i>			
345.	<i>Siganus fuscescens</i>			
346.	<i>Siganus punctatus</i>			
347.	25268 <i>Simoselaps minimus</i> (Dampierland Burrowing Snake)		P2	
348.	30948 <i>Smicromis brevisrostris</i> (Weebill)			
349.	48108 <i>Sphecotheres vieillotii</i> (Australasian Figbird)			
350.	24522 <i>Sterna bergii</i> (Crested Tern)			
351.	25640 <i>Sterna dougallii</i> (Roseate Tern)		IA	
352.	24524 <i>Sterna dougallii</i> subsp. <i>gracilis</i> (Roseate Tern)		IA	
353.	25642 <i>Sterna hirundo</i> (Common Tern)		IA	
354.	24527 <i>Sterna hirundo</i> subsp. <i>longipennis</i> (Common Tern)		IA	
355.	48593 <i>Sternula albifrons</i> (Little Tern)		IA	
356.	42347 <i>Stomiopera unicolor</i> (White-gaped Honeyeater)			
357.	42348 <i>Stomiopera unicolor</i> subsp. <i>unicolor</i> (White-gaped Honeyeater)			
358.	24924 <i>Strophurus ciliaris</i> subsp. <i>aberrans</i>			
359.	25754 <i>Sula leucogaster</i> (Brown Booby)		IA	
360.	24828 <i>Sula leucogaster</i> subsp. <i>plotus</i> (Brown Booby)		IA	
361.	24259 <i>Sus scrofa</i> (Pig)	Y		
362.	25705 <i>Tachybaptus novaehollandiae</i> (Australasian Grebe, Black-throated Grebe)			
363.	30872 <i>Taeniopygia bichenovii</i> (Double-barred Finch)			
364.	30873 <i>Taeniopygia bichenovii</i> subsp. <i>annulosa</i> (Double-barred Finch)			
365.	30870 <i>Taeniopygia guttata</i> (Zebra Finch)			
366.	30871 <i>Taeniopygia guttata</i> subsp. <i>castanotis</i> (Zebra Finch)			
367.	24175 <i>Taphozous georgianus</i> (Common Sheath-tailed Bat)			
368.	<i>Thalasseus bengalensis</i>			
369.	48597 <i>Thalasseus bergii</i> (Crested Tern)		IA	
370.	24845 <i>Threskiornis spinicollis</i> (Straw-necked Ibis)			
371.	25202 <i>Tiliqua multifasciata</i> (Central Blue-tongue)			
372.	25548 <i>Todiramphus chloris</i> (Collared Kingfisher)			
373.	42351 <i>Todiramphus pyrrhopygius</i> (Red-backed Kingfisher)			
374.	25549 <i>Todiramphus sanctus</i> (Sacred Kingfisher)			
375.	24309 <i>Todiramphus sanctus</i> subsp. <i>sanctus</i> (Sacred Kingfisher)			
376.	25723 <i>Trichoglossus haematodus</i> (Rainbow Lorikeet)			
377.	24803 <i>Tringa brevipes</i> (Grey-tailed Tattler)		P4	
378.	24806 <i>Tringa glareola</i> (Wood Sandpiper)		IA	
379.	24808 <i>Tringa nebularia</i> (Common Greenshank, greenshank)		IA	
380.	24810 <i>Tringa totanus</i> (Common Redshank, redshank)		IA	
381.	24851 <i>Turnix velox</i> (Little Button-quail)			
382.	30954 <i>Tursiops aduncus</i> (Indo-Pacific Bottlenose Dolphin)			
383.	<i>Tylosurus</i> sp.			
384.	<i>Upeneus tragula</i>			
385.	<i>Valamugil buchani</i>			
386.	<i>Valenciennesa longipinnis</i>			
387.	<i>Valenciennesa muralis</i>			
388.	25217 <i>Varanus glauerti</i> (Kimberley Rock Monitor)			
389.	25213 <i>Varanus glebopalma</i> (Black-palmed Rock Monitor)			
390.	25218 <i>Varanus gouldii</i> (Bungarra or Sand Monitor)			
391.	25214 <i>Varanus scalaris</i> (Spotted Tree Monitor)			
392.	25526 <i>Varanus tristis</i> (Racehorse Monitor)			
393.	24203 <i>Vespadelus caurinus</i> (Western Cave Bat, Northern Cave-bat)			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
394.	41351	<i>Xenus cinereus</i> (Terek Sandpiper)		IA	
395.		<i>Yongeichthys nebulosus</i>			
396.		<i>Zenarchopterus gilli</i>			
397.	24857	<i>Zosterops luteus</i> (Yellow White-eye)			
Chromista					
398.	35220	<i>Canistrocarpus cervicornis</i>			
399.	35910	<i>Canistrocarpus crispatus</i>			
400.	36197	<i>Chnoospora implexa</i>			
401.	26693	<i>Colpomenia peregrina</i>			
402.	26694	<i>Colpomenia sinuosa</i>			
403.	26764	<i>Dictyopteris australis</i>			
404.	29954	<i>Dictyopteris woodwardia</i>			
405.	26775	<i>Dictyota ciliolata</i>			
406.	48244	<i>Feldmannia mitchelliae</i>			
407.	26946	<i>Hormophysa cuneiformis</i>			
408.	26949	<i>Hydroclathrus clathratus</i>			
409.	27043	<i>Lobophora variegata</i>			
410.	27113	<i>Padina australis</i>			
411.	27115	<i>Padina boryana</i>			
412.	27225	<i>Rosenvingea intricata</i>			
413.	35908	<i>Rosenvingea nhatrangensis</i>			
414.	44573	<i>Sargassopsis decurrens</i>			
415.	42641	<i>Sargassum aquifolium</i>			
416.	27242	<i>Sargassum flavicans</i>			
417.	27255	<i>Sargassum polycystum</i>			
418.	48301	<i>Sargassum rasta</i>			
419.	42785	<i>Sirophysalis trinodis</i>			
420.	27304	<i>Sporochnus comosus</i>			
421.	27321	<i>Stypopodium flabelliforme</i>			
422.	27345	<i>Turbinaria gracilis</i>			
423.	27346	<i>Turbinaria ornata</i>			
Fungi					
424.	27585	<i>Arthopyrenia analepta</i>			Y
425.	38761	<i>Auricularia cornea</i>			
426.	27662	<i>Chrysothrix candelaris</i>			
427.		<i>Daldinia concentrica</i>			
428.	27714	<i>Diploicia canescens</i>			
429.	27728	<i>Dirinaria applanata</i>			
430.	27730	<i>Dirinaria confluens</i>			
431.	27731	<i>Dirinaria picta</i>			
432.		<i>Hypoxylon rubiginosum</i>			
433.		<i>Lecanora</i> sp.			
434.	44002	<i>Lithothelium nanosporum</i>			
435.	27948	<i>Pertusaria leioplacella</i>			
436.	27956	<i>Pertusaria thiospoda</i>			
437.		<i>Phellinus gilvus</i>			
438.	28012	<i>Pyrenula nitida</i>			
439.	28225	<i>Ramalina subfraxinea</i> var. <i>norstictica</i>			
440.	28226	<i>Ramalina subfraxinea</i> var. <i>subfraxinea</i>			
441.	28055	<i>Roccella montagnei</i>			
442.	45882	<i>Sporisorium ryleyi</i>			
443.	45846	<i>Tilletia whiteochloae</i>			
444.	48858	<i>Trametes muelleri</i>			
445.	28082	<i>Trypethelium eluteriae</i>			Y
Plantae					
446.	19262	<i>Abildgaardia schoenoides</i>			
447.	3678	<i>Abrus precatorius</i> (Crabs Eyes)			
448.	16979	<i>Abrus precatorius</i> subsp. <i>precatorius</i>			
449.	11325	<i>Abutilon indicum</i> var. <i>australiense</i>			
450.	4901	<i>Abutilon otocarpum</i> (Desert Chinese Lantern)			
451.	40300	<i>Acacia calligera</i>			
452.	17013	<i>Acacia colei</i> var. <i>colei</i>			
453.	17014	<i>Acacia colei</i> var. <i>ileocarpa</i>			
454.		<i>Acacia delibrata</i>			
455.	12085	<i>Acacia deltoidea</i> subsp. <i>deltoidea</i>			
456.	3371	<i>Acacia hippuroides</i>			
457.	3447	<i>Acacia monticola</i> (Gawar, Lilwardi)			
458.	13401	<i>Acacia neurocarpa</i>			
459.	3483	<i>Acacia pellita</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
460.	3491 <i>Acacia platycarpa</i> (Pindan Wattle)			
461.	29210 <i>Acacia</i> sp. Kununurra (Lullfitz 6195)			
462.	3579 <i>Acacia trachycarpa</i> (Minni Ritchi, Balgali)			
463.	3580 <i>Acacia translucens</i> (Poverty Bush, Banmung)			
464.	19641 <i>Acacia tumida</i> var. <i>tumida</i>			
465.	3601 <i>Acacia wickhamii</i>			
466.	16155 <i>Acacia wickhamii</i> subsp. <i>wickhamii</i>			
467.	26440 <i>Acanthopora dendroides</i>			
468.	26441 <i>Acanthopora spicifera</i>			
469.	2645 <i>Achyranthes aspera</i> (Chaff Flower)			
470.	44 <i>Acrostichum speciosum</i>			
471.	26448 <i>Actinotrichia fragilis</i>			
472.	5224 <i>Adenia heterophylla</i>			
473.	11305 <i>Adenia heterophylla</i> subsp. <i>australis</i>			
474.	6486 <i>Aegialitis annulata</i> (Club Mangrove)			
475.	6478 <i>Aegiceras corniculatum</i> (River Mangrove)			
476.	2646 <i>Aerva javanica</i> (Kapok Bush)	Y		
477.	3680 <i>Aeschynomene indica</i> (Budda Pea)			
478.	2653 <i>Alternanthera pungens</i> (Khaki Weed)	Y		
479.	17574 <i>Alysicarpus ovalifolius</i>	Y		
480.	17573 <i>Alysicarpus suffruticosus</i>		P2	
481.	20018 <i>Amaranthus undulatus</i>			
482.	5277 <i>Ammannia baccifera</i>			
483.	43680 <i>Ammannia muelleri</i>			
484.	5278 <i>Ammannia multiflora</i>			
485.	26459 <i>Amphiroa crassa</i>			
486.	26461 <i>Amphiroa foliacea</i>			
487.	26462 <i>Amphiroa fragilissima</i>			
488.	26463 <i>Amphiroa gracilis</i>			
489.	43480 <i>Amphiroa tribulus</i>			
490.	2369 <i>Amyema benthamii</i>			
491.	13700 <i>Amyema bifurcata</i>			
492.	14109 <i>Amyema mackayensis</i>			
493.	11874 <i>Amyema sanguinea</i> var. <i>sanguinea</i>			
494.	2386 <i>Amyema thalassia</i>			
495.	11392 <i>Amyema villiflora</i> subsp. <i>villiflora</i>			
496.	35872 <i>Anadyomene plicata</i>			
497.	210 <i>Aristida holathera</i>			
498.	12063 <i>Aristida holathera</i> var. <i>holathera</i>			
499.	211 <i>Aristida hygrometrica</i> (Northern Kerosene Grass)			
500.	26486 <i>Asparagopsis taxiformis</i>			
501.	8781 <i>Asparagus racemosus</i>			
502.	4742 <i>Atalaya variifolia</i> (Wingleaf Whitewood)			
503.	6828 <i>Avicennia marina</i> (White Mangrove)			
504.	48138 <i>Avrainvillea carteri</i>			
505.	36362 <i>Avrainvillea erecta</i>			
506.	26498 <i>Avrainvillea obscura</i>			
507.	7045 <i>Bacopa floribunda</i>			
508.	1743 <i>Batis argillicola</i>			
509.	12757 <i>Bauhinia cunninghamii</i>			
510.	5183 <i>Bergia ammannioides</i>			
511.	5184 <i>Bergia pedicellaris</i>			
512.	7860 <i>Blumea integrifolia</i>			
513.	7865 <i>Blumea saxatilis</i>			
514.	26507 <i>Boerghesia forbesii</i>			
515.	2769 <i>Boerhavia burbridgeana</i>			
516.	2770 <i>Boerhavia coccinea</i> (Tar Vine, Wituka)			
517.	2771 <i>Boerhavia dominii</i>			
518.	2772 <i>Boerhavia gardneri</i>			
519.	6605 <i>Bonamia linearis</i>			
520.	26508 <i>Boodlea composita</i>			
521.	26509 <i>Bornetella oligospora</i>			
522.	26516 <i>Botryocladia leptopoda</i>			
523.	13010 <i>Brachychiton diversifolius</i> subsp. <i>diversifolius</i>			
524.	4602 <i>Breynia cernua</i>			
525.	4603 <i>Bridelia tomentosa</i>			
526.	5291 <i>Bruguiera exaristata</i> (Ribbed Mangrove)			
527.	44550 <i>Bryopsis pennata</i> var. <i>secunda</i>			Y
528.	29703 <i>Buchanania oblongifolia</i>			
529.	4718 <i>Buchanania obovata</i> (Wild Mango, Walangga)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
530.	13682 <i>Buchnera asperata</i>			
531.	7047 <i>Buchnera linearis</i> (Blackrod)			
532.	7050 <i>Buchnera urticifolia</i> (Blackrod)			
533.	750 <i>Bulbostylis barbata</i>			
534.	18073 <i>Byblis filifolia</i>			
535.	17854 <i>Byblis rorida</i>			
536.	3624 <i>Caesalpinia major</i>			
537.	10852 <i>Cajanus acutifolius</i>			
538.	11055 <i>Cajanus cinereus</i>			
539.	10972 <i>Cajanus marmoratus</i>			
540.	10863 <i>Cajanus reticulatus</i>			
541.	2871 <i>Calandrinia strophiolata</i>			
542.	2872 <i>Calandrinia tepperiana</i>			
543.	14925 <i>Calotropis gigantea</i>	Y		
544.	6582 <i>Calotropis procera</i> (Calotrope)	Y		
545.	5457 <i>Calytrix exstipulata</i> (Kimberley Heather)			
546.	4997 <i>Camptostemon schultzei</i> (Kapok Mangrove)			
547.	11617 <i>Canarium australianum</i> var. <i>glabrum</i>			
548.	3749 <i>Canavalia rosea</i> (Wild Jack Bean)			
549.	2976 <i>Capparis lasiantha</i> (Split Jack, Balqarda)			
550.	2980 <i>Capparis sepiaria</i>			
551.	2981 <i>Capparis spinosa</i>			
552.	48291 <i>Capparis spinosa</i> subsp. <i>nummularia</i>			
553.	6567 <i>Carissa lanceolata</i> (Conkerberry, Marnuwiji)			
554.	1161 <i>Cartonema parviflorum</i> (Wiridjagu)			
555.	2949 <i>Cassytha capillaris</i>			
556.	2950 <i>Cassytha filiformis</i> (Love Vine, Jirawan)			
557.	6569 <i>Catharanthus roseus</i> (Pink Periwinkle)	Y		
558.	42620 <i>Caulerpa chemnitzia</i>			
559.	26559 <i>Caulerpa cupressoides</i>			
560.	44539 <i>Caulerpa cylindracea</i>			
561.	44546 <i>Caulerpa dichotoma</i>			Y
562.	44547 <i>Caulerpa lamourouxii</i>			
563.	26568 <i>Caulerpa lentillifera</i>			
564.	44551 <i>Caulerpa macrodisca</i>			
565.	26573 <i>Caulerpa racemosa</i>			
566.	35122 <i>Caulerpa racemosa</i> var. <i>racemosa</i>			
567.	26576 <i>Caulerpa serrulata</i>			
568.	26577 <i>Caulerpa sertularioides</i>			
569.	26579 <i>Caulerpa taxifolia</i>			
570.	35125 <i>Caulerpa taxifolia</i> var. <i>falcifolia</i>			
571.	35124 <i>Caulerpa taxifolia</i> var. <i>taxifolia</i>			
572.	26582 <i>Caulerpa verticillata</i>			
573.	48913 <i>Celtis strychnoides</i>			
574.	257 <i>Cenchrus biflorus</i> (Gallon's Curse)	Y		
575.	258 <i>Cenchrus ciliaris</i> (Buffel Grass)	Y		
576.	259 <i>Cenchrus echinatus</i> (Burrgrass)	Y		
577.	29721 <i>Cenchrus setiger</i> (Birdwood Grass)	Y		
578.	26606 <i>Ceratodictyon spongiosum</i>			
579.	39680 <i>Ceriops australis</i>			
580.	13713 <i>Chamaecrista absus</i>			
581.	13756 <i>Chamaecrista mimosoides</i>			
582.	14811 <i>Chamaecrista moorei</i>			
583.	20064 <i>Chamaecrista rotundifolia</i>	Y		
584.	26618 <i>Champia parvula</i>			
585.	266 <i>Chloris barbata</i> (Purpletop Chloris)	Y		
586.	26628 <i>Chondria armata</i>			
587.	273 <i>Chrysopogon fallax</i> (Golden Beard Grass)			
588.	275 <i>Chrysopogon pallidus</i> (Ribbongrass)			
589.	48838 <i>Citrullus amarus</i>	Y		
590.	36316 <i>Cladophora herpestica</i>			
591.	44726 <i>Cladophoropsis vaucheriiformis</i>			
592.	2983 <i>Cleome cleomoides</i> (Justago)			
593.	31553 <i>Cleome</i> sp. <i>Bonaparte Archipelago</i> (A.A. Mitchell 4774)			
594.	2986 <i>Cleome tetrandra</i>			
595.	11886 <i>Cleome tetrandra</i> var. <i>tetrandra</i>			
596.	2988 <i>Cleome viscosa</i> (Tickweed, Tjinduwadhu)			
597.	6729 <i>Clerodendrum floribundum</i> (Lollybush)			
598.	13693 <i>Clerodendrum floribundum</i> var. <i>coriaceum</i>			
599.	13691 <i>Clerodendrum floribundum</i> var. <i>ovatum</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
600.	3769 <i>Clitoria ternatea</i>	Y		
601.	5214 <i>Cochlospermum fraseri</i> (Kapok Bush, Malindjarr)			
602.	35917 <i>Codium arabicum</i>			
603.	35857 <i>Codium dwarkense</i>			
604.	26673 <i>Codium geppiorum</i>			
605.	26683 <i>Codium spongiosum</i>			
606.	2778 <i>Codonocarpus cotinifolius</i> (Native Poplar, Kundurangu)			
607.	26691 <i>Coelothrix irregularis</i>			
608.	32999 <i>Colocasia esculenta</i> var. <i>esculenta</i>	Y		
609.	1165 <i>Commelina ensifolia</i> (Wandering Jew, Buargu)			
610.	12767 <i>Corchorus aestuans</i>			
611.	4863 <i>Corchorus pumilio</i>			
612.	4864 <i>Corchorus sidoides</i> (Flannel Weed)			
613.	18415 <i>Corchorus sidoides</i> subsp. <i>sidoides</i>			
614.	16788 <i>Corymbia bella</i>			
615.	14650 <i>Corymbia flavescens</i>			
616.	17089 <i>Corymbia greeniana</i>			
617.	17100 <i>Corymbia polycarpa</i>			
618.	48175 <i>Crinum arenarium</i>			
619.	13466 <i>Crotalaria brevis</i>			
620.	3774 <i>Crotalaria cunninghamii</i> (Green Birdflower, Bilbun)			
621.	20176 <i>Crotalaria cunninghamii</i> subsp. <i>cunninghamii</i>			
622.	20179 <i>Crotalaria medicaginea</i> var. <i>neglecta</i>			
623.	19398 <i>Crotalaria ramosissima</i>			
624.	<i>Crotalaria</i> sp.			
625.	13711 <i>Croton habrophyllus</i>			
626.	12683 <i>Cryptostegia madagascariensis</i>	Y		
627.	17433 <i>Cullen badocanum</i>			
628.	17435 <i>Cullen candidum</i>		P1	
629.	26712 <i>Curdiea obesa</i>			
630.	31213 <i>Cuscuta chinensis</i>			
631.	1166 <i>Cyanotis axillaris</i>			
632.	45973 <i>Cyanthillium cinereum</i> var. <i>lanatum</i>	Y		
633.	54 <i>Cyclosorus interruptus</i>			
634.	1628 <i>Cymbidium canaliculatum</i>			
635.	279 <i>Cymbopogon ambiguus</i> (Scentgrass)			
636.	282 <i>Cymbopogon procerus</i> (Lemon Grass)			
637.	129 <i>Cymodocea serrulata</i>			
638.	48278 <i>Cynanchum brevipedicellatum</i>			
639.	6585 <i>Cynanchum pedunculatum</i>			
640.	48280 <i>Cynanchum viminalis</i> subsp. <i>australe</i>			
641.	48281 <i>Cynanchum viminalis</i> subsp. <i>brunonianum</i>			
642.	283 <i>Cynodon dactylon</i> (Couch)	Y		
643.	12801 <i>Cyperus blakeanus</i>			
644.	777 <i>Cyperus bulbosus</i> (Bush Onion, Tjanmata)			
645.	778 <i>Cyperus carinatus</i>			
646.	784 <i>Cyperus conicus</i>			
647.	12810 <i>Cyperus cunninghamii</i> subsp. <i>uniflorus</i>			
648.	11024 <i>Cyperus latzii</i>			
649.	803 <i>Cyperus microcephalus</i>			
650.	12807 <i>Cyperus microcephalus</i> subsp. <i>microcephalus</i>			
651.	804 <i>Cyperus nervulosus</i>			
652.	806 <i>Cyperus polystachyos</i> (Bunchy Sedge)			
653.	813 <i>Cyperus sexflorus</i>			
654.	290 <i>Dactyloctenium radulans</i> (Button Grass)			
655.	14174 <i>Decaisnina angustata</i>			
656.	11407 <i>Dendrophthoe acacioides</i> subsp. <i>acacioides</i>			
657.	3853 <i>Desmodium filiforme</i>			
658.	3858 <i>Desmodium trichostachyum</i>			
659.	18230 <i>Desmodium triflorum</i>	Y		
660.	29616 <i>Dichotomaria marginata</i>			
661.	7166 <i>Dicliptera armata</i>			
662.	26769 <i>Dictyosphaeria cavernosa</i>			
663.	26782 <i>Digenea simplex</i>			
664.	309 <i>Digitaria bicornis</i> (Finger Grass)			
665.	311 <i>Digitaria ciliaris</i> (Summer Grass)	Y		
666.	1508 <i>Dioscorea bulbifera</i> (Ganmanggu, Gunu)			
667.	1510 <i>Dioscorea transversa</i>			
668.	17902 <i>Diospyros humilis</i>			
669.	19085 <i>Diospyros rugosula</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
670.	48738 <i>Distimake dissectus</i> var. <i>dissectus</i>	Y		
671.	38461 <i>Dodonaea hispidula</i> var. <i>arida</i>			
672.	38462 <i>Dodonaea hispidula</i> var. <i>phylloptera</i>			
673.	4776 <i>Dodonaea platyptera</i>			
674.	4782 <i>Dodonaea viscosa</i> (Sticky Hopbush)			
675.	17213 <i>Drosera broomensis</i>			
676.	17215 <i>Drosera derbyensis</i>			
677.	43588 <i>Drosera serpens</i>			
678.	19079 <i>Ectrosia schultzii</i> var. <i>annua</i>			Y
679.	11453 <i>Ectrosia schultzii</i> var. <i>schultzii</i>			
680.	6682 <i>Ehretia saligna</i> (False Cedar)			
681.	14301 <i>Ehretia saligna</i> var. <i>saligna</i>			
682.	36142 <i>Endosiphonia spinuligera</i>			
683.	160 <i>Enhalus acoroides</i>			
684.	363 <i>Enneapogon pallidus</i> (Conetop Nineawn)			
685.	367 <i>Enteropogon dolichostachyus</i>			
686.	375 <i>Eragrostis cumingii</i> (Cuming's Love Grass)			
687.	381 <i>Eragrostis falcata</i> (Sickle Lovegrass)			
688.	393 <i>Eragrostis setifolia</i> (Neverfail Grass)			
689.	395 <i>Eragrostis speciosa</i> (Handsome Lovegrass)			
690.	402 <i>Eriachne avenacea</i>			
691.	404 <i>Eriachne ciliata</i> (Slender Wandarrie Grass)			
692.	410 <i>Eriachne glauca</i> (Pan Wandarrie Grass)			
693.	12055 <i>Eriachne glauca</i> var. <i>glauca</i>			
694.	412 <i>Eriachne melicacea</i>			
695.	414 <i>Eriachne obtusa</i> (Northern Wandarrie Grass)			
696.	48185 <i>Eriachne pindanica</i> (Pindan Wiregrass)			
697.	5564 <i>Eucalyptus brachyandra</i> (Tropical Red Box)			
698.	5715 <i>Eucalyptus miniata</i> (Woollybutt, Manawan)			
699.	5785 <i>Eucalyptus tectifica</i> (Darwin Box)			
700.	38182 <i>Eucheuma arnoldii</i>			
701.	26827 <i>Eucheuma denticulatum</i>			
702.	42841 <i>Euphorbia armstrongiana</i> var. <i>distans</i>			
703.	42849 <i>Euphorbia hassallii</i>			
704.	4629 <i>Euphorbia hirta</i> (Asthma Plant)	Y		
705.	42863 <i>Euphorbia mitchelliana</i> var. <i>mitchelliana</i>			
706.	4635 <i>Euphorbia myrtoides</i>			
707.	42879 <i>Euphorbia trigonosperma</i>			
708.	11416 <i>Evolvulus alsinoides</i> var. <i>decumbens</i>			
709.	15126 <i>Excoecaria ovalis</i>			
710.	11169 <i>Exocarpos latifolius</i> (Broad-leaved Cherry)			
711.	26831 <i>Exophyllum wentii</i>			
712.	31578 <i>Ficus aculeata</i> var. <i>indecora</i> (Ranji)			
713.	19643 <i>Ficus atricha</i>			
714.	19648 <i>Ficus brachypoda</i>			
715.	43508 <i>Ficus geniculata</i> var. <i>insignis</i>			
716.	1753 <i>Ficus platypoda</i> (Native Fig, Makartu)			
717.	1759 <i>Ficus virens</i> (Albayi)			
718.	12096 <i>Ficus virens</i> var. <i>virens</i>			
719.	49126 <i>Fimbristylis crosslandii</i>			
720.	847 <i>Fimbristylis cymosa</i>			
721.	850 <i>Fimbristylis depauperata</i>			
722.	855 <i>Fimbristylis ferruginea</i>			
723.	881 <i>Fimbristylis sericea</i>			
724.	890 <i>Fimbristylis tetragona</i>			
725.	1055 <i>Flagellaria indica</i> (Gadji)			
726.	12013 <i>Flueggea virosa</i> subsp. <i>melanthesoides</i> (Dogwood, Guwal)			
727.	896 <i>Fuirena ciliaris</i>			
728.	3886 <i>Galactia tenuiflora</i>			
729.	26835 <i>Galaxaura rugosa</i>			
730.	26837 <i>Ganonema farinosum</i>			
731.	26839 <i>Ganonema pinnatum</i>			
732.	15234 <i>Gardenia pyriformis</i> subsp. <i>keartlandii</i>			
733.	17560 <i>Gardenia pyriformis</i> subsp. <i>pyriformis</i>			
734.	17561 <i>Gardenia resinosa</i> subsp. <i>resinosa</i>			
735.	26842 <i>Gelidiella acerosa</i>			
736.	2836 <i>Glinus oppositifolius</i>			
737.	3942 <i>Glycine tomentella</i> (Woolly Glycine)			
738.	4485 <i>Glycosmis trifoliata</i>			
739.	18229 <i>Gomphrena brachystylis</i> subsp. <i>pindanensis</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
740.	2676 <i>Gomphrena canescens</i> (Batchelors Buttons)			
741.	2677 <i>Gomphrena celosioides</i> (Gomphrena Weed)	Y		
742.	18226 <i>Gomphrena connata</i>			
743.	2681 <i>Gomphrena diffusa</i>			
744.	18364 <i>Gomphrena diffusa</i> subsp. <i>arenicola</i>			Y
745.	2682 <i>Gomphrena flaccida</i> (Gomphrena Weed)			
746.	2687 <i>Gomphrena tenella</i>			
747.	6158 <i>Gonocarpus leptothecus</i>			
748.	7500 <i>Goodenia coronopifolia</i>			
749.	13163 <i>Goodenia sepalosa</i> var. <i>sepalosa</i>			
750.	4916 <i>Gossypium populifolium</i>			
751.	13043 <i>Gossypium rotundifolium</i>			
752.	35899 <i>Gracilaria canaliculata</i>			
753.	26873 <i>Gracilaria salicornia</i>			
754.	26874 <i>Gracilaria textorii</i>			
755.	2016 <i>Grevillea heliosperma</i> (Rock Grevillea)			
756.	2079 <i>Grevillea pyramidalis</i> (Caustic Bush, Tjungu)			
757.	15975 <i>Grevillea pyramidalis</i> subsp. <i>pyramidalis</i>			
758.	16476 <i>Grevillea refracta</i> subsp. <i>refracta</i>			
759.	13441 <i>Grevillea wickhamii</i> subsp. <i>wickhamii</i>			
760.	4868 <i>Grewia breviflora</i>			
761.	13565 <i>Grewia glabra</i>			
762.	4872 <i>Grewia retusifolia</i> (Dog's Balls)			
763.	13228 <i>Gymnanthera oblonga</i>			
764.	2960 <i>Gyrocarpus americanus</i> (Helicopter Tree, Bilangkamar)			
765.	13748 <i>Gyrocarpus americanus</i> subsp. <i>pachyphyllus</i>			
766.	2789 <i>Gyrostemon tepperi</i>			
767.	45678 <i>Haemodorum capitatum</i>		P1	
768.	2129 <i>Hakea arborescens</i> (Common Hakea)			
769.	2178 <i>Hakea macrocarpa</i> (Dyaridany, Jaradinty)			
770.	26891 <i>Halimeda cylindracea</i>			
771.	26892 <i>Halimeda discoidea</i>			
772.	26894 <i>Halimeda macroloba</i>			
773.	35906 <i>Halimeda opuntia</i>			
774.	26896 <i>Halimeda simulans</i>			
775.	26898 <i>Halimeda velasquezii</i>			
776.	47273 <i>Halimeda xishaensis</i>			
777.	131 <i>Halodule uninervis</i>			
778.	162 <i>Halophila decipiens</i>			
779.	164 <i>Halophila ovalis</i> (Sea Wrack)			
780.	38081 <i>Halymenia dilatata</i>			
781.	37642 <i>Halymenia durvillei</i>			
782.	38100 <i>Halymenia maculata</i>			
783.	5020 <i>Helicteres rhynchocarpa</i>			
784.	6708 <i>Heliotropium diversifolium</i>			
785.	10882 <i>Heliotropium foliatum</i>			
786.	10992 <i>Heliotropium glabellum</i>			
787.	13126 <i>Heliotropium leptaleum</i>			
788.	29960 <i>Heliotropium microsalsoides</i>			
789.	6713 <i>Heliotropium ovalifolium</i>			
790.	6714 <i>Heliotropium paniculatum</i>			
791.	17312 <i>Heliotropium ramullpatens</i>			
792.	26915 <i>Hennedya crispa</i>			
793.	4920 <i>Herissantia crispa</i>			
794.	443 <i>Heteropogon contortus</i> (Bunch Speargrass)			
795.	26930 <i>Heterosiphonia crassipes</i>			
796.	29358 <i>Hibiscus apodus</i>			
797.	4929 <i>Hibiscus geranioides</i>			
798.	4933 <i>Hibiscus leptocladus</i>			
799.	4934 <i>Hibiscus meraukensis</i> (Merauke Hibiscus)			
800.	4945 <i>Hibiscus vitifolius</i>			
801.	5215 <i>Hybanthus aurantiacus</i>			
802.	5219 <i>Hybanthus enneaspermus</i>			
803.	11346 <i>Hybanthus enneaspermus</i> subsp. <i>enneaspermus</i>			
804.	26956 <i>Hydrolithon reinboldii</i>			
805.	35871 <i>Hydroponia urvillei</i>			
806.	26970 <i>Hypnea pannosa</i>			
807.	26972 <i>Hypnea spinella</i>			
808.	26978 <i>Hypoglossum harveyanum</i>			Y
809.	3973 <i>Indigofera colutea</i> (Sticky Indigo)			

Name ID	Species Name	Naturalised	Conservation Code	Endemic To Query Area
810.	3976 <i>Indigofera haplophylla</i>			
811.	3978 <i>Indigofera hirsuta</i> (Hairy Indigo)			
812.	3980 <i>Indigofera linifolia</i>			
813.	3981 <i>Indigofera linnaei</i> (Birdsville Indigo)			
814.	6623 <i>Ipomoea optica</i>			
815.	6632 <i>Ipomoea macrantha</i>			
816.	6633 <i>Ipomoea muelleri</i> (Poison Morning Glory, Yumbu)			
817.	11312 <i>Ipomoea pes-caprae</i> subsp. <i>brasiliensis</i>			
818.	6637 <i>Ipomoea polymorpha</i>			
819.	13761 <i>Ischaemum australe</i> var. <i>arundinaceum</i>			
820.	6643 <i>Jacquemontia paniculata</i>			
821.	26983 <i>Jania adhaerens</i>			
822.	26985 <i>Jania micrarthrodia</i>			
823.	11315 <i>Jasminum didymum</i> subsp. <i>didymum</i>			
824.	12059 <i>Jasminum didymum</i> subsp. <i>lineare</i> (Desert Jasmine)			
825.	6502 <i>Jasminum molle</i>			
826.	4656 <i>Jatropha gossypifolia</i> (Bellyache Bush)	Y		
827.	7118 <i>Josephinia eugeniae</i> (Josephinia Burr)			
828.	6733 <i>Lantana camara</i> (Common Lantana)	Y		
829.	26998 <i>Laurencia brongniartii</i>			
830.	48408 <i>Laurencia dendroidea</i>			
831.	27004 <i>Laurencia intricata</i>			
832.	<i>Laurencia papillosa</i>			
833.	3613 <i>Leucaena leucocephala</i> (Leucaena)	Y		
834.	27021 <i>Liagora ceranoides</i>			
835.	19073 <i>Lithomyrtus retusa</i>			
836.	27037 <i>Lithophyllum kotschyannum</i>			
837.	11425 <i>Lophostemon grandiflorus</i> subsp. <i>grandiflorus</i>		P3	
838.	6136 <i>Ludwigia perennis</i>			
839.	5296 <i>Lumnitzera racemosa</i> (White-flowered Black Mangrove)			
840.	18376 <i>Luvunga monophylla</i>			
841.	11809 <i>Lysiana spathulata</i> subsp. <i>spathulata</i>			
842.	4070 <i>Macroptilium atropurpureum</i> (Purple Bean)	Y		
843.	4658 <i>Mallotus nesophilus</i>			
844.	4719 <i>Mangifera indica</i> (Mango)	Y		
845.	16540 <i>Marsdenia geminata</i>			
846.	27057 <i>Martensia fragilis</i>			Y
847.	35902 <i>Mastophora rosea</i>			
848.	20638 <i>Megathyrsus maximus</i>	Y		
849.	9178 <i>Melaleuca alsophila</i>			
850.	5875 <i>Melaleuca argentea</i> (Silver Cadjeput, Bandaran)			
851.	17791 <i>Melaleuca cajuputi</i> subsp. <i>cajuputi</i>			
852.	5901 <i>Melaleuca dealbata</i> (Karnbor)			
853.	5942 <i>Melaleuca nervosa</i> (Fibrebark)			
854.	5989 <i>Melaleuca viridiflora</i> (Broadleaf Paperbark)			
855.	5051 <i>Melhania oblongifolia</i>			
856.	4516 <i>Melia azedarach</i> (White Cedar)			
857.	5052 <i>Melochia corchorifolia</i>			
858.	38200 <i>Merremia incisa</i>			
859.	48283 <i>Mesosphaerum suaveolens</i>	Y		
860.	27074 <i>Microdictyon umbilicatum</i>			
861.	31374 <i>Microstachys chamaelea</i>			
862.	6492 <i>Mimusops elengi</i> (Walara)			
863.	6525 <i>Mitrasacme hispida</i>			
864.	6531 <i>Mitrasacme nummularia</i>			
865.	15074 <i>Mitrasacme scritchicola</i>			
866.	7335 <i>Morinda citrifolia</i>			
867.	1167 <i>Murdannia graminea</i> (Baniyu)			
868.	20774 <i>Musa acuminata</i>	Y		
869.	17158 <i>Myoporum montanum</i> (Native Myrtle)			
870.	2573 <i>Neobassia astrocarpa</i>			
871.	44525 <i>Neoziella divaricata</i>			
872.	27099 <i>Neomeris van-bosseae</i>			
873.	13340 <i>Oldenlandia corymbosa</i> var. <i>corymbosa</i>	Y		
874.	7339 <i>Oldenlandia galioides</i>			
875.	7340 <i>Oldenlandia mitrasacmoides</i>			
876.	6651 <i>Operculina aequisepala</i>			
877.	6652 <i>Operculina brownii</i> (Potato Vine, Bara)			
878.	6005 <i>Osbornia octodonta</i> (Myrtle Mangrove)			
879.	4518 <i>Owenia reticulata</i> (Native Walnut, Bandal)			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
880.	4519 <i>Owenia vernicosa</i> (Emu Apple)			
881.	36400 <i>Palisada perforata</i>			
882.	104 <i>Pandanus spiralis</i> (Screwpine, Wakirri)			
883.	11737 <i>Pandanus spiralis</i> var. <i>convexus</i>			
884.	503 <i>Panicum decompositum</i> (Native Millet, Kaltu-kaltu)			
885.	505 <i>Panicum laevinode</i>			
886.	509 <i>Panicum schinzii</i>	Y		
887.	3193 <i>Parinari nonda</i>			
888.	17362 <i>Parsonia kimberleyensis</i>		P1	
889.	523 <i>Paspalidium rarum</i> (Rare Paspalidium)			
890.	5226 <i>Passiflora foetida</i> (Stinking Passion Flower)	Y		
891.	14096 <i>Passiflora foetida</i> var. <i>hispida</i>	Y		
892.	13570 <i>Pavetta kimberleyana</i>			
893.	5284 <i>Pemphis acidula</i>			
894.	546 <i>Perotis rara</i> (Comet Grass)			
895.	2263 <i>Persoonia falcata</i> (Wild Pear, Gandala)			
896.	6734 <i>Phyla nodiflora</i> var. <i>nodiflora</i>	Y		
897.	9056 <i>Phyllanthus baccatus</i>			
898.	14462 <i>Phyllanthus exilis</i>			
899.	4680 <i>Phyllanthus maderaspatensis</i>			
900.	20652 <i>Physalis angulata</i>	Y		
901.	5290 <i>Planchonia careya</i> (Mangaloo, Yundu)			
902.	47 <i>Platyzoma microphyllum</i> (Braid Fern)			
903.	43944 <i>Pluchea longiseta</i>			
904.	8168 <i>Pluchea rubelliflora</i>			
905.	6491 <i>Plumbago zeylanica</i> (Native Plumbago)			
906.	2902 <i>Polycarpaea involucreta</i>			
907.	2903 <i>Polycarpaea longiflora</i>			
908.	4577 <i>Polygala tepperi</i>			
909.	6653 <i>Polymeria ambigua</i> (Morning Glory)			
910.	27171 <i>Polysiphonia blandii</i>			
911.	27180 <i>Polysiphonia subtilissima</i>			
912.	4691 <i>Poranthera microphylla</i> (Small Poranthera)			
913.	27186 <i>Portieria hornemannii</i>			
914.	2876 <i>Portulaca bicolor</i>			
915.	2881 <i>Portulaca filifolia</i>			
916.	2883 <i>Portulaca napiformis</i>			
917.	2884 <i>Portulaca oleracea</i> (Purslane, Wakati)			
918.	2886 <i>Portulaca pilosa</i> (Djanggara)	Y		
919.	6735 <i>Premna acuminata</i> (Ngalinginkal)			
920.	<i>Premna</i> sp.			
921.	18208 <i>Psydrax pendulina</i>			
922.	41223 <i>Pterocaulon paradoxum</i>			
923.	<i>Pterocaulon</i> sp.			
924.	8192 <i>Pterocaulon sphacelatum</i> (Apple Bush, Fruit Salad Plant)			
925.	41220 <i>Pterocaulon tricholobum</i>			
926.	2713 <i>Ptilotus corymbosus</i>			
927.	2721 <i>Ptilotus exaltatus</i> (Tall Mulla Mulla)			
928.	2725 <i>Ptilotus fusiformis</i>			
929.	37980 <i>Ptilotus giganteus</i>			
930.	2737 <i>Ptilotus lanatus</i>			
931.	2751 <i>Ptilotus polystachyus</i> (Prince of Wales Feather)			
932.	19509 <i>Pupalia micrantha</i>	Y		
933.	5295 <i>Rhizophora stylosa</i> (Spotted-leaved Red Mangrove)			
934.	4190 <i>Rhynchosia australis</i> (Rhynchosia)			
935.	<i>Rhynchospora</i> sp.			
936.	7360 <i>Richardia scabra</i> (White Eye)	Y		
937.	17890 <i>Ruellia tuberosa</i>	Y		
938.	30434 <i>Salsola australis</i>			
939.	2357 <i>Santalum lanceolatum</i> (Northern Sandalwood, Yamguli)			
940.	27230 <i>Sarconema filiforme</i>			
941.	7623 <i>Scaevola macrostachya</i>			
942.	13153 <i>Scaevola taccada</i>			
943.	1027 <i>Scleria brownii</i>			
944.	12303 <i>Senna costata</i>			
945.	12310 <i>Senna goniodes</i>			
946.	12312 <i>Senna notabilis</i>			
947.	10848 <i>Senna occidentalis</i>	Y		
948.	12316 <i>Senna surattensis</i>			
949.	12317 <i>Senna surattensis</i> subsp. <i>sulfurea</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
950.	12319 <i>Senna venusta</i>			
951.	31172 <i>Sersalisia sericea</i> (Nangi)			
952.	4196 <i>Sesbania cannabina</i> (Sesbania Pea)			
953.	4198 <i>Sesbania formosa</i> (White Dragon Tree)			
954.	2818 <i>Sesuvium portulacastrum</i>			
955.	605 <i>Setaria apiculata</i> (Pigeon Grass)			
956.	32984 <i>Sida acuta</i> subsp. <i>acuta</i>	Y		
957.	4979 <i>Sida hackettiana</i>			
958.	4981 <i>Sida intricata</i> (Tangled Sida)			
959.	18150 <i>Sida rohlenae</i> subsp. <i>occidentalis</i>			
960.	16993 <i>Sida</i> sp. Rabbit Flat (B.J. Carter 626)			
961.	7000 <i>Solanum cunninghamii</i>			
962.	7001 <i>Solanum dioicum</i> (Gilu)			
963.	7027 <i>Solanum petraeum</i>			
964.	27281 <i>Solieria robusta</i>			
965.	5288 <i>Sonneratia alba</i> (Pornupan)			
966.	12921 <i>Sorghum ecarinatum</i>			
967.	12920 <i>Sorghum interjectum</i>			
968.	619 <i>Sorghum plumosum</i> (Plume Canegrass)			
969.	620 <i>Sorghum stipoideum</i> (Annual Sorghum)			
970.	28347 <i>Spermacoce occidentalis</i>			
971.	<i>Spermacoce</i> sp.			
972.	43943 <i>Sphaeromorphaea littoralis</i>			
973.	38441 <i>Sphagnetocola trilobata</i>	Y		
974.	625 <i>Spinifex longifolius</i> (Beach Spinifex)			
975.	44523 <i>Spongophloeia tissotii</i>			
976.	635 <i>Sporobolus virginicus</i> (Marine Couch)			
977.	27310 <i>Spyridia filamentosa</i>			
978.	7101 <i>Stemodia lythriifolia</i> (Bunu Bunu)			
979.	7103 <i>Striga curviflora</i>			
980.	45717 <i>Stylidium pindanicum</i> (Pindan Triggerplant)		P3	
981.	12353 <i>Stylosanthes hamata</i> (Verano Stylo)	Y		
982.	4214 <i>Stylosanthes humilis</i> (Townsville Stylo)	Y		
983.	12354 <i>Stylosanthes scabra</i>	Y		
984.	14491 <i>Stylosanthes viscosa</i>	Y		
985.	2638 <i>Suaeda arbusculoides</i>			
986.	132 <i>Syringodium isoetifolium</i>			
987.	12113 <i>Syzygium eucalyptoides</i> subsp. <i>bleeseri</i>			
988.	6047 <i>Syzygium suborbiculare</i>			
989.	33236 <i>Tecticornia halocnemoides</i> (Shrubby Samphire)			
990.	33238 <i>Tecticornia halocnemoides</i> subsp. <i>tenuis</i>			
991.	33356 <i>Tecticornia indica</i> subsp. <i>indica</i>			
992.	4253 <i>Templetonia hookeri</i>			
993.	4266 <i>Tephrosia crocea</i> (Baynjoed)			
994.	4269 <i>Tephrosia flammea</i>			
995.	13922 <i>Tephrosia laxa</i> var. <i>angustata</i>			
996.	4272 <i>Tephrosia leptoclada</i>			
997.	4274 <i>Tephrosia oblongata</i>			
998.	4280 <i>Tephrosia rosea</i> (Flinders River Poison, Bungoo'dah)			
999.	19531 <i>Tephrosia rosea</i> var. <i>clementii</i>			
1000.	19529 <i>Tephrosia rosea</i> var. <i>rosea</i>			
1001.	4281 <i>Tephrosia simplicifolia</i>			
1002.	34716 <i>Tephrosia</i> sp. Pentecost River (I.D. Cowie 4168)			
1003.	20777 <i>Tephrosia valliculata</i>		P3	
1004.	4287 <i>Tephrosia virens</i>			
1005.	5300 <i>Terminalia canescens</i> (Joolal)			
1006.	5303 <i>Terminalia ferdinandiana</i> (Mador)			
1007.	5306 <i>Terminalia hadleyana</i>			
1008.	5307 <i>Terminalia latipes</i>			
1009.	5309 <i>Terminalia petiolaris</i> (Masroorl)			
1010.	169 <i>Thalassia hemprichii</i>			
1011.	133 <i>Thalassodendron ciliatum</i>			
1012.	4992 <i>Thespesia populneoides</i> (Laba)			
1013.	1326 <i>Thysanotus chinensis</i>			
1014.	2942 <i>Tinospora smilacina</i> (Snakevine, Oondala)			
1015.	27334 <i>Titanophora weberae</i>			
1016.	27335 <i>Tolypocladia calodictyon</i>			
1017.	27336 <i>Tolypocladia glomerulata</i>			
1018.	6270 <i>Trachymene didiscoides</i>			
1019.	1745 <i>Trema tomentosa</i>			

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
1020.	44305 <i>Trianthena pilosum</i>			
1021.	2830 <i>Trianthena portulacastrum</i> (Giant Pigweed)	Y		
1022.	4368 <i>Tribulopsis angustifolia</i>			
1023.	19138 <i>Tribulopsis</i> sp. Koolan Island (K.F. Kenneally 8278)		P1	
1024.	4375 <i>Tribulus cistoides</i>			
1025.	4383 <i>Tribulus terrestris</i> (Caltrop)	Y		
1026.	6727 <i>Trichodesma zeylanicum</i> (Camel Bush, Kumbalin)			
1027.	27339 <i>Trichogloea requienii</i>			
1028.	27340 <i>Tricleocarpa cylindrica</i>			
1029.	8252 <i>Tridax procumbens</i> (Tridax, Tridax Daisy)	Y		
1030.	17888 <i>Triodia acutispicula</i>		P3	
1031.	17883 <i>Triodia bynoei</i>			
1032.	696 <i>Triodia pungens</i> (Soft Spinifex)			
1033.	17539 <i>Triumfetta breviaculeata</i>			
1034.	14961 <i>Triumfetta carteri</i>			
1035.	4880 <i>Triumfetta micracantha</i>			
1036.	4881 <i>Triumfetta plumigera</i>			
1037.	17534 <i>Triumfetta ryeae</i>			
1038.	13469 <i>Triumfetta triandra</i>			
1039.	27348 <i>Udotea argentea</i>			
1040.	27349 <i>Udotea flabellum</i>			
1041.	35302 <i>Udotea glaucescens</i>			
1042.	35261 <i>Ulva clathrata</i>			
1043.	715 <i>Urochloa mosambicensis</i> (Sabi Grass)	Y		
1044.	10865 <i>Urochloa subquadrifera</i>			
1045.	48823 <i>Utricularia bidentata</i>		P3	
1046.	7140 <i>Utricularia kimberleyensis</i> (Kimberley Bladderwort)			
1047.	29526 <i>Vachellia pachyphloia</i> subsp. <i>brevipinnula</i>			
1048.	29525 <i>Vachellia pachyphloia</i> subsp. <i>pachyphloia</i>			
1049.	36143 <i>Valonia fastigiata</i>			
1050.	46438 <i>Valonia ventricosa</i>			
1051.	27357 <i>Valoniopsis pachynema</i>			
1052.	35904 <i>Vanvoorstia spectabilis</i>			
1053.	7663 <i>Velleia panduriformis</i> (Cabbage Poison)			
1054.	11800 <i>Vigna lanceolata</i> var. <i>filiformis</i>			
1055.	4324 <i>Vigna radiata</i> (Mung Bean)			
1056.	18503 <i>Vigna radiata</i> var. <i>sublobata</i>			
1057.	14398 <i>Vigna vexillata</i>			
1058.	48985 <i>Vincetoxicum carnosum</i>			
1059.	48983 <i>Vincetoxicum cinerascens</i>			
1060.	48987 <i>Vincetoxicum flexuosum</i>			
1061.	6742 <i>Vitex glabrata</i> (Vitex, Yauru)			
1062.	11359 <i>Vitex trifolia</i> var. <i>subtrisecta</i>	Y		
1063.	12725 <i>Wahlenbergia caryophylloides</i>			
1064.	5106 <i>Waltheria indica</i>			
1065.	725 <i>Whiteochloa airoides</i>			
1066.	6578 <i>Wrightia saligna</i>			
1067.	6661 <i>Xenostegia tridentata</i>			
1068.	730 <i>Xerochloa imberbis</i> (Rice Grass)			
1069.	1142 <i>Xyris complanata</i>			
1070.	735 <i>Yakirra pauciflora</i>			
1071.	27370 <i>Yamadaella caenomyce</i>			
1072.	4327 <i>Zornia chaetophora</i>			
1073.	12680 <i>Zornia prostrata</i> var. <i>prostrata</i>			

Conservation Codes

T - Rare or likely to become extinct
X - Presumed extinct
IA - Protected under international agreement
S - Other specially protected fauna
1 - Priority 1
2 - Priority 2
3 - Priority 3
4 - Priority 4
5 - Priority 5

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.

NatureMap Species Report - Lombadina flora

Created By Guest user on 29/10/2020

Kingdom Plantae
Conservation Status Conservation Taxon (T, X, IA, S, P1-P5)
Current Names Only Yes
Core Datasets Only Yes
Species Group All Plants
Method 'By Circle'
Centre 122° 58' 32" E, 16° 28' 05" S
Buffer 40km
Group By Conservation Status

Conservation Status	Species	Records
Priority 1	4	6
Priority 2	1	2
Priority 3	5	15
TOTAL	10	23

Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
Priority 1				
1.	17435 <i>Cullen candidum</i>		P1	
2.	45678 <i>Haemodorum capitatum</i>		P1	
3.	17362 <i>Parsonsia kimberleyensis</i>		P1	
4.	19138 <i>Tribulopsis</i> sp. Koolan Island (K.F. Kenneally 8278)		P1	
Priority 2				
5.	17573 <i>Alysicarpus suffruticosus</i>		P2	
Priority 3				
6.	11425 <i>Lophostemon grandiflorus</i> subsp. <i>grandiflorus</i>		P3	
7.	45717 <i>Stylidium pindanicum</i> (Pindan Triggerplant)		P3	
8.	20777 <i>Tephrosia valleculata</i>		P3	
9.	17888 <i>Triodia acutispicula</i>		P3	
10.	48823 <i>Utricularia bidentata</i>		P3	

Conservation Codes

T - Rare or likely to become extinct
X - Presumed extinct
IA - Protected under international agreement
S - Other specially protected fauna
1 - Priority 1
2 - Priority 2
3 - Priority 3
4 - Priority 4
5 - Priority 5

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.

Appendix 3

Atlas of Living Australia Database Search Results



Reptiles

Family	Species	Common Name	Number
Agamidae	<i>Chelosania brunnea</i>	Chameleon Dragon	1
Agamidae	<i>Chlamydosaurus kingii</i>	Friilled Lizard	4
Agamidae	<i>Diporiphora pindan</i>	Pindan Two-line Dragon	18
Agamidae	<i>Lophognathus gilberti</i>	Gilbert's Dragon	11
Agamidae	<i>Pogona minor mitchelli</i>	Dwarf Bearded Dragon	7
Cheloniidae	<i>Chelonia mydas</i>	Green Turtle	37
Cheloniidae	<i>Lepidochelys olivacea</i>	Pacific Ridley	1
Colubridae	<i>Dendrelaphis punctulatus</i>	Common Tree Snake	2
Colubridae	<i>Fordonia leucobalia</i>	White-bellied Mangrove Snake	1
Diplodactylidae	<i>Amalosia rhombifer</i>	Zigzag Velvet Gecko	1
Diplodactylidae	<i>Diplodactylus</i>	Western Fat-tailed Gecko	1
Diplodactylidae	<i>Oedura gracilis</i>	Gracile Velvet Gecko	3
Diplodactylidae	<i>Strophurus ciliaris</i>	Northern Spiny-tailed Gecko	1
Diplodactylidae	<i>Strophurus ciliaris aberrans</i>	Northern Spiny-tailed Gecko	7
Elapidae	<i>Demansia angusticeps</i>		1
Elapidae	<i>Demansia papuensis</i>	Greater Black Whipsnake	1
Elapidae	<i>Furina ornata</i>	Orange-naped Snake	1
Elapidae	<i>Pseudechis australis</i>	King Brown Snake	3
Elapidae	<i>Pseudonaja mengdeni</i>	Western Brown Snake	4
Elapidae	<i>Simoselaps minimus</i>	Dampierland Burrowing Snake	3
Gekkonidae	<i>Gehyra australis</i>	Northern Dtella	9
Gekkonidae	<i>Gehyra nana</i>	Northern Spotted Rock Dtella	6
Gekkonidae	<i>Gehyra occidentalis</i>	Kimberley Plateau Dtella	17
Gekkonidae	<i>Gehyra pilbara</i>	Pilbara Dtella	18
Gekkonidae	<i>Gehyra punctata</i>	Spotted Dtella	8
Gekkonidae	<i>Gehyra variegata</i>	Tree Dtella	1
Gekkonidae	<i>Heteronotia binoei</i>	Bynoe's Gecko	43
Pygopodidae	<i>Delma borea</i>	Rusty-topped Delma	5
Pygopodidae	<i>Delma tincta</i>	Excitable Delma	1
Pygopodidae	<i>Lialis burtonis</i>	Burton's Snake-lizard	8
Pythonidae	<i>Antaresia childreni</i>	Children's Python	3
Pythonidae	<i>Antaresia stimsoni stimsoni</i>		1
Pythonidae	<i>Liasis olivaceus olivaceus</i>		1
Scincidae	<i>Carlia amax</i>	Bauxite Rainbow-skink	2
Scincidae	<i>Carlia munda</i>	Shaded-litter Rainbow-skink	3
Scincidae	<i>Carlia triacantha</i>	Desert Rainbow-skink	11
Scincidae	<i>Cryptoblepharus tythos</i>	Pygmy Snake-eyed Skink	7
Scincidae	<i>Ctenotus colletti</i>	Buff-tailed Finesnout Ctenotus	1
Scincidae	<i>Ctenotus helenae</i>	Clay-soil Ctenotus	4
Scincidae	<i>Ctenotus inornatus</i>	Bar-shouldered Ctenotus	31
Scincidae	<i>Ctenotus serventyi</i>	North-western Sandy-loam Ctenotus	1
Scincidae	<i>Eremiascincus isolepis</i>	Northern Bar-lipped Skink	49
Scincidae	<i>Lerista apoda</i>	Dampier Land Limbless Slider	7
Scincidae	<i>Lerista bipes</i>	North-western Sandslider	57
Scincidae	<i>Lerista greeri</i>	South-eastern Kimberley Sandslider	1

Family	Species	Common Name	Number
Scincidae	<i>Lerista griffini</i>	Stout Sandslider	33
Scincidae	<i>Lerista separanda</i>	Dampierland Plain Slider	1
Scincidae	<i>Menetia maini</i>	Northern Dwarf Skink	1
Scincidae	<i>Morethia ruficauda ruficauda</i>		1
Scincidae	<i>Morethia storri</i>	Top End Firetail Skink	3
Scincidae	<i>Notoscincus ornatus</i>	Ornate Soil-crevice Skink	1
Scincidae	<i>Notoscincus ornatus wotjulum</i>		9
Scincidae	<i>Tiliqua multifasciata</i>	Centralian Blue-tongue	1
Scincidae	<i>Tiliqua scincoides intermedia</i>	Northern Blue-tongued Skink	2
Typhlopidae	<i>Anilius diversus</i>	Northern Blind Snake	4
Varanidae	<i>Varanus glauerti</i>	Kimberley Rock Monitor	2
Varanidae	<i>Varanus glebopalma</i>	Black-palmed Monitor	1
Varanidae	<i>Varanus gouldii</i>	Gould's Goanna	1
Varanidae	<i>Varanus scalaris</i>	Spotted Tree Monitor	1
Varanidae	<i>Varanus tristis</i>	Black-headed Monitor	2

Mammals

Family	Species	Common Name	Number
Balaenopteridae	<i>Megaptera novaeangliae</i>	Humpback whale	23
Dasyuridae	<i>Dasyurus hallucatus</i>	Northern Quoll	1
Dasyuridae	<i>Planigale maculata</i>	Common Planigale	1
Dasyuridae	<i>Pseudantechinus ningbing</i>	Ningbing Pseudantechinus	2
Dasyuridae	<i>Sminthopsis macroura froggatti</i>	Froggatt's Stripe-faced Dunnart	3
Delphinidae	<i>Sousa sahalensis</i>	Australian humpbacked dolphin	1
Dugongidae	<i>Dugong dugon</i>	dugong	45
Emballonuridae	<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheathtail-bat	2
Emballonuridae	<i>Taphozous georgianus</i>	Common Sheathtail-bat	3
Macropodidae	<i>Onychogalea unguifera</i>	Northern Nailtail Wallaby	1
Molossidae	<i>Ozimops cobourgianus</i>	North-western Free-tailed Bat	2
Muridae	<i>Conilurus penicillatus</i>	Brush-tailed Tree-rat	1
Muridae	<i>Melomys burtoni</i>	Grassland Melomys	24
Muridae	<i>Pseudomys delicatulus</i>	Delicate Mouse	4
Muridae	<i>Rattus exulans</i>	Pacific Rat	1
Muridae	<i>Rattus rattus</i>	Black Rat	7
Pteropodidae	<i>Macroglossus minimus</i>	Northern Blossom-bat	6
Pteropodidae	<i>Pteropus alecto</i>	Black Flying-fox	4
Pteropodidae	<i>Pteropus scapulatus</i>	Little Red Flying-fox	2
Suidae	<i>Sus scrofa</i>	Pig	1
Vespertilionidae	<i>Chalinolobus gouldii</i>	Gould's Wattled Bat	8
Vespertilionidae	<i>Chalinolobus nigrogriseus</i>	Hoary Wattled Bat	7
Vespertilionidae	<i>Nyctophilus arnhemensis</i>	Arnhem Long-eared Bat	20
Vespertilionidae	<i>Nyctophilus daedalus</i>	Pallid Long-eared Bat	3
Vespertilionidae	<i>Pipistrellus westralis</i>	Northern Pipistrelle	8
Vespertilionidae	<i>Scotorepens greyii</i>	Little Broad-nosed Bat	15
Vespertilionidae	<i>Scotorepens sanborni</i>	Northern Broad-nosed Bat	1
Vespertilionidae	<i>Vespadelus caurinus</i>	Northern Cave Bat	5

Avifauna

Family	Species	Common Name	Number
Acanthizidae	<i>Gerygone chloronota</i>		1
Acanthizidae	<i>Gerygone fusca</i>	Western Gerygone	2
Acanthizidae	<i>Gerygone levigaster</i>	Mangrove Gerygone	21
Acanthizidae	<i>Gerygone olivacea</i>	White-throated Gerygone	32
Acanthizidae	<i>Gerygone tenebrosa</i>	Dusky Gerygone	14
Acanthizidae	<i>Smicromis brevirostris</i>	Weebill	12
Accipitridae	<i>Accipiter cirrocephalus</i>	Collared Sparrowhawk	14
Accipitridae	<i>Accipiter fasciatus</i>	Brown Goshawk	26
Accipitridae	<i>Aquila audax</i>	Wedge-tailed Eagle	9
Accipitridae	<i>Circus approximans</i>	Swamp Harrier	3
Accipitridae	<i>Circus assimilis</i>	Spotted Harrier	3
Accipitridae	<i>Elanus axillaris</i>	Black-shouldered Kite	3
Accipitridae	<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	62
Accipitridae	<i>Haliastur indus</i>	Brahminy Kite	78
Accipitridae	<i>Haliastur sphenurus</i>	Whistling Kite	44
Accipitridae	<i>Hieraaetus morphnoides</i>	Little Eagle	3
Accipitridae	<i>Lophoictinia isura</i>	Square-tailed Kite	4
Accipitridae	<i>Milvus migrans</i>	Black Kite	22
Accipitridae	<i>Pandion cristatus</i>	Eastern Osprey	32
Accipitridae	<i>Pandion haliaetus</i>	Osprey	32
Aegothelidae	<i>Aegotheles cristatus</i>	Australian Owlet-nightjar	4
Alcedinidae	<i>Ceyx azureus</i>	Azure Kingfisher	1
Alcedinidae	<i>Dacelo leachii</i>	Blue-winged Kookaburra	21
Alcedinidae	<i>Todiramphus chloris</i>	Collared Kingfisher	2
Alcedinidae	<i>Todiramphus pyrrhopygius</i>	Red-backed Kingfisher	4
Alcedinidae	<i>Todiramphus sanctus</i>	Sacred Kingfisher	30
Anatidae	<i>Anas gracilis</i>	Grey Teal	9
Anatidae	<i>Anas superciliosa</i>	Pacific Black Duck	29
Anatidae	<i>Apus pacificus</i>	Pacific Swift	6
Anatidae	<i>Aythya australis</i>	Hardhead	4
Anatidae	<i>Dendrocygna arcuata</i>	Wandering Whistling-Duck	7
Anatidae	<i>Nettapus pulchellus</i>	Green Pygmy-goose	1
Anhingidae	<i>Anhinga novaehollandiae</i>	Australasian Darter	5
Ardeidae	<i>Ardea alba</i>	Great Egret	5
Ardeidae	<i>Ardea intermedia</i>	Intermediate Egret	4
Ardeidae	<i>Ardea modesta</i>		7
Ardeidae	<i>Ardea pacifica</i>	White-necked Heron	4
Ardeidae	<i>Ardea sumatrana</i>	Great-billed Heron	4
Ardeidae	<i>Butorides striatus</i>	Striated Heron	32
Ardeidae	<i>Egretta garzetta</i>	Little Egret	17
Ardeidae	<i>Egretta novaehollandiae</i>	White-faced Heron	26
Ardeidae	<i>Egretta sacra</i>	Pacific Reef-Heron	100
Ardeidae	<i>Nycticorax caledonicus</i>	Rufous Night Heron	4
Artamidae	<i>Artamus cinereus</i>	Black-faced Woodswallow	27

Family	Species	Common Name	Number
Artamidae	<i>Artamus leucorhynchus</i>	White-breasted Woodswallow	60
Artamidae	<i>Artamus minor</i>	Little Woodswallow	26
Artamidae	<i>Artamus personatus</i>	Masked Woodswallow	13
Artamidae	<i>Cracticus nigrogularis</i>	Pied Butcherbird	49
Artamidae	<i>Gymnorhina tibicen</i>	Australian Magpie	1
Burhinidae	<i>Burhinus grallarius</i>	Bush Thick-knee	8
Burhinidae	<i>Esacus magnirostris</i>	Beach Thick-knee	48
Burhinidae	<i>Esacus neglectus</i>	Beach Stone-curlew	2
Cacatuidae	<i>Cacatua sanguinea</i>	Little Corella	46
Cacatuidae	<i>Calyptorhynchus banksii</i>	Red-tailed Black-Cockatoo	62
Cacatuidae	<i>Eolophus roseicapilla</i>		1
Cacatuidae	<i>Nymphicus hollandicus</i>	Cockatiel	1
Campephagidae	<i>Coracina novaehollandiae</i>	Black-faced Cuckooshrike	118
Campephagidae	<i>Coracina papuensis</i>	White-bellied Cuckooshrike	1
Campephagidae	<i>Lalage leucomela</i>	Varied Triller	3
Campephagidae	<i>Lalage sueurii</i>	White-winged Triller	33
Caprimulgidae	<i>Eurostopodus argus</i>	Spotted Nightjar	7
Centropodidae	<i>Centropus phasianinus</i>	Pheasant Coucal	16
Charadriidae	<i>Calidris acuminata</i>		8
Charadriidae	<i>Calidris alba</i>		10
Charadriidae	<i>Calidris canutus</i>	Red Knot	2
Charadriidae	<i>Calidris ferruginea</i>		9
Charadriidae	<i>Calidris ruficollis</i>		33
Charadriidae	<i>Calidris subminuta</i>		1
Charadriidae	<i>Calidris tenuirostris</i>		15
Charadriidae	<i>Charadrius leschenaultii</i>		52
Charadriidae	<i>Charadrius mongolus</i>		18
Charadriidae	<i>Charadrius ruficapillus</i>	Red-capped Plover	70
Charadriidae	<i>Elseyornis melanops</i>	Black-fronted Dotterel	18
Charadriidae	<i>Erythrogonys cinctus</i>	Red-kneed Dotterel	3
Charadriidae	<i>Pluvialis fulva</i>	Pacific golden plover	12
Charadriidae	<i>Pluvialis squatarola</i>	Grey plover	25
Charadriidae	<i>Vanellus miles</i>	Masked Lapwing	17
Charadriiformes	<i>Tringa brevipes</i>		58
Charadriiformes	<i>Tringa glareola</i>		6
Charadriiformes	<i>Tringa nebularia</i>		39
Charadriiformes	<i>Tringa stagnatilis</i>		3
Charadriiformes	<i>Tringa totanus</i>		1
Ciconiidae	<i>Ephippiorhynchus asiaticus</i>	Black-necked Stork	18
Cisticolidae	<i>Cisticola exilis</i>	Golden-headed Cisticola	3
Climacteridae	<i>Climacteris melanura</i>	Black-tailed Treecreeper	5
Columbidae	<i>Chalcophaps indica</i>	Emerald Dove	1
Columbidae	<i>Geopelia cuneata</i>	Diamond Dove	13
Columbidae	<i>Geopelia humeralis</i>	Bar-shouldered Dove	170
Columbidae	<i>Geopelia striata</i>	Peaceful Dove	167
Columbidae	<i>Ocyphaps lophotes</i>	Crested Pigeon	5

Family	Species	Common Name	Number
Columbidae	<i>Petrophassa albipennis</i>	White-quilled Rock-Pigeon	1
Columbidae	<i>Ptilinopus regina</i>	Rose-crowned Fruit-Dove	22
Coraciidae	<i>Eurystomus orientalis</i>	Dollarbird	5
Corvidae	<i>Corvus orru</i>	Torresian Crow	62
Cuculidae	<i>Cacomantis pallidus</i>	Pallid Cuckoo	7
Cuculidae	<i>Cacomantis variolosus</i>	Brush Cuckoo	24
Cuculidae	<i>Chalcites basalis</i>	Horsfield's Bronze-Cuckoo	13
Cuculidae	<i>Chalcites minutillus</i>	Little Bronze-Cuckoo	8
Cuculidae	<i>Chalcites osculans</i>	Black-eared Cuckoo	1
Cuculidae	<i>Cuculus optatus</i>		2
Cuculidae	<i>Scythrops novaehollandiae</i>	Channel-billed Cuckoo	4
Dicruridae	<i>Dicrurus bracteatus</i>		2
Estrildidae	<i>Emblema pictum</i>	Painted Firetail	1
Estrildidae	<i>Erythrura gouldiae</i>	Gouldian Finch	35
Estrildidae	<i>Heteromunia pectoralis</i>	Pictorella Munia	1
Estrildidae	<i>Lonchura castaneothorax</i>	Chestnut-breasted Munia	10
Estrildidae	<i>Poephila acuticauda</i>	Long-tailed Finch	83
Estrildidae	<i>Stizoptera bichenovii</i>		127
Estrildidae	<i>Taeniopygia guttata</i>	Zebra Finch	19
Falconidae	<i>Falco berigora</i>	Brown Falcon	25
Falconidae	<i>Falco cenchroides</i>	Australian Kestrel	14
Falconidae	<i>Falco longipennis</i>	Australian Hobby	8
Falconidae	<i>Falco peregrinus</i>	Peregrine Falcon	2
Fregatidae	<i>Fregata ariel</i>	Lesser Frigatebird	21
Fregatidae	<i>Fregata minor</i>	Great Frigatebird	1
Glareolidae	<i>Stiltia isabella</i>	Australian Pratincole	4
Gruidae	<i>Grus rubicunda</i>	Brolga	3
Haematopodidae	<i>Haematopus fuliginosus</i>	Sooty Oystercatcher	88
Haematopodidae	<i>Haematopus longirostris</i>	Pied Oystercatcher	92
Hirundinidae	<i>Petrochelidon ariel</i>	Fairy Martin	3
Hirundinidae	<i>Petrochelidon nigricans</i>	Tree Martin	28
Laridae	<i>Anous stolidus</i>		5
Laridae	<i>Chlidonias hybrida</i>	Whiskered Tern	29
Laridae	<i>Chlidonias leucopterus</i>		2
Laridae	<i>Chroicocephalus novaehollandiae</i>	Silver Gull	154
Laridae	<i>Gelochelidon nilotica</i>	Gull-billed Tern	32
Laridae	<i>Hydroprogne caspia</i>	Caspian Tern	10
Laridae	<i>Onychoprion anaethetus</i>	Bridled Tern	9
Laridae	<i>Onychoprion fuscata</i>	Sooty Tern	3
Laridae	<i>Sterna dougallii</i>		8
Laridae	<i>Sterna hirundo</i>		7
Laridae	<i>Sternula albifrons</i>	Little Tern	12
Laridae	<i>Thalasseus bengalensis</i>	Lesser Crested Tern	23
Laridae	<i>Thalasseus bergii</i>		64
Maluridae	<i>Malurus assimilis</i>	Purple-backed Fairywren	14
Maluridae	<i>Malurus elegans</i>	Red-winged Fairywren	1

Family	Species	Common Name	Number
Maluridae	<i>Malurus lamberti</i>	Variegated Fairy-wren	8
Maluridae	<i>Malurus melanocephalus</i>	Red-backed Fairywren	97
Megaluridae	<i>Cincloramphus cruralis</i>	Brown Songlark	1
Megaluridae	<i>Cincloramphus mathewsi</i>	Rufous Songlark	4
Meliphagidae	<i>Certhionyx pectoralis</i>		18
Meliphagidae	<i>Certhionyx variegatus</i>	Pied Honeyeater	3
Meliphagidae	<i>Conopophila rufogularis</i>	Rufous-throated Honeyeater	74
Meliphagidae	<i>Epthianura crocea</i>	Yellow Chat	1
Meliphagidae	<i>Gavicalis virescens</i>	Singing Honeyeater	85
Meliphagidae	<i>Lichmera indistincta</i>	Brown Honeyeater	162
Meliphagidae	<i>Melithreptus albogularis</i>	White-throated Honeyeater	11
Meliphagidae	<i>Melithreptus gularis</i>	Black-chinned Honeyeater	12
Meliphagidae	<i>Myzomela erythrocephala</i>	Red-headed Myzomela	39
Meliphagidae	<i>Philemon argenticeps</i>	Silver-crowned Friarbird	9
Meliphagidae	<i>Philemon citreogularis</i>	Little Friarbird	98
Meliphagidae	<i>Ptilotula flavescens</i>	Yellow-tinted Honeyeater	94
Meliphagidae	<i>Ptilotula keartlandi</i>		1
Meliphagidae	<i>Ptilotula plumula</i>		1
Meliphagidae	<i>Puffinus assimilis</i>		1
Meliphagidae	<i>Ramsayornis fasciatus</i>	Bar-breasted Honeyeater	1
Meliphagidae	<i>Stomiopera unicolor</i>	White-gaped Honeyeater	146
Meropidae	<i>Merops ornatus</i>	Rainbow Bee-eater	107
Monarchidae	<i>Grallina cyanoleuca</i>	Magpie-lark	79
Monarchidae	<i>Myiagra alecto</i>	Shining Flycatcher	16
Monarchidae	<i>Myiagra inquieta</i>	Restless Flycatcher	24
Monarchidae	<i>Myiagra rubecula</i>	Leaden Flycatcher	23
Monarchidae	<i>Myiagra ruficollis</i>	Broad-billed Flycatcher	31
Motacillidae	<i>Anthus novaeseelandiae</i>	Australasian Pipit	6
Nectariniidae	<i>Dicaeum hirundinaceum</i>	Mistletoebird	122
Neosittidae	<i>Daphoenositta chrysoptera</i>	Varied Sittella	14
Oceanitidae	<i>Oceanites oceanicus</i>	Wilson's Storm Petrel	1
Oriolidae	<i>Oriolus flavocinctus</i>	Yellow Oriole	1
Oriolidae	<i>Oriolus sagittatus</i>	Olive-backed Oriole	30
Oriolidae	<i>Sphecotheres vieilloti</i>	Australasian Figbird	1
Otididae	<i>Ardeotis australis</i>	Australian Bustard	1
Pachycephalidae	<i>Colluricincla harmonica</i>	Gray Shrikethrush	39
Pachycephalidae	<i>Pachycephala lanioides</i>	White-breasted Whistler	13
Pachycephalidae	<i>Pachycephala melanura</i>	Black-tailed Whistler	33
Pachycephalidae	<i>Pachycephala pectoralis</i>	Golden Whistler	1
Pachycephalidae	<i>Pachycephala rufiventris</i>	Rufous Whistler	131
Pardalotidae	<i>Pardalotus rubricatus</i>	Red-browed Pardalote	3
Pardalotidae	<i>Pardalotus striatus</i>	Striated Pardalote	28
Pelecanidae	<i>Pelecanus conspicillatus</i>	Australian Pelican	25
Petroicidae	<i>Melanodryas cucullata</i>	Hooded Robin	18
Petroicidae	<i>Microeca fascinans</i>	Jacky-winter	63
Petroicidae	<i>Microeca flavigaster</i>		19

Family	Species	Common Name	Number
Phaethontidae	<i>Phaethon rubricauda</i>	Red-tailed Tropicbird	2
Phalacrocoracidae	<i>Microcarbo melanoleucos</i>	Little Pied Cormorant	9
Phalacrocoracidae	<i>Phalacrocorax melanoleucos</i>	Little Pied Cormorant	5
Phalacrocoracidae	<i>Phalacrocorax sulcirostris</i>	Little Black Cormorant	4
Phalacrocoracidae	<i>Phalacrocorax varius</i>	Pied Cormorant	33
Phasianidae	<i>Coturnix ypsilophora</i>	Brown Quail	11
Podargidae	<i>Podargus strigoides</i>	Tawny Frogmouth	5
Podicipedidae	<i>Poliiocephalus poliocephalus</i>	Hoary-headed Grebe	1
Podicipedidae	<i>Tachybaptus novaehollandiae</i>	Australasian Grebe	23
Pomatostomidae	<i>Pomatostomus temporalis</i>	Gray-crowned Babbler	75
Psittacidae	<i>Aprosmictus erythropterus</i>	Red-winged Parrot	83
Psittacidae	<i>Melopsittacus undulatus</i>	Budgerigar	6
Psittacidae	<i>Psitteuteles versicolor</i>	Varied Lorikeet	15
Psittacidae	<i>Trichoglossus haematodus</i>	Rainbow Lorikeet	37
Ptilonorhynchidae	<i>Ptilonorhynchus nuchalis</i>	Great Bowerbird	63
Rallidae	<i>Eulabeornis castaneiventris</i>	0	1
Rallidae	<i>Fulica atra</i>	Eurasian Coot	2
Rallidae	<i>Gallirallus philippensis</i>		1
Rallidae	<i>Porphyrio porphyrio</i>	Purple Swamphen	1
Rallidae	<i>Porzana tabuensis</i>	Spotless Crake	1
Recurvirostridae	<i>Himantopus himantopus</i>	Black-winged Stilt	9
Rhipiduridae	<i>Rhipidura albiscapa</i>	Grey Fantail	6
Rhipiduridae	<i>Rhipidura leucophrys</i>	Willie-wagtail	168
Rhipiduridae	<i>Rhipidura phasiana</i>	Mangrove Fantail	18
Rhipiduridae	<i>Rhipidura rufiventris</i>	Northern Fantail	75
Scolopacidae	<i>Actitis hypoleucos</i>	Common Sandpiper	53
Scolopacidae	<i>Arenaria interpres</i>		35
Scolopacidae	<i>Limosa lapponica</i>	Bar-tailed Godwit	33
Scolopacidae	<i>Limosa limosa</i>	Black-tailed Godwit	3
Scolopacidae	<i>Numenius madagascariensis</i>	Eastern curlew	23
Scolopacidae	<i>Numenius minutus</i>	Little Curlew	3
Scolopacidae	<i>Numenius phaeopus</i>	Whimbrel	51
Scolopacidae	<i>Xenus cinereus</i>	Terek Sandpiper	11
Strigidae	<i>Ninox connivens</i>	Barking Owl	9
Strigidae	<i>Ninox novaeseelandiae</i>	Southern Boobook	1
Sulidae	<i>Sula leucogaster</i>		58
Threskiornithidae	<i>Platalea regia</i>	Royal Spoonbill	15
Threskiornithidae	<i>Plegadis falcinellus</i>	Glossy Ibis	6
Threskiornithidae	<i>Threskiornis moluccus</i>	Australian Ibis	42
Threskiornithidae	<i>Threskiornis spinicollis</i>	Straw-necked Ibis	28
Timaliidae	<i>Zosterops lateralis</i>		2
Timaliidae	<i>Zosterops luteus</i>	Australian Yellow White-eye	72
Turnicidae	<i>Turnix castanotus</i>	Chestnut-backed Button-quail	6
Turnicidae	<i>Turnix maculosus</i>	Red-backed Buttonquail	2
Turnicidae	<i>Turnix velox</i>	Little Button-quail	1
Tytonidae	<i>Tyto javanica</i>	Eastern Barn Owl	3

Amphibians

Family	Species	Common	Number
Pelodyadidae	<i>Cyclorana australis</i>	Giant Frog	4
	<i>Litoria caerulea</i>	Green Tree Frog	4
	<i>Litoria rothii</i>	Northern Laughing Tree Frog	3
Limnodynastidae	<i>Platyplectrum ornatum</i>	Ornate Burrowing Frog	1

Invertebrates

Family	Species	Number
Camaenidae	<i>Quistrachia leptogramma</i>	6
	<i>Quistrachia monogramma</i>	1
	<i>Rhagada bulgana</i>	15
	<i>Rhagada cygna</i>	18
	<i>Rhagada dominica</i>	3
	<i>Rhagada primigena</i>	1
	<i>Rhagada reinga</i>	1
	<i>Rhagada sp.</i>	38
	<i>Westraltrachia oscarensis</i>	1
Buthidae	<i>Lychas sp.</i>	5
Urodacidae	<i>Urodacus hoplurus</i>	7
Anamidae	<i>Aname sp.</i>	7
Halonoproctidae	<i>Conothele sp.</i>	1

Appendix 4

EPBC Act Protected Matters Database Search Results



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 29/10/20 13:22:39

[Summary](#)

[Details](#)

[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



This map may contain data which are
©Commonwealth of Australia
(Geoscience Australia), ©PSMA 2015

[Coordinates](#)

[Buffer: 40.0Km](#)



This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	1
Wetlands of International Importance:	None
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	1
Listed Threatened Ecological Communities:	1
Listed Threatened Species:	31
Listed Migratory Species:	53

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	None
Commonwealth Heritage Places:	None
Listed Marine Species:	84
Whales and Other Cetaceans:	15
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Australian Marine Parks:	2

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	2
Regional Forest Agreements:	None
Invasive Species:	9
Nationally Important Wetlands:	None
Key Ecological Features (Marine)	None

National Heritage Properties

Name	State	Status
Natural		
The West Kimberley	WA	Listed place

Commonwealth Marine Area

[\[Resource Information \]](#)

Approval is required for a proposed activity that is located within the Commonwealth Marine Area which has, will have, or is likely to have a significant impact on the environment. Approval may be required for a proposed action taken outside the Commonwealth Marine Area but which has, may have or is likely to have a significant impact on the environment in the Commonwealth Marine Area. Generally the Commonwealth Marine Area stretches from three nautical miles to two hundred nautical miles from the coast.

Name

EEZ and Territorial Sea

Marine Regions

[\[Resource Information \]](#)

If you are planning to undertake action in an area in or close to the Commonwealth Marine Area, and a marine bioregional plan has been prepared for the Commonwealth Marine Area in that area, the marine bioregional plan may inform your decision as to whether to refer your proposed action under the EPBC Act.

Name

[North-west](#)

Listed Threatened Ecological Communities

[\[Resource Information \]](#)

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Name	Status	Type of Presence
Monsoon vine thickets on the coastal sand dunes of Dampier Peninsula	Endangered	Community likely to occur within area

Listed Threatened Species

[\[Resource Information \]](#)

Name	Status	Type of Presence
Birds		
Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
Erythrotriorchis radiatus Red Goshawk [942]	Vulnerable	Species or species habitat may occur within area
Erythrura gouldiae Gouldian Finch [413]	Endangered	Species or species habitat known to occur within area
Falco hypoleucos Grey Falcon [929]	Vulnerable	Species or species habitat may occur within area
Limosa lapponica baueri Bar-tailed Godwit (baueri), Western Alaskan Bar-tailed Godwit [86380]	Vulnerable	Species or species habitat may occur within area

Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat known to occur within area
Papasula abbotti Abbott's Booby [59297]	Endangered	Species or species habitat may occur within area
Pezoporus occidentalis Night Parrot [59350]	Endangered	Species or species habitat may occur within area
Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat likely to occur within area
Tyto novaehollandiae kimberli Masked Owl (northern) [26048]	Vulnerable	Species or species habitat may occur within area
Mammals		
Balaenoptera borealis Sei Whale [34]	Vulnerable	Species or species habitat may occur within area
Balaenoptera musculus Blue Whale [36]	Endangered	Species or species habitat likely to occur within area
Balaenoptera physalus Fin Whale [37]	Vulnerable	Species or species habitat may occur within area
Macroderma gigas Ghost Bat [174]	Vulnerable	Species or species habitat likely to occur within area
Macrotis lagotis Greater Bilby [282]	Vulnerable	Species or species habitat known to occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Breeding known to occur within area
Saccolaimus saccolaimus nudicluniatus Bare-rumped Sheath-tailed Bat, Bare-rumped Sheath-tail Bat [66889]	Vulnerable	Species or species habitat may occur within area
Xeromys myoides Water Mouse, False Water Rat, Yirrkoo [66]	Vulnerable	Species or species habitat may occur within area
Reptiles		
Caretta caretta Loggerhead Turtle [1763]	Endangered	Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Breeding known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Breeding likely to occur within area
Dermochelys imbricata		

Glyphis garricki Northern River Shark, New Guinea River Shark [82454]	Endangered	Breeding likely to occur within area
Pristis clavata Dwarf Sawfish, Queensland Sawfish [68447]	Vulnerable	Breeding known to occur within area
Pristis pristis Freshwater Sawfish, Largetooth Sawfish, River Sawfish, Leichhardt's Sawfish, Northern Sawfish [60756]	Vulnerable	Species or species habitat known to occur within area
Pristis zijsron Green Sawfish, Dindagubba, Narrowsnout Sawfish [68442]	Vulnerable	Breeding known to occur within area
Rhincodon typus Whale Shark [66680]	Vulnerable	Foraging, feeding or related behaviour known to occur within area

Listed Migratory Species

[[Resource Information](#)]

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Migratory Marine Birds		
Anous stolidus Common Noddy [825]		Species or species habitat likely to occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Calonectris leucomelas Streaked Shearwater [1077]		Species or species habitat known to occur within area
Fregata ariel Lesser Frigatebird, Least Frigatebird [1012]		Breeding known to occur within area
Fregata minor Great Frigatebird, Greater Frigatebird [1013]		Species or species habitat likely to occur within area
Onychoprion anaethetus Bridled Tern [82845]		Breeding known to occur within area
Sterna dougallii Roseate Tern [817]		Breeding known to occur within area
Sternula albifrons Little Tern [82849]		Breeding known to occur within area
Sula sula Red-footed Booby [1023]		Breeding known to occur within area
Migratory Marine Species		
Anoxypristis cuspidata Narrow Sawfish, Knifetooth Sawfish [68448]		Species or species habitat likely to occur within area
Balaenoptera borealis Sei Whale [34]	Vulnerable	Species or species habitat may occur within area

Fin Whale [37]	Vulnerable	Species or species habitat may occur within area
Carcharhinus longimanus Oceanic Whitetip Shark [84108]		Species or species habitat may occur within area
Carcharodon carcharias White Shark, Great White Shark [64470]	Vulnerable	Species or species habitat may occur within area
Caretta caretta Loggerhead Turtle [1763]	Endangered	Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Breeding known to occur within area
Crocodylus porosus Salt-water Crocodile, Estuarine Crocodile [1774]		Species or species habitat likely to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Breeding likely to occur within area
Dugong dugon Dugong [28]		Foraging, feeding or related behaviour likely to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Breeding likely to occur within area
Isurus oxyrinchus Shortfin Mako, Mako Shark [79073]		Species or species habitat likely to occur within area
Isurus paucus Longfin Mako [82947]		Species or species habitat likely to occur within area
Manta alfredi Reef Manta Ray, Coastal Manta Ray, Inshore Manta Ray, Prince Alfred's Ray, Resident Manta Ray [84994]		Species or species habitat known to occur within area
Manta birostris Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray, Oceanic Manta Ray [84995]		Species or species habitat likely to occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Breeding known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Breeding known to occur within area
Orcaella heinsohni Australian Snubfin Dolphin [81322]		Species or species habitat likely to occur within area
Orcinus orca Killer Whale, Orca [46]		Species or species habitat may occur within area
Pristis clavata		

Whale Shark [66680]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Sousa chinensis Indo-Pacific Humpback Dolphin [50]		Breeding known to occur within area
Tursiops aduncus (Arafura/Timor Sea populations) Spotted Bottlenose Dolphin (Arafura/Timor Sea populations) [78900]		Species or species habitat known to occur within area

Migratory Terrestrial Species

Cecropis daurica Red-rumped Swallow [80610]		Species or species habitat may occur within area
Cuculus optatus Oriental Cuckoo, Horsfield's Cuckoo [86651]		Species or species habitat may occur within area
Hirundo rustica Barn Swallow [662]		Species or species habitat may occur within area
Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat likely to occur within area

Migratory Wetlands Species

Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat known to occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Charadrius veredus Oriental Plover, Oriental Dotterel [882]		Species or species habitat may occur within area
Glareola maldivarum Oriental Pratincole [840]		Species or species habitat may occur within area
Limosa lapponica Bar-tailed Godwit [844]		Species or species habitat

[Tringa nebularia](#)

Common Greenshank, Greenshank [832]

within area

Species or species habitat likely to occur within area

Other Matters Protected by the EPBC Act

Listed Marine Species

[[Resource Information](#)]

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Birds		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat known to occur within area
Anous stolidus Common Noddy [825]		Species or species habitat likely to occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba Great Egret, White Egret [59541]		Species or species habitat known to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris canutus Red Knot, Knot [855]	Endangered	Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat known to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Calonectris leucomelas Streaked Shearwater [1077]		Species or species habitat known to occur within area

Black-eared Cuckoo [705]		Species or species habitat likely to occur within area
Fregata ariel		
Lesser Frigatebird, Least Frigatebird [1012]		Breeding known to occur within area
Fregata minor		
Great Frigatebird, Greater Frigatebird [1013]		Species or species habitat likely to occur within area
Glareola maldivarum		
Oriental Pratincole [840]		Species or species habitat may occur within area
Haliaeetus leucogaster		
White-bellied Sea-Eagle [943]		Species or species habitat known to occur within area
Hirundo daurica		
Red-rumped Swallow [59480]		Species or species habitat may occur within area
Hirundo rustica		
Barn Swallow [662]		Species or species habitat may occur within area
Larus novaehollandiae		
Silver Gull [810]		Breeding known to occur within area
Limosa lapponica		
Bar-tailed Godwit [844]		Species or species habitat known to occur within area
Merops ornatus		
Rainbow Bee-eater [670]		Species or species habitat may occur within area
Motacilla cinerea		
Grey Wagtail [642]		Species or species habitat may occur within area
Motacilla flava		
Yellow Wagtail [644]		Species or species habitat likely to occur within area
Numenius madagascariensis		
Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat known to occur within area
Pandion haliaetus		
Osprey [952]		Breeding known to occur within area
Papasula abbotti		
Abbott's Booby [59297]	Endangered	Species or species habitat may occur within area
Rostratula benghalensis (sensu lato)		
Painted Snipe [889]	Endangered*	Species or species habitat likely to occur within area
Sterna albifrons		
Little Tern [813]		Breeding known to occur within area

Red-footed Booby [1023]	Breeding known to occur within area
Tringa nebularia Common Greenshank, Greenshank [832]	Species or species habitat likely to occur within area
Fish	
Campichthys tricarinatus Three-keel Pipefish [66192]	Species or species habitat may occur within area
Choeroichthys brachysoma Pacific Short-bodied Pipefish, Short-bodied Pipefish [66194]	Species or species habitat may occur within area
Choeroichthys suillus Pig-snouted Pipefish [66198]	Species or species habitat may occur within area
Corythoichthys amplexus Fijian Banded Pipefish, Brown-banded Pipefish [66199]	Species or species habitat may occur within area
Corythoichthys flavofasciatus Reticulate Pipefish, Yellow-banded Pipefish, Network Pipefish [66200]	Species or species habitat may occur within area
Cosmocampus banneri Roughridge Pipefish [66206]	Species or species habitat may occur within area
Doryrhamphus excisus Bluestripe Pipefish, Indian Blue-stripe Pipefish, Pacific Blue-stripe Pipefish [66211]	Species or species habitat may occur within area
Doryrhamphus janssi Cleaner Pipefish, Janss' Pipefish [66212]	Species or species habitat may occur within area
Filicampus tigris Tiger Pipefish [66217]	Species or species habitat may occur within area
Halicampus brocki Brock's Pipefish [66219]	Species or species habitat may occur within area
Halicampus grayi Mud Pipefish, Gray's Pipefish [66221]	Species or species habitat may occur within area
Halicampus nitidus Glittering Pipefish [66224]	Species or species habitat may occur within area
Halicampus spinostris Spiny-snout Pipefish [66225]	Species or species habitat may occur within area
Haliichthys taeniophorus Ribbioned Pipehorse, Ribbioned Seadragon [66226]	Species or species habitat may occur within area

Spotted Seahorse, Yellow Seahorse [66237]	Species or species habitat may occur within area
Hippocampus planifrons Flat-face Seahorse [66238]	Species or species habitat may occur within area
Hippocampus spinosissimus Hedgehog Seahorse [66239]	Species or species habitat may occur within area
Hippocampus trimaculatus Three-spot Seahorse, Low-crowned Seahorse, Flat-faced Seahorse [66720]	Species or species habitat may occur within area
Micrognathus micronotus Tidepool Pipefish [66255]	Species or species habitat may occur within area
Solegnathus hardwickii Pallid Pipehorse, Hardwick's Pipehorse [66272]	Species or species habitat may occur within area
Solegnathus lettiensis Gunther's Pipehorse, Indonesian Pipefish [66273]	Species or species habitat may occur within area
Solenostomus cyanopterus Robust Ghostpipefish, Blue-finned Ghost Pipefish, [66183]	Species or species habitat may occur within area
Syngnathoides biaculeatus Double-end Pipehorse, Double-ended Pipehorse, Alligator Pipefish [66279]	Species or species habitat may occur within area
Trachyrhamphus bicoarctatus Bentstick Pipefish, Bend Stick Pipefish, Short-tailed Pipefish [66280]	Species or species habitat may occur within area
Trachyrhamphus longirostris Straightstick Pipefish, Long-nosed Pipefish, Straight Stick Pipefish [66281]	Species or species habitat may occur within area

Mammals

Dugong dugon Dugong [28]	Foraging, feeding or related behaviour likely to occur within area
---	--

Reptiles

Acalyptophis peronii Horned Seasnake [1114]	Species or species habitat may occur within area
Aipysurus duboisii Dubois' Seasnake [1116]	Species or species habitat may occur within area
Aipysurus eydouxii Spine-tailed Seasnake [1117]	Species or species habitat may occur within area

[Aipysurus laevis](#)

Olive Seasnake [1120]	Species or species habitat may occur within area
-----------------------	--

Caretta caretta Loggerhead Turtle [1763]	Endangered	Foraging, feeding or related behaviour known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Breeding known to occur within area
Crocodylus johnstoni Freshwater Crocodile, Johnston's Crocodile, Johnston's River Crocodile [1773]		Species or species habitat may occur within area
Crocodylus porosus Salt-water Crocodile, Estuarine Crocodile [1774]		Species or species habitat likely to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Breeding likely to occur within area
Disteira kingii Spectacled Seasnake [1123]		Species or species habitat may occur within area
Disteira major Olive-headed Seasnake [1124]		Species or species habitat may occur within area
Emydocephalus annulatus Turtle-headed Seasnake [1125]		Species or species habitat may occur within area
Ephalophis greyi North-western Mangrove Seasnake [1127]		Species or species habitat may occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Breeding likely to occur within area
Hydrelaps darwiniensis Black-ringed Seasnake [1100]		Species or species habitat may occur within area
Hydrophis elegans Elegant Seasnake [1104]		Species or species habitat may occur within area
Hydrophis mcdowelli null [25926]		Species or species habitat may occur within area
Hydrophis ornatus Spotted Seasnake, Ornate Reef Seasnake [1111]		Species or species habitat may occur within area
Lapemis hardwickii Spine-bellied Seasnake [1113]		Species or species habitat may occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Breeding known to occur within area
Pelamis platurus Yellow-bellied Seasnake [1091]		Species or species habitat may occur within area

Bryde's Whale [35]		Species or species habitat may occur within area
Balaenoptera musculus Blue Whale [36]	Endangered	Species or species habitat likely to occur within area
Balaenoptera physalus Fin Whale [37]	Vulnerable	Species or species habitat may occur within area
Delphinus delphis Common Dolphin, Short-beaked Common Dolphin [60]		Species or species habitat may occur within area
Grampus griseus Risso's Dolphin, Grampus [64]		Species or species habitat may occur within area
Megaptera novaeangliae Humpback Whale [38]	Vulnerable	Breeding known to occur within area
Orcaella brevirostris Irrawaddy Dolphin [45]		Species or species habitat likely to occur within area
Orcinus orca Killer Whale, Orca [46]		Species or species habitat may occur within area
Pseudorca crassidens False Killer Whale [48]		Species or species habitat likely to occur within area
Sousa chinensis Indo-Pacific Humpback Dolphin [50]		Breeding known to occur within area
Stenella attenuata Spotted Dolphin, Pantropical Spotted Dolphin [51]		Species or species habitat may occur within area
Tursiops aduncus Indian Ocean Bottlenose Dolphin, Spotted Bottlenose Dolphin [68418]		Species or species habitat likely to occur within area
Tursiops aduncus (Arafura/Timor Sea populations) Spotted Bottlenose Dolphin (Arafura/Timor Sea populations) [78900]		Species or species habitat known to occur within area
Tursiops truncatus s. str. Bottlenose Dolphin [68417]		Species or species habitat may occur within area

Australian Marine Parks

[Resource Information]

Name	Label
Kimberley	Habitat Protection Zone (IUCN IV)
Kimberley	Multiple Use Zone (IUCN VI)

Invasive Species[\[Resource Information \]](#)

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

Name	Status	Type of Presence
Frogs		
Rhinella marina Cane Toad [83218]		Species or species habitat may occur within area
Mammals		
Canis lupus familiaris Domestic Dog [82654]		Species or species habitat likely to occur within area
Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
Rattus rattus Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
Plants		
Cenchrus ciliaris Buffel-grass, Black Buffel-grass [20213]		Species or species habitat likely to occur within area
Jatropha gossypifolia Cotton-leaved Physic-Nut, Bellyache Bush, Cotton-leaf Physic Nut, Cotton-leaf Jatropha, Black Physic Nut [7507]		Species or species habitat likely to occur within area
Lantana camara Lantana, Common Lantana, Kamara Lantana, Large-leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White Sage, Wild Sage [10892]		Species or species habitat may occur within area
Parkinsonia aculeata Parkinsonia, Jerusalem Thorn, Jelly Bean Tree, Horse Bean [12301]		Species or species habitat likely to occur within area
Reptiles		
Ramphotyphlops braminus Flowerpot Blind Snake, Brahminy Blind Snake, Cacing Besi [1258]		Species or species habitat may occur within area

and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-16.468 122.9755

[-Department of Environment and Primary Industries, Victoria](#)
[-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
[-Department of Environment, Water and Natural Resources, South Australia](#)
[-Department of Land and Resource Management, Northern Territory](#)
[-Department of Environmental and Heritage Protection, Queensland](#)
[-Department of Parks and Wildlife, Western Australia](#)
[-Environment and Planning Directorate, ACT](#)
[-Birdlife Australia](#)
[-Australian Bird and Bat Banding Scheme](#)
[-Australian National Wildlife Collection](#)
[-Natural history museums of Australia](#)
[-Museum Victoria](#)
[-Australian Museum](#)
[-South Australian Museum](#)
[-Queensland Museum](#)
[-Online Zoological Collections of Australian Museums](#)
[-Queensland Herbarium](#)
[-National Herbarium of NSW](#)
[-Royal Botanic Gardens and National Herbarium of Victoria](#)
[-Tasmanian Herbarium](#)
[-State Herbarium of South Australia](#)
[-Northern Territory Herbarium](#)
[-Western Australian Herbarium](#)
[-Australian National Herbarium, Canberra](#)
[-University of New England](#)
[-Ocean Biogeographic Information System](#)
[-Australian Government, Department of Defence](#)
[Forestry Corporation, NSW](#)
[-Geoscience Australia](#)
[-CSIRO](#)
[-Australian Tropical Herbarium, Cairns](#)
[-eBird Australia](#)
[-Australian Government – Australian Antarctic Data Centre](#)
[-Museum and Art Gallery of the Northern Territory](#)
[-Australian Government National Environmental Science Program](#)
[-Australian Institute of Marine Science](#)
[-Reef Life Survey Australia](#)
[-American Museum of Natural History](#)
[-Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
[-Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
-Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.

Appendix 5

Significant Flora Species Recorded within 40 km and their Likelihood of Occurrence



Taxon	Habit and Habitat (WA Herbarium 2020)	Source			Likelihood of Occurrence in Study Area	
		Nature Map	DBCA TPFL	WA Herbarium	Initial Ranking Based on Desktop Study	Final Ranking Including Results of Field Survey
Priority 1						
<i>Cullen candidum</i>	Erect spindly shrub to 3 m. Grows in woodland; sometimes common in disturbed areas, along roadsides.	✓	✓	✓	Unlikely to occur; single record from "One Arm Point" in 1987 (location coordinates not precise); not recorded by any other surveys on Dampier Peninsula. (NR=21.1 km NE.)	Unlikely to occur.
<i>Haemodorum capitatum</i>	Bulbous perennial herb, to 50 cm tall, growing in Melaleuca woodland.	✓		✓	Unlikely to occur; no suitable habitat and only 1 record from locality (NR 13.6 km NNE).	Unlikely to occur.
<i>Parsonsia kimberleyensis</i>	Climber growing in vine thickets.	✓		✓	Unlikely to occur; does not appear to be any suitable habitat (NR 13.4 km NE).	Unlikely to occur.
<i>Tribulopsis</i> sp. Koolan Island (K.F. Kenneally 8278)	Prostrate annual herb growing in rocky sandstone areas.	✓		✓	Would not occur; no suitable habitat (NR 32.5 km NE).	Would not occur.
Priority 2						
<i>Alysicarpus suffruticosus</i>	Low shrub to 40 cm tall, growing in open woodland in valley floors and creeks over sandstone.	✓	✓	✓	Unlikely to occur; no suitable habitat (NR=32.5 km NE).	Unlikely to occur.
Priority 3						
<i>Lophostemon grandiflorus</i> subsp. <i>grandiflorus</i>	Low tree associated with creeklines and low-lying areas behind dunes.	✓		✓	Would not occur; no suitable habitat (NR 5.4 km NE).	Would not occur.
<i>Stylidium pindanicum</i>	Herb to 35 cm tall, growing in seepage areas, dry creeks through pindan and other low-lying habitats.	✓		✓	Unlikely to occur; does not appear to be any suitable habitat (NR 13.6 km NNE).	Unlikely to occur.
<i>Tephrosia valleculata</i>	Shrub to 1.6 m tall, growing on rugged rocky areas.	✓		✓	Would not occur; no suitable habitat (NR 23.6 km NE).	Would not occur.
<i>Triodia acutispicula</i>	Hummock grass, growing in sand and rocky areas.	✓		✓	May occur; suitable habitat and several records in locality (NR 2.6 km ENE).	Unlikely to occur.
<i>Utricularia bidentata</i>	Small delicate herb to 15 cm tall, growing in moist areas along creeklines.	✓		✓	Unlikely to occur; does not appear to be any suitable habitat (NR 11.8 km NNE).	Unlikely to occur.

Appendix 6

Significant Fauna Species Recorded within 40 km and their Likelihood of Occurrence



FAUNA GROUP	Conservation Status		Record Source			Habitat Preference	Habitat Potentially Available in Study Area?	Nearest Record	Record Details (if available)	Likelihood of Occurrence in Study Area	
	State	Federal	Nature Map	ALA	EPBC					Initial Ranking Based on Desktop Study	Final Ranking Including Results of Field Survey
REPTILES											
<i>Lerista separanda</i> Dampierland Plain Slider	Priority 2	–	✓	✓	–	Sandy areas of the SW Kimberley coast.	Yes	14 km SW of Lombadina	Three records from 2008.	Likely to occur.	Likely to occur.
<i>Simoselaps minimus</i> Dampierland Burrowing Snake	Priority 2	–	✓	✓	–	Dampier Peninsula endemic. Coastal dunes and adjacent <i>Acacia</i> shrubland on SW Kimberley coast.	Yes	4 km NW of Lombadina	Six records spanning 1989, 1992, 2008.	Likely to occur.	Likely to occur.
MAMMALS											
<i>Ozimops cobourgianus</i> Northern Coastal Free-tailed Bat	Priority 1	–	✓	✓	–	Mangroves and adjacent coastal habitat. Roosts in tree hollows	Yes	3 km SW of Lombadina.	11 records from 1981 and 2008.	Likely to occur.	Likely to occur
<i>Macrotis lagotis</i> Bilby	Vulnerable	Vulnerable	✓	–	✓	<i>Triodia</i> hummock grassland and <i>Acacia</i> scrub across northern Australia.	Yes	10 km NE of Lombadina.	Single record from 1975.	May occur.	May occur
<i>Macroderma gigas</i> Ghost Bat	Vulnerable	Vulnerable	–	–	✓	Patchy distribution, requires undisturbed roost caves or mineshafts.	No	No records on Dampier Peninsula.	Nearest record 85 km northeast on Bathurst Island.	Unlikely to occur.	Unlikely to occur.
<i>Dasyurus hallucatus</i> Northern Quoll	Endangered	Endangered	–	✓	–	Range of vegetation types, mostly within 200 km of coast.	No	17 km NE of Lombadina.	Single record from 2017, low accuracy co-ordinates associated with record. Nearest NatureMap record is from Broome in 2015.	Unlikely to occur.	Unlikely to occur.
BIRDS											
<i>Erythrura gouldiae</i> Gouldian Finch	Priority 4	Endangered	✓	✓	✓	Open grassy eucalypt woodlands and grasslands. Forages on native grass especially sorghum.	Yes	Multiple records within 5 km of Lombadina	Records made between 1977 and 2017.	Likely to occur.	Likely to occur.
<i>Apus pacificus</i> Pacific Swift	Migratory	Migratory/Marine	✓	✓	–	Almost exclusively aerial	Yes	9 km SE of Lombadina.	Single record from 2012.	Likely to occur.	Likely to occur (aerial only).
<i>Falco peregrinus</i> Peregrine Falcon	Other Specially Protected Species	–	✓	✓	–	Variety of habitats	Yes	14 km NNE of Lombadina.	Single record from 2014.	Likely to occur.	Likely to occur
<i>Cuculus optatus</i> Oriental Cuckoo	Migratory	Migratory/Marine	–	✓	–	Dense open woodlands and forests, riparian forests, sometimes gardens	No	7 km W of Lombadina.	Three birds seen in 2020.	May occur.	May occur
<i>Glareola maldivarum</i> Oriental Pratincole	Migratory	Migratory	–	–	✓	Mostly aerial, above open country.	Yes	60 km SW of Lombadina.	–	Unlikely to occur.	May occur
<i>Hirundo rustica</i> Barn Swallow	Migratory	Migratory	–	–	✓	Wide variety of habitats, but generally preferring open country, often near water.	Yes	90 km SW offshore of Lombadina	Many records in Broome.	Unlikely to occur.	May occur
<i>Erythrotriorchis radiatus</i> Red Goshawk	–	Vulnerable	–	–	✓	Mosaic of vegetation types, often near wetlands. They often occur at the boundary between two vegetation types, and often favour forests or woodlands dominated by eucalypts or paperbarks. They avoid very dense or very open habitats.	No	No records within 40 km of Lombadina.	Single record from 1976, 30 km north of Broome.	Unlikely to occur.	Unlikely to occur.
<i>Falco hypoleucos</i> Grey Falcon	–	Vulnerable	–	–	✓	Open plains, both timbered and treeless, open woodland, and tree-lined watercourses	Yes	No records on Dampier Peninsula.	Nearest record on Koolan Island, 80 km NE of Dampier Peninsula.	Unlikely to occur.	Unlikely to occur.

FAUNA GROUP	Conservation Status		Record Source			Habitat Preference	Habitat Potentially Available in Study Area?	Nearest Record	Record Details (if available)	Likelihood of Occurrence in Study Area	
	State	Federal	Nature Map	ALA	EPBC					Initial Ranking Based on Desktop Study	Final Ranking Including Results of Field Survey
<i>Tyto novaehollandiae kimberli</i> Masked Owl (northern)	-	Vulnerable	-	-	✓	Forests, woodlands, timbered waterways and open country on the fringe of these areas. Requires tall trees with suitable hollows for nesting and roosting and adjacent areas for foraging.	No	No records on Dampier Peninsula.	Nearest record over 200 km NE.	Unlikely to occur.	Unlikely to occur
<i>Charadrius veredus</i> Oriental Plover	Migratory	Migratory/Marine	-	-	✓	Short grasslands (natural and man-made) and sparsely vegetated plains, margins of wetlands. Less commonly tidal mudflats.	No	50 km SSW of Lombadina	One record in 2002	May occur.	Unlikely to occur
<i>Numenius minutus</i> Little Curlew	Migratory	Migratory/Marine	✓	✓	-	Coastal and inland grasslands and black soil plains in northern Australia, near swamps and flooded areas. Also feed on playing fields, paddocks and urban lawns.	Yes	2 km N of Lombadina.	Single record from 2018.	Likely to occur.	Unlikely to occur.
<i>Gelochelidon macrotarsa</i> Australian [Gull-billed] Tern <i>Gelochelidon nilotica</i> [Common] Gull-billed Tern ¹	Migratory	Migratory/Marine	✓	✓	-	Estuaries and coasts (both species), inland wetlands and grasslands (primarily Australian).	No	Four records within 3 km to the west of Lombadina in past 5 years	Records involving up to 10 individuals, both Common and Australian	May occur.	Unlikely to occur (likely to overfly).
<i>Chlidonias leucopterus</i> White-winged Tern	Migratory	Migratory/Marine	-	✓	-	Freshwater wetlands, grasslands (esp. if flooded), estuaries, and coastal seas	No	<1 km NE of Lombadina	Single bird sighted at wastewater treatment ponds in 2018.	Unlikely to occur.	Unlikely to occur (likely to overfly).
<i>Plegadis falcinellus</i> Glossy Ibis	Migratory	Migratory/Marine	-	✓	-	Freshwater wetlands and floodplains	No	Recorded in Lombadina	Single bird in 2018, precise location not indicated.	Unlikely to occur.	Unlikely to occur.
<i>Pandion cristatus</i> Eastern Osprey	Migratory	Migratory/Marine	✓	✓	-	Estuaries, coasts and offshore islands, occasionally freshwater wetlands or inland along major rivers.	No	<1 km NE of Lombadina	Single bird sighted at wastewater treatment ponds in 2016.	May occur.	Unlikely to occur (likely to overfly).
<i>Cecropis daurica</i> Red-rumped Swallow	Migratory	Migratory/Marine	-	-	✓	Rare migrant to Western Australia. Wide variety of habitats, but generally preferring open country, often near water.	Yes	Nearest records from Broome.	-	Unlikely to occur.	Unlikely to occur.
<i>Motacilla tschutschensis</i> Eastern Yellow Wagtail	Migratory	Migratory/Marine	-	-	✓	Open country, particularly well-watered grasslands or wetlands margins.	No	No records within 40 km of Lombadina	-	May occur.	Unlikely to occur.
<i>Motacilla cinerea</i> Grey Wagtail	Migratory	Migratory/Marine	-	-	✓	Rare migrant to Western Australia. Close to water, particularly fast flowing waterways. Beaches and rockpools often used on migration.	No	No records within 40 km of Lombadina	-	May occur.	Unlikely to occur.
<i>Pezoporus occidentalis</i> Night Parrot		Endangered	-	-	✓	Old growth spinifex, esp. ring-forming spinifex for roosting and nesting, samphire or higher productivity grasslands for foraging.	No	No records on Dampier Peninsula.	-	Unlikely to occur.	Would not occur.

Appendix 7

Vegetation Structural Classes and Condition Scale



Vegetation Structural Classes*

Stratum	Canopy Cover (%)				
	70-100%	30-70%	10-30%	2-10%	<2%
Trees over 30 m	Tall closed forest	Tall open forest	Tall woodland	Tall open woodland	Scattered tall trees
Trees 10-30 m	Closed forest	Open forest	Woodland	Open woodland	Scattered trees
Trees under 10 m	Low closed forest	Low open forest	Low woodland	Low open woodland	Scattered low trees
Shrubs over 2 m	Tall closed scrub	Tall open scrub	Tall shrubland	Tall open shrubland	Scattered tall shrubs
Shrubs 1-2 m	Closed heath	Open heath	Shrubland	Open shrubland	Scattered shrubs
Shrubs under 1 m	Low closed heath	Low open heath	Low shrubland	Low open shrubland	Scattered low shrubs
Hummock grasses	Closed hummock grassland	Hummock grassland	Open hummock grassland	Very open hummock grassland	Scattered hummock grasses
Grasses, Sedges, Herbs	Closed tussock grassland / bunch grassland / sedgeland / herbland	Tussock grassland / bunch grassland / sedgeland / herbland	Open tussock grassland / bunch grassland / sedgeland / herbland	Very open tussock grassland / bunch grassland / sedgeland / herbland	Scattered tussock grasses / bunch grasses / sedges / herbs

* Based on Muir (1977) , and Aplin's (1979) modification of the vegetation classification system of Specht (1970); Aplin T.E.H. (1979). The Flora. Chapter 3 In O'Brien, B.J. (ed.) (1979). *Environment and Science*. University of Western Australia Press; Muir B.G. (1977). Biological Survey of the Western Australian Wheatbelt. Part II: Vegetation and habitat of Bending Reserve. *Records of the Western Australian Museum*, Suppl. No. 3; Specht R.L. (1970). Vegetation. In: *The Australian Environment*. 4th edn (Ed. G.W. Leeper). Melbourne.

Vegetation Condition Scale for use on flora surveys*

Vegetation Condition	South West and Interzone Botanical Provinces	Eremaean and Northern Botanical Provinces
Pristine	Pristine or nearly so, no obvious signs of disturbance or damage caused by human activities since European settlement.	
Excellent	Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species. Damage to trees caused by fire, the presence of non-aggressive weeds and occasional vehicle tracks.	Pristine or nearly so, no obvious signs of damage caused by human activities since European settlement.
Very Good	Vegetation structure altered, obvious signs of disturbance. Disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.	Some relatively slight signs of damage caused by human activities since European settlement. For example, some signs of damage to tree trunks caused by repeated fire, the presence of some relatively non-aggressive weeds, or occasional vehicle tracks.
Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.	More obvious signs of damage caused by human activity since European settlement, including some obvious impact on the vegetation structure such as that caused by low levels of grazing or slightly aggressive weeds.
Poor		Still retains basic vegetation structure or ability to regenerate it after very obvious impacts of human activities since European settlement, such as grazing, partial clearing, frequent fires or aggressive weeds.
Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. Disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds at high density, partial clearing, dieback and grazing.	Severely impacted by grazing, very frequent fires, clearing or a combination of these activities. Scope for some regeneration but not to a state approaching good condition without intensive management. Usually with a number of weed species present including very aggressive species.
Completely Degraded	The structure of the vegetation is no longer intact and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees and shrubs.	Areas that are completely or almost completely without native species in the structure of their vegetation; i.e. areas that are cleared or 'parkland cleared' with their flora comprising weed or crop species with isolated native trees or shrubs.


*Taken from EPA (2016c). Adapted from:
 -Keighery, B.J. (1994). Bushland Plant Survey: a Guide to Plant Community Survey for the Community Wildflower Society of WA (Inc.), Nedlands, Western Australia; and
 -Trudgen M.E. (1988). A Report on the Flora and Vegetation of the Port Kennedy Area. Unpublished report prepared for Bowman Bishaw and Associates, West Perth.

Appendix 8




Survey Effort Map





- Study area
- Foot traverse
- Mapping note



Lombadina Campground Survey Effort



Appendix 9

Vascular Flora Species List



Family	Species	Local Aboriginal (Bardi) Name	Comments
Acanthaceae	<i>Nelsonia campestris</i>		
Aizoaceae	<i>Trianthema pilosum</i>		
Amaranthaceae	* <i>Aerva javanica</i>		Weed
	* <i>Gomphrena celosioides</i>		Weed
Anacardiaceae	* <i>Mangifera indica</i>		Planted; Mango
Apocynaceae	* <i>Catharanthus roseus</i>		Planted; Vinca
	* <i>Plumeria rubra</i>		Planted; Red Frangipani
	<i>Vincetoxicum cinerascens</i>		
Arecaceae	* <i>Cocos nucifera</i>		Planted; Coconut
Asteraceae	* <i>Cyanthillium cinereum</i>		Weed
	* <i>Tridax procumbens</i>		Weed
Boraginaceae	<i>Ehretia saligna</i> var. <i>saligna</i>		
	<i>Heliotropium glabellum</i>		
Bromeliaceae	* <i>Ananas comosus</i>		Planted; Pineapple
Burseraceae	<i>Canarium australianum</i>	Jalgir	
Cleomaceae	<i>Arivela tetrandra</i>		
	<i>Arivela viscosa</i>	Boolorrboolorr	
Combretaceae	<i>Terminalia ferdinandiana</i>	Madoor	
Convolvulaceae	<i>Bonamia linearis</i>		
	* <i>Distimake dissectus</i> var. <i>dissectus</i>		Weed
	<i>Evolvulus alsinoides</i>		
	* <i>Ipomoea batatas</i>		Planted; Sweet Potato
	<i>Ipomoea brasiliensis</i>		
	<i>Ipomoea polymorpha</i>		
Cucurbitaceae	<i>Cucumis argenteus</i>		
Cyperaceae	<i>Bulbostylis barbata</i>		
Ebenaceae	<i>Diospyros humilis</i>	Birimbiri	
Euphorbiaceae	<i>Euphorbia drummondii</i>		
	* <i>Jatropha gossypifolia</i>		Weed; WoNS
Fabaceae	<i>Acacia tumida</i>	Wanggay	
	<i>Bauhinia cunninghamii</i>	Jigal	

Family	Species	Local Aboriginal (Bardi) Name	Comments
Fabaceae (cont.)	<i>Crotalaria brevis</i>		
	<i>Crotalaria medicaginea</i>		
	<i>Glycine tomentella</i>		
	<i>Indigofera hirsuta</i>		
	<i>Indigofera linifolia</i>		
	* <i>Macroptilium atropurpureum</i>		Weed
	<i>Rhynchosia minima</i>		
	* <i>Stylosanthes hamata</i>		Weed
	* <i>Stylosanthes scabra</i>		Weed
	<i>Tephrosia leptoclada</i>		
	<i>Zornia chaetophora</i>		
<i>Zornia prostrata</i>			
Goodeniaceae	<i>Goodenia sepalosa</i>		
Gyrostemonaceae	<i>Gyrostemon tepperi</i>		
Hernandiaceae	<i>Gyrocarpus americanus</i>	Bilanggamarr	
Lamiaceae	* <i>Mesosphaerum suaveolens</i>		Weed
	* <i>Ocimum basilicum</i>		Planted; Sweet Basil
	<i>Premna acuminata</i>		
Lecythidaceae	<i>Planchonia careya</i>	Goolay	
Loranthaceae	<i>Amyema sanguinea</i>		
	<i>Lysiana spathulata</i>		
Malvaceae	<i>Abutilon indicum</i>		
	<i>Brachychiton diversifolius</i>	Gorrgorr	
	<i>Corchorus aestuans</i>		
	<i>Corchorus pumilio</i>		
	* <i>Hibiscus sabdariffa</i>		Planted; Rosella
	<i>Sida hackettiana</i>		
	<i>Sida rohlenae</i>		
	<i>Triumfetta breviaculeata</i>		
<i>Waltheria indica</i>			
Meliaceae	* <i>Azadirachta indica</i>		Weed
	* <i>Khaya senegalensis</i>		Planted
	<i>Owenia reticulata</i>	Lambilamb	

Family	Species	Local Aboriginal (Bardi) Name	Comments
Menispermaceae	<i>Tinospora smilacina</i>	Oondal	
Moraceae	<i>Ficus aculeata</i>	Ranyji	
	<i>Ficus opposita</i>		
Myrtaceae	<i>Corymbia bella</i>	Maroolal	
	<i>Corymbia greeniana</i>	Biilal	
	<i>Corymbia polycarpa</i>	Gaardga	
	<i>Eucalyptus miniata</i>	Manowan	
Nyctaginaceae	<i>Boerhavia gardneri</i>		
Passifloraceae	<i>*Passiflora foetida</i> var. <i>hispida</i>		Weed
Poaceae	<i>*Bothriochloa pertusa</i>		Planted
	<i>*Cenchrus biflorus</i>		Weed
	<i>*Cenchrus setiger</i>		Weed
	<i>*Chloris barbata</i>		Weed
	<i>Chrysopogon pallidus</i>	Irrooloo	
	<i>Dactyloctenium radulans</i>		
	<i>Digitaria bicornis</i>		
	<i>Eragrostis eriopoda</i>		
	<i>Sorghum ecarinatum</i>	Oonbi	
Portulacaceae	<i>Calandrinia strophilata</i>	Gumbin	
	<i>Portulaca napiformis</i>	Ngoorrarr	
Proteaceae	<i>Persoonia falcata</i>	Gamooloon	
Rubiaceae	<i>Gardenia pyriformis</i>		
	<i>Spermacoce occidentalis</i>		
Santalaceae	<i>Santalum lanceolatum</i>	Bilooloor	
Solanaceae	<i>Solanum cunninghamii</i>		
Verbenaceae	<i>*Phyla nodiflora</i>		Planted
Violaceae	<i>Afrohybanthus enneaspermus</i>		
Zygophyllaceae	<i>Tribulopsis angustifolia</i>		

Appendix 10

Potential Fauna Species List



Notes on Potential Fauna Species List

EPBC Act Marine Listed Birds

The EPBC Act maintains a list of fauna species recognised as Matters of National Environmental Significance that are protected under the Act. Marine species are included under the Act, including bird species that rely upon the marine environment for survival.

However, the Act also erroneously lists some species that do not actually rely upon the marine environment for survival, but instead rely upon land environments and in some cases, are often widespread and common (Garnett 2013). Eighty-eight Marine listed bird species were returned from the desktop review, of which 39 are listed solely as Marine. Many of these fall into the category of erroneous listings; that is, species that do not rely upon marine environments for survival, and all are relatively common and widespread. Given this, and that the survey area does not encompass marine environs or seabird breeding colonies, marine listed species have not been considered further in this assessment.

EPCA Act Migratory Listed Birds

Migratory species are also protected under the EPBC Act as Matters of National Environmental Significance. Fifty-two species listed as Migratory were returned from the desktop study. For clarity and conciseness, species that have a strong preference for coastal habitats or freshwater wetlands, which are available in close proximity to the study area, but are not available in the study area have not been included in the likelihood of occurrence assessment. These species may utilise adjacent habitats, and potentially may overfly or occur in proximity to the study area on occasion, but are considered very unlikely to utilise habitats within the study area and are not considered further here for the purpose of this assessment.

Reptiles

Family	Species Name	Common Name	Conservation Status			NatureMap	ALA	EPBC
			State	Federal				
Diplodactylidae	<i>Amalosia rhombifer</i>		-	-	Y	Y		
	<i>Oedura gracilis</i>		-	-	Y	Y		
	<i>Strophurus ciliaris aberrans</i>		-	-	Y	Y		
Gekkonidae	<i>Gehyra gemina</i> ¹		-	-	Y	Y		
	<i>Gehyra kimberleyi</i> ²		-	-	Y	Y		
	<i>Gehyra nana</i>		-	-	Y	Y		
	<i>Gehyra occidentalis</i>		-	-	Y	Y		
	<i>Gehyra pilbara</i>		-	-	Y	Y		
	<i>Heteronotia binoei</i>	Bynoe's Gecko	-	-	Y	Y		
Pygopodidae	<i>Delma borea</i>		-	-	Y	Y		
	<i>Delma tincta</i>		-	-	Y	Y		
	<i>Lialis burtonis</i>		-	-	Y	Y		
Agamidae	<i>Chelosania brunnea</i>	Chameleon Dragon	-	-	Y	Y		
	<i>Chlamydosaurus kingii</i>	Frill-necked Lizard	-	-	Y	Y		
	<i>Diporiphora pindan</i>	Pindan Dragon	-	-	Y	Y		
	<i>Lophognathus horneri</i> ³		-	-	Y	Y		
	<i>Pogona minor mitchelli</i>		-	-	Y	Y		
Scincidae	<i>Carlia amax</i>		-	-		Y		
	<i>Carlia munda</i>		-	-	Y	Y		
	<i>Carlia triacantha</i>		-	-	Y	Y		
	<i>Cryptoblepharus tytthos</i>		-	-	Y	Y		
	<i>Ctenotus colletti</i>		-	-	Y	Y		
	<i>Ctenotus helenae</i>		-	-		Y		
	<i>Ctenotus inornatus</i> ⁴		-	-	Y	Y		
	<i>Eremiascincus isolepis</i>		-	-	Y	Y		
	<i>Lerista apoda</i>		-	-	Y	Y		
	<i>Lerista bipes</i>		-	-	Y	Y		
	<i>Lerista griffini</i>		-	-	Y	Y		
	<i>Lerista separanda</i>		P2	-	Y	Y		
	<i>Menetia maini</i>		-	-	Y	Y		
	<i>Morethia storri</i>		-	-	Y	Y		
	<i>Notoscincus ornatus wotjulum</i>		-	-	Y	Y		
	<i>Tiliqua multifasciata</i>	Central Blue-tongue	-	-	Y	Y		
	<i>Tiliqua scincoides intermedia</i>		-	-		Y		
Varanidae	<i>Varanus glauerti</i>	Kimberley Rock Goanna	-	-	Y	Y		
	<i>Varanus glebopalma</i>	Black-palmed Rock Goanna	-	-	Y	Y		
	<i>Varanus gouldii</i>	Bungarra or Sand Goanna	-	-	Y	Y		
	<i>Varanus scalaris</i>	Spotted Tree Goanna	-	-	Y	Y		
	<i>Varanus tristis</i>	Racehorse Goanna	-	-	Y	Y		
Typhlopidae	<i>Anilius diversus</i>		-	-		Y		
	<i>Indotyphlops braminus</i>	Flowerpot Blind Snake	-	-			Y	
Pythonidae	<i>Antaresia childreni</i>	Children's Python	-	-	Y	Y		
	<i>Antaresia stimsoni stimsoni</i>		-	-	Y	Y		
	<i>Liasis olivaceus olivaceus</i>		-	-	Y	Y		

Family	Species Name	Common Name	Conservation Status			NatureMap	ALA	EPBC
			State	Federal				
Colubridae	<i>Dendrelaphis punctulatus</i>	Green Tree Snake	-	-	Y	Y		
Homalopsidae	<i>Fordonia leucobalia</i>	White-bellied Mangrove Snake	-	-	Y	Y		
Elapidae	<i>Demansia angusticeps</i>		-	-	Y	Y		
	<i>Demansia papuensis</i>	Great Black Whipsnake	-	-	Y	Y		
	<i>Furina ornata</i>	Moon Snake	-	-	Y	Y		
	<i>Pseudechis australis</i>	Mulga Snake	-	-	Y	Y		
	<i>Pseudonaja mengdeni</i>	Western Brown Snake	-	-	Y	Y		
	<i>Simoselaps minimus</i>		P2	-	Y	Y		

P2 = Priority 2

¹ Formerly included within *G. australis*, previous records listed as such

² Previous records of *G. punctata* and *G. variegata*, and possibly some records of *G. pilbara*, likely referable to this recently-described species

³ Formerly included within *L. gilberti* or *Amphibolurus gilberti*, previous records may be listed as such

⁴ previous records of *C. helenae* likely referable to this species, though taxonomy is not settled

Amphibians

Family	Species	Common	Conservation Status			NatureMap	ALA	EPBC
			State	Federal				
Pelodyadidae	<i>Cyclorana australis</i>	Giant Frog	-	-	Y	Y		
	<i>Litoria caerulea</i>	Green Tree Frog	-	-	Y	Y		
	<i>Litoria rothii</i>	Northern Laughing Tree Frog	-	-	Y	Y		
Limnodynastidae	<i>Platyplectrum ornatum</i>	Ornate Burrowing Frog	-	-	Y	Y		
Bufonidae	<i>Rhinella marina</i>	Cane Toad	-	-			Y	

Mammals

Family	Species	Common	Conservation Status			NatureMap	ALA	EPBC
			State	Federal				
Dasyuridae	<i>Dasyurus hallucatus</i>	Northern Quoll	EN	Endangered		Y		
	<i>Planigale maculata</i>	Common Planigale	-	-	Y	Y		
	<i>Pseudantechinus ningbing</i>	Ningbing Pseudantechinus	-	-	Y	Y		
	<i>Sminthopsis macroura frogattii</i>	Frogatt's Stripe-faced Dunnart	-	-		Y		
Thylacomyidae	<i>Macrotis lagotis</i>	Bilby, Dalgyte	VU	Vulnerable	Y		Y	
Macropodidae	<i>Onychogalea unguifera</i>	Northern Nailtail Wallaby, Karrabul	-	-		Y		
Muridae	<i>Melomys burtoni</i>	Grassland Melomys	-	-	Y	Y		
	<i>Pseudomys delicatulus</i>	Delicate Mouse	-	-	Y	Y		
	<i>Rattus exulans</i>	Pacific Rat	-	-		Y		
	<i>Rattus rattus</i>	Black Rat	-	-	Y	Y		
Pteropodidae	<i>Macroglossus minimus</i>	Northern Blossom-bat	-	-	Y	Y		
	<i>Pteropus alecto</i>	Black Flying-fox	-	-	Y	Y		
	<i>Pteropus scapulatus</i>	Little Red Flying-fox	-	-	Y	Y		
Macrodermatidae	<i>Macroderma gigas</i>	Ghost Bat	VU	Vulnerable			Y	
Emballonuridae	<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tailed Bat	-	-	Y	Y		
	<i>Taphozous georgianus</i>	Common Sheath-tailed Bat	-	-	Y	Y		
Molossidae	<i>Ozimops cobourgianus</i> ¹	Northern Coastal Free-tailed Bat	P1		Y	Y		

Family	Species	Common	Conservation Status				
			State	Federal	NatureMap	ALA	EPBC
Miniopteridae	<i>Miniopterus orianae orianae</i>	Large Bent-winged Bat	-	-	Y		
Vespertilionidae	<i>Chalinolobus gouldii</i>	Gould's Wattled Bat	-	-	Y	Y	
	<i>Chalinolobus nigrogriseus</i>	Hoary Wattled Bat	-	-	Y	Y	
	<i>Nyctophilus arnhemensis</i>	Arnhem Long-eared Bat	-	-	Y	Y	
	<i>Nyctophilus daedalus</i>	Pallid Long-eared Bat	-	-	Y	Y	
	<i>Pipistrellus westralis</i>	Northern Pipistrelle	-	-	Y	Y	
	<i>Scotorepens greyii</i>	Little Broad-nosed Bat	-	-	Y	Y	
	<i>Scotorepens sanborni</i>	Northern Broad-nosed Bat	-	-	Y	Y	
	<i>Vespadelus caurinus</i>	Northern Cave-bat	-	-	Y	Y	
Suidae	<i>Sus scrofa</i>	Pig	-	-	Y	Y	
Bovidae	<i>Bos taurus</i>	European Cattle	-	-	Y		
	<i>Capra hircus</i>	Goat	-	-	Y		

EN = Endangered, VU = Vulnerable, P1 = Priority 1

¹ Formerly included within *Mormopterus loriae* as *Mormopterus loriae cobourgianus*

Avifauna

Family	Species	Common	Conservation Status					Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal	Nature-Map	ALA	EPBC				
Phasianidae	<i>Coturnix ypsilophora</i>	Brown Quail			Y	Y					
Anatidae	<i>Dendrocygna arcuata</i>	Wandering Whistling Duck		MA		Y				Not marine	
	<i>Nettapus pulchellus</i>	Green Pygmy Goose		MA		Y				Not marine	
	<i>Anas superciliosa</i>	Pacific Black Duck			Y	Y					
	<i>Anas gracilis</i>	Grey Teal				Y					
	<i>Aythya australis</i>	Hardhead			Y	Y					
Podargidae	<i>Podargus strigoides</i>	Tawny Frogmouth			Y	Y					
Caprimulgidae	<i>Eurostopodus argus</i>	Spotted Nightjar		MA	Y	Y				Not marine	
Aegothelidae	<i>Aegotheles cristatus</i>	Australian Owlet-nightjar			Y	Y					
Apodidae	<i>Apus pacificus</i>	Pacific Swift	MI	MI; MA	Y	Y		Almost exclusively aerial	✓		
Otididae	<i>Ardeotis australis</i>	Australian Bustard			Y	Y					
Cuculidae	<i>Centropus phasianinus</i>	Pheasant Coucal			Y	Y					
Cuculidae	<i>Scythrops novaehollandiae</i>	Channel-billed Cuckoo		MA	Y	Y				Not marine	
Cuculidae	<i>Chrysococcyx basalis</i>	Horsfield's Bronze Cuckoo				Y					
Cuculidae	<i>Chrysococcyx osculans</i>	Black-eared Cuckoo		MA		Y				Not marine	
Cuculidae	<i>Chrysococcyx minutillus</i>	Little Bronze Cuckoo			Y	Y					
Cuculidae	<i>Cacomantis pallidus</i>	Pallid Cuckoo		MA	Y	Y				Not marine	
Cuculidae	<i>Cacomantis variolosus</i>	Brush Cuckoo			Y	Y					
Cuculidae	<i>Cuculus optatus</i>	Oriental Cuckoo	MI	MI; MA		Y		Dense open woodlands and forests, riparian forests, sometimes gardens	✓		
Columbidae	<i>Chalcophaps longirostris</i>	Pacific Emerald Dove			Y	Y					

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Columbidae	<i>Phaps chalcoptera</i>	Common Bronzewing			Y							
Columbidae	<i>Ocyphaps lophotes</i>	Crested Pigeon				Y						
Columbidae	<i>Petrophassa albipennis</i>	White-quilled Rock Pigeon				Y						
Columbidae	<i>Geopelia cuneata</i>	Diamond Dove			Y	Y						
Columbidae	<i>Geopelia placida</i>	Peaceful Dove			Y	Y						
Columbidae	<i>Geopelia humeralis</i>	Bar-shouldered Dove			Y	Y						
Columbidae	<i>Ptilinopus regina</i>	Rose-crowned Fruit Dove			Y	Y						
Rallidae	<i>Gallirallus philippensis</i>	Buff-banded Rail			Y	Y						
Rallidae	<i>Porzana tabuensis</i>	Spotless Crake		MA	Y	Y					Not marine	
Rallidae	<i>Eulabeornis castaneoventris</i>	Chestnut Rail			Y	Y						
Rallidae	<i>Porphyrio melanotus</i>	Australasian Swamphen		MA		Y					Not marine	
Rallidae	<i>Fulica atra</i>	Eurasian Coot				Y						
Gruidae	<i>Antigone rubicunda</i>	Brolga			Y	Y						
Podicipedidae	<i>Tachybaptus novaehollandiae</i>	Australasian Grebe			Y	Y						
Podicipedidae	<i>Poliiocephalus poliocephalus</i>	Hoary-headed Grebe				Y						
Turnicidae	<i>Turnix maculosus</i>	Red-backed Buttonquail				Y						
Turnicidae	<i>Turnix castanotus</i>	Chestnut-backed Buttonquail				Y						
Turnicidae	<i>Turnix velox</i>	Little Buttonquail			Y	Y						
Burhinidae	<i>Burhinus grallarius</i>	Bush Stone-curlew			Y	Y						
Burhinidae	<i>Esacus magnirostris</i>	Beach Stone-curlew			Y	Y						
Haematopodidae	<i>Haematopus longirostris</i>	Pied Oystercatcher			Y	Y						
Haematopodidae	<i>Haematopus fuliginosus</i>	Sooty Oystercatcher			Y	Y						
Recurvirostridae	<i>Himantopus leucocephalus</i>	Pied Stilt		MA		Y					Not marine	
Charadriidae	<i>Vanellus miles</i>	Masked Lapwing				Y						
Charadriidae	<i>Erythrogonyx cinctus</i>	Red-kneed Dotterel				Y						
Charadriidae	<i>Pluvialis fulva</i>	Pacific Golden Plover	MI	MI; MA	Y	Y		Muddy, rocky and sandy wetlands, shores, paddocks, saltmarsh, coastal golf courses, estuaries and lagoons	X	✓		
Charadriidae	<i>Pluvialis squatarola</i>	Grey Plover	MI	MI; MA	Y	Y		Almost entirely coastal, being found mainly on marine shores, inlets, estuaries and lagoons with large tidal mudflats or sandflats for feeding, sandy beaches for roosting, and also on rocky coasts	X	✓		
Charadriidae	<i>Charadrius ruficapillus</i>	Red-capped Plover		MA	Y	Y					Not marine	
Charadriidae	<i>Charadrius mongolus</i>	Lesser Sand Plover	EN; MI	EN; MI; MA	Y	Y		Tidal mud flats	X	✓		

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Charadriidae	<i>Charadrius leschenaultii</i>	Greater Sand Plover	VU; MI	VU; MI; MA	Y	Y		Tidal sandy flats, roosts on beaches at high tide	X	✓		
Charadriidae	<i>Charadrius veredus</i>	Oriental Plover	MI	MI; MA			Y	Generally inland; in open grasslands in arid and semi-arid zones; and less often in estuarine or littoral environments	X	X		
Charadriidae	<i>Eseyornis melanops</i>	Black-fronted Dotterel			Y	Y						
Rostratulidae	<i>Rostratula australis</i>	Australian Painted Snipe		EN			Y	Shallow, brackish or freshwater terrestrial wetlands	X	X		
Scolopacidae	<i>Numenius phaeopus</i>	Eurasian Whimbrel	MI	MI; MA	Y	Y		Coast, on tidal and estuarine mudflats, especially near mangroves, beaches and rocky shores.	X	✓		
Scolopacidae	<i>Numenius minutus</i>	Little Curlew	MI	MI; MA	Y	Y		Coastal and inland grasslands and black soil plains in northern Australia, near swamps and flooded areas. They also feed on playing fields, paddocks and urban lawns	✓			
Scolopacidae	<i>Numenius madagascariensis</i>	Far Eastern Curlew	CR; MI	CR; MI; MA	Y	Y	Y	Intertidal mudflats and sandflats, often with beds of seagrass, on sheltered coasts, especially estuaries, mangrove swamps, bays, harbours and lagoons	X	✓		
Scolopacidae	<i>Limosa lapponica</i>	Bar-tailed Godwit	CR (ssp. menzbieri)/VU (ssp. baueri); MI	EN (ssp. menzbieri)/VU (ssp. baueri); MI; MA	Y	Y	Y	Estuarine mudflats, beaches and mangroves	X	✓		
Scolopacidae	<i>Limosa limosa</i>	Black-tailed Godwit	MI	MI; MA	Y	Y		Shallow inland wetlands	X	X		
Scolopacidae	<i>Arenaria interpres</i>	Ruddy Turnstone	MI	MI; MA	Y	Y		Exposed rocks or reefs, often with shallow pools, and on beaches, mudflats	X	✓		
Scolopacidae	<i>Calidris tenuirostris</i>	Great Knot	CR; MI	CR; MI; MA	Y	Y		Intertidal mudflats and sandflats in sheltered coasts, including bays harbours and estuaries. They forage on the moist mud, and they often roost on beaches or in nearby low vegetation, such as mangroves or dune vegetation	X	✓		
Scolopacidae	<i>Calidris canutus</i>	Red Knot	EN; MI	EN; MI; MA	Y	Y	Y	Coast in sandy estuaries with tidal mudflats	X	✓		
Scolopacidae	<i>Calidris acuminata</i>	Sharp-tailed Sandpiper	MI	Migratory/ Marine	Y	Y		Grassy edges of shallow inland freshwater wetlands. It is also found around sewage farms, flooded fields, mudflats, mangroves, rocky shores and beaches.	X	✓		
Scolopacidae	<i>Calidris ferruginea</i>	Curlew Sandpiper	CR; MI	CR; MI; MA	Y	Y	Y	Intertidal mudflats of estuaries, lagoons, mangroves, as well as beaches, rocky shores and around lakes, dams and floodwaters				
Scolopacidae	<i>Calidris subminuta</i>	Long-toed Stint	MI	MI; MA		Y		Muddy fringes of fresh wetlands	X	X		
Scolopacidae	<i>Calidris ruficollis</i>	Red-necked Stint	MI	MI; MA	Y	Y		Coastal, in sheltered inlets, bays, lagoons, estuaries, intertidal mudflats and protected sandy or coralline shores. They may also be seen in saltworks, sewage farms, saltmarsh, shallow wetlands including lakes, swamps, riverbanks, waterholes, bore drains, dams, soaks and pools in saltflats, flooded paddocks or damp grasslands	X	✓		

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Scolopacidae	<i>Calidris alba</i>	Sanderling	MI	MI; MA	Y	Y		Open sandy beaches at the edge of the waves, on sandbars and spits. They roost on bare sand in the dunes or behind piles of kelp	X	✓		
Scolopacidae	<i>Calidris melanotos</i>	Pectoral Sandpiper	MI	MI; MA			Y	Shallow, grassy edges of of freshwater wetlands	X	X		
Scolopacidae	<i>Xenus cinereus</i>	Terek Sandpiper	MI	MI; MA	Y	Y		Coast in mangrove swamps, tidal mudflats and the seashore	X	✓		
Scolopacidae	<i>Actitis hypoleucos</i>	Common Sandpiper	MI	MI; MA	Y	Y		Coastal or inland wetlands with muddy edges or rocky shores	X	X		
Scolopacidae	<i>Tringa brevipes</i>	Grey-tailed Tattler	MI; P4	MI	Y	Y		Sheltered coasts with reefs and rock platforms or with intertidal mudflats, intertidal rocky, coral or stony reefs, platforms and islets that are exposed at high tide, also shores of rock, shingle, gravel and shells and on intertidal mudflats in embayments, estuaries and coastal lagoons, especially those fringed with mangroves	X	✓		
Scolopacidae	<i>Tringa totanus</i>	Common Redshank	MI	MI; MA	Y	Y		Coastal, sheltered tidal flats	X	✓		
Scolopacidae	<i>Tringa stagnatilis</i>	Marsh Sandpiper	MI	MI; MA		Y		Fresh or brackish (slightly salty) wetlands such as rivers, water meadows, sewage farms, drains, lagoons and swamps	X	X		
Scolopacidae	<i>Tringa glareola</i>	Wood Sandpiper	MI	MI; MA	Y	Y		Inland shallow freshwater wetlands, often with other waders. They prefer ponds and pools with emergent reeds and grass, surrounded by tall plants or dead trees and fallen timber	X	X		
Scolopacidae	<i>Tringa nebularia</i>	Common Greenshank	MI	MI; MA	Y	Y		Coast and inland, in estuaries and mudflats, mangrove swamps and lagoons, and in billabongs, swamps, sewage farms and flooded crops	X	✓		
Glareolidae	<i>Stiltia isabella</i>	Australian Pratincole		MA		Y					Not marine	
Glareolidae	<i>Glareola maldivarum</i>	Oriental Pratincole	MI	MI			Y	Mostly aerial, above open country	✓			
Laridae	<i>Anous stolidus</i>	Brown Noddy	MI	MI; MA	Y	Y		Off-shore tropical islands	X	X		
Laridae	<i>Chroicocephalus novaehollandiae</i>	Silver Gull		MA	Y	Y	Y				Not marine	
Laridae	<i>Gelochelidon nilotica</i>	[Common] Gull-billed Tern ¹	MI	MI; MA	Y	Y		Primarily beaches and estuarine mudflats, much less common inland or on freshwater than Australian Tern	X	✓		
Laridae	<i>Gelochelidon macrotarsa</i>	Australian [Gull-billed] Tern ¹	MI	MI; MA	Y	Y		Freshwater swamps, brackish and salt lakes, beaches and estuarine mudflats, floodwaters, sewage farms, irrigated croplands and grasslands				
Laridae	<i>Hydroprogne caspia</i>	Caspian Tern	MI	MI	Y	Y		Sheltered coastal waters, inland water bodies including large rivers, fresh to saline lakes, wetlands	X	✓		
Laridae	<i>Thalasseus bergii</i>	Greater Crested Tern	MI	MI; MA	Y	Y		Forages over coastal seas, roosts on sandy beaches, rocks and man-made structures	X	✓		

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Laridae	<i>Thalasseus bengalensis</i>	Lesser Crested Tern		MA	Y	Y		Breeds in colonies on small offshore islands. Usually a strictly coastal species, there are occasional records in the arid interior of Australia, where birds were possibly blown by passing tropical cyclones	X	✓		
Laridae	<i>Sternula albifrons</i>	Little Tern	MI	MI; MA	Y	Y		Mainly coastal, being found on beaches, sheltered inlets, estuaries, lakes, sewage farms, lagoons, river mouths and deltas	X	✓		
Laridae	<i>Onychoprion anaethetus</i>	Bridled Tern	MI	MI	Y	Y		Forages at sea, breeds on offshore islands	X	X		
Laridae	<i>Onychoprion fuscatus</i>	Sooty Tern		MA	Y	Y		Pelagic, nests on sandy or rocky offshore islands	X	X	Marine	
Laridae	<i>Sterna dougallii</i>	Roseate Tern	MI	MI; MA	Y	Y		Restricted to tropical and subtropical seas and coastlines, coral reefs and vegetated islands	X	X		
Laridae	<i>Sterna hirundo</i>	Common Tern	MI	MI; MA	Y	Y		Coastal when not breeding and found in offshore waters, ocean beaches, estuaries and large lakes. Common Terns are occasionally seen in freshwater swamps, floodwaters, sewage farms and brackish and saline lakes	X	✓		
Laridae	<i>Chlidonias hybrida</i>	Whiskered Tern		MA		Y						
Laridae	<i>Chlidonias leucopterus</i>	White-winged Tern	MI	MI; MA		Y		Coastal or sub-coastal wetlands including tidal estuaries, lagoons, grassy swamps, and sewage ponds	X	X		
Phaethontidae	<i>Phaethon rubricauda</i>	Red-tailed Tropicbird	MI; P4	MI; MA	Y	Y		Pelagic waters, rarely seen in sight of land	X	X		
Oceanitidae	<i>Oceanites oceanicus</i>	Wilson's Storm Petrel	MI	MI; MA		Y		Seabird, usually far offshore	X	X		
Procellariidae	<i>Calonectris leucomelas</i>	Streaked Shearwater	MI	MI; MA			Y	Mostly offshore waters	X	✓		
Procellariidae	<i>Puffinus assimilis</i>	Little Shearwater		MA		Y		Offshore islands mainly in the south of WA	X	X	Marine	
Ciconiidae	<i>Ephippiorhynchus asiaticus</i>	Black-necked Stork			Y	Y						
Fregatidae	<i>Fregata minor</i>	Great Frigatebird	MI	MI; MA		Y		Seabird. occasional visitor to mainland coast, but more so during tropical storms/cyclones	X	✓		
Fregatidae	<i>Fregata ariel</i>	Lesser Frigatebird	MI	MI; MA	Y	Y		Seabird. forages at sea and along coastlines, breeds on remote tropical islands	X	✓		
Sulidae	<i>Papasula abbotti</i>	Abbott's Booby		EN			Y	Tropical seas, Christmas Island	X	X		
Sulidae	<i>Sula sula</i>	Red-footed Booby	MI	MI; MA			Y	Pelagic, often found far from land	X	X		
Sulidae	<i>Sula leucogaster</i>	Brown Booby	MI	MI; MA	Y	Y		Rarely occurs on shoreline, sometimes seen on pylons/piers. Breeds on offshore islands	X	X		
Phalacrocoracidae	<i>Microcarbo melanoleucos</i>	Little Pied Cormorant			Y	Y						
Phalacrocoracidae	<i>Phalacrocorax sulcirostris</i>	Little Black Cormorant				Y						
Phalacrocoracidae	<i>Phalacrocorax varius</i>	Australian Pied Cormorant			Y	Y						
Anhingidae	<i>Anhinga novaehollandiae</i>	Australasian Darter			Y	Y						

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Threskiornithidae	<i>Threskiornis molucca</i>	Australian White Ibis		MA		Y					Not marine	
Threskiornithidae	<i>Threskiornis spinicollis</i>	Straw-necked Ibis		MA	Y	Y					Not marine	
Threskiornithidae	<i>Plegadis falcinellus</i>	Glossy Ibis	MI	MI; MA		Y		Well-vegetated wetlands, floodplains, mangroves and ricefields	X	X		
Threskiornithidae	<i>Platalea regia</i>	Royal Spoonbill			Y	Y						
Ardeidae	<i>Nycticorax caledonicus</i>	Nankeen Night Heron		MA	Y	Y					Not marine	
Ardeidae	<i>Butorides striata</i>	Striated Heron			Y	Y						
Ardeidae	<i>Bubulcus coromandus</i>	Eastern Cattle Egret		MA			Y	Grasslands and shallow open freshwater wetlands with low emergent vegetation	X	X		
Ardeidae	<i>Ardea pacifica</i>	White-necked Heron				Y						
Ardeidae	<i>Ardea sumatrana</i>	Great-billed Heron			Y	Y						
Ardeidae	<i>Ardea alba</i>	Great Egret		MA	Y	Y					Not marine	
Ardeidae	<i>Ardea intermedia</i>	Intermediate Egret		MA	Y	Y					Not marine	
Ardeidae	<i>Egretta novaehollandiae</i>	White-faced Heron			Y	Y						
Ardeidae	<i>Egretta garzetta</i>	Little Egret		MA	Y	Y					Not marine	
Ardeidae	<i>Egretta sacra</i>	Pacific Reef Heron		MA	Y	Y					Not marine	
Pelecanidae	<i>Pelecanus conspicillatus</i>	Australian Pelican		MA	Y	Y					Not marine	
Pandionidae	<i>Pandion cristatus</i>	Eastern Osprey	MI	MI; MA	Y	Y		Coast and in terrestrial wetlands of tropical and temperate Australia and off-shore islands, occasionally ranging inland along rivers	X	✓		
Accipitridae	<i>Elanus axillaris</i>	Black-shouldered Kite			Y	Y						
Accipitridae	<i>Lophoictinia isura</i>	Square-tailed Kite				Y						
Accipitridae	<i>Hieraaetus morphnoides</i>	Little Eagle				Y						
Accipitridae	<i>Aquila audax</i>	Wedge-tailed Eagle			Y	Y						
Accipitridae	<i>Erythrotriorchis radiatus</i>	Red Goshawk		VU			Y	Mosaic of vegetation types, often near wetlands. They often occur at the boundary between two vegetation types, and often favour forests or woodlands dominated by eucalypts or paperbarks. They avoid very dense or very open habitats	✓			
Accipitridae	<i>Accipiter fasciatus</i>	Brown Goshawk		MA	Y	Y					Not marine	
Accipitridae	<i>Accipiter cirrocephalus</i>	Collared Sparrowhawk			Y	Y						
Accipitridae	<i>Circus approximans</i>	Swamp Harrier		MA		Y					Not marine	
Accipitridae	<i>Circus assimilis</i>	Spotted Harrier			Y	Y						
Accipitridae	<i>Milvus migrans</i>	Black Kite			Y	Y						
Accipitridae	<i>Haliastur sphenurus</i>	Whistling Kite		MA	Y	Y					Not marine	
Accipitridae	<i>Haliastur indus</i>	Brahminy Kite		MA	Y	Y					Not marine	
Accipitridae	<i>Haliaeetus leucogaster</i>	White-bellied Sea Eagle		MA	Y	Y	Y	Marine, but also utilises inland wetlands				

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Tytonidae	<i>Tyto novaehollandiae kimberli</i>	Masked Owl (northern)		VU				Y	Forests, woodlands, timbered waterways and open country on the fringe of these areas. The main requirements are tall trees with suitable hollows for nesting and roosting and adjacent areas for foraging	✓		
Tytonidae	<i>Tyto javanica</i>	Eastern Barn Owl					Y					
Strigidae	<i>Ninox connivens</i>	Barking Owl			Y	Y						
Strigidae	<i>Ninox boobook</i>	Australian Boobook				Y						
Coraciidae	<i>Eurystomus orientalis</i>	Oriental Dollarbird		MA	Y	Y						Not marine
Alcedinidae	<i>Dacelo leachii</i>	Blue-winged Kookaburra			Y	Y						
Alcedinidae	<i>Todiramphus sordidus</i>	Torresian Kingfisher			Y	Y						
Alcedinidae	<i>Todiramphus sanctus</i>	Sacred Kingfisher		MA	Y	Y						Not marine
Alcedinidae	<i>Todiramphus pyrrhopygius</i>	Red-backed Kingfisher			Y	Y						
Alcedinidae	<i>Ceyx azureus</i>	Azure Kingfisher			Y	Y						
Meropidae	<i>Merops ornatus</i>	Rainbow Bee-eater		MA	Y	Y						Not marine
Falconidae	<i>Falco cenchroides</i>	Nankeen Kestrel		MA	Y	Y						Not marine
Falconidae	<i>Falco longipennis</i>	Australian Hobby			Y	Y						
Falconidae	<i>Falco berigora</i>	Brown Falcon			Y	Y						
Falconidae	<i>Falco hypoleucos</i>	Grey Falcon	VU	VU				Y	Open plains and treed watercourses	✓		
Falconidae	<i>Falco peregrinus</i>	Peregrine Falcon	OS		Y	Y			Variety of habitats	✓		
Cacatuidae	<i>Nymphicus hollandicus</i>	Cockatiel				Y						
Cacatuidae	<i>Calyptrorhynchus banksii</i>	Red-tailed Black Cockatoo			Y	Y						
Cacatuidae	<i>Eolophus roseicapilla</i>	Galah			Y	Y						
Cacatuidae	<i>Cacatua sanguinea</i>	Little Corella			Y	Y						
Psittaculidae	<i>Aprosmictus erythropterus</i>	Red-winged Parrot			Y	Y						
Psittaculidae	<i>Pezoporus occidentalis</i>	Night Parrot	CR	EN				Y	Grasslands that are dominated by old-growth spinifex, though they have also been recorded in shrublands dominated by samphire, bluebush and saltbush	X	X	
Psittaculidae	<i>Psitteuteles versicolor</i>	Varied Lorikeet				Y						
Psittaculidae	<i>Trichoglossus moluccanus</i>	Rainbow Lorikeet			Y	Y						
Psittaculidae	<i>Melopsittacus undulatus</i>	Budgerigar			Y	Y						
Ptilonorhynchidae	<i>Chlamydera nuchalis</i>	Great Bowerbird			Y	Y						
Climacteridae	<i>Climacteris melanurus</i>	Black-tailed Treecreeper				Y						
Maluridae	<i>Malurus assimilis</i>	Purple-backed Fairywren				Y						
Maluridae	<i>Malurus elegans</i>	Red-winged Fairywren				Y						
Maluridae	<i>Malurus melanocephalus</i>	Red-backed Fairywren			Y	Y						

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Meliphagidae	<i>Epthianura crocea</i>	Yellow Chat			Y	Y						
Meliphagidae	<i>Conopophila rufogularis</i>	Rufous-throated Honeyeater			Y	Y						
Meliphagidae	<i>Ramsayornis fasciatus</i>	Bar-breasted Honeyeater			Y	Y						
Meliphagidae	<i>Certhionyx variegatus</i>	Pied Honeyeater				Y						
Meliphagidae	<i>Myzomela erythrocephala</i>	Red-headed Myzomela			Y	Y						
Meliphagidae	<i>Philemon citreogularis</i>	Little Friarbird			Y	Y						
Meliphagidae	<i>Philemon argenticeps</i>	Silver-crowned Friarbird			Y	Y						
Meliphagidae	<i>Lichmera indistincta</i>	Brown Honeyeater			Y	Y						
Meliphagidae	<i>Cissomela pectoralis</i>	Banded Honeyeater			Y	Y						
Meliphagidae	<i>Melithreptus gularis</i>	Black-chinned Honeyeater			Y	Y						
Meliphagidae	<i>Melithreptus albogularis</i>	White-throated Honeyeater			Y	Y						
Meliphagidae	<i>Stomiopera unicolor</i>	White-gaped Honeyeater			Y	Y						
Meliphagidae	<i>Gavicalis virescens</i>	Singing Honeyeater			Y	Y						
Meliphagidae	<i>Ptilotula flavescens</i>	Yellow-tinted Honeyeater			Y	Y						
Meliphagidae	<i>Ptilotula keartlandi</i>	Grey-headed Honeyeater				Y						
Meliphagidae	<i>Ptilotula plumula</i>	Grey-fronted Honeyeater				Y						
Pardalotidae	<i>Pardalotus rubricatus</i>	Red-browed Pardalote			Y	Y						
Pardalotidae	<i>Pardalotus striatus</i>	Striated Pardalote			Y	Y						
Acanthizidae	<i>Smicronis brevirostris</i>	Weebill			Y	Y						
Acanthizidae	<i>Gerygone levigaster</i>	Mangrove Gerygone			Y	Y						
Acanthizidae	<i>Gerygone fusca</i>	Western Gerygone				Y						
Acanthizidae	<i>Gerygone tenebrosa</i>	Dusky Gerygone			Y	Y						
Acanthizidae	<i>Gerygone magnirostris</i>	Large-billed Gerygone			Y							
Acanthizidae	<i>Gerygone chloronota</i>	Green-backed Gerygone				Y						
Acanthizidae	<i>Gerygone olivacea</i>	White-throated Gerygone				Y						
Pomatostomidae	<i>Pomatostomus temporalis</i>	Grey-crowned Babbler			Y	Y						
Artamidae	<i>Artamus leucorhynchus</i>	White-breasted Woodswallow			Y	Y						
Artamidae	<i>Artamus personatus</i>	Masked Woodswallow			Y	Y						
Artamidae	<i>Artamus cinereus</i>	Black-faced Woodswallow			Y	Y						
Artamidae	<i>Artamus minor</i>	Little Woodswallow			Y	Y						
Artamidae	<i>Gymnorhina tibicen</i>	Australian Magpie				Y						
Artamidae	<i>Cracticus nigrogularis</i>	Pied Butcherbird			Y	Y						

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Campephagidae	<i>Coracina novaehollandiae</i>	Black-faced Cuckooshrike		MA		Y	Y					Not marine
Campephagidae	<i>Coracina papuensis</i>	White-bellied Cuckooshrike		MA			Y					Not marine
Campephagidae	<i>Lalage tricolor</i>	White-winged Triller					Y					
Campephagidae	<i>Lalage leucomela</i>	Varied Triller				Y	Y					
Neosittidae	<i>Daphoenositta chrysoptera</i>	Varied Sittella				Y	Y					
Pachycephalidae	<i>Pachycephala occidentalis</i>	Golden Whistler					Y					
Pachycephalidae	<i>Pachycephala melanura</i>	Mangrove Golden Whistler				Y	Y					
Pachycephalidae	<i>Pachycephala rufiventris</i>	Rufous Whistler				Y	Y					
Pachycephalidae	<i>Pachycephala lanioides</i>	White-breasted Whistler				Y	Y					
Pachycephalidae	<i>Colluricincla harmonica</i>	Grey Shrikethrush				Y	Y					
Oriolidae	<i>Sphecotheres vieilloti</i>	Australasian Figbird				Y	Y					
Oriolidae	<i>Oriolus sagittatus</i>	Olive-backed Oriole				Y	Y					
Oriolidae	<i>Oriolus flavocinctus</i>	Green Oriole				Y	Y					
Dicruridae	<i>Dicrurus bracteatus</i>	Spangled Drongo		MA		Y	Y					Not marine
Rhipiduridae	<i>Rhipidura leucophrys</i>	Willie Wagtail				Y	Y					
Rhipiduridae	<i>Rhipidura rufiventris</i>	Northern Fantail				Y	Y					
Rhipiduridae	<i>Rhipidura albiscapa</i>	Grey Fantail				Y	Y					
Rhipiduridae	<i>Rhipidura phasiana</i>	Mangrove Fantail				Y	Y					
Monarchidae	<i>Grallina cyanoleuca</i>	Magpie-lark		MA		Y	Y					Not marine
Monarchidae	<i>Myiagra rubecula</i>	Leaden Flycatcher				Y	Y					
Monarchidae	<i>Myiagra ruficollis</i>	Broad-billed Flycatcher				Y	Y					
Monarchidae	<i>Myiagra alecto</i>	Shining Flycatcher				Y	Y					
Monarchidae	<i>Myiagra inquieta</i>	Restless Flycatcher				Y	Y					
Corvidae	<i>Corvus orru</i>	Torresian Crow				Y	Y					
Petroicidae	<i>Melanodryas cucullata</i>	Hooded Robin				Y	Y					
Petroicidae	<i>Peneothello pulverulenta</i>	Mangrove Robin				Y						
Petroicidae	<i>Microeca flavigaster</i>	Lemon-bellied Flyrobin				Y	Y					
Petroicidae	<i>Microeca fascians</i>	Jacky Winter				Y	Y					
Hirundinidae	<i>Hirundo rustica</i>	Barn Swallow	MI	MI				Y	Wide variety of habitats	✓		
Hirundinidae	<i>Cecropis daurica</i>	Red-rumped Swallow	MI	MI; MA				Y	Very occasionally seen in WA overflying most habitats	✓		
Hirundinidae	<i>Petrochelidon ariel</i>	Fairy Martin				Y	Y					
Hirundinidae	<i>Petrochelidon nigricans</i>	Tree Martin		MA		Y	Y					Not marine
Locustellidae	<i>Cincloramphus cruralis</i>	Brown Songlark					Y					
Locustellidae	<i>Cincloramphus mathewsi</i>	Rufous Songlark					Y					

Family	Species	Common	Conservation Status			Nature-Map	ALA	EPBC	Habitat Preference of Conservation Significant Species	Habitat Available in Study Area?	Habitat Available Adjacent to Study Area?	Included in Likelihood Assessment?
			State	Federal								
Cisticolidae	<i>Cisticola exilis</i>	Golden-headed Cisticola				Y						
Zosteropidae	<i>Zosterops luteus</i>	Canary White-eye			Y	Y						
Zosteropidae	<i>Zosterops lateralis</i>	Silveryeye		MA		Y					Not marine	
Dicaeidae	<i>Dicaeum hirundinaceum</i>	Mistletoebird			Y	Y						
Estrildidae	<i>Emblema pictum</i>	Painted Finch				Y						
Estrildidae	<i>Poephila acuticauda</i>	Long-tailed Finch			Y	Y						
Estrildidae	<i>Taeniopygia guttata</i>	Zebra Finch			Y	Y						
Estrildidae	<i>Taeniopygia bichenovii</i>	Double-barred Finch			Y	Y						
Estrildidae	<i>Erythrura gouldiae</i>	Gouldian Finch	P4	EN	Y	Y	Y	Open grassy eucalypt woodlands and grasslands. Forages on native grass especially sorghum	✓			
Estrildidae	<i>Lonchura castaneothorax</i>	Chestnut-breasted Mannikin			Y	Y						
Estrildidae	<i>Heteromunia pectoralis</i>	Pictorella Mannikin				Y						
Motacillidae	<i>Motacilla tschutschensis</i>	Eastern Yellow Wagtail	MI	MI; MA			Y	Open, moist, grassy or muddy areas, sewage treatment plants, sportsfields, sometimes beaches	X	✓		
Motacillidae	<i>Motacilla cinerea</i>	Grey Wagtail	MI	MI; MA			Y	Offshore islands, usually close to water including beaches and rock pools	X	✓		
Motacillidae	<i>Anthus australis</i>	Australasian Pipit		MA		Y					Not marine	

CR = Critically Endangered, EN = Endangered, VU = Vulnerable, OS = Other Specially Protected, MI = Migratory, MA = Marine, P4 = Priority 4

¹ Formerly considered part of one species, Gull-billed Tern *G. nilotica*, and listed as such, but now generally considered to involve two species

Invertebrates

Group	Family	Species	ALA	SRE ?
Land Snails	Camaenidae	<i>Quistrachia leptogramma</i>	Y	No
		<i>Rhagada bulgana/Rhagada cygna</i>	Y	Uncertain
		<i>Rhagada reinga</i>	Y	Potential
Scorpions	Buthidae	<i>Lychas</i> sp.	Y	N/A
	Urodacidae	<i>Urodacus hoplurus</i>	Y	No
Mygalomorph Spider	Nemesiidae	<i>Aname</i> sp.	Y	N/A
	Ctenizidae	<i>Conothele</i> sp.	Y	N/A

Appendix 11



Weed Locations



Species	Waypoint No.	Easting	Northing	Abundance
* <i>Aerva javanica</i>	467	488881	8173663	1
* <i>Azadirachta indica</i>	460	488827	8173585	1
* <i>Azadirachta indica</i>	482	488795	8173629	1
* <i>Cenchrus biflorus</i>	499	488657	8173751	Not recorded
* <i>Cenchrus setiger</i>	466	488870	8173663	Dense
* <i>Cenchrus setiger</i>	475	488870	8173696	Dense
* <i>Cenchrus setiger</i>	476	488849	8173607	Dense
* <i>Cenchrus setiger</i>	494	488635	8173596	Dense
* <i>Cenchrus setiger</i>	497	488603	8173685	Not recorded
* <i>Chloris barbata</i>	499	488657	8173751	Not recorded
* <i>Chloris barbata</i>	500	488689	8173773	Not recorded
* <i>Cyanthillium cinereum</i>	482	488795	8173629	1
* <i>Cyanthillium cinereum</i>	489	488827	8173729	6
* <i>Distimake dissectus</i> var. <i>dissectus</i>	472	488934	8173751	2
* <i>Distimake dissectus</i> var. <i>dissectus</i>	475	488870	8173696	1
* <i>Distimake dissectus</i> var. <i>dissectus</i>	482	488795	8173629	2
* <i>Distimake dissectus</i> var. <i>dissectus</i>	497	488603	8173685	1
* <i>Gomphrena celosioides</i>	499	488657	8173751	Not recorded
* <i>Jatropha gossypifolia</i>	497	488603	8173685	30
* <i>Jatropha gossypifolia</i>	503	488742	8173762	3
* <i>Macroptilium atropurpureum</i>	494	488635	8173596	2
* <i>Macroptilium atropurpureum</i>	500	488689	8173773	Not recorded
* <i>Macroptilium atropurpureum</i>	502	488731	8173751	Not recorded
* <i>Mesosphaerum suaveolens</i>	467	488881	8173663	14
* <i>Mesosphaerum suaveolens</i>	494	488635	8173596	1
* <i>Mesosphaerum suaveolens</i>	497	488603	8173685	Many
* <i>Mesosphaerum suaveolens</i>	498	488625	8173707	Many
* <i>Passiflora foetida</i> var. <i>hispida</i>	494	488635	8173596	6
* <i>Passiflora foetida</i> var. <i>hispida</i>	499	488657	8173751	Not recorded
* <i>Passiflora foetida</i> var. <i>hispida</i>	500	488689	8173773	Not recorded
* <i>Stylosanthes hamata</i>	467	488881	8173663	1
* <i>Stylosanthes hamata</i>	494	488635	8173596	1
* <i>Stylosanthes hamata</i>	498	488625	8173707	1
* <i>Stylosanthes hamata</i>	499	488657	8173751	Not recorded
* <i>Stylosanthes scabra</i>	471	488955	8173784	1
* <i>Stylosanthes scabra</i>	497	488603	8173685	2
* <i>Tridax procumbens</i>	499	488657	8173751	Not recorded