



**Leonora (Lot 51, Mt Ida Road)**

# **Biological Survey**

**Prepared for**  
**Horizon Power**

**October 2021**

● people ● planet ● professional

| Document Reference | Revision       | Prepared by       | Reviewed by   | Admin Review | Submitted to Client |            |
|--------------------|----------------|-------------------|---------------|--------------|---------------------|------------|
|                    |                |                   |               |              | Copies              | Date       |
| 4581AA_Rev0        | Internal Draft |                   |               | -            | -                   | 18/08/2021 |
| 4581AA_Rev1        | Client Draft   | 360 Environmental | Horizon Power | -            | 1 electronic        | 18/08/2021 |
| 4581AA_Rev2        | Final          | 360 Environmental | Horizon Power | -            | 1 electronic        | 05/10/2021 |

## Disclaimer

This report is issued in accordance with, and is subject to, the terms of the contract between the Client and 360 Environmental Pty Ltd, including, without limitation, the agreed scope of the report. To the extent permitted by law, 360 Environmental Pty Ltd shall not be liable in contract, tort (including, without limitation, negligence) or otherwise for any use of, or reliance on, parts of this report without taking into account the report in its entirety and all previous and subsequent reports. 360 Environmental Pty Ltd considers the contents of this report to be current as at the date it was produced. This report, including each opinion, conclusion, and recommendation it contains, should be considered in the context of the report as a whole. The opinions, conclusions and recommendations in this report are limited by its agreed scope. More extensive, or different, investigation, sampling and testing may have produced different results and therefore different opinions, conclusions, and recommendations. Subject to the terms of the contract between the Client and 360 Environmental Pty Ltd, copying, reproducing, disclosing, or disseminating parts of this report is prohibited (except to the extent required by law) unless the report is produced in its entirety including this cover page, without the prior written consent of 360 Environmental Pty Ltd.

© Copyright 2021 360 Environmental Pty Ltd ACN 109 499 041

# Executive Summary

Horizon Power commissioned 360 Environmental Pty Ltd to undertake a biological (flora and vegetation, targeted flora and vertebrate fauna) survey and assessment to inform decisions regarding the specific lease area for the proposed construction of a new power station in Leonora, Western Australia.

The Shire of Leonora have agreed (in principle) to lease a portion of their Lot known as Lot 51 on DP 55908 located at Mt Ida Road, Leonora, Western Australia. The leased area is expected to form a development envelope within the Survey Area. The Survey Area is located 200 m southeast of the Leonora Airport and comprises 6.6 hectares.

## Flora and Vegetation

The flora desktop assessment identified 46 conservation significant taxa occurring within 100 km of the Survey Area. A pre-survey likelihood of occurrence assessment was undertaken and identified that of these, none had previously been recorded within the Survey Area, none had a high or medium likelihood of occurrence, and 45 had a low likelihood of occurrence. The remaining one taxon was not assessed due to lack of information about habitat preference and distribution.

The reconnaissance flora and vegetation and targeted flora survey recorded floristic composition and vegetation types from four relevés, mapping notes and opportunistic observations. A total of 43 taxa from 15 families across 25 genera were recorded.

No Threatened flora species pursuant to the Environment Protection and Biodiversity Conservation Act 1999 and/or gazetted as Threatened pursuant to the Biodiversity and Conservation Act 2016, or Department of Biodiversity, Conservation and Attractions listed Priority flora were recorded during the survey.

One introduced flora species, *\*Cenchrus ciliaris*, was recorded during the survey. *\*Cenchrus ciliaris* is not listed as a Weed of National Significance by the Commonwealth Department of Agriculture, Water, and the Environment and as a Declared Pest by the State Department of Primary Industries and Regional Development.

Two vegetation types were described and mapped within the Survey Area. Vegetation in the Survey Area was generally representative of existing broad scale vegetation, soil, and land system mapping for the area.

Vegetation condition within the Survey Area ranged from Very Good to Good. Evidence of disturbance included access and vehicle tracks, weeds, and rubbish.

## Vertebrate Fauna

The fauna desktop assessment identified 242 terrestrial vertebrate fauna species, of which 178 are conservation significant, comprising:

- A total of 146 bird species including 21 conservation significant species

- A total of 35 mammal species including three conservation significant species
- A total of 54 reptile species including one conservation significant species
- Seven amphibian species, none of which are conservation significant.

A post-survey likelihood of occurrence assessment was undertaken and found that one fauna species of conservation significance was considered to have a high likelihood of occurrence within the Survey Area, the Peregrine Falcon (*Falco peregrinus*). Four conservation significant fauna species were considered to have a medium likelihood of occurrence within the Survey Area:

- Oriental Plover (*Charadrius veredus*) – Migratory, Marine
- Grey Falcon (*Falco hypoleucus*) – Migratory
- Princess Parrot (*Polytelis alexandrae*) – Priority 4, Vulnerable
- Woma (*Aspidites ramsayi*) – Priority 1 (southwest subpopulation).

The remaining conservation significant species are considered to have a low likelihood of occurrence.

The field survey recorded two terrestrial vertebrate fauna species, comprising of one native bird and one introduced mammal species, none of which were of conservation significance.

One fauna habitat type was identified within the Survey Area, described as Mulga woodland.

## Table of Abbreviations

| Abbreviation      | Description   |
|-------------------|---|
| 360 Environmental | 360 Environmental Pty Ltd   |
| BAM Act           | Biosecurity and Agriculture Management Act 2007   |
| BC Act            | WA Biodiversity Conservation Act 2016   |
| BoM               | Bureau of Meteorology   |
| °C                | Degree Celsius  |
| CD                | Conservation Dependent Fauna  |
| CR                | Critically Endangered   |
| DAWE              | Department of Agriculture, Water, and the Environment   |
| DBCA              | Department of Biodiversity, Conservation and Attractions  |
| DP                | Declared Pest   |
| DWER              | Department of Water and Environmental Regulation  |
| EN                | Endangered  |
| EP Act            | WA Environmental Protection Act 1986  |
| EPA               | Environmental Protection Authority  |
| EPBC Act          | Environment Protection Biodiversity and Conservation Act 1999   |
| ESA               | Environmentally Sensitive Area  |
| GIS               | Geographic Information System   |
| ha                | Hectares  |
| IBRA              | Interim Biogeographic Regionalisation for Australia   |
| IBSA              | Index of Biodiversity Surveys for Assessments   |
| km                | Kilometres  |
| m                 | Metres  |
| mm                | Millimetres   |
| MA                | Marine  |
| MI                | Migratory   |
| MNES              | Matters of National Environmental Significance  |
| NVIS              | National Vegetation Information System  |
| OS                | Other Specially Protected Fauna   |
| P                 | Priority  |
| PEC               | Priority Ecological Community   |
| PMST              | Protected Matters Search Tool   |
| Study Area        | The database search area (varied according to each parameter)   |
| Survey Area       | The Survey Area is defined area on Lot 51 on Mt Ida Road in Leonora, Western Australia. The Survey Area is located 200 m southeast of the Leonora Airport and comprises 6.6 hectares. |
| T                 | Threatened  |

| Abbreviation | Description                                |
|--------------|--|
| TEC          | Threatened Ecological Community            |
| TPFL         | Threatened and Priority Flora Database     |
| TPFRF        | Threatened and Priority Flora Report Forms |
| VU           | Vulnerable                                 |
| WA           | Western Australia                          |
| WAH          | Western Australian Herbarium               |
| WAM          | Western Australian Museum                  |
| WoNS         | Weeds of National Significance             |

## Table of Contents

|          |   |           |
|----------|---|-----------|
| <b>1</b> | <b>Introduction.....</b>                        | <b>7</b>  |
| 1.1      | The Project.....                                | 7         |
| 1.2      | Objectives and Scope .....                      | 7         |
| <b>2</b> | <b>Background.....</b>                          | <b>8</b>  |
| 2.1      | Protection of Flora, Vegetation and Fauna ..... | 8         |
| 2.2      | Existing Environment .....                      | 9         |
| <b>3</b> | <b>Methods .....</b>                            | <b>12</b> |
| 3.1      | Desktop Assessment .....                        | 12        |
| 3.2      | Field Survey .....                              | 14        |
| 3.3      | Flora and Vegetation.....                       | 14        |
| 3.4      | Vertebrate Fauna .....                          | 16        |
| 3.5      | Limitations .....                               | 17        |
| <b>4</b> | <b>Results.....</b>                             | <b>19</b> |
| 4.1      | Flora and Vegetation.....                       | 19        |
| 4.2      | Vertebrate Fauna Results.....                   | 23        |
| <b>5</b> | <b>Discussion .....</b>                         | <b>26</b> |
| 5.1      | Flora and Vegetation.....                       | 26        |
| 5.2      | Vertebrate Fauna .....                          | 27        |
| <b>6</b> | <b>Conclusion .....</b>                         | <b>30</b> |
| <b>7</b> | <b>References .....</b>                         | <b>31</b> |
| <b>8</b> | <b>Limitations of this Report .....</b>         | <b>34</b> |

## List of Graphs

|  |          |
|--|----------|
| <b>Graph 1: Long-term Maximum and Minimum temperatures for Leonora WA (12046) and Monthly Rainfall for Leonora Aero (12241) (Bureau of Meteorology, 2021). .....</b> | <b>9</b> |
|--|----------|

## List of Tables

|   |           |
|---|-----------|
| <b>Table 1: Representation of Laverton 28 Vegetation Systems Association on a State, Regional and Local Scale (Government of Western Australia, 2019) .....</b> | <b>10</b> |
| <b>Table 2: Database Searches of the Survey Area.....</b>   | <b>13</b> |
| <b>Table 3: Likelihood of Occurrence Criteria.....</b>  | <b>14</b> |
| <b>Table 4: Limitations and constraints associated with the survey.....</b>   | <b>17</b> |
| <b>Table 5: TECs and PECs identified during the Desktop Assessment .....</b>  | <b>19</b> |
| <b>Table 6: Vegetation Types Occurring within the Survey Area.....</b>  | <b>22</b> |
| <b>Table 7: Fauna Habitat within the Survey Area .....</b>  | <b>24</b> |

## List of Figures

|                                |
|--------------------------------|
| <b>Figure 1: Survey Area</b>   |
| <b>Figure 2: Hydrography</b>   |
| <b>Figure 3: Survey Effort</b> |

**Figure 4: DBCA Threatened and Priority Flora and Communities**

**Figure 5: Introduced Flora**

**Figure 6: Vegetation Types and Condition**

**Figure 7: DBCA Threatened and Priority Fauna Locations**

**Figure 8: Fauna Habitat**

## **List of Appendices**

**Appendix A Flora Literature Review**

**Appendix B Flora Database Searches**

**Appendix C Flora Likelihood of Occurrence**

**Appendix D Inventory of Vascular Flora**

**Appendix E Flora Site Sheets**

**Appendix F Fauna Literature Review and Database Searches Results**

**Appendix G Fauna Likelihood of Occurrence**

**Appendix H Fauna Habitat Assessments**

## 1 Introduction

### 1.1 The Project

Horizon Power commissioned 360 Environmental Pty Ltd (360 Environmental) to undertake a biological survey and assessment (reconnaissance flora and vegetation survey, targeted flora survey and basic terrestrial vertebrate fauna survey) to inform decisions regarding the lease area for the proposed construction of a new power station in Leonora, Western Australia (herein referred to as the Survey Area).

The Shire of Leonora have agreed (in principle) to lease a portion of Lot 51 on DP 55908 located at Mt Ida Road, Leonora, Western Australia. The leased area is expected to form a development envelope within the Survey Area. The Survey Area is located 200 m southeast of the Leonora Airport and comprises 6.6 ha (Figure 1).

### 1.2 Objectives and Scope

The purpose of the survey was to delineate key flora and fauna values within the Survey Area and identify potential environmental sensitivities that may impact the Project.

The scope of works included:

- Complete a biological assessment pursuant to the requirements of a native vegetation clearing permit application
- A desktop assessment to determine environmental values and conservation significant flora, fauna, habitat, vegetation, or other environmental features (such as riparian areas, wetlands) relating to the Survey Area
- A field survey to assess flora, vegetation and fauna values including:
  - Reconnaissance Flora and Vegetation Survey
  - Targeted Flora Survey
  - Basic Fauna Survey
- Preparation of a combined technical flora, vegetation, and fauna survey report.
- An assessment against the Ten Clearing Principles
- Recommendations for any approvals requirements that would be required to clear within the Survey Area
- Provision of mapping and spatial data compiled in accordance with Index of Biodiversity Surveys for Assessment (IBSA).

## 2 Background

### 2.1 Protection of Flora, Vegetation and Fauna

Western Australian flora and fauna is protected formally and informally by legislative and non-legislative measures:

#### Legislative measures

- Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act)
- WA *Biodiversity Conservation Act 2016* (BC Act)
- WA *Environmental Protection Act 1986* (EP Act)
- WA *Biosecurity and Agriculture Management Act 2007* (BAM Act).

#### Non-legislative measures

- WA Department of Biodiversity Conservation and Attractions (DBCA) Priority lists for flora, fauna, and ecological communities
- Weeds of National Significance (WoNS)
- Recognition of locally significant populations by DBCA.

These protection mechanisms are supported by guidance documents published by the Environmental Protection Authority (EPA) and Department of Agriculture Water and the Environment (DAWE; formerly Department of Environment, and Department of Sustainability Environment Water Population and Communities):

- Technical Guidance - Flora and Vegetation Surveys for Environmental Impact Assessment (Environmental Protection Authority, 2016b)
- Technical Guidance - Terrestrial vertebrate fauna surveys for environmental impact assessment (Environmental Protection Authority, 2020)
- Matters of National Environmental Significance Significant impact guidelines 1.1 Environment Protection and Biodiversity Conservation Act 1999 (Department of the Environment, 2013).

Conservation codes used throughout this report are in accordance with Conservation Codes for Western Australian Flora and Fauna (Department of Biodiversity Conservation and Attractions, 2020) and Definitions, Categories and Criteria for Threatened and Priority Ecological Communities (Department of Environment and Conservation, 2013).


## 2.2 Existing Environment

### 2.2.1 Climate

The closest long-term Bureau of Meteorology (BoM) weather station with a climate dataset is Leonora WA (Station 12046), located approximately 1.3 km southeast of the Survey Area. The closest long-term Bureau of Meteorology (BoM) weather station with a rainfall dataset is Leonora Aero (Station 12241), located approximately 0.2 km west of the Survey Area.

The long-term mean minimum temperature for Leonora WA weather station ranges from 6.3°C (July) to 22.0°C (January) (1991 to 2021) and the long-term mean maximum temperature ranges from 18.8°C (July) to 36.8°C (January) (Graph 1) (Bureau of Meteorology, 2021).

The Leonora Aero weather station recorded 145.2 mm of rainfall in the 12 months prior to the survey (June 2020 to May 2021), which is 108.9 mm below the long-term average of 254.1 mm (Bureau of Meteorology, 2021). In the three months prior to the survey (March to May 2021), 60.8 mm of rainfall was recorded, which is 6.6 mm below the long-term average of 67.4 mm for the same time period (Bureau of Meteorology, 2021).



**Graph 1: Long-term Maximum and Minimum temperatures for Leonora WA (12046) and Monthly Rainfall for Leonora Aero (12241) (Bureau of Meteorology, 2021).**

### 2.2.2 Interim Biogeographic Regionalisation of Australia

The Interim Biogeographic Regionalisation of Australia (IBRA) divides Australia into 89 bioregions based on major biological, geographical, and geological attributes. These bioregions are subdivided into 419 subregions as part of a refinement of the IBRA framework (Department of the Environment and Energy, 2016). The Survey Area occurs within the Murchison bioregion and the Eastern Murchison (MUR01) subregion.

The Eastern Murchison subregion is characterised by its internal drainage, and extensive areas of elevated red desert sandplains with minimal dune development (Cowan, 2001). The subregion is characterised by broad plains of red-brown soils and breakaway complexes as well as red sandplains. Vegetation is dominated by Mulga Woodlands often rich in ephemerals; hummock grasslands, saltbush shrublands and *Tecticornia* shrublands.

### 2.2.3 Soil Landscapes and Land Systems

Soil landscape and land system mapping of Western Australia described broad soil and landscape characteristics from regional to local scales, ranging from 1:20,000 to 1:250,000 (Department of Agriculture and Food WA, 2012). The Survey Area occurs within the Gundockerta System (279Gu), which is described as extensive, gently undulating calcareous stony plains supporting bluebush shrublands.

### 2.2.4 Hydrography

The Survey Area does not intersect any major watercourses or water bodies that are mapped by State Government GIS databases (Department of Water and Environmental Regulation, 2018). The closest watercourse is a minor drainage line, located 130 m south of the Survey Area (Figure 2).

### 2.2.5 Broad Vegetation Associations

Mapping of pre-European vegetation within Western Australia was completed on a broad scale (1:1,000,000) by Beard (1976). These vegetation types were later refined by Shepherd *et al.* (2002) resulting in 819 vegetation associations.

The Laverton 28 broad vegetation association is mapped over the Survey Area. Laverton 28 is described as low woodland, open low woodland, or sparse woodland of Mulga (*Acacia aneura* and associated species). The representation on a regional, state, and local scale of Laverton 28 is shown in Table 1.

**Table 1: Representation of Laverton 28 Vegetation Systems Association on a State, Regional and Local Scale (Government of Western Australia, 2019)**

| Pre-European Extent (ha)                      | Current Extent (ha) | Remaining (%) |
|---|---------------------|---------------|
| Representation across Western Australia       |                     |               |
| 395,895.08                                    | 392,171.83          | 99.06         |
| Representation across the Murchison Bioregion |                     |               |

| Pre-European Extent (ha)                                     | Current Extent (ha) | Remaining (%) |
|--|---------------------|---------------|
| 224,291.84   | 220,583.71          | 98.35         |
| <b>Representation across the Eastern Murchison Subregion</b> |                     |               |
| 141,411.26   | 137,703.12          | 97.38         |
| <b>Representation across the Shire of Leonora</b>            |                     |               |
| 126,344.70   | 124,136.29          | 98.25         |

## 2.2.6 Environmentally Sensitive Areas

Environmentally Sensitive Areas (ESAs) are declared by the Department of Water and Environmental Regulation (DWER) to prevent the degradation of important environmental values such as Threatened flora, Threatened Ecological Communities (TECs) or significant wetlands (Department of Water and Environmental Regulation, 2014).

The Survey Area is not mapped within an ESA (Department of Water and Environmental Regulation, 2021). The nearest ESA is associated with Lake Ballard, which is located approximately 70 km south of the Survey Area.

## 2.2.7 Conservation Areas

The Survey Area does not intersect any Conservation Areas (Department of Biodiversity Conservation and Attractions, 2021a). The nearest Conservation Area is an un-named reserve (R 46847) vested under the Conservation Commission of WA, which is located 57 km south of the Survey Area.

## 3 Methods

The biological survey was undertaken in accordance with relevant EPA and DAWE guidelines (see section 2.1).

### 3.1 Desktop Assessment

#### 3.1.1 Literature Review

Background information on the Survey Area and surrounds was compiled prior to the field survey (see Section 2). Historical vegetation mapping conducted by Beard (1976) and Shepherd *et al.* (2002), land systems mapping (Department of Agriculture and Food WA, 2012), and the IBRA classification system (Department of the Environment and Energy, 2016) were consulted to provide broad contextual knowledge of vegetation units and habitat likely to be encountered within the Survey Area.

The literature review also considered eight publicly available biological reports undertaken in the vicinity of the Survey Area:

- Biological Survey of the Eastern Goldfields of Western Australia Part 10 (Hall, McKenzie and Keighery, 1994), approximately 98 km northeast of the Survey Area.
- Detailed Flora and Vegetation Survey of Tims Find (Native Vegetation Solutions, 2019), located approximately 83.6 km west of the Survey Area.
- Leonora Rail Yard Expansion Project Level 1 Flora and Vegetation Survey (Western Botanical, 2013), located approximately 0.6 km northeast of the Survey Area.
- Level 2 Fauna Risk Assessment for Granny Deeps Project Area (Terrestrial Ecosystems, 2011), approximately 100 km east of the Survey Area.
- Level 2 Vertebrate Fauna Assessment, King of the Hills Project (Terrestrial Ecosystems, 2020a), approximately 29 km north northwest of the Survey Area.
- Preliminary Environmental Impact Assessment, Flora Survey and Environmental Management Plan (GHD, 2011), approximately 51.9 km east of the Survey Area.
- Report for Gwalia Materials, Preliminary Environmental Impact Assessment, Flora Survey and Environmental Management Plan (GHD, 2011), approximately 54 km east of the Survey Area.
- Vertebrate Fauna Risk Assessment for the Granny Smith Solar Power Farm Project (Terrestrial Ecosystems, 2018), approximately 100 km east of the Survey Area.
- Vertebrate Fauna Risk Assessment Granny Smith Tailing Storage Facility Expansion (Terrestrial Ecosystems, 2020b), approximately 105 km east of the Survey Area.

### 3.1.2 Database Searches

Database searches were undertaken to identify potential conservation significant flora and fauna taxa, ecological communities, and Matters of National Environmental Significance (MNES) within or surrounding the Survey Area (Table 2). The search area for each parameter was varied to reflect distances recommended by DBCA (Table 2). The search areas are herein referred to collectively as the Study Area.

Currently listed Priority Ecological Communities (PECs) and TECs that occur in the region were examined to determine if any corresponded with the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). In addition, an EPBC Protected Matters Search (PMST) was undertaken to identify the potential for Matters of National Environmental Significance (MNES) to occur within or surrounding the Survey Area (Department of Agriculture Water and the Environment, 2021a)).

**Table 2: Database Searches of the Survey Area**

| Database Name   | Date Received | Search Target   | Search Area                          |
|---|---------------|---|--------------------------------------|
| DBCA Threatened and Priority Ecological Communities database custom search (Department of Biodiversity Conservation and Attractions, 2021c) | 19 May 2021   | TECs and PECs   | 50 km buffer around the Survey Area  |
| DBCA Threatened and Priority Flora Species List (TP list) custom search (Department of Biodiversity Conservation and Attractions, 2021e)    | 11 May 2021   | Threatened and Priority flora   | 100 km buffer around the Survey Area |
| Western Australian Herbarium flora custom search (Department of Biodiversity Conservation and Attractions, 2021f)                           |               |   |                                      |
| DBCA Threatened and Priority Fauna List custom search (Department of Biodiversity Conservation and Attractions, 2021d)                      | 19 May 2021   | Threatened Priority Fauna   | 100 km buffer around the Survey Area |
| NatureMap area search (Department of Biodiversity Conservation and Attractions, 2021b)  | 25 May 2021   | Threatened and priority flora and fauna, and inventory of potential flora and fauna | 40 km buffer around the Survey Area  |
| Protected Matters Search Tool area search (Department of Agriculture Water and the Environment, 2021a)                                      | 25 May 2021   | Commonwealth listed threatened flora and fauna, and TECs                            | 50 km buffer around the Survey Area  |

### 3.1.3 Likelihood of Occurrence

Conservation significant flora species identified from the desktop assessment were assessed to determine the likelihood of their occurrence within the Survey Area, both prior and post field survey.

Conservation significant fauna species identified from the desktop assessment were assessed following the field survey to determine the likelihood of their occurrence within the Survey Area.

Fauna species listed as Marine under the EPBC Act were not included as conservation significant as the Marine listing only applies within Commonwealth marine areas.

The assessments were completed based on the likelihood of occurrence criteria presented in Table 3.

**Table 3: Likelihood of Occurrence Criteria**

| Rank                       | Criteria   |
|----------------------------|--|
| Previously Recorded        | The species has been previously recorded in the Survey Area  |
| High<br>(Likely to occur)  | <ul style="list-style-type: none"> <li>• There are existing records of the species in close proximity to the Survey Area (within 5 km), and for fauna has been recorded in the Survey Area in the last 15 years</li> <li>• The species is strongly linked to a specific habitat, which is present in the Survey Area or</li> <li>• The species has more general habitat preferences, and suitable habitat is present.</li> </ul>   |
| Medium<br>(May occur)      | <ul style="list-style-type: none"> <li>• There are existing records of the species from the locality (within 15 km), however:           <ul style="list-style-type: none"> <li>◦ The species is strongly linked to a specific habitat, of which only a small amount is present in the Survey Area or</li> <li>◦ The species has more general habitat preferences, but only some suitable habitat is present.</li> </ul> </li> <li>• There is suitable habitat in the Survey Area, but the species is recorded infrequently in the locality.</li> </ul> |
| Low<br>(Unlikely to occur) | <ul style="list-style-type: none"> <li>• The species is linked to a specific habitat, which is absent from the Survey Area or</li> <li>• Suitable habitat is present, however there are no existing records of the species from the locality despite reasonable previous search effort in suitable habitat or</li> <li>• There is some suitable habitat in the Survey Area, however the species is very infrequently recorded in the locality.</li> </ul>  |

### 3.2 Field Survey

The reconnaissance flora and vegetation survey and basic terrestrial vertebrate fauna survey was undertaken by Principal Botanist [REDACTED] and Ecologist [REDACTED] on 24th June 2021. The survey effort is shown in Figure 3.

### 3.3 Flora and Vegetation

#### 3.3.1 Establishment of Flora sites

The Survey Area was assessed using relevés and meandering traverses to gather information to characterise and delineate vegetation and compile an inventory of vascular flora. At least one flora site was sampled in each vegetation type observed within the Survey Area. At each relevé, the following information was recorded using a Fulcrum mobile data collection device:

- Location
- Site code
- Date and personnel

- Landform and soil description
- Relevant site descriptors including, slope, aspect, litter cover, bare ground cover and fire history
- Inventory of vascular flora including the approximate average height and percentage foliar cover for each taxon recorded
- Vegetation description in accordance with National Vegetation Information System (NVIS), Level 6 ‘sub-association’, whereby the dominant growth form, height, cover, and species (up to five species) for the three traditional strata (upper, mid, and ground) are described
- Vegetation condition in accordance with the Eremaean and Northern Botanical Provinces vegetation condition scale (Environmental Protection Authority, 2016b) and evidence of disturbance (for example clearing, rubbish, feral animals, weed incursion and evidence of feral animals and dieback) where present
- Photograph of the vegetation occurring within the site.

A total of four relevés were established within the Survey Area.

### 3.3.2      Opportunistic Flora

Additional flora taxa observed opportunistically around flora sites or while traversing on foot within the Survey Area were also recorded. Where populations of conservation significant flora taxa, introduced flora, Declared Pests (DPs) or WoNS were encountered, a GPS location and a count of the individuals present was recorded.

### 3.3.3      Targeted Searching

Prior to the survey, conservation significant flora with the likelihood or potential to occur within the Survey Area was compiled (see section 3.1). Field personnel familiarised themselves with photographs, reference samples and descriptions of these taxa before conducting the survey.

The entire Survey Area was searched using meandering traverses. Personnel also actively searched for conservation significant flora species in and around flora sites, while traversing on foot within the Survey Area and in known locations or preferred habitat encountered in the Survey Area.

### 3.3.4      Taxonomy and Nomenclature

Where field identification of plant taxa was not possible in the field, specimens were collected for identification using resources at the WAH. Identification of flora collections was completed by Principal Botanist/Taxonomist Ben Eckermann.

The finalised species list was checked against FloraBase (Western Australian Herbarium, 2021) to determine the conservation status and known distribution of each taxon. Introduced species were compared against the current BAM Act Declared Plants list and the WoNS list to determine

their control status (Department of Agriculture Water and the Environment, 2021b; Department of Primary Industries and Regional Development, 2021).

### 3.3.5 Vegetation Unit and Condition Mapping

Broad vegetation and condition mapping was conducted in the field, with boundaries delineated over aerial photography, at a scale of 1:5,000. Broad vegetation units were refined based on taxonomic identification of flora collections and site notes taken during the field survey. Vegetation condition mapping was refined based on site data and mapping notes. Finalised polygons were digitised and produced as electronic mapping data using GIS software.

## 3.4 Vertebrate Fauna

### 3.4.1 Fauna Habitat Assessment

Fauna habitat assessments were undertaken throughout the Survey Area to identify fauna habitat values. Habitat assessment locations are shown in Figure 3. The following information was collected at each site using Fulcrum, a mobile data collection app:

- Site photo
- Landform
- Soil type and colour
- Rock types, surface stone cover and size classes
- Key habitat and microhabitat features including leaf litter, logs, burrows, rocky outcrops, rock crevices, hollows, water sources
- Habitat quality, fire history and evidence of disturbance
- General description of vegetation structure.

Fauna habitat mapping was based on a combination of field observations, fauna habitat assessment data and vegetation mapping.

### 3.4.2 Opportunistic Fauna Records

Throughout the field survey opportunistic observations of fauna species (including feral species) were recorded. Data collected included:

- Family, genus, species, and common name
- Conservation status
- Introduced status
- Abundance/population size
- Western Australian Museum lodgment number (if applicable)
- Sampling type
- Sampling point number

- Habitat description:
  - Brief description if species is not conservation significant
  - A full detailed description if species is conservation significant.
- Coordinates in GDA94 or GDA2020, easting and northings
- Photograph (if possible)
- Notes (any additional information, if necessary)
- Brief location.

In addition to the direct observation of fauna species, secondary evidence such as tracks, diggings and scats was noted. All fauna observations were recorded using a mobile device with GPS capability.

#### 3.4.3 Taxonomy

Terrestrial vertebrate fauna taxa were identified in the field.

Where there was doubt on species names (through subsequent name changes or taxonomic reviews), an effort was made to determine the current scientific name for each taxon. Taxonomy and nomenclature in this report follows the WA Museum checklist 2021 (Western Australian Museum, 2021) where relevant.

### 3.5 Limitations

Limitations and constraints of the flora, vegetation and fauna survey are detailed below in Table 4.

**Table 4: Limitations and constraints associated with the survey**

| Variable                    | Degree of Limitation | Potential Constraints on Survey Outcomes  |
|-----------------------------|----------------------|---|
| <b>Survey Scope</b>         | Not a limitation     | The reconnaissance flora and vegetation, targeted flora and basic vertebrate fauna surveys were undertaken in accordance with (Environmental Protection Authority, 2016b, 2020) and were considered appropriate to support approvals applications.  |
| <b>Availability of Data</b> | Not a limitation     | All data required to complete the scope of works including regional and local contextual information was available.   |
| <b>Site Access</b>          | Not a limitation     | The Survey Area was able to be accessed on foot.  |
| <b>Survey Effort</b>        | Not a limitation     | Sufficient time was allocated to the flora and vegetation and fauna surveys, given the size and complexity of the Survey Area, and the expected level of survey intensity.<br>The survey effort is displayed in Figure 3.   |
| <b>Experience</b>           | Not a limitation     | The flora and vegetation and fauna surveys were undertaken by Principal Botanist Ben Eckermann and Ecologist Lachlan Crossley, respectively.<br>Ben has 15 years' experience conducting surveys of similar scope throughout Western Australia.<br>Lachlan has three years' experience conducting surveys of similar scope throughout Western Australia. |

| Variable                               | Degree of Limitation       | Potential Constraints on Survey Outcomes  |
|--|----------------------------|---|
| <b>Timing,<br/>weather,<br/>season</b> | Not a limitation for Flora | The recommended primary survey period for flora and vegetation and fauna surveys in the Eremaean Botanical Province as per the EPA Technical Guidance is six to eight weeks post wet season (March – June). The survey was undertaken in June, within the optimal survey period for the region.   |
|  | A limitation for Fauna     | Morning temperatures were quite low when the fauna survey began, and the survey site was very dry. These limiting factors may explain the lack of fauna species observed.   |
| <b>Life forms sampled</b>              | Not a limitation           | <p>The Survey Area was traversed on foot and sites representative of all vegetation were sampled. All flora and fauna species encountered within the Survey Area were recorded.</p> <p>A total of 44 vascular flora taxa were recorded from the Survey Area, comprising 97.7% native flora taxa and 2.3% introduced flora taxa.</p> <p>Of the 44 flora taxa recorded, two taxa (4.5%), could not be identified to species level because they were sterile at the time of the survey (and may represent confirmed species). This was not considered a constraint as it represented a small portion of the flora sampled.</p> <p>None of the unknown flora taxa collected were analogous to Threatened or Priority flora taxa identified by the database searches as likely to occur within the Survey Area, nor were they representative of flora of other significance.</p> <p>A total of five vertebrate fauna taxa were recorded from the Survey Area, comprising three confirmed native taxa and two confirmed introduced taxa. All five taxa recorded were able to be identified by direct or indirect observation with high level of scientific confidence to species or genus level and were not considered Threatened or Priority Fauna.</p> |
| <b>Mapping Reliability</b>             | Not a limitation           | <p>Vegetation types were described and mapped based on relevé data.</p> <p>High resolution aerial mapping current at the time of the survey was used to differentiate all vegetation greater than 1 ha in size.</p> <p>Fauna habitat mapping was based largely on vegetation mapping.</p>   |
| <b>Disturbances (fire, flood etc.)</b> | Not a limitation           | Areas of disturbance associated with access and vehicle tracks, weeds, and rubbish were recorded but were not a constraint on the results of the survey.  |
| <b>Completeness</b>                    | Not a limitation           | <p>The survey was considered complete for a reconnaissance flora and vegetation survey, all vegetation types were surveyed and delineated within the Survey Area and a minimum of one relevé was surveyed for each vegetation type.</p> <p><b>ADD COMPLETENESS FOR TARGETED FLORA</b></p> <p>The survey was considered complete for a targeted flora survey. The Survey Area was traversed on foot via meandering traverses. None of the unconfirmed flora taxa collected were analogous to Threatened or Priority flora taxa identified by the database searches as likely to occur within the Survey Area, nor were they representative of flora of other significance.</p> <p>The survey was considered complete for a basic vertebrate fauna survey.</p>  |

## 4 Results

### 4.1 Flora and Vegetation

#### 4.1.1 Desktop Assessment

The key findings of the previous flora and vegetation reports reviews are summarised in Appendix A.

The desktop assessment identified 46 conservation significant species occurring within the vicinity of the Study Area (Figure 4; Appendix B). This list was comprised of:

- No Threatened species
- Fourteen Priority 1 species
- Three Priority 2 species
- Twenty-four Priority 3 species
- Five Priority 4 species.

One species, *Ptilotus chortophytus* (P1), was located approximately 60 km south of the Survey Area by the DBCA database searches. This record would appear to be an error as Florabase reports this taxon as being restricted to areas north of Geraldton (Western Australian Herbarium, 2021). This taxon was removed from the desktop assessment and likelihood of occurrence assessment due to the spatial disjunction as reported by DBCA and Florabase.

No State or Commonwealth listed TECs or DBCA listed PECs were mapped within the Survey Area. Two PECs occur within 50 km of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c) (Table 5, Figure 4).

**Table 5: TECs and PECs identified during the Desktop Assessment**

| Name   | State TEC /<br>DBCA PEC | Commonwealth<br>TEC | Location in<br>relation to Survey<br>Area |
|--|-------------------------|---------------------|---|
| Melita calcrete groundwater assemblage type on Raeside palaeodrainage on Melita (Sons of Gwalia) Station | Priority 1              | Not listed          | 2.9 km south of the Survey Area           |
| Sturt Meadows calcrete groundwater assemblage type on Raeside palaeodrainage on Sturt Meadows Station    | Priority 1              | Not listed          | 38.3 km west of the Survey Area           |

#### 4.1.2 Pre-Survey Likelihood of Occurrence

The pre-survey likelihood of occurrence assessment for the Survey Area identified that of the 46 conservation significant flora species identified by the desktop assessment:

- None had previously been recorded within the Survey Areas
- None were considered to have a high likelihood of occurrence

- None were considered to have a medium likelihood of occurrence
- Forty-five were considered to have a low likelihood of occurrence.

The remaining one taxon was not assessed due to lack of information about habitat preference and distribution. The likelihood of occurrence assessment is provided in Appendix C.

#### 4.1.3 Flora Composition

A total of 43 taxa from 15 families across 25 genera were recorded during the survey (Appendix D). The dominant families were Fabaceae (10 taxa), Chenopodiaceae (eight taxa) and Poaceae (seven taxa). The most dominant genus was *Acacia* (eight taxa).

#### 4.1.4 Flora of Conservation Significance

##### 4.1.4.1 Threatened and Priority Flora

No Threatened flora species pursuant to the EPBC Act and/or gazetted as Threatened flora pursuant to the BC Act were recorded during the survey.

No DBCA listed Priority flora taxa were recorded within the Survey Area.

##### 4.1.4.2 Flora of Other Conservation Significance

Flora may be considered of other conservation significance if it represents a range extension, novel taxon, species that play a keystone role in a community, has relic status, is locally endemic, or represents the extent of a species range. No flora taxa recorded from the Survey Area were considered flora of other conservation significance.

#### 4.1.5 Introduced Flora

One introduced flora species, *\*Cenchrus ciliaris*, was recorded in the Survey Area, representing 2.3% of the total taxa recorded (Figure 5, Appendix D). Approximately 500 individuals of *\*Cenchrus ciliaris* were recorded within the Survey Area. *\*Cenchrus ciliaris* is not listed as a Declared Pest under the BAM Act (Department of Primary Industries and Regional Development, 2021) and as a WoNS (Department of Agriculture Water and the Environment, 2021b).

#### 4.1.6 Unconfirmed Flora

Two specimens (4.7% of the taxa recorded) could not be identified to species level because the taxa were sterile at the time of the survey (Appendix D). These have been assigned a confirmed genus, *Maireana* sp. and *Sida* sp.

The two unconfirmed flora taxa may represent duplicates of taxa that were confirmed within the Survey Area.

None of the unconfirmed flora taxa were analogous to Priority flora taxa identified by the database searches.

#### 4.1.7 Vegetation Types

Two vegetation types were described and mapped within the Survey Area (Table 6; Figure 6).

Detailed site sheets for each flora site are provided in Appendix E.

#### 4.1.8 Vegetation Condition

Vegetation condition within the Survey Area ranged from Very Good (3.6 ha / 53.8%) to Good (3.1 ha / 46.2%) (Figure 6).

Evidence of disturbance included access and vehicle tracks, weeds, and rubbish.

**Table 6: Vegetation Types Occurring within the Survey Area**

| Vegetation Unit | Area and Proportion of the Survey Area   | Sites                             | Vegetation Condition | Photograph   |
|-----------------|--|-----------------------------------|----------------------|--|
| Plains          | P1: <i>Acacia caesaneura</i> and <i>Acacia mulganeura</i> low woodland over <i>Eremophila forestii</i> subsp. <i>forrestii</i> mid sparse shrubland over <i>Eragrostis eriopoda</i> low sparse tussock grassland | 3.6 ha<br>53.8%<br>LER01<br>LER03 | Very Good            |   |
|                 | P2: Mixed <i>Acacia</i> spp. low open woodland over <i>Eragrostis eriopoda</i> low sparse tussock grassland  | 3.1 ha<br>46.2%<br>LER02<br>LER04 | Good                 |  |

#### 4.1.9 Threatened and Priority Ecological Communities

No vegetation representative of any Commonwealth or State listed TECs or PECs were recorded within the Survey Area.

Vegetation may be of significance for a range of reasons, other than a listing as a TEC or a PEC, including (Environmental Protection Authority, 2016a):

- Vegetation extent being below a threshold level
- Scarcity
- Unusual species
- Novel combinations of species
- A role as a refuge
- A role as a key habitat for threatened species or large populations representing a significant proportion of the local to regional total population of a species
- Being representative of the range of a unit (particularly a good local and/or regional example of a unit in ‘prime’ habitat, at the extremes of range, recently discovered range extensions, or isolated outliers of the main range) and/or
- A restricted distribution.

No vegetation of other significance was recorded in the Survey Area.

#### 4.1.10 Survey Adequacy

A total of four relevés were established within the Survey Area. This was adequate to ensure at least one relevé was sampled in each vegetation type, and that coverage across the Survey Area was sufficient.

### 4.2 Vertebrate Fauna Results

#### 4.2.1 Desktop Assessment

The key findings of the fauna reports reviewed are summarised in Appendix F.

The desktop assessment identified 242 terrestrial vertebrate fauna species, of which 178 are conservation significant, comprising:

- A total of 146 bird species including 21 conservation significant species
- A total of 35 mammal species including three conservation significant species
- A total of 54 reptile species including one conservation significant species
- Seven amphibian species, none of which are conservation significant.

The results of the DBCA Threatened and Priority Fauna database search are mapped in Figure 7 and database search results are presented in Appendix F.

#### 4.2.2 Post-Survey Likelihood of Occurrence

The post-survey likelihood of occurrence assessment found that:

- One species has a high likelihood of occurrence within the Survey Area
- Four species have a medium likelihood of occurrence within the Survey Area
- The remaining species has a low likelihood of occurrence within the Survey Area.

The results of the likelihood of occurrence assessment are presented in Appendix G.

#### 4.2.3 Fauna Habitat

One broad fauna habitat was identified and mapped within the Survey Area (Figure 8). Open Mulga Woodland was continuous throughout the Survey Area with slight variation in the density of mulgas present with sparse understory of *Eremophila forrestii* subsp. *forrestii* mid sparse shrubland over *Eragrostis eriopoda* low sparse tussock grassland and is analogous with the vegetation type. The habitat quality was found to be Very Good to Good throughout the Survey Area.

A description, area of extent within the Survey Area and a representative photo is provided for the fauna habitat in Table 7. Site sheets for each habitat assessment are shown in Appendix H.

**Table 7: Fauna Habitat within the Survey Area**

| Fauna Habitat  | Area (ha) | Area (%) | Description  | Representative Photo   |
|----------------|-----------|----------|--|--|
| Mulga woodland | 6.7       | 100      | Mixed <i>Acacia</i> spp. low open woodland over <i>Eragrostis eriopoda</i> low sparse tussock grassland. |  |

#### 4.2.4 Fauna Records

##### 4.2.4.1 Sightings

One native bird species from one family, the Crested Pigeon (*Ocyphaps lophotes*), was observed within the Survey Area. One introduced mammal species from one family was also observed, the Rabbit (*Oryctolagus Cuniculus*).

#### 4.2.4.2 Secondary Evidence

One native mammal species was identified based on scat the Short-beaked Echidna (*Tachyglossus aculeatus*), and one introduced species was identified based on scat, the Cat (*Felis catus*).

One reptile genus, *Varanus* sp., was tentatively identified based on scats and diggings, but was unable to be identified at the species level due to the lack of an actual observation.

#### 4.2.5 Conservation Significant Fauna

No fauna species of conservation significance (Threatened or Priority), or evidence of these species such as tracks, scats, nest, diggings, burrows, or direct sightings were recorded within or directly surrounding the Survey Area.

## 5 Discussion

### 5.1 Flora and Vegetation

#### 5.1.1 Flora Composition

The suite of flora taxa recorded during the survey was considered typical for Eastern Murchison (MUR01) subregion and aligned with the database search results obtained.

#### 5.1.2 Survey Adequacy

The flora and vegetation survey effort was in accordance with the scope of works, and appropriate for a reconnaissance flora and vegetation survey in the region. At least one relevé was sampled within each vegetation type within the Survey Area. The inventory of vascular flora and records of weed species was compiled using site data and opportunistic observations made while sampling sites and traversing between sites.

In addition, the targeted flora survey was considered appropriate for the Survey Area and was searched using meandering traverses, however additional flora taxa may be recorded with additional survey effort due to seasonal conditions and flowering periods.

#### 5.1.3 Flora of Conservation Significance

##### 5.1.3.1 Threatened and Priority Flora

No Threatened flora species pursuant to the EPBC Act and/or gazetted as Threatened flora pursuant to the BC Act, or DBCA Priority listed flora were identified by the database searches or recorded within the Survey Area.

No Priority Flora were identified by the pre-survey likelihood of occurrence assessment as having a high or medium likelihood of occurrence. Following the survey, the likelihood of occurrence assessment was updated, and no Priority Flora taxa were considered to have a high or medium likelihood of occurrence within the Survey Area.

Two taxa encountered and/or collected during the field survey were sterile and could not be confidently identified. These were carefully checked at the WAH and none were analogous to Threatened or Priority flora taxa identified by the database searches.

##### 5.1.4 Introduced Flora

One weed species, *\*Cenchrus ciliaris*, was recorded in the Survey Area. *\*Cenchrus ciliaris* has a legal status of Permitted – s11 and does not have an assigned control category.

*\*Cenchrus ciliaris* (Buffel Grass) is a tufted perennial grass to 1.5 m, native to Asia, Africa, and the Americas. The species flowers between February to October, and the flowers are purple.

*\*Cenchrus ciliaris* grows in white, red, or brown sand, stony red loam, or black cracking clay (Western Australian Herbarium, 2021). The species is naturalised throughout mainland Australia as it is an important pasture species (Identic Pty Ltd, 2016).

### 5.1.5 Vegetation Types

The vegetation types described in the Survey Area were correlated with the Beard (1976) and Shepherd *et al.* (2002) broad vegetation systems associations by examining similarities in vegetation descriptions. Differences exist with the terminology used in the descriptions as they are based on different methods of categorising and characterising vegetation types, and the different spatial scale of the analysis (i.e. region vs. local scale).

P1 and P2 are considered to be broadly representative of the Laverton 28, low woodland, open low woodland or sparse woodland of Mulga. Laverton 28 is well represented at the State, regional and sub-regional levels all having over 90% of the pre-European extent remaining. P1 and P2 appear to be well represented beyond the Survey Area based on aerial imagery.

No vegetation representative of any Commonwealth or State TECs, or DBCA listed PECs, was recorded in the Survey Area.

## 5.2 Vertebrate Fauna

### 5.2.1 Species Recorded within the Survey Area

No fauna species of conservation significance were recorded during the survey.

### 5.2.2 Post-Survey Likelihood of Occurrence

#### 5.2.2.1 High Likelihood of Occurrence

One species of conservation significance, the Peregrine Falcon, was considered to have a high likelihood of occurrence within the Survey Area.

#### **Peregrine Falcon (*Falco peregrinus*) – Other Specially Protected Fauna**

The Peregrine Falcon is an uncommon but wide-ranging bird across Australia (Barrett *et al.*, 2003). It occurs mainly along rivers and ranges as well as wooded watercourses and lakes. It nests primarily on cliffs, granite outcrops and quarries, although is also known to occupy existing raptor and corvid stick nests (Menkhorst *et al.*, 2017). The diet of the Peregrine Falcon has been well studied and primarily includes flocking species such as parrots, pigeons and on the east coast, European Starlings (Olsen and Fuentes, 2008). Given the extent of the Peregrine Falcon's distribution, wide range of habitat preferences and high mobility, it is unlikely the species will rely on habitats within the Survey Area. However, given the presence of a record within 1.5 km of the Survey Area it is likely that it will fly over, perch or hunt in the area.

#### 5.2.2.2 Medium Likelihood of Occurrence

Four conservation significant fauna species are considered to have a medium likelihood of occurrence within the Survey Area. Each of the species listed above are discussed below.

#### **Oriental Plover (*Charadrius veredus*) – Migratory, Marine**

Oriental Plovers arrive on northern Australian soil during the wet season preferring thinly vegetated plains and grasslands. They can also be found where dense vegetation has recently

been burnt (Morcombe, 2003; Menkhorst *et al.*, 2017). The Oriental Plover is considered to have a medium likelihood of occurrence due to the presence of an existing record 17 km from the Survey Area. This species may use the habitat within the survey area for perching and foraging.

#### **Grey Falcon (*Falco hypoleucus*) – Migratory**

The Grey Falcon is an elusive and endemic bird of the arid interior (Schoenjahn, Pavey and Walter, 2019). It distributed sparsely over Australia's arid and semi-arid zones and is absent from Cape York Peninsula, south of the Great Dividing Range in Victoria, and south of 26°S in Western Australia (Johnstone and Storr, 1998; BirdLife International, 2016). The Grey Falcon is restricted largely to areas of the highest annual average temperatures where there is average annual rainfall of less than 500 mm. It favours lightly timbered and untimbered lowland plains that are crossed by tree lined watercourses, but frequents other habitats, including grassland and sand dune habitats (Johnstone and Storr, 1998; BirdLife International, 2016). The species was considered as having a medium likelihood of occurrence due to the presence of a record 60 km away and the high chance that it may fly over or perch within the survey area. There is also suitable foraging habitat nearby.

#### **Princess Parrot (*Polytelis alexandrinae*) – Priority 4, Vulnerable**

The Princess Parrot is a slim to medium-sized parrot that inhabits sand dunes and sand flats in the arid zone of western and central Australia (Higgins, 1999). It occurs in open savanna woodlands and shrublands that usually consist of scattered stands of Eucalyptus and Casuarina, an understorey of shrubs such as Acacia and Cassia amongst others and a ground cover dominated by Triodia species (Morcombe, 2003; Garnett, Szabo and Dutson, 2011). Considered medium likelihood due to the presence of a record 8 km away and the chance that it may fly over the survey area.

#### **Woma (*Aspidites ramsayi*) – Priority 1 (southwest subpopulation)**

The Woma is a largely nocturnal snake, which seeks shelter in hollow logs or under leaf debris during the day and preys upon a variety of terrestrial vertebrates during the night. The Woma is found in the west and center of Australia, from Western Australia through southern Northern Territory and northern South Australia to southern Queensland north-western New South Wales (Cogger, 2014). Throughout this region, they typically occur within woodlands, heaths and shrublands, often with spinifex. It shelters mainly in abandoned monitor and mammal burrows and in soil cracks (Wilson and Swan, 2017). Considered medium likelihood due to the presence of suitable habitat 'mammal burrows' within the survey area. The nearest record of this species is 95 km away but due to the cryptic nature of this species (being nocturnal), it could be found within the survey area if a higher level of survey effort is applied (Department of Environment and Conservation, 2012).

### 5.2.3 Fauna Habitat

All fauna habitats identified in the Survey Area during the field survey are common throughout the surrounding remnant vegetation areas (bushland surrounding the Survey Area) and also common throughout the overall bioregion and subregion.

The fauna habitats that occur within the Survey Area provide a range of values to fauna as refuge, foraging and breeding habitat. All habitats are continuous and extensive outside of the Survey Area, and habitats within the Survey Areas are not critical to maintain overall habitat connectivity.

## 6 Conclusion

### Flora and Vegetation

- No Threatened flora species pursuant to the EPBC Act 1999 and/or gazetted as Threatened pursuant to the BC Act, or DBCA listed Priority flora were recorded during the reconnaissance and targeted flora survey.
- One introduced species, *\*Cenchrus ciliaris*, was recorded during the survey. *\*Cenchrus ciliaris* is not a listed DP or WoNS
- Two vegetation types were mapped within the Survey Area. None represent any Commonwealth or State listed TECs or PECs.

### Vertebrate Fauna

- No vertebrate fauna species of conservation significance were recorded during the basic vertebrate fauna survey
- One species of conservation significance was considered to have a high likelihood of occurrence within the Survey Area and four species of conservation significance were considered to have a medium likelihood of occurrence within the Survey Area
- The Survey Area contains suitable habitat for conservation significant fauna species; however, the fauna habitat occurs in relatively small extents within the Survey Area and occurs outside the Survey Area, therefore conservation significant fauna species are unlikely to rely on habitat within the Survey Area.

## 7 References

- Barrett, G. et al. (2003) *The new atlas of Australian birds*. Melbourne, Australia: Royal Australasian Ornithologists Union.
- Beard, J. S. (1976) *Vegetation survey of Western Australia. Western Australia 1: 1 000 000 vegetation series. Design and cartography by Dept. of Geography, University of W.A.*
- BirdLife International (2016) *Falco hypoleucus*. *The IUCN Red List of Threatened Species 2016*. Available at: <https://www.iucnredlist.org/species/22696479/93566768#text-fields> (Accessed: 24 February 2020).
- Bureau of Meteorology (2021) *Monthly climate data statistics*. Available at: [www.bom.gov.au/climate/data](http://www.bom.gov.au/climate/data).
- Cogger, H. G. (2014) *Reptiles and amphibians of Australia*. 7th edn. Melbourne, Australia: CSIRO Publishing.
- Cowan, M. (2001) *Murchison 1 (MUR1 - East Murchison Subregion)*.
- Department of Agriculture and Food WA (2012) *Soil-landscape systems of Western Australia (GIS dataset)*. Perth, Australia.
- Department of Agriculture Water and the Environment (2021a) *Protected Matters Search Tool*. Canberra, Australia. Available at: <http://www.environment.gov.au/webgis-framework/apps/pmst/pmst.jsf>.
- Department of Agriculture Water and the Environment (2021b) *Weeds of National Significance*. Available at: <https://weeds.org.au/>.
- Department of Biodiversity Conservation and Attractions (2020) *Conservation Codes for Western Australian Flora*. Available at: [https://www.dpaw.wa.gov.au/images/documents/plants-animals/threatened-species/Listings/Conservation code definitions.pdf](https://www.dpaw.wa.gov.au/images/documents/plants-animals/threatened-species/Listings/Conservation%20code%20definitions.pdf).
- Department of Biodiversity Conservation and Attractions (2021a) *DBCA - Legislated Lands and Waters (DBCA-011) GIS Dataset*.
- Department of Biodiversity Conservation and Attractions (2021b) *NatureMap*. Perth, Western Australia. Available at: <https://naturemap.dpaw.wa.gov.au/>.
- Department of Biodiversity Conservation and Attractions (2021c) *Threatened and Priority Ecological Communities database request (custom search)*. Perth, Australia.
- Department of Biodiversity Conservation and Attractions (2021d) *Threatened and Priority Fauna database request (custom search)*. Perth, Australia.
- Department of Biodiversity Conservation and Attractions (2021e) *Threatened and Priority Flora List (TPFL) database request (custom search)*. Perth, Australia.
- Department of Biodiversity Conservation and Attractions (2021f) *Western Australia Herbarium Flora Database (custom search)*. Perth, Australia.
- Department of Environment and Conservation (2012) *Woma Python Aspidites ramsayi (Macleay, 1882)*. Perth, Australia. Available at: <https://library.dbca.wa.gov.au/static/FullTextFiles/925292.pdf>.
- Department of Environment and Conservation (2013) *Definitions, categories and criteria for Threatened and Priority Ecological Communities*. Perth, Australia. Available at:

[https://www.dpaw.wa.gov.au/images/plants-animals/threatened-species/definitions\\_categories\\_and\\_criteria\\_for\\_threatened\\_and\\_priority\\_ecological\\_communities.pdf](https://www.dpaw.wa.gov.au/images/plants-animals/threatened-species/definitions_categories_and_criteria_for_threatened_and_priority_ecological_communities.pdf).

Department of Primary Industries and Regional Development (2021) *Declared plants*. Available at: <https://www.agric.wa.gov.au/organisms>.

Department of the Environment (2013) *Matters of National Environmental Significance: Significant impact guidelines 1.1*. Canberra, Australia. Available at: [http://www.environment.gov.au/system/files/resources/42f84df4-720b-4dcf-b262-48679a3aba58/files/nes-guidelines\\_1.pdf](http://www.environment.gov.au/system/files/resources/42f84df4-720b-4dcf-b262-48679a3aba58/files/nes-guidelines_1.pdf).

Department of the Environment and Energy (2016) *Interim Biogeographic Regionalisation for Australia, Version 7*. Canberra, Australia. Available at: [www.environment.gov.au/land/nrs/science/ibra/](http://www.environment.gov.au/land/nrs/science/ibra/).

Department of Water and Environmental Regulation (2014) *Environmentally Sensitive Areas - Fact Sheet, Environmentally Sensitive Areas*. Available at: [https://www.der.wa.gov.au/images/documents/your-environment/native-vegetation/Fact\\_sheets/fs24-clearing-regs\\_ESAs.pdf](https://www.der.wa.gov.au/images/documents/your-environment/native-vegetation/Fact_sheets/fs24-clearing-regs_ESAs.pdf).

Department of Water and Environmental Regulation (2018) *Hydrography Linear (Hierarchy) GIS Dataset*.

Department of Water and Environmental Regulation (2021) *Clearing Regulations - Environmentally Sensitive Areas GIS Dataset*.

Environmental Protection Authority (2016a) *Environmental Factor Guideline: Flora and Vegetation*. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/Guideline-Flora-Vegetation-131216\\_4.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Guideline-Flora-Vegetation-131216_4.pdf) (Accessed: 12 April 2019).

Environmental Protection Authority (2016b) *Technical Guidance: Flora and Vegetation surveys for Environmental Impact Assessment*. Perth, Australia. Available at: [http://www.epa.wa.gov.au/sites/default/files/Policies\\_and\\_Guidance/EPA Technical Guidance - Flora and Vegetation survey\\_Dec13.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA Technical Guidance - Flora and Vegetation survey_Dec13.pdf).

Environmental Protection Authority (2020) *Technical Guidance: Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment*. Perth, Australia. Available at: <https://www.epa.wa.gov.au/policies-guidance/technical-guidance-terrestrial-vertebrate-fauna-surveys-environmental-impact>.

Garnett, S. T., Szabo, J. K. and Dutson, G. (2011) *The action plan for Australian birds 2010*. Collingwood, Victoria.

GHD (2011) *Preliminary Environmental Impact Assessment, Flora Survey and Environmental Management Plan*.

Government of Western Australia (2019) *2018 Statewide Vegetation Statistics - Full Report*. Available at: <https://catalogue.data.wa.gov.au/dataset/dbca-statewide-vegetation-statistics/resource/b7bd60c2-bff6-4637-b213-aee4706412c7>.

Hall, N. J., McKenzie, N. L. and Keighery, G. J. (1994) 'The Biological Survey of the Eastern Goldfields of Western Australia - Part 10 - Sandstone Sir Samuel and Laverton Leonora Study Areas', *Records of the Western Australian Museum*.

Higgins, P. J. (1999) *Handbook of Australian, New Zealand and Antarctic Birds. Volume Four -*

- Parrots to Dollarbird, Handbook of Australian, New Zealand & Antarctic Birds.*
- Identic Pty Ltd (2016) *Cenchrus ciliaris - Weeds of Western Australia Fact Sheet*.
- Johnstone, R. E. and Storr, G. M. (1998) *Handbook of Western Australian birds*. Edited by D. Louise. Perth, Australia: Western Australian Museum.
- Menkhorst, P. et al. (2017) *The Australian bird guide*. Australia: CSIRO Publishing.
- Morcombe, M. (2003) *Field Guide to Australian Birds*. Archerfield: Steve Parish Publishing Pty Ltd.
- Native Vegetation Solutions (2019) *Detailed Flora and Vegetation Survey of Tims Find*.
- Olsen, J. and Fuentes, E. (2008) 'Dietary shifts based upon prey availability in Peregrine Falcons and Australian Hobbies breeding near Canberra, Australia', *Journal of Raptor Research*, 42, pp. 125–137.
- Schoenjahn, J., Pavey, C. R. and Walter, G. H. (2019) 'Ecology of the Grey Falcon Falco hypoleucus—current and required knowledge', *Emu*. doi: 10.1080/01584197.2019.1654393.
- Shepherd, D. P., Beeston, G. R. and Hopkins, A. J. M. (2002) *Native Vegetation in Western Australia Technical Report 249*. Perth, Australia.
- Terrestrial Ecosystems (2011) *Level 2 Fauna Risk Assessment for Granny Deep Project Area*.
- Terrestrial Ecosystems (2018) *Vertebrate Fauna Risk Assessment for the Granny Smith Solar Power Farm Project*.
- Terrestrial Ecosystems (2020a) *Level 2 Vertebrate Fauna Assessment - King of the Hills Project*.
- Terrestrial Ecosystems (2020b) *Vertebrate Fauna Risk Assessment - Granny Smith Tailing Storage Facility Expansion*.
- Western Australian Herbarium (2021) *FloraBase - The Western Australian Flora*. Perth, Australia. Available at: <https://florabase.dpaw.wa.gov.au>.
- Western Australian Museum (2021) *Checklist of the Terrestrial Vertebrate Fauna of Western Australia*. Available at: <http://museum.wa.gov.au/research/departments/terrestrial-zoology/checklist-terrestrial-vertebrate-fauna-western-australia>.
- Western Botanical (2013) *Leonora Rail Yard Expansion Project Level 1 Flora and Vegetation Survey*.
- Wilson, S. and Swan, G. (2017) *A complete guide to reptiles of Australia*. 5th edn, *Reptiles of Australia*. 5th edn. Australia: New Holland Publishers.

## 8 Limitations of this Report

This report is produced strictly in accordance with the scope of services set out in the contract or otherwise agreed in accordance with the contract. 360 Environmental makes no representations or warranties in relation to the nature and quality of soil and water other than the visual observation and analytical data in this report.

In the preparation of this report, 360 Environmental has relied upon documents, information, data, and analyses ('client's information') provided by the client and other individuals and entities. In most cases where client's information has been relied upon, such reliance has been indicated in this report. Unless expressly set out in this report, 360 Environmental has not verified that the client's information is accurate, exhaustive, or current and the validity and accuracy of any aspect of the report including, or based upon, any part of the client's information is contingent upon the accuracy, exhaustiveness, and currency of the client's information. 360 Environmental shall not be liable to the client or any other person in connection with any invalid or inaccurate aspect of this report where that invalidity or inaccuracy arose because the client's information was not accurate, exhaustive, and current or arose because of any information or condition that was concealed, withheld, misrepresented, or otherwise not fully disclosed or available to 360 Environmental.


Aspects of this report, including the opinions, conclusions, and recommendations it contains, are based on the results of the investigation, sampling and testing set out in the contract and otherwise in accordance with normal practices and standards. The investigation, sampling and testing are designed to produce results that represent a reasonable interpretation of the general conditions of the site that is the subject of this report. However, due to the characteristics of the site, including natural variations in site conditions, the results of the investigation, sampling and testing may not accurately represent the actual state of the whole site at all points.

It is important to recognise that site conditions, including the extent and concentration of contaminants, can change with time. This is particularly relevant if this report, including the data, opinions, conclusions, and recommendations it contains, are to be used a considerable time after it was prepared. In these circumstances, further investigation of the site may be necessary.

Subject to the terms of the contract between the Client and 360 Environmental Pty Ltd, copying, reproducing, disclosing, or disseminating parts of this report is prohibited (except to the extent required by law) unless the report is produced in its entirety including this page, without the prior written consent of 360 Environmental Pty Ltd.



# Figures



### Legend

- Survey Area
- Cadastral Lines

|   |   |   |  |
|---|---|---|--|
| <br>• LOCALITY MAP SOURCED FROM LANDGATE 2020<br>• OTHER DATA SOURCED LANDGATE 2020<br>• AERIAL PHOTOGRAPHY SOURCED LANDGATE 2020<br>(© Western Australian Land Information Authority 2020) | 0 25 50 75<br>Meters<br>1:2,500 @ A4<br>N<br><br>- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS | <b>360</b><br>environmental<br>a 10 Bermondsey St, West Leederville, 6007 WA<br>t (08) 9388 9360<br>f (08) 9381 2360<br>w www.360environmental.com.au | PROJECT ID<br>4581<br>DATE<br>04/08/2021 |
|   |   | HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51   |  |
|   |   | CREATED<br>[REDACTED]   | CHECKED<br>[REDACTED]                    |
|   |   | APPROVED<br>[REDACTED]  | REVISION<br>0                            |
| Horizon Power<br>Mt Ida Road, Leonora   |   |   |  |
| <b>Leonora Biological Survey</b>  |   |   |  |

**Figure 1**  
**Survey Area**



COPYRIGHT IN THIS DOCUMENT ENDOWS THE OWNER OF THE DOCUMENT THE EXCLUSIVE RIGHT TO REPRODUCE OR PUBLISH THE DOCUMENT AND THE ENVIRONMENTAL ACT DOES NOT HOLD ANY RESPONSIBILITY FOR THE USE OF THE DOCUMENT.

### Legend


- Survey Area
- Watercourse - minor
- Lake
- Water Reservoir

|  |  |  |                    |
|--|--|--|--------------------|
| <br>1:50,000 @ A4                                    |  | 360 environmental<br>a 10 Bernondsay St, West Leederville, 6007 WA<br>t (08) 9388 8360<br>f (08) 9381 2360<br><a href="http://www.360environmental.com.au">w www.360environmental.com.au</a> | DATE<br>04/08/2021 |
| - NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS |  |  |                    |
| <b>LOCALITY MAP</b>                                  |  |  |                    |
|  |  |  |                    |
|  |  | PROJECT ID<br>4581   |                    |
|  |  | HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51  |                    |
|  |  | CREATED   CHECKED   APPROVED   REVISION<br>[REDACTED]   [REDACTED]   [REDACTED]   0  |                    |
| Horizon Power<br>Mt Ida Road, Leonora                |  |  |                    |
| Leonora Biological Survey                            |  |  |                    |

### SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2020  
 - OTHER DATA SOURCED LANDGATE 2020  
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2020  
 (© Western Australian Land Information Authority 2020)

**Figure 2**  
**Hydrography**





### Legend


- Survey Area
- Cadastral Lines
- Relevé
- Fauna Habitat Assessment
- GPS Tracks

|  |   |          |
|--|---|----------|
|  | a 10 Bernondsay St, West Leederville, 6007 WA<br>(08) 9388 8360<br>(08) 9381 2360<br><a href="http://www.360environmental.com.au">www.360environmental.com.au</a> |          |
| DATE 17/08/2021                                      |   |          |
| PROJECT ID 4581                                      |   |          |
| HORIZONTAL DATUM AND PROJECTION GDA 1994 MGA Zone 51 |   |          |
| CREATED  | CHECKED   | APPROVED |
| REVISION 0   |   |          |
| Horizon Power<br>Mt Ida Road, Leonora                |   |          |
| Leonora Biological Survey                            |   |          |

- LOCALITY MAP SOURCED FROM LANDGATE 2020  
- OTHER DATA SOURCED LANDGATE 2020  
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2020  
(© Western Australian Land Information Authority 2020)

Figure 3  
Survey Effort






### Legend

- Survey Area
- Cadastral Lines
- Introduced Flora
  - \**Cenchrus ciliaris*

|  |  |  |  |
|--|--|--|--|
| <br>SLIP ENABLER   |  | a 10 Bernondsay St, West Leederville, 6007 WA<br>t (08) 9388 8360   f (08) 9381 2360<br><a href="http://www.360environmental.com.au">www.360environmental.com.au</a> |  |
| 0 25 50 75 N<br>Meters<br>1:2,500 @ A4   |  | PROJECT ID<br>4581      DATE<br>04/08/2021   |  |
| - NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS   |  |  |  |
| <b>LOCALITY MAP</b>  |  |  |  |
|  |  |  |  |
| HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51                                      |  |  |  |
| CREATED   CHECKED   APPROVED   REVISION<br>[Redacted]   [Redacted]   [Redacted]   [Redacted] |  |  |  |
| Horizon Power<br>Mt Ida Road, Leonora  |  |  |  |
| Leonora Biological Survey  |  |  |  |

**Figure 5**  
Introduced Flora



### Legend


- Survey Area
- Cadastral Lines
- Vegetation Types and Condition
  - P1 (Very Good Condition)
  - P2 (Good Condition)

**SLIP ENABLER**

- LOCALITY MAP SOURCED FROM LANDGATE 2020  
 - OTHER DATA SOURCED LANDGATE 2020  
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2020  
 (© Western Australian Land Information Authority 2020)

|   |  |   |
|---|--|---|
| <br>Meters<br>1:2,500 @ A4  |  | N   |
| - NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS  |  |   |
| <b>LOCALITY MAP</b>   |  |   |
|   |  |   |
| <b>360</b><br>environmental a 10 Bernondsay St, West Leederville, 6007 WA<br>t (08) 9388 8360<br>f (08) 9381 2360<br><a href="http://www.360environmental.com.au">w www.360environmental.com.au</a> |  | DATE<br>04/08/2021                                      |
| PROJECT ID<br><b>4581</b>   |  | HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51 |
| CREATED<br>CHECKED<br>APPROVED<br>REVISION  |  |   |
| Horizon Power<br>Mt Ida Road, Leonora   |  |   |
| Leonora Biological Survey   |  |   |

**Figure 6**  
Vegetation Types and Condition




### Legend

- Survey Area
- DBCA Threatened and Priority Fauna**
- Brush-tailed mulgara (P4)
- Common Sandpiper (MI)
- Common greenshank, greenshank (MI)
- Glossy ibis (MI)
- Gull-billed tern (MI)
- Hooded plover, hooded dotterel (P4)
- Long-tailed dunnart (P4)
- Malleefowl (VU)
- Pacific golden plover (MI)
- Peregrine falcon (OS)
- Poseidon slater (P1)
- Red knot (EN)
- Sharp-tailed sandpiper (MI)
- Streaked shearwater (MI)
- Woma (southwest subpop.) (P1)
- Wood sandpiper (MI)

|   |            |  |
|---|------------|--|
| <br>Meters<br>1:1,000,000 @ A4                          |            | <b>360</b><br>environmental<br>a 10 Bermondsey St, West Leederville, 6007 WA<br>(08) 9388 8360<br>(08) 9381 2360<br><a href="http://www.360environmental.com.au">www.360environmental.com.au</a> |
| PROJECT ID  | DATE       |  |
| 4581  | 04/08/2021 |  |
| HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51 |            |  |
| CREATED   | CHECKED    | APPROVED   |
| REVISION  |            |  |
| Horizon Power<br>Mt Ida Road, Leonora                   |            |  |
| Leonora Biological Survey                               |            |  |

**LOCALITY MAP**

**Figure 7**  
DBCA Threatened and Priority Fauna Locations



### Legend

- Survey Area
- Cadastral Lines
- Fauna Habitat
- Mulga woodland

|   |   |
|---|---|
| <br>PROJECT ID<br>4581<br>HORIZONTAL DATUM AND PROJECTION<br>GDA 1994 MGA Zone 51<br>CREATED<br>CHECKED<br>APPROVED<br>REVISION | 360 Environmental a 10 Bernondsay St, West Leederville, 6007 WA<br>t (08) 9388 8360<br>f (08) 9381 2360<br><a href="http://www.360environmental.com.au">w www.360environmental.com.au</a><br>DATE<br>17/08/2021<br><br><b>LOCALITY MAP</b><br><br>Leonora Biological Survey |
|---|---|

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2020  
 - OTHER DATA SOURCED LANDGATE 2020  
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2020  
 (© Western Australian Land Information Authority 2020)

Figure 8  
Fauna Habitat



# Appendices

## **Appendix A**

### **Flora Literature Review**

| Report  | Project Area                        | Survey Timing | Survey Effort  | Conservation Significant Ecological Communities | Conservation Significant Flora | Introduced Flora   |
|---|-------------------------------------|---------------|--|---|--------------------------------|--|
| Detailed Flora and Vegetation Survey of Tims Find (Native Vegetation Solutions, 2019)                   | 83.6 km west of the Survey Area     | July 2019     | Detailed Flora and Vegetation Survey:<br>• Thirty quadrats | None recorded.                                  | None recorded.                 | • <i>Lactuca serriola</i><br>• <i>Sisymbrium irio</i>  |
| Leonora Rail Yard Expansion Project Level 1 Flora and Vegetation Survey (Western Botanical, 2013)       | 0.6 km northeast of the Survey Area | January 2013  | Reconnaissance Flora and Vegetation Survey                 | None recorded.                                  | None recorded.                 | • <i>Cenchrus ciliaris</i><br>• <i>Centauraea melitensis</i><br>• <i>Citrullus amarus</i><br>• <i>Eragrostis curvula</i><br>• <i>Erigeron bonariensis</i><br>• <i>Salvia verbenaca</i> |
| Preliminary Environmental Impact Assessment, Flora Survey and Environmental Management Plan (GHD, 2011) | 51.9 km east of the Survey Area     | November 2010 | Reconnaissance Flora and Vegetation Survey                 | None recorded                                   | None recorded.                 | • <i>Cenchrus ciliaris</i><br>• <i>Citrullus colocynthis</i><br>• <i>Rumex vesicarius</i><br>• <i>Salvia verbenaca</i>   |

| Conservation significant flora or vegetation          |                                 |                                     |                                 |
|---|---------------------------------|-------------------------------------|---------------------------------|
|   | 33.6 km west of the Survey Area | 0.6 km northeast of the Survey Area | 51.9 km east of the Survey Area |
| T   | ★                               |                                     |                                 |
| Ricinocarpos brevis                                   |                                 |                                     |                                 |
| P1  | ★                               |                                     |                                 |
| Drosera eremaea                                       | ★                               |                                     |                                 |
| Frankenia georgei                                     |                                 | ★                                   |                                 |
| Jacksonia lanicarpa                                   | ★                               |                                     |                                 |
| Ptilotus tetrandrus                                   | ★                               |                                     |                                 |
| Stenanthemum patens                                   | ★                               |                                     |                                 |
| Tecticornia sp. Lake Way (P. Armstrong 05/961)        |                                 | ★                                   | ★                               |
| P2  |                                 | ★                                   |                                 |
| Eremophila mirabilis                                  |                                 | ★                                   |                                 |
| Eucalyptus educta                                     |                                 | ★                                   |                                 |
| Hyalosperma stoevae                                   |                                 | ★                                   |                                 |
| P3  |                                 |                                     | ★                               |
| Angianthus prostratus                                 |                                 |                                     | ★                               |
| Calotis sp. Perrin Vale Station (R.J. Cranfield 7096) |                                 | ★                                   |                                 |
| Calytrix hislopii                                     |                                 | ★                                   |                                 |

| Conservation significant flora or vegetation                |                           |             |                                     |
|---|---------------------------|-------------|-------------------------------------|
| (Native Vegetation Solutions, 2019)                         | (Western Botanical, 2013) | (GHD, 2011) |                                     |
|   |                           |             | 0.6 km northeast of the Survey Area |
|   |                           |             | 51.9 km east of the Survey Area     |
|   |                           |             | 83.6 km west of the Survey Area     |
| <i>Calytrix praecipua</i>                                   | ★                         |             |                                     |
| <i>Chrysocephalum apiculatum</i> subsp. <i>norsemanense</i> |                           | ★           |                                     |
| <i>Cratystylis centralis</i>                                | ★                         |             |                                     |
| <i>Eremophila annosicaulis</i>                              | ★                         |             |                                     |
| <i>Eremophila simulans</i> subsp. <i>megacalyx</i>          | ★                         |             |                                     |
| <i>Eremophila veronica</i>                                  |                           | ★           |                                     |
| <i>Goodenia lyrate</i>                                      |                           |             | ★                                   |
| <i>Grevillea subterlineata</i>                              | ★                         |             |                                     |
| <i>Hybanthus floribundus</i> subsp. <i>chloroxanthus</i>    |                           | ★           |                                     |
| <i>Micromyrtus serrulata</i>                                |                           | ★           |                                     |
| <i>Notisia intonsa</i>                                      |                           | ★           |                                     |
| <i>Olearia mucronata</i>                                    |                           | ★           |                                     |
| <i>Philotheca coateana</i>                                  | ★                         |             |                                     |
| <i>Philotheca deserti</i> subsp. <i>brevifolia</i>          |                           | ★           |                                     |
| <i>Pterostylis virens</i>                                   |                           |             | ★                                   |
| <i>Tecticornia cymbiformis</i>                              |                           |             | ★                                   |
| <i>Triglochin protuberans</i>                               |                           | ★           |                                     |

### Conservation significant flora or vegetation

|                                     | 83.6 km west of the Survey Area | 0.6 km northeast of the Survey Area | 51.9 km east of the Survey Area |
|-------------------------------------|---------------------------------|-------------------------------------|---------------------------------|
| (Native Vegetation Solutions, 2019) | ★                               |                                     |                                 |
| (Western Botanical, 2013)           |                                 |                                     | ★                               |
| (GHD, 2011)                         |                                 |                                     |                                 |

✓ Denotes species was found during survey

★ Denotes species was identified by database searches during desktop assessment, which typically include an additional buffer around the Project Area, but were not found during survey

## **Appendix B**

### **Flora Database Searches**





|                              |   |   |   |  |  |            |
|------------------------------|---|---|---|--|--|------------|
| <i>Carex sanguinalis</i>     | 3 | low dome-shaped shrub 0.3 to 1 m tall by 0.4 to 1 m in diameter. Plants in late bud.  | Open low plain, fed sandy loam to clayey loam with ironstone gravel on surface.                   | Emergent (canopy cover <2%) Acacia parvocarpa growing to 4 m tall over Dwart Scrub (D) (canopy cover 10 to 30%) mixed species. | ca. 320 m S of Leeton Laverton Road, between Leeton Laverton Road, the proposed Hwy Mine. Mine access is via track ca. 5.7 km E of Leeton.                       | 15/09/2002 |
| <i>Carex sanguinalis</i>     | 3 | low shrub growing 0.5 to 0.8 m tall with a spread of 0.5 m diameter.  | Flat plain, Red sandy loam with ironstone gravel.   | Mulga Open Scrub 3-4 m tall over mixed Dwart Scrub (Mair 1977).  | 47.3 km by road E of Leeton by road to mine turn off, 730 m from Laverton Leonna Road, then 150 m E along Eddine   | 20/04/2007 |
| <i>Carex sanguinalis</i>     | 3 | low dome-shaped shrub 0.3 to 1 m tall by 0.4 to 1 m in diameter.  | Open low plain, fed sandy loam to clayey loam with ironstone gravel on surface.                   | Emergent (canopy cover <2%) Acacia parvocarpa growing to 4 m tall over Dwart Scrub (D) (canopy cover 10 to 30%) mixed species. | located ca. 320 m S of the Leeton Laverton Road, between cleared exploration gullies for the proposed Hwy Mine. Mine access is via track ca. 5.7 km E of Leeton. | 16/08/2002 |
| <i>Carex sanguinalis</i>     | 3 | low dome-shaped shrub 0.3 to 1 m tall by 0.4 to 1 m in diameter. Plants in late bud.  | Open low plain, fed sandy loam to clayey loam with ironstone gravel on surface.                   | Emergent (canopy cover <2%) Acacia parvocarpa growing to 4 m tall over Dwart Scrub (D) (canopy cover 10 to 30%) mixed species. | ca. 350 m S of Leeton Laverton Road, between cleared exploration gullies for the proposed Hwy Mine. Mine access is via track ca. 5.7 km E of Leeton.             | 16/08/2002 |
| <i>Carex sanguinalis</i>     | 3 | low dome-shaped shrub 0.3 to 1 m tall by 0.4 to 1 m in diameter. Plants in late bud.  | Open low plain, fed sandy loam to clayey loam with ironstone gravel on surface.                   | Emergent (canopy cover <2%) Acacia parvocarpa growing to 4 m tall over Dwart Scrub (D) (canopy cover 10 to 30%) mixed species. | Adjacent to gullies, 700 m S of Laverton Lemon Road, 1.1 km E of Peak Shell Well, 0.5 km S of Leeton   | 17/10/2007 |
| <i>Carex sanguinalis</i>     | 3 | Shrub to 1.2 m tall with a spread to 2 m.   | Low plain, Brown sandy loam.  | Mulga scrub with Acacia parvocarpa over Dwart Scrub (Mair 1977).   | ca. 350 m S of Leeton Laverton Road, between cleared exploration gullies for the proposed Hwy Mine. Mine access is via track ca. 5.7 km E of Leeton.             | 16/08/2002 |
| <i>Carex sanguinalis</i>     | 3 | Compact shrub to 80 cm tall x 80 cm in diameter.  | Dwarf Scrub.  | Dwarf Scrub.   | Adjacent to gullies, 700 m S of Laverton Lemon Road, 1.1 km E of Peak Shell Well, 0.5 km S of Leeton   | 17/10/2007 |
| <i>Dioclea eremaea</i>       | 1 |   |   | In low mulga shrubland (2-4 m tall) with <i>Pithecellobium obliquum</i> .  | 1000 plants scattered over 500 m by 400 m area.  | 14/08/2013 |
| <i>Eremophila emodioides</i> | 3 |   |   | In low mulga shrubland (2-4 m tall) with <i>Pithecellobium obliquum</i> .  | 200 plants in 200 m area inspected - restricted habitat.   | 10/10/2008 |
| <i>Eremophila emodioides</i> | 3 |   |   | On stony banks (ironstone laterite).   | In full flower or fruit.   |            |
| <i>Eremophila emodioides</i> | 1 | Small rounded shrub 0.75 m x 1 m, calyx/canines outside, inside calmine, but base tube unsplotted.  | On stony brown dry loam.  | In open mixed <i>Acacia shublindiana</i> and over <i>Eremophila myrtacea</i> low shrubland.                                    | Same location as previously recorded. Drove macramtha subsp. <i>eremaeae</i> [P].  |            |
| <i>Eremophila emodioides</i> | 2 | Shrub to 1.5 m high, most to 0.5 m.   | On clayey sand in granite country.  | Occasional.  | Within 5 m cleared gully, 250 m N of Mt. Ascano, 9.96 km N of Copperfield, 9.1 km N of Mt. Beacon, 2.7 km W of Leeton  | 29/09/2002 |
| <i>Eremophila emodioides</i> | 2 | Ever compact shrub 2 m x 75 cm. Flowers cream, calyx pink.  | Brown sandy dry loam.   | Abundance: occasional.   | 5 km S of Leeton, Leeton road, c. 2.1 km SW of Mount Morgan's Mine turnoff   | 14/08/2013 |
| <i>Eremophila emodioides</i> | 2 | Spreading shrub 1m x 80 cm. Leaves green/grey, flowers yellow with purple spots.  | Woodland, <i>Acacia aneura</i> , <i>A. quadiimagine</i> .   | Abundance: occasional  | 121.1 m. S of Leeton, Austin Botanical District.   | //         |
| <i>Eremophila emodioides</i> | 2 | Spreading shrub to 1.5 x 2 m, but most ca. 0.8 m high.  | Stripped granite surface, red/brown boulders and over granite.                                    | Common.  | 36/10/1993   |            |
| <i>Eremophila emodioides</i> | 2 | 2 feet high.  | Rocky soil.   | Niagara Dam  | 13/09/1966   |            |
| <i>Eremophila emodioides</i> | 2 | Shrub to 1.5 m high, most to 0.5 m.   | Nature Reserve, Drainage line, laterite. Brown dry gravelly loam, clayey sand on granite country. | Healthy population. Area unlikely to ever burn due to rock soil. Some plants extend outside reserve which is fenced.           | Niagara Dam Nature Reserve, Leeton Shire   | 15/02/2006 |
| <i>Eremophila emodioides</i> | 2 | low shrubs 0.3-1 m x 0.4-1.6 m. Calyx dull carmine more rarely yellowish outside, inside calmine pale yellow in the lower part or yellowish corolla pale yellow, prominently purple spotted outside on the tubes and lobes, unspotted inside. | On stony brown dry loams, In rocky soil.  | Abundance: occasional. Population sample: 1 branch per individual.   | Niagara Dam, 1.3 km from boundary fence entity, Austin Botanical District  | 29/09/1990 |
| <i>Eremophila emodioides</i> | 2 | 60 cm high.   |   | Niagara Dam  | 13/09/1966   |            |
| <i>Eremophila emodioides</i> | 2 | Erect shrub 3 m high.   |   | Near Lake Besside, 30 miles S of Leeton  | /80/1974   |            |
| <i>Eremophila emodioides</i> | 3 | Erect open straggly shrub. Flowers purple, leaves linear, resinous.   | Stony yellow sand associated with granite.  | 4 km NW of (Hedens Weel), Niambu Station   | 16/03/1989   |            |
| <i>Eremophila emodioides</i> | 3 | Shrub 2-4 ft. flowers pale to dark violet.  |   | repent hair faun (not at all rare).  | Ca. 20 miles N of Leeton   | 5/09/1990  |
| <i>Eremophila emodioides</i> | 3 |   | Red sand.   | Abundance: occasional.   | Gibraltar Rock, Western Australian Goldfields  | 7/1885     |
| <i>Eremophila emodioides</i> | 4 | Erect male 4 m high. Stems orange.  | Flat, Red-orange clay-loam.   | Site 159, Tamroda, 40 km N of Leeton   | 7/04/2006  |            |
| <i>Eremophila emodioides</i> | 1 | Shrub 40 cm high.   | Red sand.   | Site 159, Tamroda, 30 km N of Leeton   | 7/04/2007  |            |
| <i>Frankenia georgei</i>     | 1 | Shrub 30 cm high.   | Flow line, Most damp - clay.  | 500 metres ESE of Alexander Bore, Leedamya Station   | 16/08/1988   |            |
| <i>Goodenia rotunda</i>      | 3 | Prostrate, flowers yellow.  | In red sandy loam, near claypan.  | 20 miles W of Laverton   | 23/09/1961   |            |
| <i>Grevillea ericifolia</i>  | 4 |   |   | 1.6 miles S of Menzies Road  | 28/10/1980   |            |
| <i>Grevillea incognita</i>   | 4 |   |   | Abundance: occurring in drainage line, ca 1000 plants.   | 23/08/1997   |            |
| <i>Grevillea incognita</i>   | 4 |   | Greenstone outcrop.   | Mount Roberts, 100 m W of track running N-S, along range   | 19/12/1990   |            |

|  |   |  |   |                  |  |  |            |
|--|---|--|---|------------------|--|--|------------|
| <i>Grevillea incrassata</i>                          | 4 | Bush 1m.   | Along small creek.  | In Acacia scrub. | Abundance: common.   | 2 km W of Mount Clifford, 5° E of Tonic Lake, Bellard, 20 km NW of Menzies   | 16/03/1981 |
| <i>Grevillea subeflexa</i>                           | 3 |  | Iland in salt lake. White sand over limestone.  |                  |  | E-side of Lake Bellard, boundary of Riverine Station and Adelie Station  | 14/08/1988 |
| <i>Grevillea subeflexa</i>                           | 3 | Erect compact shrub 3 m high. Pollen presenter oblique.  | White-grey clv. / Cop. dune.  |                  | Abundance: frequent.   | Population structure: mature, 75% flowering, condition of plants: healthy, Ecological disturbance: isolated Avicula contexts and Eadiinae sp. bushy grasses. |            |
| <i>Hemigelia exilis</i>                              | 4 |  | Isolated Acacia gaudichiana, A. aneura tall shrubs. Melia to tall Barringtonia polis isolated shrubs. Isolated low Solanum scabiosifolium and Pittosporum oblongifolium shrubs. | 50+ plants.      | Population: HE 4.  | Murin Murin ca 60 km leonora   | 9/03/2012  |
| <i>Hemigelia exilis</i>                              | 4 | Erect multistemmed shrub 0.75 m high. Fruiting.  | Hill slope, red laterite over ultramafic.   | abundant.        | Population: HE 4.  | Gleichenia/Maria Station, E of Leonora   | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect multistemmed shrub 2 m high. Flowers mauve, fruit developing.  | Edge of brachauka. Red laterite, ultramafic.  | occasional.      | Less than 50 plants. Population: M 9.  | Comprefield Bara Station (Newcrest) Mount Mba, 5.5 km SW of Leonora  | 15/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect, multistemmed shrub 0.75 m high. Flowers mauve, fruit developing.  | Above creek on mixed bank. Feldspar-quartz conglomerate over granite. Red ferric clay.  |                  | Growing in the shade of trees. Some plants as tall as 2 m. Growing, others not. Rooted in riverbank, others not. |  |            |
| <i>Hemigelia exilis</i>                              | 4 | Erect, multistemmed shrub 0.75 m high. Flowers mauve, young fruits, mature fruits brown.                                     | Fair ground above a creek. Quartz-feldspar gravel conglomerate over granite. Red ferric clay.   |                  | Population: HE 8.  | Poison Creek, on Leonora - Agnew road  | 13/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Compact shrub 1.2-1.5 m. Grows very pale whilst blue, base of medial lobe of lower lip whitish-yellow spotted brownish tail. | Red brown clay flats. Brown clay/leam.  |                  | Population: HE 8. Often in small non-estuarine streams along the population.                                     | Ridge to Melville Station, 8 km from Leonora - Leekin Road   | 26/10/1993 |
| <i>Hemigelia exilis</i>                              | 4 | Woody shrub 1.25 m high, flowers pale cream, many stamens.   | Unseen, lateritic chert.  |                  | 17 plants on 22/9/1995; recovery more than 200 plants in population.   | Ridge to Melville Station - Agnew Road   | 13/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect, multistemmed shrub 1.25 m high.   | Red sandy clay over laterite, young ultramafic soil.  |                  | Most of the plants were sterile, few with immature fruits. Population: HE 7.                                     | Minara Station   | 20/09/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Low plain gently sloping into drainage gullies and creeks. Residual laterite formation. Ferruginous lateritic colluvium.  |                  | Wilson Creek, W of Tonic/Bore Mining Centre, Leonora, Goldfields-Esperance.                                      | Wilson Creek, W of Tonic/Bore Mining Centre, Leonora, Goldfields-Esperance.  | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Low plains, gently sloping into drainage gullies and creeks.  |                  | Edge of population has track through it.   | Minara Station, between Leonora and Laverton   | 28/05/1996 |
| <i>Hemigelia exilis</i>                              | 4 | Erect shrub 1-1.5 m high. Flowers mauve, occasionally white. Multistemmed shrub brown base.                                  | Hedge, fire-resistant belt, saline.   | 20 plants.       | Wilson Creek, N of Minimine Well, Leonora  | Wilson Creek, N of Minimine Well, Leonora  | 19/08/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect compact shrub.   | Unseen, lateritic, greenstone and chert.  |                  | C. 16 seedlings and 25 along track margin or in centre.  | 1 km N of Bullock Hole Creek, Leonora  | 20/09/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect multistemmed shrub 75 cm high. Fruits immature and mature.   | Plain.  |                  | Station near Yenda Station Boundary  | Wilson Creek, N of Minimine Well, Leonora  | 21/07/1989 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Slope, red laterite, ultramafic.  |                  | Minara Station, E of Leonora   | Minara Station, E of Leonora   | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect multistemmed shrub 1 m high. Fruiting.   | Low plain adjacent to creek, sloping into drainage gullies and creeks. Ferruginous lateritic colluvium.   |                  | Wilson Creek, W of Tonic/Bore Mining Centre, Leonora, Goldfields-Esperance.                                      | Wilson Creek, W of Tonic/Bore Mining Centre, Leonora, Goldfields-Esperance.  | 19/08/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Hill slope, red laterite over ultramafic.   |                  | Population: HE 6.  | Minara Station   | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Spreading shrub with pink to white flowers growing up to 1.5 m tall. Fruiting.   | Low hard pan residue with thin gravelly soil.   |                  | "Keep it dark" gold prospect, c. 10 km E of Minara Station Home-track.   | Minara Station, E of Leonora   | 22/09/1998 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Hilly, red laterite, ultramafic.  |                  | Population: HE 3.  | Gleichenia/Maria Station, E of Leonora   | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Low plains gently sloping into drainage gullies and creeks.   |                  | Population: HE 6.  | Marshall Creek, E of Hebron Well, Leonora  | 10/06/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub 1-1.5 m.   | Low plains, gently sloping into drainage gullies and creeks.  |                  | Population: HE 5.  | Wilson Creek, N of Minimine Well, Leonora  | 08/06/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Erect, multistemmed shrub 1.25 m high. Flowers purple.   | Hill slope, red laterite, ultramafic.   |                  | Population: HE 5.  | Gleichenia Station, E of Leonora   | 14/11/1995 |
| <i>Hemigelia exilis</i>                              | 4 | Multi stemmed shrub 1.2 m tall.  | Outcropping laterite.   |                  | ca 0.5 km E of Lake Carey on Gleichenia Station  | ca 0.5 km E of Lake Carey  | 27/10/1997 |
| <i>Hemigelia exilis</i>                              | 4 | Multi stemmed shrub 1.2 m tall.  | Crest of slope with massive laterite outcropping.   |                  | Lovrise above Lake Carey.  |  | 29/10/1997 |
| <i>Hemigelia exilis</i>                              | 4 | Multi stemmed shrub, 1.2 m tall.   | Crest of slope with massive laterite outcropping.   |                  | Sturt Meadow Station, Coys Creek   | Sturt Meadow Station, Coys Creek   | 30/10/1994 |
| <i>Hemigelia exilis</i>                              | 4 | Multi stemmed shrub growth 30 cm tall.   | Rocky pavement, adjacent to mid slopes of low hills. Shallow red sandy banks on pavements.  |                  | Minara Murin lease, Leonora Station  | Minara Murin lease, Leonora Station  | 30/10/1997 |
| <i>Hemigelia exilis</i>                              | 4 | Multi stemmed shrub 1 m tall. Flowers purple.  | Rocky lower slopes of hill side. Red sandy loam.  |                  | Adjacent to gidgee, 1.3 km S of Laverton   | 1.7 km by road E of Leonora to mine carrford, 920 m from Laverton road to Eridgee, then 280 m E along gidgee road, then 40 m S on edge of pavement           | 19/04/2007 |
| <i>Hemigelia exilis</i>                              | 4 | Shrub with pale blue flowers.  | Creek bed with ironstone boulders.  |                  | Lots more laterite than in previous population.  | Minara Station   | 21/09/1995 |
| <i>Hemigelia exilis</i>                              | 4 |  | Rocky pavement, adjacent to mid slopes of low hills. Shallow red sandy banks on pavements.  |                  | Lucalynta lease, ca 2 km E of Lake Carey   |  | 29/12/1997 |
| <i>Hemigelia exilis</i>                              | 4 |  | Scattered mulga to 3 m tall over mixed Open Dwarf Scrub (Mar 1977).   |                  | Sturt Meadow Station, Coys Creek   | Sturt Meadow Station, Coys Creek   | 10/2004    |
| <i>Hemigelia exilis</i>                              | 4 |  | Scattered mulga over 50 m area of pavement.   |                  |  |  |            |
| <i>Hemigelia exilis</i>                              | 4 |  | 10 scattered over 50 m area.  |                  |  |  |            |
| <i>Hemigelia exilis</i>                              | 4 |  | Adjacet to gidgee, 1.3 km S of Laverton   |                  |  |  |            |
| <i>Hemigelia exilis</i>                              | 4 |  | 10 scattered over 50 m area.  |                  |  |  |            |
| <i>Hemigelia exilis</i>                              | 4 |  | Is out of regrowth side and down centre of track, more than 50 plants.  |                  |  |  |            |
| <i>Hybanthus floribundus</i> subsp. <i>charcotii</i> | 3 | Compact rounded shrub 60 cm high x 80 cm wide. White flowers.  | Slope of a low hill. Red clay/leam. Ironstone rocks. Burn > 10 years.   |                  |  |  |            |
| <i>Hybanthus floribundus</i> subsp. <i>charcotii</i> | 3 | Multi stemmed shrub 70 cm tall.  | Laterite caprock under/ing ultramafic bedrock.  |                  |  |  |            |
| <i>Hybanthus floribundus</i> subsp. <i>charcotii</i> | 3 |  | 2 km E of Hage Bone, Yundanna station, E of Leonora   |                  |  |  |            |
| <i>Hybanthus floribundus</i> subsp. <i>charcotii</i> | 3 |  | 1.5 km W of Mount Clifford, 5° E of Tonic   |                  |  |  |            |

|   |   |  |  |  |  |            |
|---|---|--|--|--|--|------------|
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 70 cm tall.  | Lateral caprock underlying ultramafic bedrock.   | 2 km E of Haig Bone, Yundamundra Station, E of Leonora.  | 1/08/1997  |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | Outcropping limestone.   | Eucalyptus lese Glencoe Station, E of Leonora.   | 29/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 60 cm tall. Flowers white.   | Acacia aenea woodland. Solanum sisymbriifolium, Pilosella oblongata, P. hispidipes.  | Murin Murin, E of Leonora.   | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | Hakea prostrata, Acacia camilloa. Very open shrubland.   | Eucalyptus lese Glencoe Station.   | 29/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 1.2 m tall.  | On crest with massive laterite outcropping.  | Murin Murin lese Glencoe Station.  | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 70 cm tall.  | In creek bed.  | Murin Murin lese Glencoe Station.  | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 1 m tall.  | Cutting paper, <i>Canthium hispidum</i> .  | Murin Murin lese Glencoe Station.  | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 70 cm tall with white flowers.   | Very open Acacia aenea woodland. <i>Scaevola spinescens</i> , <i>Acacia aenea</i> , Pilosella oblongata.   | Glenroy Station.   | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | Pilosella oblongata, <i>Aristida contorta</i> , <i>Acacia aenea</i> , <i>A. scutinaria</i> . Woodland of <i>Acacia schaffneri</i> and <i>A. aculeata</i> .   | Crest slope with scattered heathlets.  | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall. Flowers white.   | Pilosella oblongata, <i>Hemigenia palauensis</i> , <i>Acacia tetragonophylla</i> , <i>A. aenea</i> , <i>A. camilloa</i> .  | Murin Murin lese Glencoe Station.  | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 60 cm tall. Flowers white.   | Very open woodland of <i>Acacia camilloa</i> and <i>A. tetragonophylla</i> .   | Glenroy Station.   | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | <i>Solanum sisymbriifolium</i> , Pilosella oblongata, <i>T. hebetulaeoides</i> , <i>Acacia aenea</i> woodland.   | Eucalyptus lese Glencoe Station, E of Leonora.   | 29/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | <i>Eucalyptus elaeagnifolia</i> , <i>Acacia aenea</i> woodland. <i>Eremophila oppositifolia</i> woodland. Pilosella oblongata.   | Murin Murin lese Glencoe Station, E of Leonora.  | 29/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Multi stemmed shrub 40 cm tall.  | Dodonaea viscosa, Pilosella heterophyllodes, <i>Acacia aenea</i> , <i>A. tetragonophylla</i> very open shrubland.  | Glenroy Station.   | 30/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 10 cm tall x 20 cm in diameter.  | Low slopes of hills. Orange sandy loamy with rocky lacertite.  | 50 m N of main track; 2.2 km E of Eucla Bore; 1.5 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore.                    | 10/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 30 cm tall x 20 cm in diameter.  | Outcropping laterite, quartz and dolomite.   | 50 m N of main track; 2.2 km E of Eucla Bore; 1.5 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore.   | 10/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 30 cm tall x 20 cm in diameter.  | Outcropping laterite.  | 50 m N of main track; 2.2 km E of Eucla Bore; 1.5 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore.   | 10/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 30 cm tall x 20 cm in diameter.  | Low hills. Brown red sandy loamy.  | 20 plants in 20 m area inspected.  | 20/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Shrub, multi stemmed 60 cm. Flowers white. Capsule immature greenish blue colour.                                      | Mid slopes of hills. Orange sandy clay loamy.  | 10 plants in 50 m area inspected.  | 20/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 30 cm tall x 20 cm in diameter.  | Pinchish red clayey loam along creek.  | 50 plants in 20 m area inspected.  | 20/09/2008   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub 3 m tall over mixed Dwart Scrub (Muir 1977).   | Scrub Mulliga 3 m tall (Muir 1977).  | Adjacent to fine exploration gridline; 1.8 km SE of Eucla Bore; 18.3 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore. | 29/10/1997   |            |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Shrub, multi stemmed 60 cm. Flowers white. Capsule immature greenish blue colour.                                      | <i>Acacia aenea</i> , <i>A. burkartii</i> woodland. Pilosella oblongata, <i>Acacia ramulosa</i> , Pilosella oblongata, <i>Hemigenia palauensis</i> , <i>Podocarpus falcatus</i> , <i>Pithecellobium ligustrifolium</i> .                 | Scrub Mulliga 3 m tall (Muir 1977).  | Adjacent to fine exploration gridline; 1.8 km SE of Eucla Bore; 18.3 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore; 6.7 km S.E. of Eucla Bore. | 29/10/1997 |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub growing up to 0.5 m tall with a spread of 0.4 m diam.  | Upper slopes of low hills. Red sandy loam with some rocky outcropping.   | ca 100 plants in 50 m diam. area.  | ca 100 plants in 50 m diam. area.  | 9/09/2008  |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Low shrub, multi stemmed 60 cm. Flowers white. Capsule immature greenish blue colour.                                  | Drainage line. Well drained dry red clay/cane over ironstone/quartz; 0-10% loose rock; site: pastoral lease/mining lease.  | 100'2 - 100'5.   | Population structure: mature; 100% flowering. Condition of plants: healthy. Threats: mining.   | 7/03/2012  |
| <i>Hyphantus floribundus</i> subsp. <i>chloroxanthus</i>      | 3 | Shrub 2-3.5 ft.  | Emergent Eucalyptus gongolopha growing to 6 m tall over scrub dominated by <i>Acacia</i> area 3-4 m tall over low scrub of mixed <i>Acacia</i> and <i>Fremontia</i> species; 1-2 m tall over Hummock Grass of <i>Trodia</i> based sward. | 47.3 km by road E of Leonora by road to mine turn off; 520 m from Laverton Leonora Waterhole.  | 47.3 km by road E of Leonora by road to mine turn off; 520 m from Laverton Leonora Waterhole.  | 19/09/2008 |
| <i>Microseris thysanodes</i>                                  | 1 |  | In brown sandy clay on exposed breakaway platform (degraded).  | Ca 40 km E of Lester to Lester to Leonora Road and 1 km to road along the Waterhole.   | Ca 40 km E of Lester to Lester to Leonora Road and 1 km to road along the Waterhole.   | 11/03/2004 |
| <i>Microseris sericea</i>                                     | 3 | Erect, open shrub 40 cm high with white flowers.   | Clayey loam.   | Waterhole, Melba station.  | Nugara Dam Nature Reserve.   | 12/09/1988 |
| <i>Microseris sericea</i>                                     | 3 | Low shrub to 0.5 m high.   | With <i>Fremontia</i> sp.  | abundant.  | 15.5 km W of Negera Dam.   | 6/07/1995  |
| <i>Microseris sericea</i>                                     | 3 | Erect compact shrub 50 x 40 cm. Flowers white.   | Open scrub.  | Abundance: frequent.   | 1.5 km S.E. of Alexandra Bone, Agedama Station.  | 16/09/1988 |
| <i>Microseris sericea</i>                                     | 3 | Erect, compact shrub 37 cm high with white flowers.  | In red sand.   | Golffield's, S.E. Menies.  | Golffield's, S.E. Menies.  | 10/10/1991 |
| <i>Newcastelia marginis</i>                                   | 2 |  | In aerox.  | Near Menies.   | Near Menies.   | 1/07/1991  |
| <i>Newcastelia marginis</i>                                   | 2 | Flowers violet-yellow.   |  | Checkered W.F. Blackall's collecting book - M.A. Lewington 11.03.2015.   | Mount Margaret.  | 12/09/2013 |
| <i>Obione macrocarpa</i>                                      | 3 | Shrub 2-3 ft. leaves yellow.   |  | 24 km E of Lichman Well, Vast Crown Vindifiable Rock Hole.   | Mount Margaret.  | 12/09/2013 |
| <i>Obione macrocarpa</i>                                      | 3 | Shrub 2-3 feet tall, erect and densely branched.   | On chotsoe hills.  | 300 m NW of the Menies - Sandstone Road.   | Mount Margaret.  | 12/09/2013 |
| <i>Philettia coerulea</i>                                     | 3 |  |  | 8.3 km N of the road to Mt. Ma.  | Menies.  | 1/09/1997  |
| <i>Philettia tubiflora</i>                                    | 1 | Low shrub. Flowers white, anthers exerted, leaves globular, glandular.   | On breakaway plateau screete on kaolinic decaying granite.   | 800 m by track SW of the Menies - Sandstone Road.  | This is the population Weber collected from in 1975, growing on and around an old area of mining/prospecting.  | 24/10/1989 |
| <i>Pterostylis vires</i>                                      | 3 |  | low granite sheet complex, red sandy loam margins.   | 500 m by track SW of the Menies - Sandstone Road.  | Menies.  | 2/09/1997  |
| <i>Ptilotus sp. Kooleyanie (J. Jackson &amp; B. Moye 261)</i> | 1 | Small perennial herb to 12 cm high, 12 cm wide, green flowers, small succulent basal leaves.                           | Area of dense quartz.  | 800 m by track SW of the Menies - Sandstone Road.  | Menies.  | 2/09/1997  |
| <i>Ptilotus tetrandrus</i>                                    | 1 | Flowers hermaphrodite, 4 stamens, shortly united, 2-ocular.  | On rock on rocky hill.   | 8 km SW of Negera Dam (ca 50 km N of Menies).  | Menies.  | 2/09/1997  |
| <i>Seringap exstiga</i>                                       | 1 | More or less compact shrub ca 45 cm tall. Flowers - Calyx segments blue-purple when young, pinkish-purple when mature. | team.  | At junction of Wondarra - Leonora road with Laverton road; ca 19 km from Laverton.   | Wondarra - Leonora association.  | 9/10/1993  |

|  |   |   |  |  |  |  |            |
|--|---|---|--|--|--|--|------------|
| <i>Seringapexita</i>   | T | Dwarf shrub to 30 cm.   | Flat sandplain of deep red brown sandy soil.                                 | Open tall shrubland and woodland of <i>Acacia effusa</i> and <i>Eucalyptus youngiana</i> over hummock grassland or Tridia basedowii.   | Isolated plants (<1%).                     | SWATT Sandplain Survey, Weero Pastoral Lease, survey sites SWA0902C, ca. 35.62 Km ESE (12.4 km SSW) 219.22 (Region) of Leinster and ca. Homehead                     | 30/08/2013 |
| <i>Stemonium patens</i>  | 1 | Erect shrub 100 cm high.  | Slope, Dry red sand/soil, fire 10+ Years, Collection site: mining lease.     | Shrubland, With <i>Acacia rhodophloia</i> , <i>Calytrix tetragona</i> , <i>Grevillea insignis</i> [?], <i>Acacia seminervia</i> , <i>Senna artemisioides</i> , <i>Peltoscheila oblonga</i> .   | 6-20 plants.                               | Ca. 40 km NW Leinster, Ca. 8 km NW Tamroa Miresite   | 3/05/2007  |
| <i>Stemonium patens</i>  | 1 |   |  |  |  | Site 38, Marshall Peat, 70 km N of Leinster  | 20/07/1997 |
| <i>Stemonium patens</i>  | 1 | Tangled shrub, very round shape 0.8 m x 0.7 m.                              | Iwo Beach hills.   | Open shrubland with <i>Hedysarum edulis</i> , <i>Grevillea laurina</i> (rare), very little understorey, <i>Pithecellobium obliquum</i> , <i>Pithecellobium aculeatum</i> , <i>Pithecellobium stroblii</i> , <i>Fremontia cuneata</i> , <i>Eriochne galericulata</i> sp., <i>Commelinaceae</i> , <i>Acacia rigidifolia</i> subsp. <i>rigida</i> , <i>Acaena</i> area near a salt flat, fuliginea. | Infrrequent, 50-100 plants noted.          | Also found in drainage line adjacent to this site.   | 18/06/1997 |
| <i>Stemonium patens</i>  | 1 | 100 cm high, flowers white.   | Slope, loose, clay.  |  |  | Site 39, Marshall Peat, 70 km N of Leinster  | 18/06/1997 |
| <i>Stemonium patens</i>  | 1 | Bush 0.5 m high.  | On rocky hillside.   |  |  | Tamroa   | 7/04/2006  |
| <i>Tectaria cylindrica</i>   | 3 | Bounded 30 cm high shrub. Flatly, pbs-shaped segments.                      | Calcrete area.   |  |  | To. of Mount Morgan, near Lake Carey   | 16/09/1981 |
| <i>Tectaria mediterraneum</i>  | 1 |   | Cyperosidous dunes, growing close to salt lake.                              |  |  | Map Ref. L 28589 K 6813199   | 26/09/1996 |
| <i>Tectaria sp. Lake Way (P. Armstrong 05/96)</i>                    | 1 | Shrub to 50 cm, dense succulent, foliage yellow and green.                  | Flat, clay, salt lake on playa surface at edge of lake.                      |  |  | L 2. Km SE from Mount Margaret trig on the ridge of Lake Carey   | 23/03/1996 |
| <i>Thryptomene cuneata</i>   | 2 | Shrub 0.5 ft, flowers pale pink.  | Samphires.   |  |  | Near Menies, between Kalgoorlie and Leonora  | 15/03/2004 |
| <i>Thryptomene cuneata</i>   | 2 | Shrub 1-2 m tall.   | Red sand.  |  |  | 1.5 miles N of Menies.   | 10/09/1927 |
| <i>Thryptomene cuneata</i>   | 2 | Erect, compact, perennial shrub 45 cm high x 35 cm wide. Flowers pale pink. | Plain, Brown sand, stony.  |  |  | Niagara Dam  | 11/04/2006 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | Shrub 1 m tall.   | Flat top of breakaway.   | Open mulga with scattered tall shrubs to 2 m tall over open heaths.  | Uncommon - 10 plants in suitable habitats. | Adjacent to proposed Bairdine Mine site, 80%.  | 28/09/2003 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | Shrub 1.8 m tall.   | Plateau amongst steep rocky outcrops.  | Scattered tall shrubs in deeper soil pockets, with small shrubs in other areas.  | Uncommon - 20 plants in suitable habitats. | Same population as PA.3.   | 13/10/2004 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | Shrub 1.2 m tall.   | Flat top of breakaway.   | Open mulga with scattered tall shrubs to 2 m tall over open heaths.  | Uncommon - 10 plants in suitable habitats. | Unadjusted to proposed Bairdine Mine site.   | 13/10/2004 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | Small shrub to 0.3 m tall.  |  |  |  | Approximately 36-38 km WSW mine front, 0.5-1.5 Km WSW Old Goldfields railway and Darchi Webo Road turnoff and 3.4 Km NE of Goldfields Highway, 37 Km SE of Leinster. | 13/10/2004 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | low compact shrub to 1.0 m tall.  | On top and adjacent to breakaway, Shallow red sandy soil.                    |  |  | Adj. to proposed Bairdine Mine site.   | 13/10/2004 |
| <i>Thryptomene sp. Leinster (B.L. Lipsch &amp; L.A. Craven 4362)</i> | 3 | Fruit green.  | Edge of low breakaway, Rocky brown sandy/clay loam, in red sand along creek. |  |  | 3.8 Km W of Goldfields Highway and Webster Woods a Road intersection 4.5 km S of White Bell, 30.7 km SW of Leinster  | 16/03/2005 |
| <i>Triptilothrix prothecata</i>                                      | 3 |   |  |  |  | 1.5 miles E of Malcolm   | 22/08/1961 |

**DBCA Threatened and Priority Ecological Communities Database Search Results**

| COM_ID                 | COM_NAME   | STATE_CATG | COMM_CATG | BUFFER |
|------------------------|--|------------|-----------|--------|
| Sturt Meadows Calcrete | Sturt Meadows calcrite groundwater assemblage type on Raeside palaeodrainage on Sturt Meadows Station    | Priority 1 |           | 2000   |
| Melita Calcrete        | Melita calcrite groundwater assemblage type on Raeside palaeodrainage on Melita (Sons of Gwalia) Station | Priority 1 |           | 2000   |

# NatureMap Species Report

Created By Guest user on 25/05/2021

**Current Names Only** Yes  
**Core Datasets Only** Yes  
**Species Group** All Plants  
**Method** 'By Circle'  
**Centre** 121° 19' 14" E, 28° 52' 49" S  
**Buffer** 40km

| Name ID | Species Name  | Naturalised | Conservation Code | Endemic To Query Area |
|---------|---|-------------|-------------------|-----------------------|
| 1.      | 3217 <i>Acacia aneura</i> ( <i>Mulga, Wanari</i> )                        |             |                   |                       |
| 2.      | 37260 <i>Acacia aptaneura</i>   |             |                   |                       |
| 3.      | 3248 <i>Acacia burkittii</i> ( <i>Sandhill Wattle</i> )                   |             |                   |                       |
| 4.      | 36417 <i>Acacia caesaneura</i>  |             |                   |                       |
| 5.      | 3273 <i>Acacia craspedocarpa</i> ( <i>Hop Mulga</i> )                     |             |                   |                       |
| 6.      | 32118 <i>Acacia effusifolia</i>   |             |                   |                       |
| 7.      | 36781 <i>Acacia fuscaneura</i>  |             |                   |                       |
| 8.      | 36418 <i>Acacia incurvaneura</i>  |             |                   |                       |
| 9.      | 3419 <i>Acacia ligulata</i> ( <i>Umbrella Bush, Watarka</i> )             |             |                   |                       |
| 10.     | 37240 <i>Acacia macranera</i>   |             |                   |                       |
| 11.     | 36416 <i>Acacia mulganeura</i>  |             |                   |                       |
| 12.     | 3452 <i>Acacia murrayana</i> ( <i>Sandplain Wattle</i> )                  |             |                   |                       |
| 13.     | 3473 <i>Acacia oswaldii</i> ( <i>Miljee, Nelia</i> )                      |             |                   |                       |
| 14.     | 36800 <i>Acacia pteraneura</i>  |             |                   |                       |
| 15.     | 3507 <i>Acacia quadrimarginea</i>   |             |                   |                       |
| 16.     | 3510 <i>Acacia ramulosa</i> ( <i>Horse Mulga</i> )                        |             |                   |                       |
| 17.     | 19483 <i>Acacia ramulosa</i> var. <i>linophylla</i>                       |             |                   |                       |
| 18.     | 19499 <i>Acacia ramulosa</i> var. <i>ramulosa</i>                         |             |                   |                       |
| 19.     | 3513 <i>Acacia resinimarginnea</i>  |             |                   |                       |
| 20.     | 8949 <i>Acacia sibirica</i> ( <i>Bastard Mulga</i> )                      |             |                   |                       |
| 21.     | 18424 <i>Acacia</i> sp. <i>Marshall Pool</i> (G. Cockerton 3024)          |             | P3                |                       |
| 22.     | 3577 <i>Acacia tetragonophylla</i> ( <i>Kurara, Wakalpuka</i> )           |             |                   |                       |
| 23.     | 31511 <i>Acacia victoriae</i> subsp. <i>victoriae</i>                     |             |                   |                       |
| 24.     | 1505 <i>Agave americana</i> ( <i>Century Plant</i> )                      |             |                   | Y                     |
| 25.     | 11487 <i>Alectryon oleifolius</i> subsp. <i>oleifolius</i>                |             |                   |                       |
| 26.     | 19470 <i>Aluta maisonneuvei</i> subsp. <i>auriculata</i>                  |             |                   |                       |
| 27.     | 4907 <i>Alyogyne pinoniana</i> ( <i>Sand Hibiscus</i> )                   |             |                   |                       |
| 28.     | 2372 <i>Amyema fitzgeraldii</i> ( <i>Pincushion Mistletoe</i> )           |             |                   |                       |
| 29.     | 13265 <i>Amyema miraculosa</i> subsp. <i>boormanii</i>                    |             |                   |                       |
| 30.     | 40910 <i>Androcalva luteiflora</i> ( <i>Yellow-flowered Rulingia</i> )    |             |                   |                       |
| 31.     | 7826 <i>Angianthus cornutus</i>   |             |                   |                       |
| 32.     | 7834 <i>Angianthus prostratus</i>   |             | P3                |                       |
| 33.     | 7836 <i>Angianthus tomentosus</i> ( <i>Camel-grass</i> )                  |             |                   |                       |
| 34.     | 2333 <i>Anthobolus leptomeroides</i>                                      |             |                   |                       |
| 35.     | 207 <i>Aristida contorta</i> ( <i>Bunched Kerosene Grass</i> )            |             |                   |                       |
| 36.     | 7846 <i>Asteridea atrixioides</i>   |             |                   |                       |
| 37.     | 2451 <i>Atriplex bunburyana</i> ( <i>Silver Saltbush</i> )                |             |                   |                       |
| 38.     | 17801 <i>Atriplex cephalantha</i>   |             |                   |                       |
| 39.     | 2453 <i>Atriplex codonocarpa</i> ( <i>Flat-topped Saltbush</i> )          |             |                   |                       |
| 40.     | 2476 <i>Atriplex semilunaris</i> ( <i>Annual Saltbush</i> )               |             |                   |                       |
| 41.     | 2478 <i>Atriplex spongiosa</i> ( <i>Pop Saltbush</i> )                    |             |                   |                       |
| 42.     | 2481 <i>Atriplex vesicaria</i> ( <i>Bladder Saltbush</i> )                |             |                   |                       |
| 43.     | 17246 <i>Austrostipa nitida</i>   |             |                   |                       |
| 44.     | 17251 <i>Austrostipa scabra</i>   |             |                   |                       |
| 45.     | 11726 <i>Bergia perennis</i> subsp. <i>exigua</i>                         |             |                   |                       |
| 46.     | 3722 <i>Bossiaea walkeri</i>  |             |                   |                       |
| 47.     | 4999 <i>Brachychiton gregorii</i> ( <i>Desert Kurrajong, Ngalta</i> )     |             |                   |                       |
| 48.     | 7871 <i>Brachyscome ciliaris</i>  |             |                   |                       |
| 49.     | 20170 <i>Calandrinia pleiopetala</i>                                      |             |                   |                       |
| 50.     | 48773 <i>Calandrinia quartzitica</i>                                      |             | P1                |                       |
| 51.     | 5395 <i>Callistemon phoeniceus</i> ( <i>Lesser Bottlebrush, Dubarda</i> ) |             |                   |                       |
| 52.     | 7895 <i>Calocephalus multiflorus</i> ( <i>Yellow-top</i> )                |             |                   |                       |

| Name ID | Species Name  | Naturalised | Conservation Code | 'Endemic To Query Area' |
|---------|---|-------------|-------------------|-------------------------|
| 53.     | 5398 <i>Calothamnus aridus</i>  |             |                   |                         |
| 54.     | 7903 <i>Calotis hispidula</i> ( <i>Bindy Eye</i> )                                  |             |                   |                         |
| 55.     | 7905 <i>Calotis multicaulis</i> ( <i>Many-stemmed Burr-daisy</i> )                  |             |                   |                         |
| 56.     | 5456 <i>Calytrix erosipetala</i>  |             |                   |                         |
| 57.     | 43546 <i>Calytrix hislopia</i>  | P3          |                   |                         |
| 58.     | 5473 <i>Calytrix praecipua</i>  | P3          |                   |                         |
| 59.     | 12373 <i>Calytrix uncinata</i>  |             |                   |                         |
| 60.     | 3008 <i>Carrichtera annua</i> ( <i>Ward's Weed</i> )                                |             | Y                 |                         |
| 61.     | 1742 <i>Casuarina obesa</i> ( <i>Swamp Sheoak, Kuli</i> )                           |             |                   |                         |
| 62.     | 7916 <i>Centaurea melitensis</i> ( <i>Maltese Cockspur, Malta Thistle</i> )         |             | Y                 |                         |
| 63.     | 7921 <i>Centipeda thespidioides</i> ( <i>Desert Sneezewood</i> )                    |             |                   |                         |
| 64.     | 7922 <i>Cephalipterum drummondii</i> ( <i>Pompom Head</i> )                         |             |                   |                         |
| 65.     | 1216 <i>Chamaexeros macranthera</i>   |             |                   |                         |
| 66.     | 2489 <i>Chenopodium gaudichaudianum</i> ( <i>Cottony Saltbush</i> )                 |             |                   |                         |
| 67.     | 6614 <i>Convolvulus remotus</i>   |             |                   |                         |
| 68.     | 48865 <i>Cucumis myriocarpus</i> subsp. <i>myriocarpus</i>                          |             | Y                 |                         |
| 69.     | 6747 <i>Cyanostegia angustifolia</i> ( <i>Tinsel-flower</i> )                       |             |                   |                         |
| 70.     | 20759 <i>Cylindropuntia fulgida</i> var. <i>mammillata</i>                          |             | Y                 |                         |
| 71.     | 33077 <i>Cylindropuntia imbricata</i>   |             | Y                 |                         |
| 72.     | 279 <i>Cymbopogon ambiguus</i> ( <i>Scentgrass</i> )                                |             |                   |                         |
| 73.     | 281 <i>Cymbopogon obtectus</i> ( <i>Silkyheads</i> )                                |             |                   |                         |
| 74.     | 12034 <i>Dodonaea microzyga</i> var. <i>acrolobata</i>                              |             |                   |                         |
| 75.     | 4773 <i>Dodonaea petiolaris</i>   |             |                   |                         |
| 76.     | 4775 <i>Dodonaea pinifolia</i>  |             |                   |                         |
| 77.     | 4779 <i>Dodonaea rigida</i>   |             |                   |                         |
| 78.     | 11674 <i>Dodonaea viscosa</i> subsp. <i>mucronata</i>                               |             |                   |                         |
| 79.     | 33501 <i>Dysphania cristata</i> ( <i>Crested Goosefoot</i> )                        |             |                   |                         |
| 80.     | 2500 <i>Dysphania glandulosa</i>  |             |                   |                         |
| 81.     | 2502 <i>Dysphania kalpari</i> ( <i>Rat's Tail, Kalpari</i> )                        |             |                   |                         |
| 82.     | 33479 <i>Dysphania melanocarpa</i> ( <i>Black Crumbweed</i> )                       |             |                   |                         |
| 83.     | 2506 <i>Dysphania rhadinostachya</i>  |             |                   |                         |
| 84.     | 33483 <i>Dysphania saxatilis</i>  |             |                   |                         |
| 85.     | 2511 <i>Enchytraea tomentosa</i> ( <i>Barrier Saltbush</i> )                        |             |                   |                         |
| 86.     | 12064 <i>Enchytraea tomentosa</i> var. <i>tomentosa</i> ( <i>Barrier Saltbush</i> ) |             |                   |                         |
| 87.     | 19846 <i>Enebatus eremaeus</i>  |             |                   |                         |
| 88.     | 357 <i>Enneapogon caerulescens</i> ( <i>Limestone Grass</i> )                       |             |                   |                         |
| 89.     | 365 <i>Enneapogon polyphyllus</i> ( <i>Leafy Nineawn</i> )                          |             |                   |                         |
| 90.     | 378 <i>Eragrostis dielsii</i> ( <i>Mallee Lovegrass</i> )                           |             |                   |                         |
| 91.     | 387 <i>Eragrostis lanipes</i> ( <i>Creeping Wanderrie</i> )                         |             |                   |                         |
| 92.     | 7180 <i>Eremophila alternifolia</i> ( <i>Poverty Bush</i> )                         |             |                   |                         |
| 93.     | 7189 <i>Eremophila clarkei</i> ( <i>Turpentine Bush</i> )                           |             |                   |                         |
| 94.     | 17157 <i>Eremophila compacta</i> subsp. <i>compacta</i>                             |             |                   |                         |
| 95.     | 7204 <i>Eremophila eriocalyx</i> ( <i>Desert Pride</i> )                            |             |                   |                         |
| 96.     | 7207 <i>Eremophila foliosissima</i>   |             |                   |                         |
| 97.     | 7208 <i>Eremophila Forrestii</i> ( <i>Wilcox Bush</i> )                             |             |                   |                         |
| 98.     | 15052 <i>Eremophila Forrestii</i> subsp. <i>forrestii</i>                           |             |                   |                         |
| 99.     | 29532 <i>Eremophila galeata</i>   |             |                   |                         |
| 100.    | 7211 <i>Eremophila georgei</i>  |             |                   |                         |
| 101.    | 17176 <i>Eremophila Gilesii</i> subsp. <i>variabilis</i>                            |             |                   |                         |
| 102.    | 14340 <i>Eremophila glabra</i> subsp. <i>glabra</i>                                 |             |                   |                         |
| 103.    | 14191 <i>Eremophila glabra</i> subsp. <i>tomentosa</i>                              |             |                   |                         |
| 104.    | 19560 <i>Eremophila glabra</i> subsp. <i>verrucosa</i>                              |             |                   |                         |
| 105.    | 16475 <i>Eremophila glandulifera</i>  |             |                   |                         |
| 106.    | 7219 <i>Eremophila granitica</i> ( <i>Thin-leaved Poverty Bush</i> )                |             |                   |                         |
| 107.    | 7221 <i>Eremophila homoplastica</i>   |             |                   |                         |
| 108.    | 17189 <i>Eremophila hygrophana</i>  |             |                   |                         |
| 109.    | 7230 <i>Eremophila latrobei</i> ( <i>Warty Fuchsia Bush, Mintjingka</i> )           |             |                   |                         |
| 110.    | 17576 <i>Eremophila latrobei</i> subsp. <i>latrobei</i>                             |             |                   |                         |
| 111.    | 7234 <i>Eremophila longifolia</i> ( <i>Berrigan, Tulyupurpa</i> )                   |             |                   |                         |
| 112.    | 15158 <i>Eremophila mackinlayi</i> subsp. <i>spathulata</i>                         |             |                   |                         |
| 113.    | 16362 <i>Eremophila maculata</i> subsp. <i>maculata</i>                             |             |                   |                         |
| 114.    | 7239 <i>Eremophila margaretha</i> ( <i>Sandbank Poverty Bush</i> )                  |             |                   |                         |
| 115.    | 7240 <i>Eremophila metallicorum</i>   |             |                   |                         |
| 116.    | 7242 <i>Eremophila miniata</i> ( <i>Kopi Poverty Bush</i> )                         |             |                   |                         |
| 117.    | 15003 <i>Eremophila Oldfieldii</i> subsp. <i>angustifolia</i>                       |             |                   |                         |
| 118.    | 18570 <i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>                    |             |                   |                         |
| 119.    | 7250 <i>Eremophila pantoni</i>  |             |                   |                         |
| 120.    | 48951 <i>Eremophila platycalyx</i> subsp. <i>Leonora</i> ( <i>J. Morrisey 252</i> ) |             |                   |                         |
| 121.    | 7257 <i>Eremophila punicea</i> ( <i>Crimson Eremophila</i> )                        |             |                   |                         |
| 122.    | 7261 <i>Eremophila ramiflora</i>  |             |                   |                         |

| Name ID | Species Name  | Naturalised | Conservation Code | 'Endemic To Query Area' |
|---------|---|-------------|-------------------|-------------------------|
| 123.    | 7267 <i>Eremophila scoparia</i> (Broom Bush)  |             |                   |                         |
| 124.    | 7269 <i>Eremophila serrulata</i> (Serrate-leaved Eremophila)                            |             |                   |                         |
| 125.    | 17165 <i>Eremophila simulans</i> subsp. <i>megacalyx</i>                                |             | P3                |                         |
| 126.    | 7278 <i>Eremophila veronica</i>   |             | P3                |                         |
| 127.    | 15155 <i>Eremophila youngii</i> subsp. <i>youngii</i>                                   |             |                   |                         |
| 128.    | 16485 <i>Eriachne pulchella</i> subsp. <i>dominii</i>                                   |             |                   |                         |
| 129.    | 16486 <i>Eriachne pulchella</i> subsp. <i>pulchella</i>                                 |             |                   |                         |
| 130.    | 2514 <i>Eriochiton scleroaenoides</i> (Woolly Bindii)                                   |             |                   |                         |
| 131.    | 7970 <i>Erodiocephalum acanthocephalum</i>  |             |                   |                         |
| 132.    | 4334 <i>Erodium crinitum</i> (Corkscrew)  |             |                   |                         |
| 133.    | 14377 <i>Erymophyllum ramosum</i> subsp. <i>ramosum</i>                                 |             |                   |                         |
| 134.    | 35344 <i>Eucalyptus camaldulensis</i> subsp. <i>arida</i>                               |             |                   |                         |
| 135.    | 35345 <i>Eucalyptus camaldulensis</i> subsp. <i>obtusa</i> (Blunt-budded River Red Gum) |             |                   |                         |
| 136.    | 5583 <i>Eucalyptus carnei</i> (Carne's Blackbutt)                                       |             |                   |                         |
| 137.    | 5660 <i>Eucalyptus gongylocarpa</i> (Marble Gum, Baarla)                                |             |                   |                         |
| 138.    | 5684 <i>Eucalyptus kingsmillii</i> (Kingsmill's Mallee)                                 |             |                   |                         |
| 139.    | 13058 <i>Eucalyptus leptopoda</i> subsp. <i>elevata</i>                                 |             |                   |                         |
| 140.    | 5697 <i>Eucalyptus lesouefii</i> (Goldfields Blackbutt)                                 |             |                   |                         |
| 141.    | 13037 <i>Eucalyptus loxophleba</i> subsp. <i>lissophloia</i>                            |             |                   |                         |
| 142.    | 5703 <i>Eucalyptus lucasii</i> (Barlee Box)   |             |                   |                         |
| 143.    | 5803 <i>Eucalyptus youngiana</i> (Large-fruited Mallee, Yarlariba)                      |             |                   |                         |
| 144.    | 35303 <i>Euphorbia australis</i> var. <i>subtomentosa</i>                               |             |                   |                         |
| 145.    | 42869 <i>Euphorbia porcata</i>  |             |                   |                         |
| 146.    | 12097 <i>Euphorbia tannensis</i> subsp. <i>eremophila</i> (Desert Spurge)               |             |                   |                         |
| 147.    | 16722 <i>Euryomyrtus maidenii</i>   |             |                   |                         |
| 148.    | 5191 <i>Frankenia cinerea</i>   |             |                   |                         |
| 149.    | 5200 <i>Frankenia fecunda</i>   |             |                   |                         |
| 150.    | 5201 <i>Frankenia georgei</i>   |             | P1                |                         |
| 151.    | 5206 <i>Frankenia laxiflora</i> (Loose Flowered Frankenia)                              |             |                   |                         |
| 152.    | 14297 <i>Frankenia pauciflora</i> var. <i>pauciflora</i>                                |             |                   |                         |
| 153.    | 5212 <i>Frankenia setosa</i> (Bristly Frankenia)  |             |                   |                         |
| 154.    | 11008 <i>Gilruthia osbornii</i>   |             |                   |                         |
| 155.    | 7060 <i>Glossostigma diandrum</i>   |             |                   |                         |
| 156.    | 12624 <i>Gnephosis angianthoides</i>  |             |                   |                         |
| 157.    | 7988 <i>Gnephosis arachnoidea</i> (Cobwebby-headed Gnephosis)                           |             |                   |                         |
| 158.    | 7989 <i>Gnephosis brevifolia</i> (Short-leaved Gnephosis)                               |             |                   |                         |
| 159.    | 7998 <i>Gnephosis macrocephala</i>  |             |                   |                         |
| 160.    | 8002 <i>Gnephosis tenuissima</i>  |             |                   |                         |
| 161.    | 7514 <i>Goodenia havilandii</i>   |             |                   |                         |
| 162.    | 12530 <i>Goodenia macroplectra</i>  |             |                   |                         |
| 163.    | 7527 <i>Goodenia mimuloides</i>   |             |                   |                         |
| 164.    | 7529 <i>Goodenia mueckeana</i>  |             |                   |                         |
| 165.    | 7531 <i>Goodenia occidentalis</i>   |             |                   |                         |
| 166.    | 1949 <i>Grevillea acicularis</i>  |             |                   |                         |
| 167.    | 1963 <i>Grevillea berryana</i>  |             |                   |                         |
| 168.    | 2004 <i>Grevillea extorris</i>  |             |                   |                         |
| 169.    | 2019 <i>Grevillea inconspicua</i> (Cue Grevillea)                                       |             | P4                |                         |
| 170.    | 19542 <i>Grevillea nematophylla</i> subsp. <i>supraplana</i>                            |             |                   |                         |
| 171.    | 15978 <i>Grevillea oligomera</i>  |             |                   |                         |
| 172.    | 2807 <i>Gunniosia quadrifida</i> (Sturt's Pigface)                                      |             |                   |                         |
| 173.    | 19137 <i>Hakea lorea</i> subsp. <i>loreia</i>   |             |                   |                         |
| 174.    | 2182 <i>Hakea minyma</i>  |             |                   |                         |
| 175.    | 2196 <i>Hakea preissii</i> (Needle Tree, Dandjin)                                       |             |                   |                         |
| 176.    | 17557 <i>Hakea recurva</i> subsp. <i>recurva</i>  |             |                   |                         |
| 177.    | 6176 <i>Haloragis odontocarpa</i> (Mulga Nettle)  |             |                   |                         |
| 178.    | 6180 <i>Haloragis trigonocarpa</i>  |             |                   |                         |
| 179.    | 8045 <i>Helipterum craspedioides</i> (Yellow Billy Buttons)                             |             |                   |                         |
| 180.    | 43022 <i>Hibiscus</i> sp. <i>Gardneri</i> (A.L. Payne PRP 1435)                         |             |                   |                         |
| 181.    | 15448 <i>Hyalosperma glutinosum</i> subsp. <i>venustum</i>                              |             |                   |                         |
| 182.    | 11973 <i>Hybanthus floribundus</i> subsp. <i>curvifolius</i>                            |             |                   |                         |
| 183.    | 48648 <i>Hysterobaeckea ocellata</i>  |             |                   |                         |
| 184.    | 3974 <i>Indigofera georgei</i> (Bovine Indigo)  |             |                   |                         |
| 185.    | 459 <i>Iseilema eremaeum</i>  |             |                   |                         |
| 186.    | 7397 <i>Isotoma petraea</i> (Rock Isotope, Tundiwan)                                    |             |                   |                         |
| 187.    | 4953 <i>Lawrenzia densiflora</i>  |             |                   |                         |
| 188.    | 4956 <i>Lawrenzia helmsii</i> (Dunna Dunna)   |             |                   |                         |
| 189.    | 4959 <i>Lawrenzia squamata</i>  |             |                   |                         |
| 190.    | 19727 <i>Leiocarpa semicalva</i> subsp. <i>semicalva</i>                                |             |                   |                         |
| 191.    | 19237 <i>Leiocarpa websteri</i>   |             |                   |                         |
| 192.    | 3032 <i>Lepidium muelleri-ferdinandi</i>  |             |                   |                         |

| Name ID | Species Name  | Naturalised | Conservation Code | 'Endemic To Query Area' |
|---------|---|-------------|-------------------|-------------------------|
| 193.    | 3033 <i>Lepidium oxytrichum</i>                                   |             |                   |                         |
| 194.    | 3039 <i>Lepidium platypetalum</i> ( <i>Slender Peppercress</i> )  |             |                   |                         |
| 195.    | 4061 <i>Lotus cruentus</i> ( <i>Redflower Lotus</i> )             |             |                   |                         |
| 196.    | 2396 <i>Lysiana casuarinae</i>                                    |             |                   |                         |
| 197.    | 2398 <i>Lysiana murrayi</i> ( <i>Mistletoe, Parka-Parka</i> )     |             |                   |                         |
| 198.    | 36375 <i>Lysimachia arvensis</i> ( <i>Pimpernel</i> )             |             |                   |                         |
| 199.    | 2533 <i>Maireana amoena</i>                                       | Y           |                   |                         |
| 200.    | 2536 <i>Maireana atkinsiana</i> ( <i>Bronze Bluebush</i> )        |             |                   |                         |
| 201.    | 2538 <i>Maireana carnosa</i> ( <i>Cottony Bluebush</i> )          |             |                   |                         |
| 202.    | 2539 <i>Maireana convexa</i> ( <i>Mulga Bluebush</i> )            |             |                   |                         |
| 203.    | 2543 <i>Maireana eriosphaera</i>                                  |             |                   |                         |
| 204.    | 2544 <i>Maireana georgei</i> ( <i>Satin Bluebush</i> )            |             |                   |                         |
| 205.    | 2545 <i>Maireana glomerifolia</i> ( <i>Ball Leaf Bluebush</i> )   |             |                   |                         |
| 206.    | 2556 <i>Maireana planifolia</i> ( <i>Low Bluebush</i> )           |             |                   |                         |
| 207.    | 2560 <i>Maireana pyramidata</i> ( <i>Sago Bush</i> )              |             |                   |                         |
| 208.    | 2563 <i>Maireana sedifolia</i> ( <i>Pearl Bluebush, Myall</i> )   |             |                   |                         |
| 209.    | 2566 <i>Maireana thesioides</i> ( <i>Lax Bluebush</i> )           |             |                   |                         |
| 210.    | 11662 <i>Maireana tomentosa</i> subsp. <i>tomentosa</i>           |             |                   |                         |
| 211.    | 2568 <i>Maireana trichoptera</i> ( <i>Downy Bluebush</i> )        |             |                   |                         |
| 212.    | 2569 <i>Maireana triptera</i> ( <i>Threewinged Bluebush</i> )     |             |                   |                         |
| 213.    | 2571 <i>Maireana villosa</i>                                      |             |                   |                         |
| 214.    | 12949 <i>Marsdenia australis</i>                                  |             |                   |                         |
| 215.    | <i>Marsilea</i> sp.   |             |                   |                         |
| 216.    | 20288 <i>Melaleuca interioris</i>                                 |             |                   |                         |
| 217.    | 5991 <i>Melaleuca xerophila</i>                                   |             |                   |                         |
| 218.    | 3054 <i>Menkea villosula</i>                                      |             |                   |                         |
| 219.    | 6001 <i>Micromyrtus serrulata</i>                                 |             | P3                |                         |
| 220.    | 8107 <i>Minuria cunninghamii</i> ( <i>Bush Minuria</i> )          |             |                   |                         |
| 221.    | 8110 <i>Minuria leptophylla</i> ( <i>Minnie Daisy</i> )           |             |                   |                         |
| 222.    | 4098 <i>Mirbelia rhagodioides</i>                                 |             |                   |                         |
| 223.    | 490 <i>Monachather paradoxus</i>                                  |             |                   |                         |
| 224.    | 29418 <i>Monoculus monstrosus</i>                                 | Y           |                   |                         |
| 225.    | 17925 <i>Myriocephalus oldfieldii</i>                             |             |                   |                         |
| 226.    | 14186 <i>Myriocephalus pygmaeus</i>                               |             |                   |                         |
| 227.    | 11327 <i>Nicotiana occidentalis</i> subsp. <i>hesperis</i>        |             |                   |                         |
| 228.    | 11331 <i>Nicotiana occidentalis</i> subsp. <i>obliqua</i>         |             |                   |                         |
| 229.    | 11734 <i>Nicotiana rosulata</i> subsp. <i>rosulata</i>            |             |                   |                         |
| 230.    | 17 <i>Ophioglossum lusitanicum</i> ( <i>Adders Tongue</i> )       |             |                   |                         |
| 231.    | 31799 <i>Opuntia elata</i>  | Y           |                   |                         |
| 232.    | 46205 <i>Opuntia microdasys</i>                                   | Y           |                   |                         |
| 233.    | 29276 <i>Opuntia monacantha</i> ( <i>Barbary Fig</i> )            | Y           |                   |                         |
| 234.    | 12642 <i>Ozothamnus cassiopicus</i>                               |             |                   |                         |
| 235.    | 12670 <i>Parietaria cardiostegia</i>                              |             |                   |                         |
| 236.    | 10975 <i>Paspalidium basicladum</i>                               |             |                   |                         |
| 237.    | 17206 <i>Physopsis viscidula</i>                                  |             |                   |                         |
| 238.    | 19744 <i>Pittosporum angustifolium</i>                            |             |                   |                         |
| 239.    | 7299 <i>Plantago debilis</i>                                      |             |                   |                         |
| 240.    | 7300 <i>Plantago drummondii</i> ( <i>Sago Weed</i> )              |             |                   |                         |
| 241.    | 8167 <i>Pluchea dentex</i>  |             |                   |                         |
| 242.    | 45238 <i>Podolepis aristata</i> subsp. <i>affinis</i>             |             |                   |                         |
| 243.    | 8173 <i>Podolepis capillaris</i> ( <i>Wiry Podolepis</i> )        |             |                   |                         |
| 244.    | 8174 <i>Podolepis gardneri</i>                                    |             |                   |                         |
| 245.    | 8176 <i>Podolepis kendallii</i>                                   |             |                   |                         |
| 246.    | 8177 <i>Podolepis lessonii</i>                                    |             |                   |                         |
| 247.    | 8188 <i>Pogonolepis stricta</i>                                   |             |                   |                         |
| 248.    | 581 <i>Polypogon maritimus</i> ( <i>Coast Beardgrass</i> )        | Y           |                   |                         |
| 249.    | 15822 <i>Prostanthera althoferi</i> subsp. <i>althoferi</i>       |             |                   |                         |
| 250.    | 41650 <i>Prostanthera prostantheroides</i>                        |             |                   |                         |
| 251.    | 18210 <i>Psydrax rigidula</i>                                     |             |                   |                         |
| 252.    | 18155 <i>Psydrax suaveolens</i>                                   |             |                   |                         |
| 253.    | 2690 <i>Ptilotus aervoides</i>                                    |             |                   |                         |
| 254.    | 2717 <i>Ptilotus divaricatus</i> ( <i>Climbing Mulla Mulla</i> )  |             |                   |                         |
| 255.    | 2718 <i>Ptilotus drummondii</i> ( <i>Narrowleaf Mulla Mulla</i> ) |             |                   |                         |
| 256.    | 11797 <i>Ptilotus drummondii</i> var. <i>minor</i>                |             |                   |                         |
| 257.    | 2721 <i>Ptilotus exaltatus</i> ( <i>Tall Mulla Mulla</i> )        |             |                   |                         |
| 258.    | 2727 <i>Ptilotus gaudichaudii</i>                                 |             |                   |                         |
| 259.    | 2731 <i>Ptilotus helipterooides</i> ( <i>Hairy Mulla Mulla</i> )  |             |                   |                         |
| 260.    | 2747 <i>Ptilotus obovatus</i> ( <i>Cotton Bush</i> )              |             |                   |                         |
| 261.    | 11396 <i>Ptilotus obovatus</i> var. <i>obovatus</i>               |             |                   |                         |
| 262.    | 2754 <i>Ptilotus roei</i>   |             |                   |                         |

| Name ID | Species Name  | Naturalised | Conservation Code | Endemic To Query Area |
|---------|---|-------------|-------------------|-----------------------|
| 263.    | <i>Ptilotus schwartzii</i>  |             |                   |                       |
| 264.    | <i>Ptilotus schwartzii</i> var. <i>schwartzii</i>                           |             |                   |                       |
| 265.    | <i>Quinqueremulus linearis</i>  |             |                   |                       |
| 266.    | <i>Ranunculus sessiliflorus</i> (Smallflower Buttercup)                     |             |                   |                       |
| 267.    | <i>Ranunculus sessiliflorus</i> var. <i>sessiliflorus</i>                   |             |                   |                       |
| 268.    | <i>Rhagodia eremaea</i> (Thorny Saltbush)                                   |             |                   |                       |
| 269.    | <i>Rhodanthe battii</i>   |             |                   |                       |
| 270.    | <i>Rhodanthe chrysanthemoides</i>   |             |                   |                       |
| 271.    | <i>Rhodanthe chlorocephala</i> subsp. <i>rosea</i>                          |             |                   |                       |
| 272.    | <i>Rhodanthe chlorocephala</i> subsp. <i>splendida</i>                      |             |                   |                       |
| 273.    | <i>Rhodanthe manglesii</i>  |             |                   |                       |
| 274.    | <i>Rhodanthe maryonii</i>   |             |                   |                       |
| 275.    | <i>Rhodanthe propinqua</i>  |             |                   |                       |
| 276.    | <i>Rhodanthe pygmaea</i>  |             |                   |                       |
| 277.    | <i>Rhodanthe stricta</i>  |             |                   |                       |
| 278.    | <i>Roebuckiella ciliocarpa</i>  |             |                   |                       |
| 279.    | <i>Rumex vesicarius</i> (Ruby Dock)   |             | Y                 |                       |
| 280.    | <i>Rutidosis helichrysoides</i> subsp. <i>helichrysoides</i>                |             |                   |                       |
| 281.    | <i>Samolus repens</i> (Creeping Brookweed)                                  |             |                   |                       |
| 282.    | <i>Santalum lanceolatum</i> (Northern Sandalwood, Yarnguli)                 |             |                   |                       |
| 283.    | <i>Santalum spicatum</i> (Sandalwood, Wilarak)                              |             |                   |                       |
| 284.    | <i>Scaevola spinescens</i> (Currant Bush, Maroon)                           |             |                   |                       |
| 285.    | <i>Schoenia cassiniiana</i> (Schoenia)                                      |             |                   |                       |
| 286.    | <i>Schoenoplectiella dissachantha</i>                                       |             |                   |                       |
| 287.    | <i>Sclerolaena cuneata</i> (Yellow Bindii)                                  |             |                   |                       |
| 288.    | <i>Sclerolaena densiflora</i>   |             |                   |                       |
| 289.    | <i>Sclerolaena deserticola</i>  |             |                   |                       |
| 290.    | <i>Sclerolaena eriacantha</i> (Tall Bindii)                                 |             |                   |                       |
| 291.    | <i>Sclerolaena eurotioides</i> (Fluffy Bindii)                              |             |                   |                       |
| 292.    | <i>Sclerolaena gardneri</i>   |             |                   |                       |
| 293.    | <i>Sclerolaena lanicuspis</i> (Spinach Burr)                                |             |                   |                       |
| 294.    | <i>Senecio glossanthus</i> (Slender Groundsel)                              |             |                   |                       |
| 295.    | <i>Senecio gregorii</i> (Fleshy Groundsel)                                  |             |                   |                       |
| 296.    | <i>Senecio lacustrinus</i>  |             |                   |                       |
| 297.    | <i>Senecio magnificus</i> (Showy Groundsel)                                 |             |                   |                       |
| 298.    | <i>Senna artemisioides</i>  |             |                   |                       |
| 299.    | <i>Senna artemisioides</i> subsp. <i>filifolia</i>                          |             |                   |                       |
| 300.    | <i>Senna artemisioides</i> subsp. <i>helmsii</i>                            |             |                   |                       |
| 301.    | <i>Senna artemisioides</i> subsp. <i>oligophylla</i>                        |             |                   |                       |
| 302.    | <i>Senna artemisioides</i> subsp. <i>x sturtii</i>                          |             |                   |                       |
| 303.    | <i>Senna cardiosperma</i>   |             |                   |                       |
| 304.    | <i>Senna charlesiana</i>  |             |                   |                       |
| 305.    | <i>Senna glutinosa</i> subsp. <i>chateleainiana</i>                         |             |                   |                       |
| 306.    | <i>Senna manicula</i>   |             |                   |                       |
| 307.    | <i>Senna</i> sp. Meekatharra (E. Bailey 1-26)                               |             |                   |                       |
| 308.    | <i>Sida ectogama</i>  |             |                   |                       |
| 309.    | <i>Sida</i> sp. dark green fruits (S. van Leeuwen 2260)                     |             |                   |                       |
| 310.    | <i>Solanum cleistogamum</i>   |             |                   |                       |
| 311.    | <i>Solanum ferocissimum</i>   |             |                   |                       |
| 312.    | <i>Solanum lasiophyllum</i> (Flannel Bush, Mindjulu)                        |             |                   |                       |
| 313.    | <i>Solanum orbiculatum</i> subsp. <i>orbiculatum</i> (Round-leaved Solanum) |             |                   |                       |
| 314.    | <i>Stenantherium patens</i>   |             | P1                |                       |
| 315.    | <i>Stenopetalum anfractum</i>   |             |                   |                       |
| 316.    | <i>Stenopetalum filifolium</i>  |             |                   |                       |
| 317.    | <i>Streptoglossa liatroides</i>   |             |                   |                       |
| 318.    | <i>Stylidium induratum</i> (Desert Triggerplant)                            |             |                   |                       |
| 319.    | <i>Stylidium longibracteatum</i> (Long-bracted Trigger Plant)               |             |                   |                       |
| 320.    | <i>Swainsona affinis</i>  |             |                   |                       |
| 321.    | <i>Swainsona canescens</i> (Grey Swainsona)                                 |             |                   |                       |
| 322.    | <i>Swainsona formosa</i>  |             |                   |                       |
| 323.    | <i>Swainsona rostellata</i>   |             |                   |                       |
| 324.    | <i>Synaptontha tillaeacea</i> var. <i>tillaeacea</i>                        |             |                   |                       |
| 325.    | <i>Tecticornia indica</i> subsp. <i>bidens</i>                              |             |                   |                       |
| 326.    | <i>Templetonia incrassata</i>   |             |                   |                       |
| 327.    | <i>Tetragonia eremaea</i>   |             |                   |                       |
| 328.    | <i>Teucrium teucriifolium</i>   |             |                   |                       |
| 329.    | <i>Threlkeldia diffusa</i> (Coast Bonefruit)                                |             |                   |                       |
| 330.    | <i>Trachymene ornata</i> (Spongefruit)                                      |             |                   |                       |
| 331.    | <i>Tragus australianus</i> (Small Burgrass)                                 |             |                   |                       |
| 332.    | <i>Trichanthodium skirrophorum</i>  |             |                   |                       |

| Name ID | Species Name   | Naturalised | Conservation Code | <sup>1</sup> Endemic To Query Area |
|---------|--|-------------|-------------------|------------------------------------|
| 333.    | 6727 <i>Trichodesma zeylanicum</i> ( <i>Camel Bush, Kumbalin</i> ) |             |                   |                                    |
| 334.    | 18587 <i>Triglochin nana</i>                                       |             |                   |                                    |
| 335.    | 19038 <i>Triglochin protuberans</i>                                |             | P3                |                                    |
| 336.    | 7661 <i>Velleia hispida</i> ( <i>Hispid Velleia</i> )              |             |                   |                                    |
| 337.    | 7664 <i>Velleia rosea</i> ( <i>Pink Velleia</i> )                  |             |                   |                                    |
| 338.    | 15725 <i>Verbesina encelioides</i>                                 |             | Y                 |                                    |
| 339.    | 12436 <i>Verticordia interioris</i>                                |             |                   |                                    |
| 340.    | 8265 <i>Vittadinia eremaea</i>                                     |             |                   |                                    |
| 341.    | 8268 <i>Vittadinia humerata</i>                                    |             |                   |                                    |
| 342.    | 20153 <i>Vittadinia</i> sp. <i>Earaheedy</i> (D.J. Edinger 3106)   |             |                   |                                    |
| 343.    | <i>Wahlenbergia</i> sp.  |             |                   |                                    |
| 344.    | 7393 <i>Wahlenbergia tumidifructa</i>                              |             |                   |                                    |
| 345.    | 1391 <i>Wurmbea densiflora</i>                                     |             |                   |                                    |

**Conservation Codes**

I - Rare or likely to become extinct  
 X - Presumed extinct  
 IA - Protected under international agreement  
 S - Other specially protected fauna  
 1 - Priority 1  
 2 - Priority 2  
 3 - Priority 3  
 4 - Priority 4  
 5 - Priority 5

<sup>1</sup> For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.



# EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 25/05/21 19:00:46

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



This map may contain data which are  
©Commonwealth of Australia  
(Geoscience Australia), ©PSMA 2015

[Coordinates](#)

Buffer: 50.0Km



# Summary

## Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

|   |      |
|---|------|
| <a href="#">World Heritage Properties:</a>                | None |
| <a href="#">National Heritage Places:</a>                 | None |
| <a href="#">Wetlands of International Importance:</a>     | None |
| <a href="#">Great Barrier Reef Marine Park:</a>           | None |
| <a href="#">Commonwealth Marine Area:</a>                 | None |
| <a href="#">Listed Threatened Ecological Communities:</a> | None |
| <a href="#">Listed Threatened Species:</a>                | 5    |
| <a href="#">Listed Migratory Species:</a>                 | 8    |

## Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

|  |      |
|--|------|
| <a href="#">Commonwealth Land:</a>                 | 1    |
| <a href="#">Commonwealth Heritage Places:</a>      | None |
| <a href="#">Listed Marine Species:</a>             | 11   |
| <a href="#">Whales and Other Cetaceans:</a>        | None |
| <a href="#">Critical Habitats:</a>                 | None |
| <a href="#">Commonwealth Reserves Terrestrial:</a> | None |
| <a href="#">Australian Marine Parks:</a>           | None |

## Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

|  |      |
|--|------|
| <a href="#">State and Territory Reserves:</a>    | None |
| <a href="#">Regional Forest Agreements:</a>      | None |
| <a href="#">Invasive Species:</a>                | 15   |
| <a href="#">Nationally Important Wetlands:</a>   | None |
| <a href="#">Key Ecological Features (Marine)</a> | None |

## Details

### Matters of National Environmental Significance

| Listed Threatened Species  | [ Resource Information ] |  |
|--|--------------------------|--|
| Name   | Status                   | Type of Presence                                       |
| Birds  |                          |  |
| <a href="#"><u>Falco hypoleucus</u></a><br>Grey Falcon [929]                                     | Vulnerable               | Species or species habitat may occur within area       |
| <a href="#"><u>Leipoa ocellata</u></a><br>Malleefowl [934]                                       | Vulnerable               | Species or species habitat known to occur within area  |
| <a href="#"><u>Pezoporus occidentalis</u></a><br>Night Parrot [59350]                            | Endangered               | Species or species habitat may occur within area       |
| <a href="#"><u>Polytelis alexandrae</u></a><br>Princess Parrot, Alexandra's Parrot [758]         | Vulnerable               | Species or species habitat known to occur within area  |
| Mammals  |                          |  |
| <a href="#"><u>Dasyurus geoffroii</u></a><br>Chuditch, Western Quoll [330]                       | Vulnerable               | Species or species habitat may occur within area       |
| Listed Migratory Species   |                          |  |
| * Species is listed under a different scientific name on the EPBC Act - Threatened Species list. |                          |  |
| Name   | Threatened               | Type of Presence                                       |
| Migratory Marine Birds   |                          |  |
| <a href="#"><u>Apus pacificus</u></a><br>Fork-tailed Swift [678]                                 |                          | Species or species habitat likely to occur within area |
| Migratory Terrestrial Species  |                          |  |
| <a href="#"><u>Motacilla cinerea</u></a><br>Grey Wagtail [642]                                   |                          | Species or species habitat may occur within area       |
| <a href="#"><u>Motacilla flava</u></a><br>Yellow Wagtail [644]                                   |                          | Species or species habitat may occur within area       |
| Migratory Wetlands Species   |                          |  |
| <a href="#"><u>Actitis hypoleucus</u></a><br>Common Sandpiper [59309]                            |                          | Species or species habitat known to occur within area  |
| <a href="#"><u>Calidris acuminata</u></a><br>Sharp-tailed Sandpiper [874]                        |                          | Species or species habitat may occur within area       |

| Name  | Threatened | Type of Presence                                       |
|---|------------|--|
| <a href="#"><u>Calidris melanotos</u></a><br>Pectoral Sandpiper [858]                 |            | Species or species habitat may occur within area       |
| <a href="#"><u>Charadrius veredus</u></a><br>Oriental Plover, Oriental Dotterel [882] |            | Species or species habitat may occur within area       |
| <a href="#"><u>Tringa nebularia</u></a><br>Common Greenshank, Greenshank [832]        |            | Species or species habitat likely to occur within area |

## Other Matters Protected by the EPBC Act

| Commonwealth Land [ <a href="#">Resource Information</a> ]                                       |            |  |
|--|------------|--|
| Name   | Threatened | Type of Presence                                       |
| Commonwealth Land -  |            |  |
| Listed Marine Species [ <a href="#">Resource Information</a> ]                                   |            |  |
| * Species is listed under a different scientific name on the EPBC Act - Threatened Species list. |            |  |
| Name   | Threatened | Type of Presence                                       |
| Birds  |            |  |
| <a href="#"><u>Actitis hypoleucus</u></a><br>Common Sandpiper [59309]                            |            | Species or species habitat known to occur within area  |
| <a href="#"><u>Apus pacificus</u></a><br>Fork-tailed Swift [678]                                 |            | Species or species habitat likely to occur within area |
| <a href="#"><u>Calidris acuminata</u></a><br>Sharp-tailed Sandpiper [874]                        |            | Species or species habitat may occur within area       |
| <a href="#"><u>Calidris melanotos</u></a><br>Pectoral Sandpiper [858]                            |            | Species or species habitat may occur within area       |
| <a href="#"><u>Charadrius veredus</u></a><br>Oriental Plover, Oriental Dotterel [882]            |            | Species or species habitat may occur within area       |
| <a href="#"><u>Chrysococcyx osculans</u></a><br>Black-eared Cuckoo [705]                         |            | Species or species habitat known to occur within area  |
| <a href="#"><u>Merops ornatus</u></a><br>Rainbow Bee-eater [670]                                 |            | Species or species habitat may occur within area       |
| <a href="#"><u>Motacilla cinerea</u></a><br>Grey Wagtail [642]                                   |            | Species or species habitat may occur within area       |
| <a href="#"><u>Motacilla flava</u></a><br>Yellow Wagtail [644]                                   |            | Species or species habitat may occur within            |

| Name   | Threatened | Type of Presence area                                  |
|--|------------|--|
| <a href="#"><i>Thinornis rubricollis</i></a><br>Hooded Plover [59510]          |            | Species or species habitat known to occur within area  |
| <a href="#"><i>Tringa nebularia</i></a><br>Common Greenshank, Greenshank [832] |            | Species or species habitat likely to occur within area |

## Extra Information

### Invasive Species [ Resource Information ]

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

| Name  | Status | Type of Presence                                       |
|---|--------|--|
| <b>Birds</b>  |        |  |
| <i>Columba livia</i><br>Rock Pigeon, Rock Dove, Domestic Pigeon [803]         |        | Species or species habitat likely to occur within area |
| <i>Streptopelia senegalensis</i><br>Laughing Turtle-dove, Laughing Dove [781] |        | Species or species habitat likely to occur within area |
| <b>Mammals</b>  |        |  |
| <i>Camelus dromedarius</i><br>Dromedary, Camel [7]                            |        | Species or species habitat likely to occur within area |
| <i>Canis lupus familiaris</i><br>Domestic Dog [82654]                         |        | Species or species habitat likely to occur within area |
| <i>Capra hircus</i><br>Goat [2]   |        | Species or species habitat likely to occur within area |
| <i>Equus asinus</i><br>Donkey, Ass [4]  |        | Species or species habitat likely to occur within area |
| <i>Felis catus</i><br>Cat, House Cat, Domestic Cat [19]                       |        | Species or species habitat likely to occur within area |
| <i>Mus musculus</i><br>House Mouse [120]                                      |        | Species or species habitat likely to occur within area |
| <i>Oryctolagus cuniculus</i><br>Rabbit, European Rabbit [128]                 |        | Species or species habitat likely to occur within area |
| <i>Vulpes vulpes</i><br>Red Fox, Fox [18]                                     |        | Species or species                                     |

| Name  | Status | Type of Presence<br>habitat likely to occur within<br>area |
|---|--------|--|
| <b>Plants</b>   |        |  |
| <i>Carrichtera annua</i><br>Ward's Weed [9511]  |        | Species or species habitat<br>may occur within area        |
| <i>Cenchrus ciliaris</i><br>Buffel-grass, Black Buffel-grass [20213]  |        | Species or species habitat<br>may occur within area        |
| <i>Cylindropuntia</i> spp.<br>Prickly Pears [85131]   |        | Species or species habitat<br>likely to occur within area  |
| <i>Opuntia</i> spp.<br>Prickly Pears [82753]  |        | Species or species habitat<br>likely to occur within area  |
| <i>Tamarix aphylla</i><br>Athel Pine, Athel Tree, Tamarisk, Athel Tamarisk,<br>Athel Tamarix, Desert Tamarisk, Flowering Cypress,<br>Salt Cedar [16018] |        | Species or species habitat<br>likely to occur within area  |

## Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

## Coordinates

-28.88024 121.32071

## Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [Office of Environment and Heritage, New South Wales](#)
- [Department of Environment and Primary Industries, Victoria](#)
- [Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [Department of Environment, Water and Natural Resources, South Australia](#)
- [Department of Land and Resource Management, Northern Territory](#)
- [Department of Environmental and Heritage Protection, Queensland](#)
- [Department of Parks and Wildlife, Western Australia](#)
- [Environment and Planning Directorate, ACT](#)
- [Birdlife Australia](#)
- [Australian Bird and Bat Banding Scheme](#)
- [Australian National Wildlife Collection](#)
- Natural history museums of Australia
- [Museum Victoria](#)
- [Australian Museum](#)
- [South Australian Museum](#)
- [Queensland Museum](#)
- [Online Zoological Collections of Australian Museums](#)
- [Queensland Herbarium](#)
- [National Herbarium of NSW](#)
- [Royal Botanic Gardens and National Herbarium of Victoria](#)
- [Tasmanian Herbarium](#)
- [State Herbarium of South Australia](#)
- [Northern Territory Herbarium](#)
- [Western Australian Herbarium](#)
- [Australian National Herbarium, Canberra](#)
- [University of New England](#)
- [Ocean Biogeographic Information System](#)
- [Australian Government, Department of Defence](#)
- [Forestry Corporation, NSW](#)
- [Geoscience Australia](#)
- [CSIRO](#)
- [Australian Tropical Herbarium, Cairns](#)
- [eBird Australia](#)
- [Australian Government – Australian Antarctic Data Centre](#)
- [Museum and Art Gallery of the Northern Territory](#)
- [Australian Government National Environmental Science Program](#)
- [Australian Institute of Marine Science](#)
- [Reef Life Survey Australia](#)
- [American Museum of Natural History](#)
- [Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.

---

## **Appendix C**

### **Flora Likelihood of Occurrence**

**Appendix C: Assessment of the Likelihood of Occurrence of Threatened and Priority Flora as per Desktop Assessment Database Searches surrounding the Survey Area**

Distance to Nearest Record from the Survey Area is based on a distance analysis undertaken against 2020 DBCA database. High = Suitable habitat present and records less than 5 km from the Survey Area. Medium = Suitable habitat present and records greater than 5 km from the Survey Area. Low = No suitable habitat present and/or records greater than 15 km from the Survey Area. Unknown = Insufficient information available to classify . Cr= Listed as Critically Endangered under the EPBC Act. EN = Listed as Endangered under the EPBC Act. VU = Listed as Vulnerable under the EPBC Act. T = Threatened under the BC Act. P = Priority Listed. Ranked and Listed by the DBCA. Likelihoods are assessed both pre and post survey based on knowledge of the Survey Area, nearest known records, known flowering period of flora taxa and knowledge gained from the survey effort during ground truthing.

| Species   | Conservation Status |      | Source | Distance to Nearest Record (km) | Flowering Period | Preferred Habitat | Habitat occurs within the Survey Area  | Pre-Survey Likelihood of Occurrence | Post-Survey Likelihood of Occurrence |
|---|---------------------|------|--------|---------------------------------|------------------|-------------------|--|-------------------------------------|--------------------------------------|
|   | DBCA                | EPBC |        |                                 |                  |                   |  |                                     |                                      |
| <i>Acacia websteri</i>  | P1                  |      |        | X                               | 50.9             | Jun and Dec - Jan | Red sand, clay or loam. Low-lying areas, flats.  | No                                  | Low                                  |
| <i>Anacampseros</i> sp. Eremaeana (F. Hirt, J. Hort & J. Shanks 3248) | P1                  |      |        | X                               | 58.6             | Sep               | Sand patches inside rocks, brown sandy clay, granite. Depressions in rock outcrops, breakaways, flats.     | No                                  | Low                                  |
| <i>Calandrinia quadriflora</i>  | P1                  | X    |        | X                               | 26.6             | Sep - Oct         | Brown silty sands, red-brown silty loams, Quartz derived geology. Margins of salt lakes and lake channels. | No                                  | Low                                  |
| <i>Drosera eremaea</i>  | P1                  |      |        | X                               | 96.8             | Jun - Sep         | Quartz grit and loam, winter wet creeks.   | No                                  | Low                                  |
| <i>Eremophila eversa</i>  | P1                  |      |        | X                               | 80.8             | Sep               | Only known from type specimen on Yerilla Station.  | Unknown                             | Unknown                              |
| <i>Frankenia georgii</i>  | P1                  | X    |        | X                               | 29.5             | Dec               | Rocky slopes.  | No                                  | Low                                  |
| <i>Korthalsella leucathrix</i>  | P1                  |      |        | X                               | 72.3             | Aug               | Parasitic shrub, on Acacia acuminata and A. crassipedocarpa. <sup>2</sup>                                  | No                                  | Low                                  |
| <i>Micromyrtus chrysodema</i>   | P1                  |      |        | X                               | 88.5             | Mar               | Red sands. Sandplains.   | No                                  | Low                                  |
| <i>Philotricha tubiflora</i>  | P1                  |      |        | X                               | 83.2             | Jun - Oct         | Rocky rises and hills, outcrops. <sup>2</sup>  | No                                  | Low                                  |
| <i>Ptilotus</i> sp. Kookynie (J. Jackson & B. Moyle 261)              | P1                  |      |        | X                               | 58.4             | Sep               | Rocky hills, quartz.   | No                                  | Low                                  |
| <i>Ptilotus tetrandrus</i>  | P1                  |      |        | X                               | 49.1             | Aug and Oct       | Loamy red sand, swales between dunes, low in landscape.  | No                                  | Low                                  |
| <i>Stenanthemum patens</i>  | P1                  | X    |        | X                               | 30.3             | Apr - Oct         | Red/orange stony-sandy loam. Rocky slopes and hillsides.   | No                                  | Low                                  |
| <i>Tecticornia mellinaria</i>   | P1                  |      |        | X                               | 89.9             | Sep - Oct         | Red/gypseous dunes or gypseous clay pans on margins of salt lakes.   | No                                  | Low                                  |
| <i>Tecticornia</i> sp. Lake Way (P. Armstrong 05/961)                 | P1                  |      |        | X                               | 87.1             | Unknown           | Outwash plains on margins of salt lake.  | No                                  | Low                                  |
| <i>Eremophila mirabilis</i>   | P2                  |      |        | X                               | 44.1             | Jul - Sep         | Clay sand, stony clayey loam. Granite country, lateritic breakaways.                                       | No                                  | Low                                  |
| <i>Newcastelia insignis</i>   | P2                  |      |        | X                               | 93.9             | Sep - Nov         | Red or yellow sandy soils.   | No                                  | Low                                  |
| <i>Thryptomene eremaea</i>  | P2                  |      |        | X                               | 57.3             | Jul - Sep         | Red or yellow sand. Sandplains.  | No                                  | Low                                  |

| Species   | Conservation Status |      |           | Source | Distance to Nearest Record (km) | Flowering Period       | Preferred Habitat  | Habitat occurs within the Survey Area | Pre-Survey Likelihood of Occurrence | Post-Survey Likelihood of Occurrence |
|---|---------------------|------|-----------|--------|---------------------------------|------------------------|--|---------------------------------------|-------------------------------------|--------------------------------------|
|   | DBCA                | EPBC | NatureMap |        |                                 |                        |  |                                       |                                     |                                      |
| <i>Acacia</i> sp. Marshall Pool (G. Cockerton 3024)           | P3                  |      | X         |        | 0.9                             | May                    | Low hills of gabbro, basalt and calcite.   | No                                    | Low                                 | Low                                  |
| <i>Angianthus prostratus</i>                                  | P3                  |      | X         |        | 16.1                            | Jul - Sep              | Red clay or loamy soils. Saline depressions.   | No                                    | Low                                 | Low                                  |
| <i>Baeckea</i> sp. Sandstone (C.A. Gardner s.n. 26 Oct. 1963) | P3                  |      |           |        | 94.4                            | Oct                    | Orange sand, flats, sandplains. <sup>2</sup>   | No                                    | Low                                 | Low                                  |
| <i>Calandrinia</i> sp. Menzies (F. Hort et al. FH 4100)       | P3                  |      |           | X      | 53.9                            | Apr and Aug - Oct      | Red-brown/orange clayey sands. Flat stony or hardpan plains.   | Possibly                              | Low                                 | Low                                  |
| <i>Calotis</i> sp. Pennington Station (R.J. Cranfield 7096)   | P3                  |      |           | X      | 91.9                            | Aug - Sep              | Red loam, red/orange sandy clay/loam, red clay/loam over calcareous. Plains, Banded ironstone formation outcrops, granite slopes. <sup>2</sup> | No                                    | Low                                 | Low                                  |
| <i>Calytrix hislopii</i>                                      | P3                  |      | X         |        | 36.5                            | Sep - Nov              | Latent ridges, top of breakaways and granites.   | No                                    | Low                                 | Low                                  |
| <i>Calytrix praecipua</i>                                     | P3                  |      | X         |        | 22.5                            | Jun - Jul or Sep - Nov | Skeletal sandy soils over granite or laterite. Breakaways, outcrops. <sup>2</sup>  | No                                    | Low                                 | Low                                  |
| <i>Cratystylis centralis</i>                                  | P3                  |      |           | X      | 46.3                            | Aug - Oct              | Red sandy loam with ironstone gravel. Flat plains, breakaway country.  | Possibly                              | Low                                 | Low                                  |
| <i>Eremophila amosicaulis</i>                                 | P3                  |      |           | X      | 71.4                            | Jun - Sep              | Stony ironstone soils on slopes of low rocky hills.  | No                                    | Low                                 | Low                                  |
| <i>Eremophila shoneae</i> subsp. <i>diffusa</i>               | P3                  |      |           | X      | 65                              | Aug - Oct              | Stony yellow sands associated with granites, rocky slopes.   | No                                    | Low                                 | Low                                  |
| <i>Eremophila simulans</i> subsp. <i>megacylix</i>            | P3                  |      | X         |        | 32.1                            | Aug - Oct              | Rocky and sandy clay soils. Crests and slopes of banded ironstone, sandy plains  | No                                    | Low                                 | Low                                  |
| <i>Eremophila veronica</i>                                    | P3                  |      | X         |        | 38                              | Oct - Dec              | Stony clay, clay loam. Slopes of low lateritic hills and breakaways.   | No                                    | Low                                 | Low                                  |
| <i>Goodenia lyra</i>  | P3                  |      |           | X      | 76.6                            | Aug                    | Red sandy loam. Near claypan. <sup>2</sup>   | No                                    | Low                                 | Low                                  |
| <i>Grevillea subteriffineata</i>                              | P3                  |      |           | X      | 73.9                            | May - Sep              | Red-brown gravelly clayey sands. Low limestone rises, gypsumous dunes, drainage lines.   | No                                    | Low                                 | Low                                  |
| <i>Hybanthus floribundus</i> subsp. <i>chloranthus</i>        | P3                  |      |           | X      | 45.7                            | Aug - Oct              | Dark red-brown soil, never sandy, rich in iron oxide, laterite. Rocky areas, creek banks, along drainage lines. <sup>2</sup>                   | No                                    | Low                                 | Low                                  |
| <i>Hysterobaeckea ochropetala</i> subsp. <i>cometes</i>       | P3                  |      |           | X      | 93.9                            | Jul - Sep              | Red-yellow sandy soils.  | No                                    | Low                                 | Low                                  |
| <i>Microseris serrulata</i>                                   | P3                  |      | X         |        | 34.3                            | Jun - Nov              | Brownish sandy and clayey soils over granite.  | No                                    | Low                                 | Low                                  |
| <i>Olearia mucronata</i>                                      | P3                  |      |           | X      | 84.7                            | Aug - Jan              | Schistose hills, along drainage channels. <sup>2</sup>   | No                                    | Low                                 | Low                                  |
| <i>Phlœotheca coateana</i>                                    | P3                  |      |           | X      | 93.9                            | Aug - Sep              | Red-yellow/brown sands, sandy clay loam. Plains, sandplains, slopes, breakaways.   | Possibly                              | Low                                 | Low                                  |

| Species  | Conservation Status |      |           | Source | Distance to Nearest Record (km) | Flowering Period | Preferred Habitat       | Habitat occurs within the Survey Area  | Pre-Survey Likelihood of Occurrence | Post-Survey Likelihood of Occurrence |
|--|---------------------|------|-----------|--------|---------------------------------|------------------|-------------------------|--|-------------------------------------|--------------------------------------|
|  | DBCA                | EPBC | NatureMap |        |                                 |                  |                         |  |                                     |                                      |
| <i>Phyllanthus baccharoides</i>                                    | P3                  |      |           |        | X                               | 54.5             | Jul - Sep               | Red latentic and sandy clay soils Granite outcrops. <sup>2</sup>                             | No                                  | Low                                  |
| <i>Pterostylis virens</i>  | P3                  |      |           |        | X                               | 95.7             | Sep                     | Red sandy, gritty loams. Slopes and margins of granites, No banded ironstone hills.          | No                                  | Low                                  |
| <i>Tecticornia cymbiformis</i>                                     | P3                  |      |           |        | X                               | 84.9             | Mar - May               | Saline soils. Along the edge of creek lines. <sup>2</sup>                                    | No                                  | Low                                  |
| <i>Thryptomene sp. Leinsteri</i> (B.J. Lepschi & L.A. Craven 4362) | P3                  |      |           |        | X                               | 88.3             | Oct - Dec               | Red sandy loam. Granite breakaways, stony rises and outcrops.                                | No                                  | Low                                  |
| <i>Triglochin protuberans</i>                                      | P3                  |      | X         |        | X                               | 33.9             | Aug - Nov               | Red loam, grey mud over clay. Winter-wet sites, claypans, near salt lakes, margins of pools. | No                                  | Low                                  |
| <i>Conospermum todii</i>   | P4                  |      |           |        | X                               | 57.2             | Jul - Oct               | Yellow sand. Sand dunes.   | No                                  | Low                                  |
| <i>Eucalyptus jutsonii</i> subsp. <i>jutsonii</i>                  | P4                  |      |           |        | X                               | 57.5             | Feb - Mar and Nov - Dec | Red to pale orange deep sands. Undulating areas and on dunes.                                | No                                  | Low                                  |
| <i>Grevillea ericifolia</i>  | P4                  |      |           |        | X                               | 94.7             | Sep - Oct               | Gravelly loam. Lateritic ridges.   | No                                  | Low                                  |
| <i>Grevillea inconspicua</i>                                       | P4                  |      | X         |        | X                               | 38.7             | Jun - Aug               | Loam, gravel. Along drainage lines on rocky outcrops, creeklines. <sup>2</sup>               | No                                  | Low                                  |

## **Appendix D**

### **Inventory of Vascular Flora**

| Family           | Species   |
|------------------|---|
| Amaranthaceae    | <i>Ptilotus obovatus</i><br><i>Ptilotus schwartzii</i> var. <i>schwartzii</i>   |
| Apocynaceae      | <i>Marsdenia australis</i>  |
| Chenopodiaceae   | <i>Dysphania melanocarpa</i><br><i>Enchytraea tomentosa</i> var. <i>tomentosa</i><br><i>Maireana planifolia</i><br><i>Maireana</i> sp.<br><i>Maireana tomentosa</i> subsp. <i>tomentosa</i><br><i>Maireana triptera</i><br><i>Maireana villosa</i><br><i>Rhagodia eremaea</i>   |
| Convolvulaceae   | <i>Duperreya commixta</i>   |
| Euphorbiaceae    | <i>Euphorbia tannensis</i> subsp. <i>eremophila</i>   |
| Fabaceae         | <i>Acacia aptaneura</i><br><i>Acacia ayersiana</i><br><i>Acacia caesaneura</i><br><i>Acacia craspedocarpa</i><br><i>Acacia incurvaneura</i><br><i>Acacia mulganeura</i><br><i>Acacia ramulosa</i> var. <i>ramulosa</i><br><i>Acacia tetragonophylla</i><br><i>Senna artemisioides</i> subsp. <i>x artemisioides</i><br><i>Senna charlesiana</i> |
| Geraniaceae      | <i>Erodium cygnorum</i>   |
| Malvaceae        | <i>Hibiscus</i> sp. Gardneri (A.L. Payne PRP 1435)<br><i>Sida</i> sp.<br><i>Sida</i> sp. dark green fruits (S. van Leeuwen 2260)  |
| Poaceae          | * <i>Cenchrus ciliaris</i><br><i>Enneapogon caeruleuscens</i><br><i>Enneapogon polyphyllus</i><br><i>Eragrostis eriopoda</i><br><i>Eriachne pulchella</i> subsp. <i>dominii</i><br><i>Monachather paradoxus</i><br><i>Thyridolepis mitchelliana</i>   |
| Portulacaceae    | <i>Portulaca oleracea</i>   |
| Proteaceae       | <i>Hakea recurva</i> subsp. <i>recurva</i>  |
| Pteridaceae      | <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>  |
| Santalaceae      | <i>Santalum spicatum</i>  |
| Scrophulariaceae | <i>Eremophila forrestii</i> subsp. <i>forrestii</i><br><i>Eremophila latrobei</i> subsp. <i>latrobei</i><br><i>Eremophila platycalyx</i> subsp. <i>Leonora</i> (J. Morrisey 252)<br><i>Eremophila serrulata</i>   |
| Solanaceae       | <i>Solanum lasiophyllum</i>   |



## **Appendix E**

## **Flora Site Sheets**

## FLORA SITE SHEET

**Project Name** 4581 Leonora Biological Survey  
**Site:** LER01  
**Location** MGA 50      336056 mE      6804099 mN

**Described by:** BE, LC  
**Date:** 24/06/2021  
**Type:** Relevé

**Landform:** Gravelly Plain  
**Slope:** Flat  
**Rock Type:** Ironstone, Quartz  
**Soil Type:** Clay, Loam, Sand  
**Soil Colour:** Brown, Orange



**Vegetation:** *Acacia mulganeura* and *Acacia caesaneura* low woodland over *Eremophila forrestii* subsp. *forrestii* and *Maireana planifolia* mid sparse shrubland over *Eragrostis eriopoda* low sparse tussock grassland over *Ptilotus schwartzii* var. *schwartzii* low sparse herland  
**Condition:** Very Good      **Disturbance Type:** Rubbish, Vehicle track  
**Fire Age:** >10 years

### SPECIES LIST

| Taxon   | Height (cm) | Cover (%) | Notes                   |
|---|-------------|-----------|-------------------------|
| <i>Acacia incurvaneura</i>                          | 600         | 0,1       |                         |
| <i>Acacia mulganeura</i>                            | 500         | 8         | Variant 2               |
| <i>Acacia caesaneura</i>                            | 380         | 6         | Narrow phyllode variant |
| <i>Maireana planifolia</i>                          | 150         | 1         |                         |
| <i>Eremophila forrestii</i> subsp. <i>forrestii</i> | 130         | 3         |                         |
| <i>Marsdenia australis</i>                          | 130         | 0,1       |                         |
| <i>Maireana</i> sp.                                 | 60          | 0,1       |                         |
| <i>Ptilotus schwartzii</i> var. <i>schwartzii</i>   | 50          | 4         |                         |
| <i>Acacia tetragonophylla</i>                       | 40          | 0,1       |                         |
| <i>Eragrostis eriopoda</i>                          | 30          | 1         |                         |
| <i>Ptilotus obovatus</i>                            | 30          | 0,1       |                         |
| <i>Solanum lasiophyllum</i>                         | 30          | 0,1       |                         |
| <i>Thyridolepis mitchelliana</i>                    | 20          | 0,1       |                         |
| <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>    | 10          | 0,1       |                         |
| <i>Eriachne pulchella</i> subsp. <i>dominii</i>     | 5           | 0,1       |                         |
| <i>Erodium cygnorum</i>                             | 2           | 0,1       |                         |

## FLORA SITE SHEET

**Project Name** 4581 Leonora Biological Survey  
**Site:** LER02  
**Location** MGA 50      336180 mE      6803941 mN

**Described by:** BE, LC  
**Date:** 24/06/2021  
**Type:** Relevé

**Landform:** Gravelly Plain  
**Slope:** Flat  
**Rock Type:** Ironstone, Quartz  
**Soil Type:** Clay, Loam, Sand  
**Soil Colour:** Brown, Orange



**Vegetation:** *Acacia caesaneura* and *Acacia mulganeura* low open woodland over *Eragrostis eriopoda* low sparse tussock grassland

**Condition:** Good                          **Disturbance Type:** Weeds, Rubbish, Tracks  
**Fire Age:** >10 years

### SPECIES LIST

| Taxon   | Height (cm) | Cover (%) | Notes                   |
|---|-------------|-----------|-------------------------|
| <i>Acacia caesaneura</i>  | 350         | 5         | Narrow phyllode variant |
| <i>Acacia mulganeura</i>  | 350         | 3         | Variant 2               |
| <i>Rhagodia eremaea</i>   | 100         | 0,1       |                         |
| <i>Eremophila platycalyx</i> subsp. <i>Leonora</i> (J. Morrisey 2 | 70          | 0,1       |                         |
| <i>Maireana planifolia</i>  | 50          | 0,1       |                         |
| <i>Ptilotus obovatus</i>  | 30          | 0,1       |                         |
| <i>Eragrostis eriopoda</i>  | 10          | 2         |                         |
| <i>Enchyalaena tomentosa</i> var. <i>tomentosa</i>                | 10          | 0,1       |                         |
| * <i>Cenchrus ciliaris</i>  | 5           | 0,1       |                         |
| <i>Sida</i> sp.   | 3           | 0,1       |                         |
| <i>Erodium cygnorum</i>   | 2           | 0,1       |                         |

## FLORA SITE SHEET

**Project Name** 4581 Leonora Biological Survey  
**Site:** LER03  
**Location** MGA 50      336266 mE      6804160 mN

**Described by:** BE, LC  
**Date:** 24/06/2021  
**Type:** Relevé

**Landform:** Gravelly Plain  
**Slope:** Flat  
**Rock Type:** Ironstone, Quartz  
**Soil Type:** Clay, Loam, Sand  
**Soil Colour:** Brown, Orange



**Vegetation:** *Acacia caesaneura*, *Acacia mulganeura*, *Acacia ayersiana* and *Acacia aptaneura* low woodland over *Eremophila forestii* subsp. *forestii* mid sparse shrubland over *Eragrostis eriopoda* low sparse tussock grassland over *Ptilotus obovatus* low sparse hermland  
**Condition:** Very Good      **Disturbance Type:** Rubbish  
**Fire Age:** >10 years

### SPECIES LIST

| Taxon   | Height (cm) | Cover (%) | Notes                   |
|---|-------------|-----------|-------------------------|
| <i>Acacia caesaneura</i>                                  | 500         | 8         | Narrow phyllode variant |
| <i>Acacia mulganeura</i>                                  | 500         | 2         | Variant 2               |
| <i>Acacia ayersiana</i>                                   | 480         | 2         |                         |
| <i>Acacia aptaneura</i>                                   | 400         | 2         |                         |
| <i>Eremophila forestii</i> subsp. <i>forestii</i>         | 120         | 3         |                         |
| <i>Santalum spicatum</i>                                  | 110         | 0.1       |                         |
| <i>Enchytraea tomentosa</i> var. <i>tomentosa</i>         | 80          | 0.1       |                         |
| <i>Ptilotus obovatus</i>                                  | 50          | 2         |                         |
| <i>Eragrostis eriopoda</i>                                | 40          | 6         |                         |
| <i>Maireana</i> sp.                                       | 40          | 0.1       |                         |
| <i>Ptilotus schwartzii</i> var. <i>schwartzii</i>         | 40          | 0.1       |                         |
| <i>Maireana tomentosa</i> subsp. <i>tomentosa</i>         | 40          | 0.1       |                         |
| <i>Eremophila latrobei</i> subsp. <i>latrobei</i>         | 40          | 0.1       |                         |
| <i>Maireana villosa</i>                                   | 30          | 0.1       |                         |
| <i>Hibiscus</i> sp. <i>Gardneri</i> (A.L. Payne PRP 1435) | 30          | 0.1       |                         |
| <i>Monachather paradoxus</i>                              | 30          | 0.1       |                         |
| <i>Thyridolepis mitchelliana</i>                          | 20          | 0.1       |                         |
| <i>Eriachne pulchella</i> subsp. <i>dominii</i>           | 10          | 0.1       |                         |
| <i>Sida</i> sp. dark green fruits (S. van Leeuwen 2260)   | 10          | 0.1       |                         |
| <i>Solanum lasiophyllum</i>                               | 10          | 0.1       |                         |
| <i>Euphorbia tannensis</i> subsp. <i>eremophila</i>       | 10          | 0.1       |                         |
| <i>Erodium cygnorum</i>                                   | 2           | 0.1       |                         |

## FLORA SITE SHEET

**Project Name** 4581 Leonora Biological Survey  
**Site:** LER04  
**Location** MGA 50      336368 mE      6804138 mN

**Described by:** BE, LC  
**Date:** 24/06/2021  
**Type:** Relevé

**Landform:** Gravelly Plain  
**Slope:** Flat  
**Rock Type:** Ironstone, Quartz  
**Soil Type:** Clay, Loam, Sand  
**Soil Colour:** Brown, Orange



**Vegetation:** *Acacia incurvaneura* low open woodland over *Eragrostis eriopoda*, *Eriachne pulchella* subsp. *dominii* and \**Cenchrus ciliaris* low sparse grassland

**Condition:** Good      **Disturbance Type:** Rubbish  
**Fire Age:** >10 years

### SPECIES LIST

| Taxon   | Height (cm) | Cover (%) | Notes |
|---|-------------|-----------|-------|
| <i>Acacia tetragonophylla</i>                             | 600         | 0,1       |       |
| <i>Acacia incurvaneura</i>                                | 320         | 7         |       |
| * <i>Cenchrus ciliaris</i>                                | 100         | 1         |       |
| <i>Maireana planifolia</i>                                | 100         | 0,1       |       |
| <i>Maireana tomentosa</i> subsp. <i>tomentosa</i>         | 60          | 0,1       |       |
| <i>Eragrostis eriopoda</i>                                | 50          | 2         |       |
| <i>Solanum lasiophyllum</i>                               | 40          | 0,1       |       |
| <i>Monachather paradoxus</i>                              | 40          | 0,1       |       |
| <i>Marsdenia australis</i>                                | 40          | 0,1       |       |
| <i>Ptilotus obovatus</i>                                  | 30          | 0,1       |       |
| <i>Ptilotus schwartzii</i> var. <i>schwartzii</i>         | 30          | 0,1       |       |
| <i>Thyridolepis mitcelliana</i>                           | 30          | 0,1       |       |
| <i>Maireana</i> sp.                                       | 20          | 0,1       |       |
| <i>Maireana villosa</i>                                   | 20          | 0,1       |       |
| <i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>          | 10          | 0,1       |       |
| <i>Eriachne pulchella</i> subsp. <i>dominii</i>           | 5           | 2         |       |
| <i>Enneapogon polypylus</i>                               | 5           | 0,1       |       |
| <i>Hibiscus</i> sp. <i>Gardneri</i> (A.L. Payne PRP 1435) | 5           | 0,1       |       |
| <i>Erodium cygnorum</i>                                   | 3           | 0,1       |       |

## **Appendix F**

### **Fauna Literature Review and Database Searches Results**

**Appendix: Database Search and Literature Review Fauna Inventory**

\*State – Conservation status under BC Act or DBCA priority list, Federal – Conservation status under EPBC Act  
 A: Level 2 Vertebrate Fauna Assessment, King of the Hills Project (Terrestrial Ecosystems, 2020), B: Level 2 Fauna Risk Assessment for Granny Deeps Project Area. (Terrestrial Ecosystems, 2011), C: Biological Survey of the Eastern Goldfields of Western Australia Part 10 (Hall et al., 1994)

| Family           | Scientific Name                  | Common Name                               | Conservation Status <sup>+</sup> |       |           | Database | DBCA | DBCA 15yr | Field Survey | A | B | C | Literature |
|------------------|----------------------------------|---|----------------------------------|-------|-----------|----------|------|-----------|--------------|---|---|---|------------|
|                  |                                  |   | Federal                          | State | NatureMap |          |      |           |              |   |   |   |            |
| <b>AMPHIBIAN</b> |                                  |   |                                  |       |           |          |      |           |              |   |   |   |            |
| Limnodynastidae  | <i>Neobatrachus kunapalari</i>   | Kunapalari Frog                           | -                                | -     | x         |          |      |           |              |   |   |   | x          |
|                  | <i>Neobatrachus sutor</i>        | Shoemaker Frog                            | -                                | -     | x         |          |      |           |              | x | x |   |            |
|                  | <i>Platyplectrum spenceri</i>    | Centralian Burrowing Frog                 | -                                | -     | x         |          |      |           |              |   |   |   |            |
| Pelodryadidae    | <i>Cyclorana maini</i>           | Sheep Frog                                | -                                | -     | x         |          |      |           |              | x |   |   |            |
|                  | <i>Cyclorana occidentalis</i>    | Western Water-holding Frog                | -                                | -     | x         |          |      |           |              | x |   |   |            |
|                  | <i>Litoria rubella</i>           | Little Red Tree Frog                      | -                                | -     |           |          |      |           |              | x |   |   |            |
| Myobatrachidae   | <i>Pseudophryne occidentalis</i> | Western Toadlet                           | -                                | -     |           |          |      |           |              | x |   |   |            |
| <b>AVIAN</b>     |                                  |   |                                  |       |           |          |      |           |              |   |   |   |            |
| Acanthizidae     | <i>Acanthiza apicalis</i>        | Inland Thornbill (Broad-tailed Thornbill) | -                                | -     | x         |          |      |           |              | x | x |   |            |
|                  | <i>Acanthiza chrysorrhoa</i>     | Yellow-rumped Thornbill                   | -                                | -     | x         |          |      |           |              | x | x |   |            |
|                  | <i>Acanthiza robustirostris</i>  | Slaty-backed Thornbill                    | -                                | -     | x         |          |      |           |              | x | x |   |            |
|                  | <i>Acanthiza uropygialis</i>     | Chestnut-rumped Thornbill                 | -                                | -     | x         |          |      |           |              | x |   |   |            |
|                  | <i>Aphelocephala leucopsis</i>   | Southern Whiteface                        | -                                | -     | x         |          |      |           |              | x | x |   |            |
|                  | <i>Gerygone fusca</i>            | Western Gerygone                          | -                                | -     | x         |          |      |           |              | x |   |   |            |
|                  | <i>Pyrrholaemus brunneus</i>     | Redthroat                                 | -                                | -     |           |          |      |           |              | x |   |   |            |
|                  | <i>Smicronyx brevirostris</i>    | Weebill                                   | -                                | -     | x         |          |      |           |              | x |   |   |            |
| Accipitridae     | <i>Accipiter cirrocephalus</i>   | Collared Sparrowhawk                      | -                                | -     | x         |          |      |           |              | x |   |   |            |
|                  | <i>Accipiter fasciatus</i>       | Brown Goshawk                             | -                                | -     | x         |          |      |           |              | x |   |   |            |

| Family       | Scientific Name                    | Common Name                                  | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|--------------|------------------------------------|--|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|              |                                    |  |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
| Accipitridae | <i>Aquila audax</i>                | Wedge-tailed Eagle                           | -     | -                                | x    | -    |          |                     |                 | x          | x |   |
|              | <i>Circus approximans</i>          | Swamp Harrier                                | -     | -                                | x    | -    |          |                     |                 |            |   |   |
|              | <i>Haliastur sphenurus</i>         | Whistling Kite                               | -     | -                                | x    | -    |          |                     |                 |            |   |   |
|              | <i>Hieraetus morphnoides</i>       | Little Eagle                                 | -     | -                                | x    | -    |          |                     |                 |            |   |   |
|              | <i>Milvus migrans</i>              | Black Kite                                   | -     | -                                | x    | -    |          |                     |                 |            |   |   |
| Aegothelidae | <i>Aegotheles cristatus</i>        | Australian Owllet-nightjar                   | -     | -                                |      |      |          |                     | x               |            |   |   |
| Anatidae     | <i>Anas gracilis</i>               | Grey Teal                                    | -     | -                                | x    | -    |          |                     | x               | x          | x |   |
|              | <i>Anas superciliosa</i>           | Pacific Black Duck                           | -     | -                                | x    | -    |          |                     | x               |            |   |   |
|              | <i>Aythya australis</i>            | Hardhead                                     | -     | -                                | x    | -    |          |                     | x               | x          | x |   |
|              | <i>Biziura lobata</i>              | Musk Duck                                    | -     | -                                | x    | -    |          |                     | x               | x          | x |   |
|              | <i>Chenonetta jubata</i>           | Australian Wood Duck (Wood Duck, Maned Duck) | -     | -                                | x    | -    |          |                     | x               | x          | x |   |
|              | <i>Cygnus atratus</i>              | Black Swan                                   | -     | -                                | x    | -    |          |                     | x               |            |   |   |
|              | <i>Malacorhynchus membranaceus</i> | Pink-eared Duck                              | -     | -                                | x    | -    |          |                     | x               |            |   |   |
|              | <i>Tadorna tadornoides</i>         | Australian Shelduck (Mountain Duck)          | -     | -                                | x    | -    |          |                     | x               |            |   |   |
| Anhingidae   | <i>Anhinga novaehollandiae</i>     | Australasian Darter                          | -     | -                                | x    | -    |          |                     |                 |            |   |   |
| Apodidae     | <i>Apus pacificus</i>              | Pacific Swift (Fork-tailed Swift)            | MI    | -                                |      | x    |          |                     |                 |            |   |   |
| Ardeidae     | <i>Ardea modesta</i>               | Eastern Great Egret                          | -     | -                                | x    | -    |          |                     |                 |            |   |   |
|              | <i>Ardea pacifica</i>              | White-necked Heron                           | -     | -                                | x    | -    |          |                     |                 |            |   |   |
|              | <i>Egretta novaehollandiae</i>     | White-faced Heron                            | -     | -                                | x    | -    |          |                     | x               |            |   |   |
| Artamidae    | <i>Artamus cinereus</i>            | Black-faced Woodswallow                      | -     | -                                | x    | -    |          |                     | x               | x          | x |   |

| Family          | Scientific Name                 | Common Name                         | State | Conservation Status <sup>+</sup> |      |           | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|-----------------|---------------------------------|-------------------------------------|-------|----------------------------------|------|-----------|----------|---------------------|-----------------|------------|---|---|
|                 |                                 |                                     |       | Federal                          | PMST | NatureMap |          |                     |                 | A          | B | C |
|                 | <i>Artamus personatus</i>       | Masked Woodswallow                  | -     | -                                | x    |           |          |                     |                 | x          |   |   |
|                 | <i>Artamus minor</i>            | Little Woodswallow                  | -     | -                                |      |           |          |                     |                 | x          |   |   |
| Burhinidae      | <i>Burhinus grallarius</i>      | Bush Stone-curlew (Bush Thick-knee) | -     | -                                | x    |           |          |                     |                 |            |   |   |
| Cacatuidae      | <i>Cacatua roseicapilla</i>     | Galah                               | -     | -                                | x    |           |          |                     |                 | x          |   |   |
|                 | <i>Nymphicus hollandicus</i>    | Cockatiel                           | -     | -                                | x    |           |          |                     |                 |            |   |   |
| Campetherigidae | <i>Coracina maxima</i>          | Ground Cuckoo-shrike                | -     | -                                | x    |           |          |                     |                 | x          | x |   |
|                 | <i>Coracina novaehollandiae</i> | Black-faced Cuckoo-shrike           | -     | -                                | x    |           |          |                     |                 | x          | x |   |
|                 | <i>Lalage tricolor</i>          | White-winged Triller                | -     | -                                |      |           |          |                     |                 | x          |   |   |
| Caprimulgidae   | <i>Eurostopodus argus</i>       | Spotted Nightjar                    | -     | -                                | x    |           |          |                     |                 |            |   |   |
| Casuariidae     | <i>Dromaius novaehollandiae</i> | Emu                                 | -     | -                                | x    |           |          |                     |                 | x          | x |   |
| Charadriidae    | <i>Charadrius ruficollis</i>    | Red-capped Plover                   | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                 | <i>Charadrius veredus</i>       | Oriental Plover                     | MI    | MII & MA                         | x    |           |          |                     |                 |            |   |   |
|                 | <i>Elseyornis melanops</i>      | Black-fronted Dotterel              | -     | -                                | x    |           |          |                     |                 | x          | x |   |
|                 | <i>Erythrogenys cinctus</i>     | Red-kneed Dotterel                  | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                 | <i>Pluvialis fulva</i>          | Pacific Golden Plover               | MI    | MII & MA                         | x    |           |          |                     |                 |            |   |   |
|                 | <i>Thinornis cucullatus</i>     | Hooded Plover (Hooded Dotterel)     | P4    | MA                               | x    | x         | x        |                     |                 |            |   |   |
|                 | <i>Vanellus tricolor</i>        | Banded Lapwing                      | -     | -                                | x    |           |          |                     |                 |            |   |   |
| Cinclosomatidae | <i>Cinclosoma marginatum</i>    | Western Quail-thrush                | -     | -                                | x    |           |          |                     |                 |            |   |   |
| Climacteridae   | <i>Climacteris affinis</i>      | White-browed Treecreeper            | -     | -                                |      |           |          |                     |                 | x          |   |   |
| Columbidae      | <i>Columba livia</i>            | Domestic Pigeon (Rock Dove)         | -     | -                                | x    | x         |          |                     |                 |            |   |   |
|                 | <i>Geopelia cuneata</i>         | Diamond Dove                        | -     | -                                | x    |           |          |                     |                 |            |   |   |

| Family       | Scientific Name                      | Common Name                          | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|--------------|--------------------------------------|--------------------------------------|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|              |                                      |                                      |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
| Columbidae   | <i>Ocyphaps lophotes</i>             | Crested Pigeon                       | -     | -                                | x    |      | x        | x                   | x               | x          | x |   |
|              | <i>Phaps chalcoptera</i>             | Common Bronzewing                    | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Spilopelia senegalensis</i>       | Laughing Turtle Dove                 | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Corvidae     | <i>Corvus bennetti</i>               | Little Crow                          | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Corvus coronoides</i>             | Australian Raven                     | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Corvus orru</i>                   | Torresian Crow                       | -     | -                                | x    |      |          |                     | x               | x          |   |   |
| Cracticidae  | <i>Cracticus nigrogularis</i>        | Pied Butcherbird                     | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Cracticus tibicen</i>             | Australian Magpie                    | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Cracticus torquatus</i>           | Grey Butcherbird                     | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Strepera versicolor</i>           | Grey Currawong                       | -     | -                                | x    |      |          |                     | x               | x          |   |   |
| Cuculidae    | <i>Cacomantis pallidus</i>           | Pallid Cuckoo                        | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Chrysococcyx basalis</i>          | Horsfield's Bronze Cuckoo            | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Chrysococcyx osculans</i>         | Black-eared Cuckoo                   | -     | -                                | x    |      |          |                     | x               | x          |   |   |
| Dicaeidae    | <i>Dicaeum hirundinaceum</i>         | Mistletoebird                        | -     | -                                | x    |      |          |                     | x               | x          |   |   |
| Falconidae   | <i>Falco berigora</i>                | Brown Falcon                         | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Falco cenchroides</i>             | Australian Kestrel (Nankeen Kestrel) | -     | -                                | x    |      |          |                     | x               | x          |   |   |
|              | <i>Falco cenchroides cenchroides</i> |                                      | -     | -                                | x    |      |          |                     |                 |            |   |   |
|              | <i>Falco hypoleucus</i>              | Grey Falcon                          | VU    | VU                               | x    | x    |          |                     |                 |            |   |   |
|              | <i>Falco longipennis</i>             | Australian Hobby                     | -     | -                                | x    |      |          |                     |                 |            |   |   |
|              | <i>Falco peregrinus</i>              | Peregrine Falcon                     | OS    | -                                | x    | x    | 6        |                     |                 |            |   |   |
| Hirundinidae | <i>Cheramoeca leucosterna</i>        | White-backed Swallow                 | -     | -                                | x    |      |          |                     | x               | x          |   |   |

| Family       | Scientific Name                 | Common Name              | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|--------------|---------------------------------|--------------------------|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|              |                                 |                          |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
| Hirundinidae | <i>Hirundo neoxena</i>          | Welcome Swallow          | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|              | <i>Petrochelidon ariel</i>      | Fairy Martin             | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|              | <i>Petrochelidon nigricans</i>  | Tree Martin              | -     | -                                | x    |      |          |                     |                 | x          |   |   |
| Laridae      | <i>Larus novaehollandiae</i>    | Silver Gull              | -     | MA                               | x    |      |          |                     |                 |            |   |   |
|              | <i>Sterna hybrida javanica</i>  |                          | -     | -                                | x    |      |          |                     |                 |            |   |   |
|              | <i>Sterna nilotica</i>          | Gull-billed Tern         | MI    | -                                | x    |      |          |                     |                 |            |   |   |
| Maluridae    | <i>Malurus leucopterus</i>      | White-winged Fairywren   | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Malurus pulcherrimus</i>     | Blue-breasted Fairywren  | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Malurus splendens</i>        | Splendid Fairywren       | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Megapodiidae | <i>Leipoa ocellata</i>          | Malleefowl               | VU    | VU                               | x    | x    | x        | 10                  |                 | x          | x |   |
| Meliphagidae | <i>Acanthagenys rufogularis</i> | Spiny-cheeked Honeyeater | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Anthochaera carunculata</i>  | Red Wattlebird           | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Certhionyx variegatus</i>    | Pied Honeyeater          | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Epthianura albifrons</i>     | White-fronted Chat       | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Epthianura aurifrons</i>     | Orange Chat              | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Epthianura tricolor</i>      | Crimson Chat             | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Gavicalis virescens</i>      | Singing Honeyeater       | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Lichenura indistincta</i>    | Brown Honeyeater         | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Manorina flavigula</i>       | Yellow-throated Miner    | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|              | <i>Purnella albifrons</i>       | White-fronted Honeyeater | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Meropidae    | <i>Merops ornatus</i>           | Rainbow Bee-eater        | -     | -                                | x    | x    |          |                     |                 | x          | x |   |
| Monarchidae  | <i>Grallina cyanoleuca</i>      | Magpie-lark              | -     | -                                | x    |      |          |                     |                 | x          | x |   |

| Family            | Scientific Name                          | Common Name            | State | Conservation Status <sup>+</sup> |      |           | Database | DBCA 15yr<br>Survey | Field<br>Survey | A | B | C | Literature |
|-------------------|--|------------------------|-------|----------------------------------|------|-----------|----------|---------------------|-----------------|---|---|---|------------|
|                   |  |                        |       | Federal                          | PMST | NatureMap |          |                     |                 |   |   |   |            |
| Motacillidae      | <i>Anthus australis australis</i>        |                        | -     | -                                |      |           |          |                     |                 | x | x |   |            |
|                   | <i>Motacilla cinerea</i>                 | Grey Wagtail           | MI    | MI & MA                          | x    |           |          |                     |                 |   |   |   |            |
|                   | <i>Motacilla flava</i>                   | Yellow Wagtail         | MI    | MI & MA                          | x    |           |          |                     |                 |   |   |   |            |
| Neosittidae       | <i>Daphoenositta chrysoptera</i>         | Varied Sittella        | -     | -                                |      |           |          |                     | x               |   |   |   |            |
| Oreocitidae       | <i>Oreoica gutturalis</i>                | Crested Bellbird       | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
| Otididae          | <i>Ardeotis australis</i>                | Australian Bustard     | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
| Pachycephalidae   | <i>Colluricinclia harmonica</i>          | Grey Shrike-thrush     | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
|                   | <i>Pachycephala rufiventris</i>          | Rufous Whistler        | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
| Pardalotidae      | <i>Pardalotus striatus</i>               | Striated Pardalote     | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
|                   | <i>Pardalotus striatus westraliensis</i> |                        | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
| Pelecanidae       | <i>Pelecanus conspicillatus</i>          | Australian Pelican     | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
| Petroicidae       | <i>Melanodryas cucullata</i>             | Hooded Robin           | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
|                   | <i>Microeca fascinans</i>                | Jacky Winter           | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
|                   | <i>Petroica goodenovii</i>               | Red-capped Robin       | -     | -                                | x    |           |          |                     | x               | x |   |   |            |
| Phalacrocoracidae | <i>Phalacrocorax carbo</i>               | Great Cormorant        | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
|                   | <i>Phalacrocorax melanoleucos</i>        | Little Pied Cormorant  | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
|                   | <i>Phalacrocorax sulcirostris</i>        | Little Black Cormorant | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
| Podargidae        | <i>Podargus strigoides brachypterus</i>  |                        | -     | -                                | x    |           |          |                     |                 |   |   |   |            |
| Podicipedidae     | <i>Podiceps cristatus</i>                | Great Crested Grebe    | -     | -                                | x    |           |          |                     |                 |   |   |   |            |

| Family           | Scientific Name                      | Common Name                               | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|------------------|--------------------------------------|---|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|                  |                                      |   |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
|                  | <i>Poliocephalus poliocephalus</i>   | Hoary-headed Grebe                        | -     | -                                | x    |      |          |                     |                 | x          |   |   |
| Podicipedidae    | <i>Tachybaptus novaehollandiae</i>   | Australasian Grebe (Black-throated Grebe) | -     | -                                | x    |      |          |                     |                 | x          |   |   |
| Pomatostomidae   | <i>Pomatostomus superciliosus</i>    | White-browed Babbler                      | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Procellariidae   | <i>Calonectris leucomelas</i>        | Streaked Shearwater                       | MI    | -                                |      |      | x        | 6                   |                 |            |   |   |
| Psittacidae      | <i>Melopsittacus undulatus</i>       | Budgerigar                                | -     | -                                | x    |      |          |                     |                 |            |   |   |
|                  | <i>Neophema bourkii</i>              | Bourke's Parrot                           | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|                  | <i>Pezoporus occidentalis</i>        | Night Parrot                              | CR    | EN                               | x    |      |          |                     |                 |            |   |   |
|                  | <i>Platycercus varius</i>            | Mulga Parrot                              | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|                  | <i>Platycercus zonarius</i>          | Australian Ringneck                       | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|                  | <i>Polytelis alexandri</i>           | Princess Parrot                           | P4    | VU                               | x    |      |          |                     |                 | x          | x |   |
| Rallidae         | <i>Fulica atra</i>                   | Eurasian Coot                             | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|                  | <i>Tribonyx ventralis</i>            | Black-tailed Nativehen                    | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Recurvirostridae | <i>Himantopus himantopus</i>         | Black-winged Stilt                        | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|                  | <i>Recurvirostra novaehollandiae</i> | Red-necked Avocet                         | -     | -                                | x    |      |          |                     |                 |            |   |   |
|                  | <i>Cladorhynchus leucocephalus</i>   | Banded Stilt                              | -     | -                                |      |      |          |                     |                 | x          |   |   |
| Scolopacidae     | <i>Calidris acuminata</i>            | Sharp-tailed Sandpiper                    | MI    | MI & MA                          |      |      |          |                     |                 |            |   |   |
|                  | <i>Calidris canutus</i>              | Red Knot                                  | EN    | EN, MI & MA                      |      |      |          |                     |                 |            |   |   |
|                  | <i>Calidris melanotos</i>            | Pectoral Sandpiper                        | MI    | -                                |      |      |          |                     |                 |            | x |   |

| Family            | Scientific Name                           | Common Name                           | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|-------------------|---|---------------------------------------|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|                   |   |                                       |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
|                   | <i>Tringa glareola</i>                    | Wood Sandpiper                        | MI    | MI & MA                          | x    | x    | x        | 2                   |                 |            |   |   |
|                   | <i>Tringa hypoleucos</i>                  | Common Sandpiper                      | MI    | MI & MA                          | x    | x    | x        | 3                   |                 |            |   |   |
| Scolopacidae      | <i>Tringa nebularia</i>                   | Common Greenshank                     | MI    | MI & MA                          | x    | x    | x        | 5                   |                 |            |   |   |
| Threskiornithidae | <i>Platalea flavipes</i>                  | Yellow-billed Spoonbill               | -     | -                                | x    |      |          |                     |                 |            |   |   |
|                   | <i>Plegadis falcinellus</i>               | Glossy Ibis                           | MI    | -                                |      | x    |          |                     |                 |            |   |   |
|                   | <i>Threskiornis spinicollis</i>           | Straw-necked Ibis                     | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Alcedinidae       | <i>Todiramphus pyrrhopygius</i>           | Red-backed Kingfisher                 | -     | -                                | x    |      |          |                     |                 | x          | x |   |
|                   | <i>Todiramphus sanctus</i>                | Sacred Kingfisher                     | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Estrildidae       | <i>Taeniopygia guttata</i>                | Zebra Finch                           | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Psophodidae       | <i>Psophodes occidentalis</i>             | Western Wedgebill (Chiming Wedgebill) | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Ptilonorhynchidae | <i>Ptilonorhynchus maculatus guttatus</i> | Western Bowerbird                     | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Rhipiduridae      | <i>Rhipidura albiscapa</i>                | Grey Fantail                          | -     | -                                | x    |      |          |                     |                 | x          |   |   |
|                   | <i>Rhipidura albicauda</i>                | White-tailed Fantail                  | -     | -                                | x    |      |          |                     |                 |            |   |   |
|                   | <i>Rhipidura leucophrys</i>               | Willie Wagtail                        | -     | -                                | x    |      |          |                     |                 | x          | x |   |
| Tytonidae         | <i>Tyto javanica delicatula</i>           |                                       | -     | -                                | x    |      |          |                     |                 |            |   |   |
| <b>MAMMALIA</b>   |   |                                       |       |                                  |      |      |          |                     |                 |            |   |   |
| Bovidae           | <i>Bos primigenius taurus</i>             | European Cattle                       | -     | -                                |      |      |          |                     |                 | x          |   |   |
|                   | <i>Capra aegagrus hircus</i>              | Goat                                  | -     | -                                |      | x    |          |                     |                 | x          |   |   |
| Camelidae         | <i>Camelus dromedarius</i>                | Dromedary, Camel                      | -     | -                                | x    |      |          |                     |                 | x          |   |   |
| Canidae           | <i>Canis familiaris familiaris</i>        | Dog                                   | -     | -                                | x    |      |          |                     |                 | x          |   |   |

| Family       | Scientific Name                       | Common Name                   | State | Conservation Status <sup>+</sup> |      |      | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|--------------|---------------------------------------|-------------------------------|-------|----------------------------------|------|------|----------|---------------------|-----------------|------------|---|---|
|              |                                       |                               |       | Federal                          | PMST | DBCA |          |                     |                 | A          | B | C |
| Dasyuridae   | <i>Vulpes vulpes</i>                  | Red Fox                       | -     | -                                | x    | x    |          |                     |                 | x          | x | x |
|              | <i>Dasyurus blythii</i>               | Brush-tailed Mulgara, Ampurta | P4    | -                                | x    | x    | x        | x                   | x               | x          | x | x |
| Dasyuridae   | <i>Dasyurus geoffroii fortis</i>      | Western Quoll, Chuditch       | VU    | VU (at sp.<br>level)             | x    |      |          |                     |                 | x          | x | x |
|              | <i>Ningauia ridei</i>                 | Wongai Ningauai               | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Pseudantechinus woolleyae</i>      | Woolley's Pseudantechinus     | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Sminthopsis crassicaudata</i>      | Fat-tailed Dunnart            | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Sminthopsis dolichura</i>          | Little long-tailed Dunnart    | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Sminthopsis hirtipes</i>           | Hairy-footed Dunnart          | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Sminthopsis longicaudata</i>       | Long-tailed Dunnart           | P4    | -                                | x    | x    | x        | x                   | x               | x          | x | x |
|              | <i>Sminthopsis macroura</i>           | Stripe-faced Dunnart          | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Sminthopsis oolea</i>              | Ooldea Dunnart                | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Antechinomys laniger</i>           | Kultarr                       | -     | -                                |      |      |          |                     |                 | x          | x | x |
| Equidae      | <i>Equus africanus asinus</i>         | Donkey                        | -     | -                                | x    |      |          |                     |                 |            |   |   |
| Felidae      | <i>Felis catus</i>                    | Cat                           | -     | -                                | x    |      |          |                     |                 | x          | x | x |
| Leporidae    | <i>Oryctolagus cuniculus</i>          | Rabbit                        | -     | -                                | x    |      |          |                     |                 | x          | x | x |
| Macropodidae | <i>Osphranter robustus erubescens</i> | Euro, Biggada                 | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Osphranter rufus</i>               | Red Kangaroo, Marlu           | -     | -                                | x    |      |          |                     |                 | x          | x | x |
| Molossidae   | <i>Austronomus australis</i>          | White-striped Free-tailed Bat | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Ozimops kitcheneri</i>             | Western Free-tailed Bat       | -     | -                                |      |      |          |                     |                 | x          | x | x |
|              | <i>Ozimops pettersi</i>               | Inland Free-tailed Bat        | -     | -                                |      |      |          |                     |                 | x          | x | x |

| Family           | Scientific Name                         | Common Name                | State | Conservation Status <sup>+</sup> |      |           | Database | DBCA 15yr<br>Survey | Field<br>Survey | Literature |   |   |
|------------------|---|----------------------------|-------|----------------------------------|------|-----------|----------|---------------------|-----------------|------------|---|---|
|                  |   |                            |       | Federal                          | PMST | NatureMap |          |                     |                 | A          | B | C |
| Muridae          | <i>Mus musculus</i>                     | House Mouse                | -     | -                                | x    | x         |          |                     |                 | x          | x |   |
|                  | <i>Notomys alexis alexis</i>            | Spinifex Hopping-mouse     | -     | -                                |      |           |          |                     |                 | x          | x |   |
| Muridae          | <i>Pseudomys hermannsburgensis</i>      | Sandy Inland Mouse         | -     | -                                | x    |           |          |                     | x               | x          | x |   |
| Vespertilionidae | <i>Chalinolobus gouldii</i>             | Gould's Wattled Bat        | -     | -                                |      |           |          |                     | x               | x          | x |   |
|                  | <i>Nyctophilus geoffroyi geoffroyi</i>  | Lesser Long-eared Bat      | -     | -                                | x    |           |          |                     | x               |            | x |   |
|                  | <i>Nyctophilus sp.</i>                  |                            | -     | -                                |      |           |          |                     | x               |            |   |   |
|                  | <i>Scotorepens balstoni</i>             | Inland Broad-nosed Bat     | -     | -                                |      |           |          |                     | x               | x          | x |   |
|                  | <i>Vespadelus baverstocki</i>           | Inland Forest Bat          | -     | -                                |      |           |          |                     | x               |            |   |   |
|                  | <i>Vespadelus finlaysoni</i>            | Finlayson's Cave Bat       | -     | -                                |      |           |          |                     | x               | x          |   |   |
| Suidae           | <i>Sus scrofa</i>                       | Pig                        | -     | -                                | x    |           |          |                     | x               | x          | x |   |
| Tachyglossidae   | <i>Tachyglossus aculeatus acanthion</i> | Short-beaked Echidna       | -     | -                                |      |           |          |                     | x               | x          | x |   |
| <b>REPTILA</b>   |   |                            |       |                                  |      |           |          |                     |                 |            |   |   |
| Agamidae         | <i>Ctenophorus caudicinctus</i>         | Western Ring-tailed Dragon | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                  | <i>Ctenophorus fordi</i>                | Mallee Sand Dragon         | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                  | <i>Ctenophorus nuchalis</i>             | Central Netted Dragon      | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                  | <i>Ctenophorus reticulatus</i>          | Western Netted Dragon      | -     | -                                | x    |           |          |                     |                 |            |   |   |
|                  | <i>Ctenophorus scutulatus</i>           |                            | -     | -                                |      |           |          |                     |                 |            |   |   |
|                  | <i>Diporiphora amphiboluroides</i>      | Mulga Dragon               | -     | -                                |      |           |          |                     | x               |            | x |   |
|                  | <i>Pogona minor minor</i>               | Western Bearded Dragon     | -     | -                                |      |           |          |                     | x               |            | x |   |

| Family           | Scientific Name                       | Common Name                       | State | Conservation Status <sup>+</sup> |      |           | Database | DBCA 15yr | Field Survey | A | B | C | Literature |
|------------------|---------------------------------------|-----------------------------------|-------|----------------------------------|------|-----------|----------|-----------|--------------|---|---|---|------------|
|                  |                                       |                                   |       | Federal                          | PMST | NatureMap |          |           |              |   |   |   |            |
| Carphodactylidae | <i>Tymanocryptis cephalus</i>         | Coastal pebble-nimic dragons      | -     | -                                | -    | x         |          |           |              | x |   |   |            |
| Carphodactylidae | <i>Nephrurus vertebralis</i>          | Southern Banded Knob-tailed Gecko | -     | -                                | -    | x         |          |           |              | x |   |   |            |
| Diplodactylidae  | <i>Nephrurus wheeleri</i>             |                                   |       |                                  |      |           |          |           |              |   |   |   |            |
|                  | <i>Underwoodisaurus milii</i>         | Southern Barking Gecko            | -     | -                                | -    | x         |          |           |              | x |   |   |            |
| Diplodactylidae  | <i>Diplodactylus conspicillatus</i>   | Variable Fat-tailed Gecko         | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Diplodactylus granariensis</i>     | Wheat-Belt Stone Gecko            | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Diplodactylus granariensis rex</i> |                                   | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Diplodactylus pulcher</i>          |                                   | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Rhynchoedura ornata</i>            | Western Beaked Gecko              | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Strophurus assimilis</i>           | Goldfields Spiny-tailed Gecko     | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Strophurus strophurus</i>          |                                   | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Strophurus wellingtonae</i>        |                                   | -     | -                                | -    | x         |          |           |              | x |   |   |            |
| Elapidae         | <i>Brachyurophis semifasciatus</i>    |                                   | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Parasuta monachus</i>              | Inland Hooded Snake               | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Pseudechis australis</i>           | Mulga Snake                       | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Pseudechis butleri</i>             | Spotted Mulga Snake               | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Pseudonaja mengdeni</i>            | Western Brown Snake               | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Pseudonaja modesta</i>             | Ringed Brown Snake                | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Simoselaps bertholdi</i>           | Jan's Banded Snake                | -     | -                                | -    | x         |          |           |              | x |   |   |            |
|                  | <i>Suta fasciata</i>                  | Rosen's Snake                     | -     | -                                | -    | x         |          |           |              | x |   |   |            |

| Family      | Scientific Name                   | Common Name                       | State                        | Conservation Status <sup>+</sup> | Literature |      |          |
|-------------|-----------------------------------|-----------------------------------|------------------------------|----------------------------------|------------|------|----------|
|             |                                   |                                   |                              |                                  | DBCA       | PMST | Database |
| Gekkonidae  | <i>Suta punctata</i>              | Spotted Snake                     | -                            | -                                |            |      | x        |
| Gekkonidae  | <i>Gehyra variegata</i>           | Variegated gehyra                 | -                            | -                                | x          | x    | x        |
| Gekkonidae  | <i>Heteronotia binoei</i>         | Bynoe's Gecko                     | -                            | -                                | x          | x    | x        |
| Pygopodidae | <i>Pygopus nigriceps</i>          |                                   | -                            | -                                | x          | x    | x        |
| Pythonidae  | <i>Antaresia stimsoni</i>         |                                   | -                            | -                                |            | x    | x        |
|             | <i>Aspidites ramsayi</i>          | Woma                              | P1<br>(southwest<br>subpop.) | -                                |            | x    |          |
| Scincidae   | <i>Cryptoblepharus australis</i>  |                                   | -                            | -                                | x          |      |          |
|             | <i>Cryptoblepharus buchananii</i> |                                   | -                            | -                                | x          | x    |          |
|             | <i>Ctenotus schomburgkii</i>      |                                   | -                            | -                                |            | x    |          |
|             | <i>Ctenotus severus</i>           |                                   | -                            | -                                | x          | x    |          |
|             | <i>Ctenotus uber uber</i>         |                                   | -                            | -                                |            | x    |          |
|             | <i>Egernia depressa</i>           | Southern Pygmy Spiny-tailed Skink | -                            | -                                | x          | x    | x        |
|             | <i>Eremiascincus richardsonii</i> | Broad-banded Sand Swimmer         | -                            | -                                |            | x    | x        |
|             | <i>Lerista desertorum</i>         |                                   | -                            | -                                | x          | x    | x        |
|             | <i>Lerista muelleri</i>           |                                   | -                            | -                                |            | x    |          |
|             | <i>Lerista timida</i>             |                                   | -                            | -                                | x          |      |          |
|             | <i>Liopholis inornata</i>         |                                   | -                            | -                                | x          |      |          |
|             | <i>Menetia greyii</i>             |                                   | -                            | -                                |            | x    | x        |
|             | <i>Morethia butleri</i>           |                                   | -                            | -                                | x          | x    | x        |

| Family      | Scientific Name                 | Common Name         | Conservation Status <sup>+</sup> | State | Federal | Database |   |   | Field Survey |   |   | Literature |   |   |
|-------------|---------------------------------|---------------------|----------------------------------|-------|---------|----------|---|---|--------------|---|---|------------|---|---|
|             |                                 |                     |                                  |       |         | A        | B | C | A            | B | C | A          | B | C |
|             | <i>Ctenotus leonhardii</i>      |                     | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
|             | <i>Lerista distinguenda</i>     |                     | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
|             | <i>Tiliqua multifasciata</i>    | Central Blue-tongue | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
| Typhlopidae | <i>Anilios hamatus</i>          |                     | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
|             | <i>Anilios australis</i>        |                     | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
|             | <i>Anilios bicolor</i>          |                     | -                                | -     | -       |          |   |   |              |   |   | x          |   |   |
| Varanidae   | <i>Varanus caudolineatus</i>    |                     | -                                | -     | -       |          |   |   |              |   |   | x          | x |   |
|             | <i>Varanus panoptes rubidus</i> |                     | -                                | -     | -       |          |   |   |              |   |   | x          | x |   |

| Report   | Project Area  | Survey Timing and limitations    | Survey Effort  | Conservation Significant                                  | Fauna Recorded Onsite | Fauna Habitats   |
|--|---|----------------------------------|--|---|-----------------------|--|
| Level 2 Vertebrate Fauna Assessment, King of the Hills Project (Terrestrial Ecosystems, 2020)  | King of the Hills Gold Mine<br>29km North Northwest | Nov 2019<br>March 2020           | • Desktop<br>• Level 2 vertebrate fauna risk assessment                      | -   | -                     | Two fauna habitats:<br>• Open Mulga Woodland over mixed shrubs<br>• Woodland of large eucalypts over mixed shrubs along ephemeral creekline  |
| Report for Gwalia Materials, Preliminary Environmental Impact Assessment, Flora Survey and Environmental Management Plan (GHD, 2011) | Leonora-Laverton Road<br>54km East                  | Nov 2010                         | • Desktop<br>• Opportunistic vertebrate fauna (no fauna inventory available) | -   | -                     | -  |
| Biological Survey of the Eastern Goldfields of Western Australia Part 10 (Hall et al, 1994)  | Erliston Station<br>98km Northeast                  | Feb 1979<br>May 1980<br>Aug 1981 | • Vertebrate fauna survey  | -   | -                     | -  |
| Vertebrate Fauna Risk Assessment for the Granny Smith Solar Power Farm Project (Terrestrial Ecosystems, 2018)                        | Granny Smith Gold Mine<br>100km East                | Oct 2018                         | • Desktop<br>• Level 1 vertebrate fauna risk assessment                      | -   | -                     | Four fauna habitats:<br>• Open mulga woodland over scattered low shrubs (stony sandy-clay or sandy-clay substrate)<br>• Open Chenopod shrubland (stony sandy-clay) or sandy-clay substrate<br>• Chenopod and Mulga shrubland (stony sandy-clay or sandy-clay substrate)<br>• Banded ironstone rocky ridgeline with scattered Mulga and shrubs. |
| Level 2 Fauna Risk Assessment for Granny Deeps Project Area. (Terrestrial Ecosystems, 2011)  | Granny Smith Gold Mine<br>100km East                | Jan 2011                         | • Desktop<br>• Level 2 vertebrate fauna survey                               | • Long-tailed Dunnart ( <i>Sminthopsis longicaudata</i> ) | -                     | -  |
| Vertebrate Fauna Risk Assessment Granny Smith Tailing Storage Facility Expansion (Terrestrial Ecosystems, 2020)                      | Granny Smith Gold Mine<br>105km East                | April 2020                       | • Desktop<br>• Level 1 vertebrate fauna risk assessment                      | -   | -                     | Four fauna habitats:<br>• Open mulga woodland over scattered low shrubs<br>• Mulga Woodland along creekline over grasses and shrubs<br>• Mulga over chenopod shrubland<br>• Mulga Woodland over scattered low on banded ironstone formation  |

## **Appendix G**

### **Fauna Likelihood of Occurrence**

**Appendix: Conservation Significant Fauna Likelihood of Occurrence**

High = Previously recorded or Suitable habitat present and records less than 5 km from the Survey Area within the last 10 years, Medium = Suitable habitat present and records between 5 km and 15km from the Survey Area, and Low = No suitable habitat present, records greater than 15km from the Survey Area and/or records are historical

\*State: Conservation status under EC Act or DBCA priority list, Federal: Conservation status under EPBC Act

| Family       | Scientific Name             | Common Name                       | Conservation Code |         | Source | Habitat   | Likelihood of Occurrence | Distance and Justification                                       |
|--------------|-----------------------------|-----------------------------------|-------------------|---------|--------|---|--------------------------|--|
|              |                             |                                   | Federal           | State   |        |   |                          |  |
| AVIAN        |                             |                                   |                   |         |        |   |                          |  |
| Apodidae     | <i>Apus pacificus</i>       | Pacific Swift (Fork-tailed Swift) | M1                |         | X      | Low to very high airspace over varied habitat, rainforest to semi-desert <sup>2</sup>   | Low                      | Nearest record 158 kms away. Medium due to distance and habitat. |
| Charadriidae | <i>Charadrius veredus</i>   | Oriental Plover                   | M1                | M1 & MA | X      | Mainly on grasslands and thinly vegetated plains, preferring open areas. <sup>1</sup>   | Medium                   | Nearest record 17 kms away. Medium due to distance and habitat.  |
|              | <i>Pluvialis fulva</i>      | Pacific Golden Plover             | M1                | M1 & MA |        | Migrant from north Siberia. Mainly coastal in Australia, most commonly associated with tidal flats but also in other tidal settings like beaches and reefs, especially those with sea-weed <sup>1</sup> | Low                      | Nearest record 10 kms away. Low due to habitat.                  |
|              | <i>Thinornis cucullatus</i> | Hooded Plover (Hooded Dotterel)   | P4                | MA      | X      | Ocean beaches and margins of inland salt lakes <sup>1</sup>   | Low                      | Nearest record 11 kms. Low due to habitat.                       |

| Family       | Scientific Name                 | Common Name      | Conservation Code | Source  | Likelihood of Occurrence   | Distance and Justification |
|--------------|---------------------------------|------------------|-------------------|---------|--|----------------------------|
|              |                                 |                  |                   |         |  |                            |
|              |                                 |                  |                   |         |  |                            |
| Falconidae   | <i>Falco hypoleucus</i>         | Grey Falcon      | VU                | VU      | Open plains with treed watercourses in arid inland <sup>1</sup>  | Medium                     |
|              | <i>Falco peregrinus</i>         | Peregrine Falcon | OS                | X       | Most environments with suitable nest sites; cliff faces preferred, including man-made ones, commonly used; stick nests built by other species <sup>1</sup>   | High                       |
| Laridae      | <i>Larus novaehollandiae</i>    | Silver Gull      | MA                | X       | Waters near coast; beaches, reefs, jetties and piers; town lakes; public gardens; sewage ponds etc, to which many gulls commute long distances. <sup>6</sup> | Low                        |
|              | <i>Gelochelidon nilotica</i>    | Gull-billed Tern | ML                |         | Strictly coastal, at high tide often roosts with other terns/shorebirds <sup>1</sup>   | Low                        |
| Megapodiidae | <i>Leipoa ocellata</i>          | Malleefowl       | VU                | VU      | Unburned mallee and woodland with abundant litter and low scrub <sup>2</sup>   | Low                        |
| Motacillidae | <i>Motacilla cinerea</i>        | Grey Wagtail     | ML                | ML & MA | Usually near fresh sandy or rocky streams, but also on mown grass, ploughed land, sewage ponds <sup>2</sup>  | Low                        |
| Motacillidae | <i>Motacilla tschutschensis</i> | Yellow Wagtail   | ML                | ML & MA | Short grass and bare ground; swamp-margins, sewage ponds, saltmarshes, playing fields, airfields, ploughed lands, town lands. <sup>6</sup>                   | Low                        |

| Family         | Scientific Name               | Common Name            | Conservation Code | Source      | Habitat | Likelihood of Occurrence | Distance and Justification  |
|----------------|-------------------------------|------------------------|-------------------|-------------|---------|--------------------------|---|
|                |                               |                        |                   |             |         |                          |   |
| Procellariidae | <i>Calonectris leucomelas</i> | Streaked Shearwater    | M1                |             | x       |                          | In Aust., abundant off n. coast less common further s. <sup>6</sup> . Appears often to have been associated with spinifex, or among sapphire bushes on margins of salt lakes <sup>2</sup> .                                 |
| Psittacidae    | <i>Pezoporus occidentalis</i> | Night Parrot           | CR                | EN          | x       |                          | Nearest record 407 kms away. Low due to distance and lack of spinifex.  |
|                | <i>Polytelis alexandrae</i>   | Princess Parrot        | P4                | VU          | x       |                          | Nearest record 8 kms away. Medium due to distance and habitat. Species could fly over site.   |
|                |                               |                        |                   |             |         |                          | Nearest record 10 kms away. Low due to habitat.   |
| Scolopacidae   | <i>Actitis hypoleucus</i>     | Common Sandpiper       | M1                | MA & MA     | x x x   |                          | Varied coastal and interior wetlands – narrow muddy edges of billabongs, river pools, mangroves, among rocks and snags, reefs or rocky beaches; avoids wide open mudflats. Perches on branches, posts, boats <sup>1</sup> . |
|                | <i>Calidris acuminata</i>     | Sharp-tailed Sandpiper | M1                | MI & MA     | x x     |                          | Fresh or salt wetlands – muddy edges of lagoons, swamps, lakes, dams, soaks, sewage farms, temporary floodwaters <sup>2</sup> .   |
|                | <i>Calidris canutus</i>       | Red Knot               | EN                | EN, MI & MA | x x     |                          | Nearest record 45 kms away. Low due to distance and habitat.  |

| Family                  | Scientific Name           | Common Name        | Conservation Code | Source  | Habitat   | Likelihood of Occurrence | Distance and Justification                                    |
|-------------------------|---------------------------|--------------------|-------------------|---------|---|--------------------------|---|
|                         |                           |                    |                   |         |   |                          |   |
| Scolopacidae            | <i>Calidris melanotos</i> | Pectoral Sandpiper | MI                |         | Usually coastal wetlands, both fresh and saline, but also inland on permanent and temporary wetlands. Uses sites with mudflats, fringing vegetation, swamps with heavy overgrowth of vegetation <sup>2</sup>  | Low                      | Nearest Record 275 kms away. Low due to distance and habitat. |
|                         |                           | Wood Sandpiper     |                   |         | Uses freshwater wetlands, especially those with emergent sedges and taller fringing vegetation <sup>1</sup>   |                          |   |
|                         | <i>Tringa glareola</i>    |                    | MI                | MI & MA | X   | X                        | Nearest record 0.78 kms away. Low due to habitat.             |
| <i>Tringa nebularia</i> |                           |                    | MI                |         | Diverse inland and coastal spots. Away from the coast – uses both permanent and temporary wetlands – billabongs, swamps, lakes, floodplains, sewage farms and salt works ponds, flooded irrigated crops. On the coast – uses sheltered estuaries and bays with extensive mudflats, mangrove swamps, muddy shallows of harbours and lagoons, occasionally rocky tidal ledges. Prefers wet and flooded mud and clay rather than sand <sup>2</sup> | Low                      | Nearest record 10 kms away. Low due to habitat.               |
|                         |                           | Common Greenshank  |                   |         |   |                          |   |

| Family            | Scientific Name                  | Common Name                   | Conservation Code | Source            | Likelihood of Occurrence  | Distance and Justification  |
|-------------------|----------------------------------|-------------------------------|-------------------|-------------------|---|---|
|                   |                                  |                               |                   |                   |   |   |
|                   |                                  |                               |                   |                   |   |   |
| Threskiornithidae | <i>Plegadis falcinellus</i>      | Glossy Ibis                   | MI                | X                 | Shallow, fresh water, occasionally estuarine waters or dry grasslands <sup>1</sup>  | Low<br>Nearest record 92 kms away. Low due to distance and habitat.                             |
| <b>MAMMALIA</b>   |                                  |                               |                   |                   |   |   |
| Dasyuridae        | <i>Dasyurus blythi</i>           | Brush-tailed Mulgara, Ampurta | P4                | X                 | Mulgara predominantly occur in hummock grasslands (e.g. <i>Triodia</i> spp.) and shrublands on sandy soils. <sup>7</sup>  | Low<br>Nearest record 80 kms away. Low due to distance.   |
|                   | <i>Dasyurus geoffroii fortis</i> | Western Quoll, Chuditch       | VU                | VU (at sp. level) | Areas dominated by sclerophyll forest or drier woodland, heath and mallee shrubland <sup>4</sup>  | Low<br>Nearest record 322 kms away. Low due to distance and habitat.                            |
|                   | <i>Sminthopsis longicaudata</i>  | Long-tailed Dunnart           | P4                | X                 | Rugged, rocky areas in the grid zone; scree slopes, boulder and stony plateaus and stony adjacent stony plains with shrubs over spinifex hummock grasslands. <sup>5</sup> | Low<br>Nearest record 45 kms away. Low due to suitable adjacent habitat.                        |
| <b>REPTILIA</b>   |                                  |                               |                   |                   |   |   |
| Pythonidae        | <i>Aspidites ramsayi</i>         | Woma                          | P1 (SW subpop.)   | X                 | Woodlands, heaths and shrublands, often with spinifex. Shelters mainly in abandoned monitor and mammal burrows and in soil cracks. <sup>3</sup>                           | Medium<br>Nearest record 95kms away. Medium due to habitat similarities. Burrows found on site. |

## **Appendix H**

### **Fauna Habitat Assessments**